

An Application of a Theorem of Orthohomological Triangles

Prof. Ion Pătrașcu
Frații Buzești College, Craiova, Romania

Dr. Florentin Smarandache
University of New Mexico, Gallup Campus, USA

Abstract.

In this note we prove a problem given at a Romanian student mathematical competition, and we obtain an interesting result by using a *Theorem of Orthohomological Triangles*¹.


Problem L. 176 (from [1])

Let D, E, F be the projections of the centroid G of the triangle ABC on the lines BC, CA , and respectively AB . Prove that the Cevian lines AD, BE , and CF meet in an unique point if and only if the triangle is isosceles. {Proposed by Temistocle Bîrsan.}

Proof

Applying the generalized Pythagorean theorem in the triangle BGC , we obtain:

$$CG^2 = BG^2 + BC^2 - 2BD \cdot BC \quad (1)$$



Because $CG = \frac{2}{3} m_c$, $BG = \frac{2}{3} m_b$ and from the median's theorem it results:

$$4m_b^2 = 2(a^2 + c^2) - b^2 \quad \text{and} \quad 4m_c^2 = 2(a^2 + b^2) - c^2$$

From (1) we get: $BD = \frac{3a^2 - b^2 + c^2}{6a}$.

¹ It has been called the *Smarandache-Pătrașcu Theorem of Orthohomological Triangles* (see [2], [3], [4]).

From $BC = a$ and $BC = BD + DC$, we get that:

$$DC = \frac{3a^2 + b^2 - c^2}{6a}$$

Similarly we find:

$$CE = \frac{3b^2 - c^2 + a^2}{6b}, \quad EA = \frac{3b^2 + c^2 - a^2}{6b}$$

$$FA = \frac{3c^2 - a^2 + b^2}{6c}, \quad FB = \frac{3c^2 + a^2 - b^2}{6c}.$$

Applying Ceva's theorem it results that AD, BE, CF are concurrent if and only if

$$(3a^2 - b^2 + c^2)(3b^2 - c^2 + a^2)(3c^2 - a^2 + b^2) = (3a^2 + b^2 - c^2)(3b^2 + c^2 - a^2)(3c^2 + a^2 - b^2) \quad (2)$$

Let's consider the following notations:

$$a^2 + b^2 + c^2 = T, \quad 2a^2 - 2b^2 = \alpha, \quad 2b^2 - 2c^2 = \beta, \quad 2c^2 - 2a^2 = \gamma$$

From (2) it results:

$$(T + \alpha)(T + \beta)(T + \gamma) = (T - \alpha)(T - \beta)(T - \gamma).$$

And from here:

$$T^3 + (\alpha + \beta + \gamma)T^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)T + \alpha\beta\gamma = T^3 - (\alpha + \beta + \gamma)T^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)T - \alpha\beta\gamma.$$

Because $\alpha + \beta + \gamma = 0$, we obtain that $2\alpha\beta\gamma = 0$, therefore $\alpha = 0$ or $\beta = 0$ or $\gamma = 0$, thus $a = b$ or $b = c$ or $a = c$; consequently the triangle ABC is isosceles.

The reverse: If ABC is an isosceles triangle, then it is obvious that AD, BE , and CF are concurrent.

Observations

1. The proved problem asserts that:
"A triangle ABC and the pedal triangle of its weight center are orthomological triangles if and only if the triangle ABC is isosceles."
2. Using the previous result and the Smarandache-Pătrășcu Theorem (see [2], [3], [4]) we deduce that:
"A triangle ABC and the pedal triangle of its simedian center are orthomological triangles if and only if the triangle ABC is isosceles."

References

- [1] Temistocle Bîrsan, Training problems for mathematical contest, B. College Level – L. 176, *Recreații Matematice* journal, Iași, Romania, Year XII, No. 1, 2010.
- [2] Ion Pătrășcu & Florentin Smarandache, *A Theorem about Simultaneous Orthological and Homological Triangles*, in arXiv.org, Cornell University, NY, USA.

- [3] Mihai Dicu, *The Smarandache- Pătraşcu Theorem of Orthohomological Triangles*, <http://www.scribd.com/doc/28311880/Smarandache-Patrascu-Theorem-of-Orthohomological-Triangles>.
- [4] Claudiu Coandă, *A Proof in Barycentric Coordinates of the Smarandache-Pătraşcu Theorem*, Sfera journal, 2010.