

A Theorem about Simultaneous Orthological and Homological Triangles

Ion Pătrașcu
Frații Buzești College, Craiova, Romania

Florentin Smarandache
University of New Mexico, Gallup Campus, USA

Abstract. In this paper we prove that if P_1, P_2 are isogonal points in the triangle ABC , and if $A_1B_1C_1$ and $A_2B_2C_2$ are their ponder triangle such that the triangles ABC and $A_1B_1C_1$ are homological (the lines AA_1, BB_1, CC_1 are concurrent), then the triangles ABC and $A_2B_2C_2$ are also homological.

Introduction.

In order for the paper to be self-contained, we recall below the main definitions and theorems needed in solving this theorem.

Also, we introduce the notion of *Orthohomological Triangle*, which means a triangle that is simultaneously orthological and homological.

Definition 1

In a triangle ABC the Cevians AA_1 and AA_2 which are symmetric with respect to the angle's BAC bisector are called isogonal Cevians.




Fig. 1

Observation 1

If $A_1, A_2 \in BC$ and AA_1, AA_2 are isogonal Cevians then $\sphericalangle BAA_1 \equiv \sphericalangle BAA_2$. (See Fig.1.)

Theorem 1 (Steiner)

If in the triangle ABC , AA_1 and AA_2 are isogonal Cevians, A_1, A_2 are points on BC then:

$$\frac{A_1B}{A_1C} \cdot \frac{A_2B}{A_2C} = \left(\frac{AB}{AC}\right)^2$$

Proof

We have:

$$\frac{A_1B}{A_1C} = \frac{\text{area}\Delta BAA_1}{\text{area}\Delta CAA_1} = \frac{\frac{1}{2} AB \cdot AA_1 \sin(\sphericalangle BAA_1)}{\frac{1}{2} AC \cdot AA_1 \sin(\sphericalangle CAA_1)} \tag{1}$$

$$\frac{A_2B}{A_2C} = \frac{\text{area}\Delta BAA_2}{\text{area}\Delta CAA_2} = \frac{\frac{1}{2} AB \cdot AA_2 \sin(\sphericalangle BAA_2)}{\frac{1}{2} AC \cdot AA_2 \sin(\sphericalangle CAA_2)} \tag{2}$$

Because $\sin(\sphericalangle BAA_1) = \sin(\sphericalangle BAA_2)$ and $\sin(\sphericalangle CAA_1) = \sin(\sphericalangle CAA_2)$

by multiplying the relations (1) and (2) side by side we obtain the Steiner relation:

$$\frac{A_1B}{A_1C} \cdot \frac{A_2B}{A_2C} = \left(\frac{AB}{AC}\right)^2 \tag{3}$$

Theorem 2

In a given triangle, the isogonal Cevians of the concurrent Cevians are concurrent.

Proof

We'll use the Ceva's theorem which states that the triangle's ABC Cevians AA_1, BB_1, CC_1 ($A_1 \in BC, B_1 \in AC, C_1 \in AB$) are concurrent if and only if the following relation takes place:

$$\frac{A_1B}{A_1C} \cdot \frac{B_1C}{B_1A} \cdot \frac{C_1A}{C_1B} = 1 \tag{4}$$




Fig. 2

We suppose that AA_1, BB_1, CC_1 are concurrent Cevians in the point P_1 and we'll prove that their isogonal AA_2, BB_2, CC_2 are concurrent in the point P_2 . (See Fig. 2).

From the relations (3) and (4) we find:

$$\frac{A_2B}{A_2C} = \left(\frac{AB}{AC}\right)^2 \cdot \frac{A_1C}{A_1B} \quad (5)$$

$$\frac{B_2C}{B_2A} = \left(\frac{BC}{AB}\right)^2 \cdot \frac{B_1A}{B_1C} \quad (6)$$

$$\frac{C_2A}{C_2B} = \left(\frac{AC}{BC}\right)^2 \cdot \frac{C_1B}{C_1A} \quad (7)$$

By multiplying side by side the relations (5), (6) and (7) and taking into account the relation (4) we obtain:

$$\frac{A_2B}{A_2C} \cdot \frac{B_2C}{B_2A} \cdot \frac{C_2A}{C_2B} = 1,$$

which along with Ceva's theorem proves the proposed intersection.

Definition 2

The intersection point of certain Cevians and the point of intersection of their isogonal Cevians are called isogonal conjugated points or isogonal points.

Observation 2

The points P_1 and P_2 from Fig. 2 are isogonal conjugated points.

In a non right triangle its orthocenter and the circumscribed circle's center are isogonal points.

Definition 3

If P is a point in the plane of the triangle ABC , which is not on the triangle's circumscribed circle, and A', B', C' are the orthogonal projections of the point P respectively on BC, AC , and AB , we call the triangle $A'B'C'$ the podaire triangle of the point P .

Definition 4

The podaire triangle of the center of the inscribed circle in the triangle is called the contact triangle of the given triangle.




Fig. 3

Observation 3

In figure 3, $A'B'C'$ is the contact triangle of the triangle ABC . The name is connected to the fact that its vertexes are the contact points (of tangency) with the sides of the inscribed circle in the triangle ABC .

Definition 5

The podaire triangle of the orthocenter of a triangle is called orthic triangle.

Definition 6

Two triangles are called orthological if the perpendiculars constructed from the vertexes of one of the triangle on the sides of the other triangle are concurrent.

Definition 7

The intersection point of the perpendiculars constructed from the vertexes of a triangle on the sides of another triangle (the triangles being orthological) is called the triangles' orthology center.

Theorem 3 (The Orthological Triangles Theorem)

If the triangles ABC and $A'B'C'$ are such that the perpendiculars constructed from A on $B'C'$, from B on $A'C'$ and from C on $A'B'$ are concurrent (the triangles ABC and $A'B'C'$ being orthological), then the perpendiculars constructed from A' on BC , from B' on AC , and from C' on AB are also concurrent.

To prove this theorem firstly will prove the following:

Lemma 1 (Carnot)

If ABC is a triangle and A_1, B_1, C_1 are points on BC, AC, AB respectively, then the perpendiculars constructed from A_1 on BC , from B_1 on AC and from C_1 on AB are concurrent if and only if the following relation takes place:

$$A_1B^2 - A_1C^2 + B_1C^2 - B_1A^2 + C_1A^2 - C_1B^2 = 0 \tag{8}$$

Proof

If the perpendiculars in A_1, B_1, C_1 are concurrent in the point M (see Fig. 4), then from Pythagoras theorem applied in the formed right triangles we find:




Fig. 4

$$A_1B^2 = MB^2 - MA_1^2 \tag{9}$$

$$A_1C^2 = MC^2 - MA_1^2 \tag{10}$$

hence

$$A_1B^2 - A_1C^2 = MB^2 - MC^2 \tag{11}$$

Similarly it results

$$B_1C^2 - B_1A^2 = MC^2 - MA^2 \tag{12}$$

$$C_1A^2 - C_1B^2 = MA^2 - MC^2 \tag{13}$$

By adding these relations side by side it results the relation (8).

Reciprocally

We suppose that relation (8) is verified, and let's consider the point M being the intersection of the perpendiculars constructed in A_1 on BC and in B_1 on AC . We also note with C' the projection of M on AC . We have that:

$$A_1B^2 - A_1C^2 + B_1C^2 - B_1A^2 + C_1A^2 + C'A^2 - C'B^2 = 0 \tag{14}$$

Comparing (8) and (14) we find that

$$C_1A^2 - C_1B^2 = C'A^2 - C'B^2$$

and

$$(C_1A - C_1B)(C_1A + C_1B) = (C'A - C'B)(C'A + C'B)$$

and because

$$C_1A - C_1B = C'A + C'B = AB$$

we obtain that $C' = C_1$, therefore the perpendicular in C_1 passes through M also.

Observation 4

The triangle ABC and the podaire triangle of a point from its plane are orthological triangles.

The proof of Theorem 3

Let's consider ABC and $A'B'C'$ two orthological triangles (see Fig. 5). We note with M the intersection of the perpendiculars constructed from A on $B'C'$, from B on $A'C'$ and from C on $A'B'$, also we'll note with A_1, B_1, C_1 the intersections of these perpendiculars with $B'C', A'C'$ and $A'B'$ respectively.




Fig. 5

In conformity with lemma 1, we have:

$$A_1B'^2 - A_1C'^2 + B_1C'^2 - B_1A'^2 + C_1A'^2 - C_1B'^2 = 0 \quad (15)$$

From this relation using the Pythagoras theorem we obtain:

$$B'A'^2 - C'A'^2 + C'B'^2 - A'B'^2 + A'C'^2 - B'C'^2 = 0 \quad (16)$$

We note with A'_1, B'_1, C'_1 the orthogonal projections of A', B', C' respectively on BC, CA, AB . From the Pythagoras theorem and the relation (16) we obtain:

$$A'_1B'^2 - A'_1C'^2 + B'_1C'^2 - B'_1A'^2 + C'_1A'^2 - C'_1B'^2 = 0 \quad (17)$$

This relation along with Lemma 1 shows that the perpendiculars drawn from A' on BC , from B' on AC and from C' on AB are concurrent in the point M' .

The point M' is also an orthological center of triangles $A'B'C'$ and ABC .

Definition 8

The triangles ABC and $A'B'C'$ are called bylogical if they are orthological and they have the same orthological center.

Definition 9

Two triangles ABC and $A'B'C'$ are called homological if the lines AA', BB', CC' are concurrent. Their intersection point is called the homology point of triangles ABC and $A'B'C'$

Observation 6

In figure 6 the triangles AA' , BB' , CC' are homological and the homology point being O




Fig.6

If ABC is a triangle and $A'B'C'$ is its podaire triangle, then the triangles ABC and $A'B'C'$ are homological and the homology center is the orthocenter H of the triangle ABC

Definition 10

A number of n points ($n \geq 3$) are called concyclic if there exist a circle that contains all of these points.

Theorem 5 (The circle of 6 points)

If ABC is a triangle, P_1, P_2 are isogonal points on its interior and $A_1B_1C_1$ respectively




Fig. 7

$A_2B_2C_2$ the podaire triangles of P_1 and P_2 , then the points $A_1, A_2, B_1, B_2, C_1, C_2$ are concyclic.

Proof

We will prove that the 6 points are concyclic by showing that these are at the same distance of the middle point P of the line segment P_1P_2 .

It is obvious that the medians of the segments $(A_1A_2), (B_1B_2), (C_1C_2)$ pass through the point P , which is the middle of the segment (P_1P_2) . The trapezoid $A_1P_1P_2A_2$ is right angle and the mediator of the segment (A_1A_2) will be the middle line, therefore it will pass through P , (see Fig. 7).

Therefore we have:

$$PA_1 = PA_2, PB_1 = PB_2, PC_1 = PC_2 \quad (18)$$

We'll prove that $PB_1 = PC_2$ by computing the length of these segments using the median's theorem applied in the triangles $P_1B_1P_2$ and $P_1C_2P_2$.

We have:

$$4PB_1^2 = 2(P_1B_1^2 + P_2B_1^2) - P_1P_2^2 \quad (19)$$

We note

$$AP_1 = x_1, AP_2 = x_2, m(\sphericalangle BAP_1) = m(\sphericalangle CAP_2) = \alpha.$$

In the right triangle $P_2B_2B_1$ applying the Pythagoras theorem we obtain:

$$P_2B_1^2 = P_2B_2^2 + B_1B_2^2 \quad (20)$$

From the right triangle AB_2P_2 we obtain:

$$P_2B_2 = AP_2 \sin \alpha = x_2 \sin \alpha \text{ and } AB_2 = x_2 \cos \alpha$$

From the right triangle AP_1B_1 it results $AB_1 = AP_1 \cos(A - \alpha)$, therefore

$$AB_1 = x_1 \cos(A - \alpha) \text{ and } P_1B_1 = x_1 \sin(A - \alpha),$$

thus

$$B_1B_2 = |AB_2 - AB_1| = |x_2 \cos \alpha - x_1 \cos(A - \alpha)| \quad (21)$$

Substituting back in relation (17), we obtain:

$$P_2B_1^2 = x_2^2 \sin^2 \alpha + [x_2 \cos \alpha - x_1 \cos(A - \alpha)]^2 \quad (22)$$

From the relation (16), it results:

$$4PB_1^2 = 2[x_1^2 + x_2^2 - 2x_1x_2 \cos \alpha \cos(A - \alpha)] P_1P_2^2 \quad (23)$$

The median's theorem in the triangle $P_1C_2P_2$ will give:

$$4PC_2^2 = 2(P_1C_2^2 + P_2C_2^2) - P_1P_2^2 \quad (24)$$

Because $P_1C_1 = x_1 \sin \alpha$, $AC_1 = x_1 \cos \alpha$, $AC_2 = x_2 \cos(A - \alpha)$, $P_1C_2^2 = P_1C_1^2 + C_1C_2^2$,

we find that

$$4PC_2^2 = 2[x_1^2 + x_2^2 - 2x_1x_2 \cos \alpha \cos(A - \alpha)] - P_1P_2^2 \quad (25)$$

The relations (23) and (25) show that

$$PB_1 = PC_2 \quad (26)$$

Using the same method we find that :

$$PA_1 = PC_1 \quad (27)$$

The relations (18), (26) and (27) imply that:

$$PA_1 = PA_2 = PB_1 = PB_2 = PC_1 = PC_2$$

From which we can conclude that $A_1, A_2, B_1, B_2, C_1, C_2$ are concyclic.

Lemma 2 (The power of an exterior point with respect to a circle)

If the point A is exterior to circle $C(O, r)$ and d_1, d_2 are two secants constructed from A that intersect the circle in the points B, C respectively E, D , then:

$$AB \cdot AC = AE \cdot AD = cons. \quad (28)$$

Proof

The triangles ADB and ACE are similar triangles (they have each two congruent angles respectively), it results:

$$\frac{AB}{AE} = \frac{AD}{AC}$$




Fig. 8

and from here:

$$AB \cdot AC = AE \cdot AD \quad (29)$$

We construct the tangent from A to circle $C(O, r)$ (see Fig. 8). The triangles ATE and ADT are similar (the angles from the vertex A are common and $\sphericalangle ATE \equiv \sphericalangle ADT = \frac{1}{2}m(\widehat{TE})$).

We have:

$$\frac{AE}{AT} = \frac{AT}{AD},$$

it results

$$AE \cdot AD = AT^2 \quad (30)$$

By noting $AO = a$, from the right triangle ATO (the radius is perpendicular on the tangent in the contact point), we find that:

$$AT^2 = AO^2 - OT^2,$$

therefore

$$AT^2 = a^2 - r^2 = const. \quad (31)$$

The relations (29), (30) and (31) are conducive to relation (28).

Theorem 6 (Terquem)

If AA_1, BB_1, CC_1 are concurrent Cevians in the triangle ABC and A_2, B_2, C_2 are intersections of the circle circumscribed to the triangle A_1, B_1, C_1 cu $(BC), (CA), (AB)$, then the lines AA_2, BB_2, CC_2 are concurrent.

Proof

Let's consider F_1 the concurrence point of the Cevians AA_1, BB_1, CC_1 .

From Ceva's theorem it results that:

$$A_1B \cdot B_1C \cdot C_1A = A_1C \cdot B_1A \cdot C_1B \tag{32}$$




Fig 9

Considering the vertexes A, B, C 's power with respect to the circle circumscribed to the triangle $A_1B_1C_1$, we obtain the following relations:

$$AC_1 \cdot AC_2 = AB_1 \cdot AB_2 \tag{33}$$

$$BA_1 \cdot BA_2 = BC_1 \cdot BC_2 \tag{34}$$

$$CB_1 \cdot CB_2 = CA_1 \cdot CA_2 \tag{35}$$

Multiplying these relations side by side and taking into consideration the relation (32), we obtain

$$AC_2 \cdot BA_2 \cdot CB_2 = AB_2 \cdot BC_2 \cdot CA_2 \tag{36}$$

This relation can be written under the following equivalent format

$$\frac{A_2B}{A_2C} \cdot \frac{B_2C}{B_2A} \cdot \frac{C_2A}{C_2B} = 1 \tag{37}$$

From Ceva's theorem and the relation (37) we obtain that the lines AA_2, BB_2, CC_2 are concurrent in a point noted in figure 9 by F_2 .

Note 1

The points F_1 and F_2 have been named the Terquem's points by Candido of Pisa – 1900.

For example in a non right triangle the orthocenter H and the center of the circumscribed circle O are Terquem's points.

Definition 11

Two triangles are called orthohomological if they are in the simultaneously orthological and homological.

Theorem 7 (Smarandache-Pătraşcu Theorem of Orthohomological Triangles)

If P_1, P_2 are two conjugated isogonal points in the triangle ABC , and $A_1B_1C_1$ and $A_2B_2C_2$ are their respectively podaire triangles such that the triangles ABC and $A_1B_1C_1$ are homological, then the triangles ABC and $A_2B_2C_2$ are also homological.

Proof

Let's consider that F_1 is the concurrence point of the Cevians AA_1, BB_1, CC_1 (the center of homology of the triangles ABC and $A_1B_1C_1$). In conformity with Theorem 6 the circumscribed circle to triangle $A_1B_1C_1$ intersects the sides $(BC), (CA), (AB)$ in the points A_2, B_2, C_2 , these points are exactly the vertexes of the podaire triangle of P_2 , because if two circles have in common three points, then the two circles coincide; practically, the circle circumscribed to the triangle $A_1B_1C_1$ is the circle of the 6 points (Theorem 5).

Terquem's theorem implies the fact that the triangles ABC and $A_2B_2C_2$ are homological. Their homological center is F_2 , the second Terquem's point of the triangle ABC .

Observation 7

If the points P_1 and P_2 isogonal conjugated in the triangle ABC coincide, then the triangles ABC and $A_2B_2C_2$, the podaire of $P_1 = P_2$ are homological.

Proof

From $P_1 = P_2$ and the fact that P_1, P_2 are isogonal conjugate, it results that $P_1 = P_2 = I$ - the center of the inscribed circle in the triangle ABC . The podaire triangle of I is the contact triangle. In this case the lines AA_1, BB_1, CC_1 are concurrent in Γ , Gergonne's point, which is the homological center of these triangles.

Observation 8

The reciprocal of Theorem 7 for orthohomological triangles is not true.

To prove this will present a counterexample in which the triangle ABC and the podaire triangles $A_1B_1C_1, A_2B_2C_2$ of the points P_1 and P_2 are homological, but the points P_1 and P_2 are not isogonal conjugated; for this we need several results.

Definition 12

In a triangle two points on one of its side and symmetric with respect to its middle are called isometrics.

Definition 13

The circle tangent to a side of a triangle and to the other two sides' extensions of the triangle is called exterior inscribed circle to the triangle.

Observation 9

In figure 10 we constructed the extended circle tangent to the side (BC). We note its center with I_a . A triangle ABC has, in general, three exinscribed circles

Definition 14

The triangle determined by the contact points with the sides (of a triangle) of the exinscribed circle is called the cotangent triangle of the given triangle.




Fig. 10

Theorem 8

The isometric Cevians of the concurrent Cevians are concurrent.

The proof of this theorem results from the definition 14 and Ceva's theorem

Definition 15

The contact points of the Cevians and of their isometric Cevians are called conjugated isotomic points.

Lemma 3

In a triangle ABC the contact points with a side of the inscribed circle and of the exinscribed circle are isotomic points.

Proof

The proof of this lemma can be done computational, therefore using the tangents' property constructed from an exterior point to a circle to be equal, we compute the CD and BD_a (see Fig. 10) in function of the length a, b, c of the sides of the triangle ABC .

We find that $CD = p - c = BD_a$, which shows that the Cevians AD and AD_a are isogonal (p is the semi-perimeter of triangle ABC , $2p = a + b + c$).

Theorem 9

The triangle ABC and its cotangent triangle are isogonal.

We'll use theorem 8 and taking into account lemma 3, and the fact that the contact triangle and the triangle ABC are homological, the homological center being the Gergonne's point.

Observation 10

The homological center of the triangle ABC and its cotangent triangle is called Nagel's point (N).

Observation 11

The Gergonne's point (Γ) and Nagel's point (N) are isogonal conjugated points.

Theorem 10

The perpendiculars constructed on the sides of a triangle in the vertexes of the cotangent triangle are concurrent.

The proof of this theorem results immediately using lemma 1 (Carnot)

Definition 12

The concurrence point of the perpendiculars constructed in the vertexes of the cotangent triangle on the sides of the given triangle is called the Bevan's point (V).

We will prove now that the reciprocal of the theorem of the orthohomological triangles is false

We consider in a given triangle ABC its contact triangle and also its cotangent triangle. The contact triangle and the triangle ABC are homological, the homology center being the Gergonne's point (Γ). The given triangle and its cotangent triangle are homological, their homological center being Nagel's point (N). Bevan's point and the center of the inscribed circle have as podaire triangles the cotangent triangles and of contact, but these points are not isogonal conjugated (the point I is its own isogonal conjugate).

References:

1. C. Mihalescu, *Geometria elementelor remarcabile*, Ed. Tehnică, București, 1959.
2. R.A. Johnson, *Modern Geometry: An Elementary Treatise on the Geometry of the Triangle and the Circle*, 1929.