

**What can be said about
(massive) Graviton Stability?
& Is there DM generated DE ?**

**Rudn 10, June 30, Moscow,
Russia**

Dr. Andrew Beckwith
beckwith@aibep.org
abeckwith@uh.edu

Abstract

A thought experiment:

1st : Can gravitons be stable if they have mass ?

2nd : If so, did a DM generated KK particle, with tiny 4 D mass contribute to DE a billion years ago ?

- (indirectly touching upon)
- To what degree is gravity an emergent field that is partly/largely classical with extreme nonlinearity, or a QM/quantum field theory phenomenon?

Math – Physics representation of core issues of higher dimensional contribution

- Start off with a basic statement of strength of matter - graviton interaction, assuming KK graviton

- $$\mathfrak{J} = -\frac{K}{2} \cdot \sum_n \int d^4x h_{uv}^n \cdot T^{uv} \sim 1 / M_{PL}^2 \quad (1)$$

The stress energy tensor comes from the standard model, and the h term is from using a KK graviton interactions model , up to the n th mode.

Does the last slide hold if we make the following modification of a KK tower of gravitons ?

- Modification put in, as seen in later to mimic DE
- Suggestion to look at, here, is to consider what if
- $m_n(\textit{Graviton}) = \frac{n}{L} + 10^{-65}$ grams? **(2)**

Issue to raise

- ***Work presented by Maggiore, specifically delineated for non zero graviton mass, due to Fiertz Lagrangian , obtaining as a limit***

$$-3m_{\text{graviton}}^2 h = \frac{\kappa}{2} \cdot T \quad (3)$$

- **Does the above Eqn(3) permit a frequency range for massive Gravitons which permits stability for Massive Gravitons ?**

From working with Eqn (1) and from Vissers treatment of (massive) gravitons

- This is Vissers (massive) Graviton Stress energy tensor, as a start

$$T_{uv}|_{m \neq 0} = \left[\left(\frac{\hbar}{l_P^2 \lambda_g^2} \right) \cdot \left(\frac{GM}{r} \right) \cdot \exp\left(\frac{r}{\lambda_g} \right) + \left(\frac{GM}{r} \right)^2 \right] \times \begin{bmatrix} 4 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (4)$$

Eqn (1), Eqn (4), with work leads to a positive, real frequency squared value ,


- In order to obtain a non imaginary, real (massive graviton frequency), **need Eqn(5) below:**

We claim that application leads to the result of the next page 1 billion years ago as put in slide 8 (next slide)

$$0 < \frac{1}{6m_g c^2} \left(\frac{\hbar^2}{l_P^2 \lambda_g^2} \cdot \exp \left[-\frac{r}{\lambda_g} + \frac{m_g \cdot r}{\hbar} \right] + \left(\frac{MG}{r} \right) \cdot \exp \left(\frac{m_g r}{\hbar} \right) \right) < 1$$

Jerk calculation leads to, if one has a stable, non zero 4-D graviton mass

- If we define the jerk $q = -\frac{\ddot{a}a}{\dot{a}^2}$



Assuming a brane world

Z (red shift value). Change in sign for Z $\sim .42$ is almost one billion years ago, corresponding to reacceleration of the universe, i.e

Basic results of [Alves](#), et al. (2009), using their parameter values, with an additional term of "dark flow" added, corresponding to one KK additional dimensions.

**Issue is that a 4 dimensional Graviton
with small mass violates principle of
correspondence- complimentarity**

- How important is such a
violation ?**

**Explanation may show reason for G.
Smoots values of information, initially**

- See the following for the G. Smoot lecture

http://chalonge.obspm.fr/Paris07_Smoot.pdf

For brane world, use these evolution equations

Friedman equation, subsequently modified

$$\dot{a}^2 = \left[\left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{\rho^2}{36M_{Planck}^2} \right) a^2 - \kappa + \frac{C}{a^2} \right] \quad (6)$$

Density equation, with non- zero graviton mass

$$\rho \equiv \rho_0 \cdot \left(\frac{a_0}{a} \right)^3 - \left[\frac{m_g c^6}{8\pi G \hbar^2} \right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2} \right) \quad (7)$$

For LQG, use these evolution equations

Friedman equations, assuming 'constant' momentum , a.k.a. Eqn(8)

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa}{3} \cdot \rho \quad \left(\frac{\dot{a}}{a}\right)^2 \equiv \frac{\kappa}{6} \cdot \frac{p_\phi^2}{a^6} \quad \left(\frac{\ddot{a}}{a}\right) = -\frac{2 \cdot \kappa}{3} \cdot \rho$$

Density equation is the same

$$\rho \equiv \rho_0 \cdot \left(\frac{a_0}{a}\right)^3 - \left[\frac{m_g c^6}{8\pi G \hbar^2} \right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2} \right)$$

Infinite Quantum statistics. From the work presented in the Paris observatory, July 2009

Start with

$$Z_N \sim \left(\frac{1}{N!}\right) \cdot \left(\frac{V}{\lambda^3}\right)^N \quad S \approx N \cdot (\log[V/N\lambda^3] + 5/2)$$

$$S \approx N \cdot (\log[V/\lambda^3] + 5/2) \quad V \approx R_H^3 \approx \lambda^3$$

V stands for volume of nucleation regime space.
“particles” nucleate from ‘vacuum’ in QM

For DM. V for nucleation is HUGE. Graviton space V for nucleation is tiny , well inside inflation/ Therefore, the log factor drops OUT of entropy S if V chosen properly. For small V, then

$$\Delta S \approx \Delta N_{gravitons} \quad (9)$$

Some considerations about the partition function

Glinka (2007): if we identify $\Omega = \frac{1}{2|u|^2 - 1}$

- as a partition function (with u part of a Bogoliubov transformation) due to a graviton-quintessence gas, to get information theory-based entropy $S \equiv \ln \Omega$

1. Derivation by Glinka explicitly uses the Wheeler De Witt equation
2. 2. Is there in any sense a linkage of Wheeler De Witt equation with String theory results ?

PROBLEM TO CONSIDER:

Ng's result quantum counting algorithm is a **STRING theory** result. Glinka is **Wheeler De Witt equation. Equivalent ?**

Questions to raise.

Can we make a linkage between Glinka's quantum gas argument, and a small space version/ application of Ng's Quantum infinite statistics ?

In addition, if the quantum graviton gas is correct, can we model emergent structure of gravity via linkage between Ng particle count, and Q.G.G argument?

More on the Glinka quantum gas hypothesis, part 1

- Starting point to Glinka's Quantum gas, in terms of numerical count


- $$n_f = [1/4] \cdot \left[\sqrt{\frac{v(a_{initial})}{v(a)}} - \sqrt{\frac{v(a)}{v(a_{final})}} \right] \quad (10)$$

More on the Glinka Quantum gas hypothesis, part 2

- Relevant to understanding the role of

- $$\Omega_{gw}(\nu) \cong \frac{3.6}{h_0^2} \cdot \left[\frac{n_f}{10^{37}} \right] \cdot \left(\frac{\nu}{1kHz} \right)^4 \quad (11)$$

See B. P. Abbott et al.



Breakdown of field theory with respect to massive gravitons in limit

$$m_{\text{graviton}} \rightarrow 0$$

The massless equation of the graviton evolution equation takes the form

$$\partial_{\mu} \partial^{\mu} h_{\mu\nu} = \sqrt{32\pi G} \cdot \left(T_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} T^{\mu}_{\mu} \right)$$

Consider what happens with a graviton mass

$$m_{\text{graviton}} \neq 0$$

From Maggiore (2008):

$$\left(\partial_{\mu}\partial^{\mu} - m_{\text{graviton}}\right) \cdot h_{\mu\nu} = \left[\sqrt{32\pi G} + \delta^{+}\right] \cdot \left(T_{\mu\nu} - \frac{1}{3}\eta_{\mu\nu}T^{\mu}_{\mu} + \frac{\partial_{\mu}\partial_{\nu}T^{\mu}_{\mu}}{3m_{\text{graviton}}}\right)$$

The mismatch between these two equations when

$$m_{\text{graviton}} \rightarrow 0$$

Is largely due to, even if graviton mass goes to zero

$$m_{\text{graviton}} h_{\mu}^{\mu} \neq 0$$

$$m_{\text{graviton}} \cdot h_{\mu}^{\mu} = -\left[\sqrt{32\pi G} + \delta^+ \right] \cdot T_{\mu}^{\mu}$$


Try semiclassical model of graviton, as kink-anti kink pair

- How does this fit in with t'Hooft's deterministic QM?
- From a 1+ dimensional kink-antikink

$$\Psi_{i,f} [\phi(\mathbf{x})] \Big|_{\phi=\phi_{ci,cf}} = c_{i,f} \cdot \exp \left\{ - \int d\mathbf{x} \alpha \left[\phi_{Ci,f}(\mathbf{x}) - \phi_0(\mathbf{x}) \right]^2 \right\},$$

From density wave physics, 1+ dimensions

Kink-antikinks lead to a vacuum wave function. The LHS is a kink; the RHS is an antikink.



The wave functional could have t'Hooft equivalence class structure added, in 4 to 5 dimensions

- T'Hooft used in 2006 an equivalence class argument as an embedding space for simple harmonic oscillators, as given in his Figure 2, on page 8 of his 2006 article.
- “Beneath Quantum Mechanics, there may be a deterministic theory with (local) information loss. This may lead to a sufficiently complex vacuum state.” - t'Hooft
- The author submits, that a kink-anti kink formulation of the graviton, when sufficiently refined, may indeed create such a vacuum state, as a generalization of Fig 2.

The small mass of the graviton
would be for energy in

$$\Delta E \Delta t \geq \hbar$$

- Having said this, the author is fully aware of the String theory HUP variant

$$\Delta x \geq \frac{\hbar}{\Delta p} + \frac{l_s^2}{\hbar} \Delta p$$

- The idea would be to possibly obtain a way to look at counting for GW detectors

$$h_0^2 \Omega_{gw}(f) \cong \frac{3.6}{2} \cdot \left[\frac{n_f [\textit{graviton}] + n_f [\textit{neutrino}]}{10^{37}} \right] \cdot \left(\frac{\langle f \rangle}{1\text{kHz}} \right)^4$$

The following is claimed:

If n (graviton) is obtained, then higher dimensional geometry may be relevant to transmitting information via gravitons from prior to present universes

- How much information can be carried by an individual graviton?
- Assume $\Delta S \approx \Delta N_{gravitons}$
- Use Seth Lloyd's

$$I = S_{total} / k_B \ln 2 = [\#operations]^{3/4} = [\rho \cdot c^5 \cdot t^4 / \hbar]^{3/4}$$

10^7 relic gravitons yields at or more than 10^7 operations!

This value implies that per graviton, as nucleated at least 4 dimensions, there is at least **one unit** of information associated with the graviton (assuming there is at least **some relationship** between an operation and information)

10^7 or higher amounts of prior universe information transmitted to our cosmos?

Resolutions of questions about cosmological constants ?

- **1st Conclusion**, one needs a reliable information packing algorithm! I.e. for a wave length , as input into the fine structure constant, we need spatial / information limits defined for geometry
- $\Delta S \approx \Delta N_{gravitons} \approx 10^7$ is only a beginning
- **2nd Conclusion**, assumed GW detector sensitivity limits need a comprehensive look over, re do

Final inquiry, making sense of the supposed
“radius of the Universe” calculation

- Matt Roos, has put in a foundational way of testing, via experiment, how to calculate a supposed ‘radius of the universe’

- $$r_U \equiv \frac{1}{H \cdot \sqrt{|\Omega - 1|}} \quad (12)$$

Tweaking parameters of H , and

$\Omega \equiv \rho(t) / \rho_{critical}$ from our inquiry

- **The choice of H , and of density , as in the**
- **equation below will allow the dynamics of**
- **how the universe expands mesh with a fuller**
- **understanding of structure formation.**

$$\Omega \equiv \rho(t) / \rho_{critical} \quad (13)$$

Acknowledgements

- The author thanks
 - **Dr. Glinka** of Friendship university is thanked for encouraging the author to think of semi classical / WdW style issues, and their relevance to cosmology. He also told the author of the RUDN 10 conference
 - **Dr. Fangyu Li** of Chongqing University is thanked for lending his personal notes to give substance to the content of page 10 of the companion document that complements this PPT, plus his week- long hospitality to the author in China
 - **Dr. Christian Corda** is thanked for encouraging the author to explore semiclassical issues in GW and in GR, which lead to a meeting on foundational issues of gravitation in a symposium in Crete, September 2009; and forwarded the Alves et al. document to the author to expand the author's scientific horizons .

Bibliography, with 1st , 2nd , 3rd , references

- M. Alves, O. Miranda. and J. de Araujo, J. ,” Can Massive Gravitons be an Alternative to Dark Energy?”, arXiv:0907.5190 (July 2009)
- A. Beckwith “Deceleration Parameter $Q(z)$ and the Role of Nucleated GW 'graviton Gas' in the Development of DE Alternatives”; <http://vixra.org/abs/1005.0012>
- A. Beckwith “ Can a massive graviton be a stable particle ?” <http://vixra.org/abs/1006.0022>

Bibliography , Contd.

M. Visser,

[http://arxiv.org/pdf/gr-qc/9705051v2.](http://arxiv.org/pdf/gr-qc/9705051v2)

G. Smoot as in the link given below

[http://chalonge.obspm.fr/Paris07 Smoot.pdf](http://chalonge.obspm.fr/Paris07_Smoot.pdf)

General frequency reference, plus
Glinka's Quantum gas reference

- **B.P. Abbott et al., *Nature* 460, 990-993 (2009)**
- **L. A. Glinka ;**
http://arxiv.org/PS_cache/arxiv/pdf/0711/0711.1380v4.pdf

Bibliography, continued

- A.W. Beckwith, " Entropy Growth in the Early Universe and Confirmation of Initial Big Bang Condition (Wheeler de Witt Eqn. Results Vs. String Theory?)", (Submitted to Rencontres De Blois 2009 conference proceedings), <http://vixra.org/abs/0908.0109> (2009)
- A.W.Beckwith, " Gravitinos, the Lithium problem, and DM production: Is there a corresponding neutrino physics linkage?", presented in 12 Marcel Grossman conference, Paris, 2009, submitted as a confirmed conference entry <http://vixra.org/abs/0910.0052>; **published in Progress in nuclear & particle physics**

Bibliography, contd.

- A.W. Beckwith, "Stretched Neutrinos, and the Supposed Linkage to Gravitons/ HFGW Data Sets", submitted to the 12 Marcel Grossman conference, as a conference proceedings entry <http://vixra.org/abs/0911.0067>
- A.W. Beckwith, "Classical and Quantum Models of Density Wave Transport, a comparative study", PhD dissertation, December 2001
- **R. Brandenberger, "Matter bounce in Hořava-Lifshitz cosmology", *Phys. Rev. D* 80, 043516 (2009); <http://arxiv.org/abs/0904.2835>**

Bibliography, contd

- A. Buonanno, “Gravitational waves”, pp 10-52, from the Les Houches Section LXXXVI, ‘Particle physics and Cosmology, the fabric of spacetime’ which has F. Bernardeau, C. Grojean, and J. Dalibard as editors
- M. Gasperini and G. *Veneziano*, Mod. Phys. Lett. A 8, 3701 (1993)
- M. Giovannini, “Non Topological Gravitating defects in 5 dimensional space”, Class. Quantum. Grav. 23, 2006 , L73-80,
- Fangyu Li, Nan Yang, Zhenyun Fang, Robert M. Baker, Gary V. Stephenson, and Hao Wen, “[Signal photon flux and background noise in a coupling electromagnetic detecting system for high-frequency gravitational waves](http://vixra.org/abs/0907.0030)”, [Phys. Rev. D 80, 064013] Published Wed Sep 9, 2009 ; <http://vixra.org/abs/0907.0030>

Bibliography, contd

- Ng, Y.Jack, "Article: [Spacetime Foam: From Entropy and Holography to Infinite Statistics and Nonlocality](#)" Entropy 2008, 10(4), 441-461; DOI: [10.3390/e10040441](#)
- G. t'Hooft., "The mathematical basis for Deterministic Quantum Mechanics," http://arxiv.org/PS_cache/quant-ph/pdf/0604/0604008v2.pdf, (2006).
- G.t'Hooft, "Determinism beneath Quantum Mechanics", http://arxiv.org/PS_cache/quant-ph/pdf/0212/0212095v1.pdf
- [Dimitar Valev](#)," Neutrino and graviton rest mass estimations by a phenomenological approach", Aerospace Res. Bulg. 22:68-82, 2008; <http://arxiv.org/abs/hep-ph/0507255>

Finally a good book summary with up to date summaries

- From “ Series in High Energy physics, Cosmology and Gravitation” - Taylor and Francis (publishers)
- - Particle and Astroparticle Physics (2008)
- By Utpal Sarkar