

THEORY OF ELECTRON

WU SHENG-PING

ABSTRACT. The solution with no singularity of wave equation for E-M fields is solved not to Bessel function, which's geometrical size is little enough to explain all effects in matter's structure: strong, weak effect or even other new ones. The mathematic calculation leded by quantum theory reveals the weak or strong decay and static properties of elementary particles, all coincide with experimental data, and a covariant equation comprising bent space is proposed to explain mass.

CONTENTS

1. Unit Dimension of sch	1
2. Quantization	2
3. Self-consistent Electrical-magnetic Fields	3
4. Stable Particle	3
5. Radium Function	4
6. Soluton	4
7. Electrons and Their Symmetries	5
8. Mechanic Feature	6
9. Propagation and Movement	7
10. Antiparticle and Radiation	10
11. Conservation Law and Balance Formula	11
12. Muon	11
13. Pion Positive	11
14. Pion Neutral	12
15. tau	12
16. Proton	12
17. Magic Numbers	13
18. Collision and Coupling	13
19. Conclusion	13
References	14

1. UNIT DIMENSION OF sch

A rebuilding of units and physical dimensions is needed. Time s is fundamental. The velocity of light is set to 1

$$Velocity : c = 1$$

Date: Nov 10, 2011.

Key words and phrases. Maxwell equations, Decay, Antiparticle.

Hence the dimension of length is

$$L : c(s)$$

The \hbar is set to 1

$$Energy : \hbar(s^{-1})$$

In Maxwell equations the following is set

$$c\epsilon = 1, c\mu = 1$$

One can have

$$\epsilon : \frac{Q^2}{\epsilon L};$$

$$\mu : \frac{\epsilon L}{c^2 Q^2}$$

$$UnitiveElectricalCharge : \sigma = \sqrt{\hbar}$$

It's very strange that the charge is analyzed as space and mass. Charge Q is then defined as Q/σ here, without unit.

$$\sigma = 1.03 \times 10^{-17} C = 64e, e/\sigma = e/\sigma = 1/64 = 1.56 \times 10^{-2}$$

$$H : Q/(LT) : \sqrt{\hbar}/c(s^{-2})$$

$$E : \epsilon/(LQ) : \sqrt{\hbar}/c(s^{-2})$$

If \hbar, c is taken as a number instead of unit, then all physical units is described as the powers of the second: s^n .

The unit of charge can be reset by *linear variation of charge-unit*

$$Q \rightarrow CQ, Q : \sigma/C$$

We define *all physical is the number as the unit of charge is $|e|$* . In order to recover to unit σ the physical, *when do this*, each physical has factor e^n/σ with n the power of charge. We will use it without detailed explanation to set the unit electron charge $Q_e = 1$.

2. QUANTIZATION

All discussion base on a explanation of quantization, or *real* probability explanation for quantum theory, which bases on a Transfer Probability Matrix (TPM)

$$P_i(x)M = P_f(x)$$

As a fact, that a particle appears in a point at rate 1 is independent with appearing at anther point at rate 1. There still another pairs of independent states

$$S_1 = e^{ipx}, S_2 = e^{ip'x}$$

because

$$\langle s_1, s_2 \rangle = \int dV s_1 s_2^* = N\delta(p - p')$$

In fact in the TPM formulation, it's been accepted for granted that the Hermitian inner-product is the measure of the dependence of two states, and it is also implied by the formula

$$P_1 M P_2^*$$

Depending on this view point one can constructs a wave

$$e^{ipx}$$

and gifts it with the momentum explanation p , Then all quantum theory is set up.

3. SELF-CONSISTENT ELECTRICAL-MAGNETIC FIELDS

The Maxwell equations are

$$\begin{aligned}\frac{\partial H}{\partial t} + \nabla \times E &= 0 \\ \frac{\partial E}{\partial t} - \nabla \times H + \mathbf{j} &= 0\end{aligned}$$

it's discussed that plat and straight space.

Try equation the free E-M field

$$(3.1) \quad \begin{aligned}A_i^{;j} - A_j^{;i} &= -iA^{\nu*} \cdot \partial_i A_\nu, Q = 1 \\ (A_i) &:= (-V, \mathbf{A}), j = (\rho, J) \\ \partial &:= (\partial_i) := (-\partial_t, \partial_{x_1}, \partial_{x_2}, \partial_{x_3})\end{aligned}$$

The equation 3.1 have symmetries

$$CPT, CcT$$

When choose gauge

$$\begin{aligned}\partial' A &= 0 \\ \partial' &:= (\partial^i) := (\partial_t, \partial_{x_1}, \partial_{x_2}, \partial_{x_3})\end{aligned}$$

this is valid that the continuous charge current meets

$$\partial^\mu \cdot (A^{\nu*} \cdot \partial_\mu A_\nu) = \partial' j = 0$$

and

$$\partial_\mu \partial^\mu (A^{\nu*} \cdot A_\nu) = 0$$

Then

$$\begin{aligned}2\nabla_\mu A^{\nu*} \nabla^\mu A_\nu &= 2\partial_t A_\nu \partial_t A^{\nu*} - A \partial_\mu \partial^\mu A_\nu - A^{\nu*} \partial_\mu \partial^\mu A_\nu \\ \nabla_\mu A_\nu \nabla^\mu A^{\nu*} &= \partial_t A_\nu \partial_t A^{\nu*} \\ U(t)U^*(t)\mathbf{p}^2 &\approx U^*(t)_t U(t)_t \cdot A = U(t)e^{i\mathbf{p}\mathbf{r}} \\ U(t) &= e^{-|\mathbf{p}|t} \\ \frac{\langle \Delta_\infty^t A | \Delta_\infty^t A \rangle}{\langle \Delta_\infty^t A_0 | \Delta_\infty^t A_0 \rangle} &\approx e^{-t\Delta_\infty^t \varepsilon}\end{aligned}$$

ε is the gross energy. This means exponent decay with respect to time.

4. STABLE PARTICLE

All particles are elementarily E-M fields is presumed. It's trying to find stable solution of the Maxwell equations *in complex domain*. One can write down the solution initially and correct it by re-substitution. Here is the initial state

$$V = V_1 e^{ikt}, A_1 = V$$

The static fields E_0, H_0

$$(4.1) \quad \begin{aligned}\nabla \cdot E_0 &= iA_1^{\nu*} \cdot \partial_t A_{1\nu} = -\rho_0 \\ \nabla \times H_0 &= -iA_1^{\nu*} \cdot \nabla A_{1\nu} = J_0\end{aligned}$$

In the first round of substitution

$$J_1 = -i(A_0^{\nu*} \cdot \partial A_{1\nu}) - i(\partial A_0^{\nu*} \cdot A_{1\nu})$$

5. RADIUM FUNCTION

Firstly

$$\nabla^2 A = -k^2 A$$

is solved. Exactly, it's solved in spherical coordinate

$$0 = r^2 \nabla^2 f + k^2 f = (r^2 f_r)_r + k^2 r^2 f + \frac{1}{\sin \theta} (\sin \theta f_\theta)_\theta + \frac{1}{\sin^2 \theta} (f_\phi)_\phi$$

Its solution is

$$\begin{aligned} f &= R\Theta\Phi = R_l Y_{lm} \\ \Theta &= P_l^m(\cos \theta), \Phi = \cos(\alpha + m\phi) \\ R_l &= N \eta_l(kr), \eta_l(r) = r^l \int_0^\infty \frac{(1-\lambda)^l}{(1+\lambda)^{l+2}} \cos(\lambda r) d\lambda \\ &\int_0^\infty dr \cdot r^2 R^2 = 1 \end{aligned}$$

R is solved like

$$\begin{aligned} (r^2 R_r)_r &= -k^2 r^2 R + l(l+1)R, l \geq 0 \\ R &\rightarrow rR' \\ (r^2 R')_{rr} &= -k^2 r^2 R' + l(l+1)R' \\ R' &\rightarrow r^{l-1} R' \\ rR'_{rr} + 2(l+1)R'_r + k^2 rR' &= 0 \\ r &\rightarrow r/k \\ (s^2 F)' + 2(l+1)F + F' &= 0, F = F(R') \end{aligned}$$

$F()$ is the Fourier transform

$$R' = \int_0^\infty \frac{(1-\lambda)^l}{(1+\lambda)^{l+2}} \cos(\lambda r) d\lambda$$

The function R' has zero derivative at $r = 0$ and is zero as $r \rightarrow \infty$.

6. SOLUTION

The derivatives of the function of electron has a strange breaking point in coordinate origin hence without normal convenience of Fourier transform. The following are some proximation of the first rank. The solution of $l = 1, m = 1, Q = e/\sigma$ is calculated or tested for electron.

$$A_1 = N R_1(kr) Y_{1,1},$$


The curve of R_1 is like the one in the figure 1.

The magnetic dipole moment μ_z is calculated as the first rank of proximation

$$\begin{aligned} \mu_z &= \langle A^\nu | -i\partial_\phi | A_\nu \rangle / 2 \\ &= 1/2, k_e = 1 \end{aligned}$$

The power of unit of charge is not equal, but it's valid for unit $Q = e$.

$$\frac{Q}{2k} = \mu_B$$

FIGURE 1. the shape of radium function R_1 by DFT

7. ELECTRONS AND THEIR SYMMETRIES

Some states of electrical field A are defined as the core of the electron

$$e_r^+ : NR_1(-kr)Y_{1,1}e^{-ikt},$$

$$e_r^- : NR_1(kr)Y_{1,1}e^{ikt}, (CPT)$$

$$e_l^+ = NR_{-z}(e_r^+) : R_1(-kr)Y_{1,-1}e^{-ikt}$$

$$e_l^- = NR_{-z}(e_l^-) : R_1(kr)Y_{1,-1}e^{ikt}$$

$$R_{-z} : \text{Rotation} : z \rightarrow -z, x \rightarrow x, y \rightarrow -y$$

We use these symbols e -s to express the complete field (E, M)

$$\langle R_1(kr)Y | R_1(kr)Y \rangle = 1$$

By charge normalization (the equation 3.1)

$$\rho = A^{\nu*} \cdot A_{\nu,t}$$

we get

$$N = \sqrt{1/(ke_{/\sigma})}$$

On the energy of electron, $\nabla \cdot A = 0$

$$\langle \nabla A^\nu | \nabla A_\nu \rangle / 2 = \langle A^\nu | \nabla^2 A_\nu \rangle / 2 = k_e$$

1) Energy of static E-field crossing

$$\begin{aligned} \varepsilon_e &= -k_e \int dV DV' \rho(\mathbf{r}) \rho(\mathbf{r}') / |4\pi(\mathbf{r} - \mathbf{r}')| \\ &= -e_{/\sigma} k_e \int dV \rho(\mathbf{r}) / r = -\frac{1}{4.876 \times 10^{-16} s} \end{aligned}$$

2) Energy of the static M-field crossing

$$\varepsilon_m = \varepsilon_e$$

because

$$\begin{aligned} &\int dV_1 dV_2 \cdot A^{\nu'}(\mathbf{r}_1) \partial_1^\mu A_{\nu'}^*(\mathbf{r}_1) A^{\nu*}(\mathbf{r}_2 - \mathbf{r}_1) \partial_{1\mu} A_\nu(\mathbf{r}_2 - \mathbf{r}_1) / |r_2| \\ &+ A^{\nu'}(\mathbf{r}_1) \partial_{1\mu} A_{\nu'}^*(\mathbf{r}_1) A^{\nu*}(\mathbf{r}_2 - \mathbf{r}_1) \partial_1^\mu A_\nu(\mathbf{r}_2 - \mathbf{r}_1) / |r_2| \end{aligned}$$

$$\begin{aligned}
&= A^{\nu'}(\mathbf{r}_1)A^{\nu*}(\mathbf{r}_2 - \mathbf{r}_1)(\partial_{1\mu}\partial_1^\mu(A_{\nu'}^*(\mathbf{r}_1))A_\nu(\mathbf{r}_2 - \mathbf{r}_1) - A_{\nu'*}(\mathbf{r}_1)\partial_{1\mu}\partial_1^\mu A_\nu(\mathbf{r}_2 - \mathbf{r}_1) \\
&\quad - A_\nu(\mathbf{r}_2 - \mathbf{r}_1)\partial_{1\mu}\partial_1^\mu A_{\nu'*}(\mathbf{r}_1))/|r_2| \\
&= 0
\end{aligned}$$

The value of a crossing term generated by static fields between electrons are

$$\begin{array}{ccccc}
n(\cdot 2\varepsilon_e) & e_r^+ & e_r^- & e_l^+ & e_l^- \\
e_r^+ & + & - & 0 & 0 \\
e_r^- & - & + & 0 & 0 \\
e_l^+ & 0 & 0 & + & - \\
e_l^- & 0 & 0 & - & +
\end{array}$$

As two electrons fold their crossing term generated by dynamic correction (of electron function) is calculated like

$$\partial^\mu \partial_\mu \partial V \approx \partial^2 A_0 \cdot V$$

Use the gauge $\partial' A = 0$

$$\begin{aligned}
\partial^\mu \partial_\mu \partial V &= -2\rho_t \\
\varepsilon_x &= -\frac{1}{k_e} \int dV \cdot V_1^* \cdot V_{1,t} \cdot V_1 \cdot V_1^* \\
&= -\frac{1}{4.50 \times 10^{-8} s}
\end{aligned}$$

The value of a crossing term generated by this correction between electrons are

$$\begin{array}{ccccc}
n(\cdot 2\varepsilon_x) & e_r^+ & e_r^- & e_l^+ & e_l^- \\
e_r^+ & - & 0 & 0 & - \\
e_r^- & 0 & - & - & 0 \\
e_l^+ & 0 & - & - & 0 \\
e_l^- & - & 0 & 0 & -
\end{array}$$

The first correction would affect the static energy ε_e , it's

$$\varepsilon_{ex} = -\frac{2k_e Q_e \varepsilon_x}{\varepsilon_e} = -\frac{1}{4.46 \times 10^{-6} s}$$

This effect also exists between two electrons in stable particle.

8. MECHANIC FEATURE

As two electrons meet and effect each other, their phases of the vibrations are also key, but the effect of phase is not observed. Considering two electrons with the same phase start from the same place and meet at the other one, the relative theory give a result that their phase are the same as they meet. If one defines unitary and orthogonal frame field P_i

$$DP_i = 0, P_i(O) \cdot P_j(O) = 0, |P_i| = 1$$

And the frame

$$g_i^i dx_i = P_i$$

and the free and orthogonal harmonic waves

$$e^{i \sum_i p_i g^i}$$

In fact *under this base P_i all differential is good as covariant and can be operated like in straight and flat space.* More over we have the covariant spectrum indexed by p .

One can guess that all the electrons in this cosmos are generated in the same place and the same time.

If the equation that connects space and E-M fields is written down for cosmos of electrons, it's the following:

$$(8.1) \quad R_{ij} - \frac{1}{2}Rg_{ij} = 8\pi GT_{ij}$$

$$e_{/\sigma}^2 T_{ij} = F_i^{k*} F_{kj} - \delta_{ij} F_{\mu\nu} F^{\mu\nu*} / 4$$

All tensors are expressed in base P_i . This equation give mass because the space is decided by E-M fields instantly. the factor $e_{/\sigma}^2$ is to balances the physical unit.

The Einstein's Theory of space and gravity is compatible with this theory and explains the energy of space and the looking mass k (generated by moving coordinate system) of particle.

In the group of electrons

$$A = \sum_i f_i e_i, \langle f_i | f_i \rangle = 1$$

It's called *propagation*. Each f_i is normalized to 1. The complete coupling electrons is

$$f \sum_i e_i$$

When the mechanical physical is discussed, the field must be normalized to energy. To say in details it's

$$\langle \nabla A^\nu | \nabla A_\nu \rangle / 2 = \langle C \nabla A^\nu | i \partial_t | C \nabla A_\nu \rangle, \nabla \cdot \mathbf{A} = 0$$

C is normalization factor.

The spin of electron is calculated as

$$\varepsilon = \langle \nabla A^\nu | \nabla A_\nu \rangle / 2 = k_e, \nabla \cdot \mathbf{A} = 0$$

$$\langle \nabla A^\nu | \partial_t | \nabla A_\nu \rangle = 2k_e^2, \nabla \cdot \mathbf{A} = 0$$

Its spin is

$$\langle C \nabla A^\nu | -i \partial_\phi | C \nabla A_\nu \rangle = 1/2$$

9. PROPAGATION AND MOVEMENT

Define symbols

$$e_{xr} := N \cdot R_1(k_x r) Y(1, 1) e^{iK_x t},$$

$$e_{xx} := (e_{xl} + e_{xr}) / \sqrt{2}$$

The following are also (stable) classical propagations.

<i>particle</i>	<i>electron</i>	<i>neutino</i>	<i>photon</i>
<i>notation</i>	e_r^+	ν_r	γ_r
<i>structure</i>	e_r^+	$(e_r^+ + e_l^-)$	$(e_r^+ + e_r^-)$

We have results by mathematic

$$\varsigma_{k,l,m}(x) := R_l(kr) Y_{l,m},$$

meets




FIGURE 2. the shape of distribution of momenta of electron fields in one direction, calculated through spherical Bessel functions

Theorem 9.1. C_A is a global area with its center in A and its diameter is r_A

$$\lim_{r_o=r_y \rightarrow 0} \int_{I-\Sigma C_i} dV \varsigma_{k,l,m}(x) \varsigma_{k,l,m}^*(x-y) = 0, y \neq O$$

$$\int dV \varsigma_{k,l,m}(x) \varsigma_{k',l',m'}(x) = 0, k \neq k' \text{ or } l \neq l' \text{ or } m \neq m'$$

Proof. Use the limit

$$\lim_{k' \rightarrow k} \lim_{r_o=r_y \rightarrow 0} \left(\int_{I-\Sigma C_i} dV \varsigma_{k,l,m}(x) - \varsigma_{k',l,m}(x-y) \right)$$

□

Theorem 9.2. if $e^{i\mathbf{pr}}, \varsigma_{k,l,m}$ is normalized to 1,

$$e^{i\mathbf{pr}} * \varsigma_{k,l,m} = e^{i\mathbf{pr}}$$

Definition 9.3.

$$e_{x\nu}^+ = e_x^{\nu*}$$

$$\langle A|A \rangle = \langle A^\nu | A_\nu \rangle$$

Proof. because

$$\int dV e^{i\mathbf{pr}} * \varsigma_{k,l,m} \cdot (e^{i\mathbf{pr}} * \varsigma_{k,l,m})^*$$

$$= \int dV e^{i\mathbf{pr}} (e^{i\mathbf{pr}})^* \cdot \int dV \varsigma_{k,l,m} (\varsigma_{k,l,m})^* = 1$$

□

The figure 2 is the shape of distribution of momenta of electron function e_x .

The movement of the propagation is called *Movement*, ie. the third level wave, for example

$$e^{i\mathbf{pr}-ikt} * \delta(\mathbf{r}) * e$$

Theorem 9.4.

$$\nabla(e_x * e) = (\nabla e_x) * e, k_x \gg k_e$$

$$k_e (\nabla e_x) * e = k_x e_x * (\nabla e)$$

Proof. Calculate

$$\int dV_x \nabla_x (e_x(x) * e(x) \cdot \delta(x - x')) = \int dV_x \nabla_x (e_x(x) (e(x) * \delta(x - x'))),$$

$$\langle (\nabla e_x) * e | (\nabla e_x) * e \rangle$$

□

Definition 9.5.

$$\langle f_1(x_1) + f_2(x_2) | O(x) | f_1(x_1) + f_2(x_2) \rangle$$

$$= \lim_{V \rightarrow I} \left(\int_V dV_1 \int_V dV_2 \cdot (f_1(x_1) + f_2(x_2))^* (O(x_1) + O(x_2)) (f_1(x_1) + f_2(x_2)) \right) / V$$

The field of two decoupling system

$$F = F_1 + F_2$$

$$F_1 = \sum_i f_i^* \nabla e_i(x), F_2 = \sum_i g_i^* \nabla e_i(x')$$

The additive physical of Einstein tensor is adopted to express mechanics, which has probability expression. The energy

$$\epsilon = \langle F_1 + F_2 | F_1 + F_2 \rangle / e_{j\sigma}^2 = \langle C_1 F_1 + C_2 F_2 | i \partial_t | C_1 F_1 + C_2 F_2 \rangle$$

$$\langle C_1 F_1 | C_1 F_1 \rangle = 1, \langle C_2 F_2 | C_2 F_2 \rangle = 1$$

Energy is coupling at its elementary property.

Its static MDM (magnetic dipole moment) for wave F_1

$$\int d\mathbf{r} \cdot \sum_i f_i * \nabla e_i = \sqrt{\frac{k_{e_i}}{k_x}} \sum_i f_i * e_i$$

$$\mu = \langle \sum_i C_i^2 f_i * e_i(x_i) | \sum_i C_i^2 (-i \partial_\phi f_i) * e_i(x_i) \rangle$$

$$\langle \sum_i C_i f_i * e_i(x_i) | \sum_i C_i f_i * (-i \partial_\phi e_i(x_i)) \rangle$$

$$C_i = \sqrt{\frac{k_{e_i}}{k_{x_i}}}$$

Electrons is decoupled. Coupling part is interaction. The two parts of currents are j_0 and j_{int} that marks the interaction of electron that causes collapse of the system. Its spin (decoupling) is

$$S = \langle C_1 F_1 | \mathbf{r} \times \nabla | C_1 F_1 \rangle$$

F_1 is decoupled.

$$S_z = \langle \sum_i f_i * \nabla e_i(x_i) | \sum_i (-i \partial_\phi f_i) * \nabla e_i(x_i) \rangle / (2k_x)$$

$$+ \langle \sum_i f_i * \nabla e_i(x_i) / k_{e_i} | \sum_i f_i * (-i \partial_\phi \nabla e_i(x_i)) \rangle / (2k_e)$$

For a coupling electrons system x

$$F = f * \sum_i \nabla e_i, \langle f | f \rangle = 1$$

Set

$$f = U(\mathbf{r}) e^{-iKt}$$

Substitute this into

$$\partial' \cdot \partial A = 0, A = \int d\mathbf{r} \cdot F, \nabla \cdot A = 0$$

It's solved to

$$f = R_1(k_x r) Y e^{-iKt}, K^2 + k_e^2 + 2k_e K Q / (Q_e) = k_x^2$$

10. ANTIPARTICLE AND RADIATION

The radiation of photon is derive from this reaction

$$e^{ip_1x} * e_r^+ + e^{ip_2x} * e_r^- \rightarrow e^{ip_3x} * \gamma_r$$

The emission (of E-M fields), that's the reason to react forward but is not the all energy variation related, is

$$4\varepsilon_e = \frac{1}{1.219 \times 10^{-16} s}$$

this energy marks the intension of electromagnet effect.

The wave of photon

$$e^{i\mathbf{pr}+ikt} * (e_r^+ + e_r^-)$$

has a mechanic field that describes a movement of a mass

$$k_e - k_e = 0$$

The equivalent reaction is like

$$e^{ip_1x} * e_r^+ \rightarrow e^{-ip_2x} * \overline{e_r^-} + e^{ip_3x} * \gamma_r$$

$\overline{e_l^+}$ is just the equivalent for the equilibrium after the particle e_r^- is shifted to the other side of the reaction. In fact the shift is a transform of conjugation

$$\overline{e_r^-} = (e_r^-)^* \approx -e_l^+$$

The normal matter is called positive matter and this kind above is called antiparticle conventionally. (this term is different from the one derived by *CPT*)

Antimatter happens by reversing the world's line, with the same map of the event.

The radiation of neutrino depends the reaction

$$e_r^+ + e_l^- \rightarrow \nu_r$$

This reaction is with emission of an energy

$$2\varepsilon_x = \frac{1}{2.25 \times 10^{-8} s}$$

this energy marks the intension of weak effect (of this kind). As a testifying one can have

$$4\varepsilon_e : 2\varepsilon_x = 8.46 \times 10^7$$

This is the difference of the intension between electromagnetic effect and weak effect.

The antiparticle meets

$$F(A^*) = 0$$

F is the formula 3.1. The mixing of antiparticle field A and particle field B have the law

$$(10.1) \quad F(A^* + P) = 0$$

A is

$$A_1 \xrightarrow{it} A_2$$

Get the conjugation

$$A_2^* \xrightarrow{it} A_1^*$$

Another explanation is

$$\sum e^{ipr} e^{ikt}, A_i \rightarrow A_f$$

It has conjugation

$$\sum e^{-ipr} e^{ik(-t)}, A_f^* \rightarrow A_i^*$$

The formula 10.1 is reasonable expansion of the law for real number fields to that of complex fields $A + P$.

11. CONSERVATION LAW AND BALANCE FORMULA

No matter in E-M fields (the elementary) level or in movement (the third) level, the conservation law is *conservation of momentum and conservation of angular momentum*. A *balance formula* for a reaction is the equivalent formula in positive matter, ie. after all anti-matter is shifted to the other side of the reaction formula. Balance formula is suitable for the analysis of the energy transition of E-M fields in the reaction. The *invariance of electron itself* in reaction is also a conservation law.

12. MUON

μ^+ is composed of

$$\mu_r^+ : e_{\mu x} * (e_r^+ - \nu_l)$$

μ is with mass $3k_e/e/\sigma = 3 \times 64k_e$, ($k_\mu \approx m_\mu/3$) spin 1/2, MDM $\mu_B k_e/k_\mu$.

The main channel of decay

$$\mu_r^+ \rightarrow -e_r^+ + \nu_r - \nu_l$$

is with balance formula

$$e_{\mu x} * e_r^+ + e^{-ip_1 x} * e_l^- + e^{-ip_2 x} * \nu_l \rightarrow e_{\mu x}^* * \nu_l + e^{ip_3 x} * \nu_r$$

From the theorem 9.2 we can find the only gap of crossing energy between the two sides is in the first static energy correction

$$2\varepsilon_{ex} = \frac{1}{2.23 \times 10^{-6} s} [2.1970 \times 10^{-6} s][1]$$

The data in square bracket is experimental data of the full width.

13. PION POSITIVE

Pion positive is

$$\pi_r^+ : e_{\pi y} * (-\nu_l + e_r^+) + e_{\pi x} * (\nu_r)$$

It's with mass $3 - 5 \times 64k_e$, spin 1/2 and MDM $\mu_B k_e/k_{\pi^+}$.

Decay Channels:

$$\pi^+ \rightarrow \mu_r^+ + \nu_r$$

It's with balance formula

$$e_{\pi x} * \nu_r + e^{-ip_1 x} * e_{\mu x}^* * \nu_l + e_{\pi y} * e_r^+ \rightarrow e_{\pi y}^* * \nu_l + e^{ip_2 x} * \nu_r + e^{ip_1 x} * e_{\mu x} * e_r^+$$

Wave e^{ipx} is subjected by $\delta(x)$. The emission of energy is weak interaction in strong interaction term

$$2\varepsilon_x = \frac{1}{2.25 \times 10^{-8}s} \quad [(2.603 \times 10^{-8}s)[1]]$$

The referenced data is the full width.

14. PION NEUTRAL

Pion neutral is atom-like particle

$$\pi^0 : f^+ * \nu_r + f^- * \nu_l, \langle f^+, f^- \rangle = 0$$

It has mass $4k_e$, zero spin and zero MDM. Its decay modes are

$$\pi^0 \rightarrow \gamma_r + \gamma_l$$

The loss of energy is from static field

$$8\varepsilon_e = \frac{1}{6.1 \times 10^{-17}s} \quad [8.4 \times 10^{-17}s][1]$$

15. TAU

τ maybe that

$$\tau^- : 5e_r^+ - 5e_l^- + e_r^-$$

Its mass $51 \times 64k_e$, spin 1/2, MDM $50\mu_B/k_\mu$. It has decay mode

$$\tau^- \rightarrow -\mu_r^- + \nu_l - \nu_r$$

$$e_{\tau x} * 5e_r^+ + e_{\tau x} * e_r^- + e^{-ip_1 x} * e_{\mu x}^* * e_l^+ + e^{-ip_2 x} * \nu_r \rightarrow e_{\tau x}^* * 5e_r^+ + e^{ip_1 x} * e_{\mu x} * \nu_r + e^{ip_3 x} * \nu_l$$

The loss of energy is the difference of the static fields

$$e_{\tau x} * 5e_r^+ + e_{\tau x} * e_r^- \leftrightarrow e_{\tau x}^* * 5e_r^+ + e^{ip_3 x} * \nu_l$$

$$e_{\tau x} * (5e_r^+ + e_r^-) \leftrightarrow (5e_r^+ + e_r^-)$$

Calculating the difference between $X = \tau$ and $X = \delta$ we can find the emission of static E-M fields

$$\Gamma_1 = \frac{5\varepsilon_e}{k_\tau/k_e}$$

$$= \frac{1}{1.7 \times 10^{-13}s} \quad [2.9 \times 10^{-13}s, BR.0.17][1]$$

From the shape of momentum distribution I can find many experimental data has a shift of initial velocity, I judge many resonance states is evaluated with larger mass than the real. With zero initial velocity the momentum distribution is like the figure 2, with the steep edge crosses grid origin directly.

16. PROTON

Proton may be like

$$p^+ : e_{px} * (-4e_r^+ + 3e_l^+ + e_r^- - e_l^-)$$

The mass is $27 \times 64k_e$ that's very close to the real mass. The MDM is calculated as $3\mu_N$, spin is 1/2. The proton thus designed is eternal because even if decay to the finest small parts the emission is negative.

17. MAGIC NUMBERS

We define an unit: Mass-number Unite

$$m := m_e \sigma / e \approx 64 k_e$$

And we presume the Mass-number (in fact relates theoretical electron number) in a particle for the four kinds of electrons are

$$e_r^+ : i, e_r^- : j, e_l^+ : k, e_l^- : l$$

The the designation of a particle is an equation

$$\begin{cases} i^2 + j^2 + k^2 + l^2 = M/m \\ i - j + k - l = Q \\ \pm i \pm j \pm k \pm l = 2S \end{cases}$$

According to Lagrange's four Square theorem, Any integer can be sum of some four square of integers. But after adding the constraints of charge number or spin number the conditions are not so simple as the Lagrange's theorem.

If consider more complicated design like

$$i' e_r^+ + \bar{i} e_l^-, i' - \bar{i} = i$$

The equations for charge, mass and spin are

$$\begin{cases} (i' - \bar{i})^2 + (j' - \bar{j})^2 + (k' - \bar{k})^2 + (l' - \bar{l})^2 = M/m \\ (i' + \bar{i}) - (j' + \bar{j}) + (k' + \bar{k}) - (l' + \bar{l}) = Q \\ (i' - \bar{i}) + (j' - \bar{j}) - (k' - \bar{k}) - (l' - \bar{l}) = 2S \end{cases}$$

18. COLLISION AND COUPLING

The key part is to calculate the coupling that's calculated by the cross part in space-time domain for the two beams.

19. CONCLUSION

The relative theory is applied to electromagnetic wave to give the looking mass of the fields which does expresses mass, for example the solved electron function in this article. In my view point the sum-up of the grains (as electrons) of electromagnetic field is a mechanic movement with diverse effect. Fortunately this model will explain all the effects in the known world: strong, weak and electromagnetic effects, and even subclassify them further if not add new ones. In this model the only field is electromagnetic field except space, this stands for the philosophical with the point of that unified world from unique source. All depend on a simple fact: the gross momentum in a system is time-invariant, and we can devise the E-M momentum to analysis current.

The inertial mass is deduced by mechanical operator $i\partial_t$. But the gravitational mass (by the equation of 8.1) of the naked electron is 64 time of the inertial and mechanical mass, the photon and neutrino has zero mechanical mass but their gravitational mass is not zero obviously. this is hard problem unsettled by this article. For atom the inertial mass less then gravitational mass by 1/50 approximately.

The energy of matter would happen in this process, the hot matter distilled to protons as got cold with their wave functions dependent each others. the harmony between bent space and electromagnetic fields explain them all.

Except electron function my description of particles in fact has the same form with Quantum Electromagnetic Mechanics, and they two should reach the same result except for precision. But my theory isn't compatible to the theory of quarks, the upper part of standard model, if not it is calculated in the style of Quantum Electromagnetic Mechanics. In fact, The electron function is a good promotion for the experimental model of proton that went up very early.

I found these presumptions on some days of 1994-1995 and soon I grossly testify this theory the year. At that time a few people studied in HUST China knew of it. But in the following teen years I nearly forgot of it except now and several years ago a round of submission of it.

REFERENCES

- [1] K. Nakamura et al. (Particle Data Group), *JPG* 37, 075021 (2010) (URL: <http://pdg.lbl.gov>)

WUHAN UNIVERSITY, WUHAN, HUBEI PROVINCE, THE PEOPLE'S REPUBLIC OF CHINA. POST-CODE: 430074

E-mail address: hiyaho@126.com