

Programming Planck units from a virtual electron; a Simulation Hypothesis

Malcolm J. Macleod

macleod@platoscode.com

The simulation hypothesis proposes that all of reality is an artificial simulation. In this article I describe a method for programming the Planck units as simple geometrical forms within a virtual universe simulation environment. These forms are constructs of 2 mathematical (dimensionless) constants; the fine structure constant α and $\Omega = 2.0071349496...$ such that mass $M=1$, time $T=2\pi$, velocity $V = 2\pi\Omega^2$, length $L=2\pi^2\Omega^2 ...$ The 5 SI units kg, m, s, A, k are replaced by a single unit u that defines the relationships between the SI units. To convert from these base geometries to their respective SI numerical values requires 2 scalars, for example the SI values of Planck mass and the speed of light c . Using these 2 scalars with α and Ω , we can solve the remaining Planck units and the physical constants G, h, e, m_e, k_B . Results are consistent with CODATA 2014 (table). Solving for the electron f_e , we find that the scalars and u cancel leaving the formula ($f_e = 4\pi^2 r^3; r = 2^6 3\pi^2 \alpha \Omega^5$, units = 1). Thus although we can construct the electron from Planck units, the electron itself is a dimensionless mathematical constant (independent of the system of units used). Inversely we can take this virtual electron and use it to construct the Planck units. It is the dimensionless geometry of the Planck units which confers the functionality of the unit, this u is a rule set rather than a physical entity. This resolves the principal difficulty of the mathematical universe as being how to relate mathematical forms to their corresponding physical units. It is these relationships between the units and not any inherent 'physical-ness' that gives the perception of a physical reality. The rationale for the electron was derived via the sqrt of Planck momentum and a black-hole electron model as a function of magnetic-monopoles and time.

| Table 1 | Calculated* (α, Ω, k, v) | CODATA 2014 |
|----------------------|--|--|
| Speed of light | $V = 299792458 u^{17}$ | $c = 299792458$ |
| Permeability | $\mu_0^* = 4\pi/10^7 u^{56}$ | $\mu_0 = 4\pi/10^7$ |
| Rydberg constant | $R_\infty^* = 10973731.568 508 u^{13}$ | $R_\infty = 10973731.568 508(65) [15]$ |
| Planck constant | $h^* = 6.626 069 134 e-34 u^{19}$ | $h = 6.626 070 040(81) e-34 [16]$ |
| Elementary charge | $e^* = 1.602 176 511 30 e-19 u^{-27}$ | $e = 1.602 176 6208(98) e-19 [19]$ |
| Electron mass | $m_e^* = 9.109 382 312 56 e-31 u^{15}$ | $m_e = 9.109 383 56(11) e-31 [17]$ |
| Boltzmann's constant | $k_B^* = 1.379 510 147 52 e-23 u^{29}$ | $k_B = 1.380 648 52(79) e-23 [22]$ |
| Gravitation constant | $G^* = 6.672 497 192 29 e-11 u^6$ | $G = 6.674 08(31) e-11 [21]$ |

keywords: computer universe, virtual universe, mathematical universe, simulated universe, sqrt Planck momentum, Planck unit, magnetic-monopole, fine structure constant alpha, Omega, black-hole electron;

1 Background

The general universe simulation hypothesis proposes that all of reality, including the earth and the universe, is in fact an artificial simulation, analogous to a computer simulation, and as such our reality is an illusion [2].

Mathematical platonism is a metaphysical view that there are abstract mathematical objects whose existence is independent of us [1]. Mathematical realism holds that mathematical entities exist independently of the human mind. Thus humans do not invent mathematics, but rather discover it. Triangles, for example, are real entities, not the creations of the human mind [3].

Max Tegmark's Mathematical Universe Hypothesis: Our external physical reality is a mathematical structure. That is, the physical universe is mathematics in a well-defined sense, and "in those [worlds] complex enough to contain self-aware substructures [they] will subjectively perceive themselves as existing in a physically 'real' world" [10].

Planck units (defined here Planck mass m_P , Planck time t_P , Planck length l_P , Planck charge A_Q , Planck temperature T_P) are a set of units of measurement also known as natural units because the origin of their definition comes only from properties of nature and not from any human construct.

2 The virtual universe

Mathematical universe hypotheses presume that our physical universe has an underlying mathematical origin. The principal difficulty of such hypotheses is the problem of constructing physical units such as mass, space and time from their respective mathematical forms.

This article describes a mathematical universe model that uses geometrical forms to duplicate the Planck units. These geometrical forms are constructed from 2 mathematical constants; the fine structure constant α and a presumed constant Ω such that $M=1$, time $T=2\pi$, velocity $V = 2\pi\Omega^2$, length $L=2\pi^2\Omega^2 ...$ eq(42-47).

The 5 SI units kg, m, s, A, k are replaced by a single unit u .

This u defines the relationships between the SI units, it is thus a rule-base and not a descriptive of an actual physical entity. The functionality of the unit is embedded into the geometry.

To solve the physical constants in SI terms also requires 2 scalars to convert from the base geometries to their respective SI values (4.2). For example, using the following values for α, Ω, k, v gives the results in the table for G, h, c, e, m_e, k_B ;

$$\begin{aligned} k &= m_p = .2176728175... \times 10^{-7} u^{15} \text{ (kg)} \\ v &= (2\pi\Omega^2)/c = 11843707.9... u^{17} \text{ (m/s)} \\ \alpha &= 137.035999139 \text{ CODATA mean (4.4)} \\ \Omega &= 2.0071349496...; \text{ (4.5)} \end{aligned}$$

Solving for the electron (4.3) gives

$$f_e = 4\pi^2 r^3, k^0 v^0 u^0; (r = 2^6 3\pi^2 \alpha \Omega^5)$$

The electron is thus a mathematical formula constructed from dimensionless constants α, Ω ($k^0 v^0 u^0 = 1$).

3 The virtual electron

In this section I introduce the sqrt of momentum as a distinct Planck unit and suggest how this could be used as a link between the mass and charge domains.

From the formulas for the charge constants I then derive a formula for a magnetic monopole (ampere-meter AL) and from this subsequently a formula for an electron function f_e .

The electron formula is constructed from monopoles (AL) and from time T yet it is also dimensionless as the charge and time units are not independent but rather are related, collapsing within the electron whereby; $f_e = (AL)^3/T$, units = 1. Being dimensionless and so independent of any system of units, this electron formula is a mathematical constant.

Note: for convenience I use the commonly recognized value for alpha as $\alpha \sim 137.036$.

Defining Q as the sqrt of Planck momentum where Planck momentum = $m_p c = 2\pi Q^2 = 6.52485... \text{ kg.m/s}$, and a unit q whereby $q^2 = \text{kg.m/s}$ giving;

$$Q = 1.019 \ 113 \ 411..., \text{ unit} = q \quad (1)$$

$$\begin{aligned} \text{Planck momentum; } &2\pi Q^2, \text{ units} = q^2, \\ \text{Planck length; } &l_p, \text{ units} = m = q^2 s / \text{kg}, \\ &c, \text{ units} = m/s = q^2 / \text{kg}; \end{aligned}$$

3.1. In Planck terms the mass constants are typically defined in terms of Planck mass, here I use Planck momentum;

$$m_p = \frac{2\pi Q^2}{c}, \text{ unit} = \text{kg} \quad (2)$$

$$E_p = m_p c^2 = 2\pi Q^2 c, \text{ units} = \frac{\text{kg.m}^2}{\text{s}^2} = \frac{q^4}{\text{kg}} \quad (3)$$

$$t_p = \frac{2l_p}{c}, \text{ unit} = \text{s} \quad (4)$$

$$F_p = \frac{2\pi Q^2}{t_p}, \text{ units} = \frac{q^2}{\text{s}} \quad (5)$$

3.2. The charge constants in terms of Q^3, c, α, l_p ;

$$A_Q = \frac{8c^3}{\alpha Q^3}, \text{ unit } A = \frac{m^3}{q^3 s^3} = \frac{q^3}{\text{kg}^3} \quad (6)$$

$$e = A_Q t_p = \frac{8c^3}{\alpha Q^3} \cdot \frac{2l_p}{c} = \frac{16l_p c^2}{\alpha Q^3}, \text{ units} = A.s = \frac{q^3 s}{\text{kg}^3} \quad (7)$$

$$T_p = \frac{A_Q c}{\pi} = \frac{8c^3}{\alpha Q^3} \cdot \frac{c}{\pi} = \frac{8c^4}{\pi \alpha Q^3}, \text{ units} = \frac{q^5}{\text{kg}^4} \quad (8)$$

$$k_B = \frac{E_p}{T_p} = \frac{\pi^2 \alpha Q^5}{4c^3}, \text{ units} = \frac{\text{kg}^3}{q} \quad (9)$$

3.3. As with c , the permeability of vacuum μ_0 has been assigned an exact numerical value so it is our next target. The ampere is that constant current which, if maintained in two straight parallel conductors of infinite length, of negligible circular cross section, and placed 1 meter apart in vacuum, would produce between these conductors a force equal to exactly 2.10^{-7} newton per meter of length.

$$\frac{F_{electric}}{A_Q^2} = \frac{F_p}{\alpha} \cdot \frac{1}{A_Q^2} = \frac{2\pi Q^2}{\alpha t_p} \cdot \left(\frac{\alpha Q^3}{8c^3}\right)^2 = \frac{\pi \alpha Q^8}{64l_p c^5} = \frac{2}{10^7} \quad (10)$$

$$\mu_0 = \frac{\pi^2 \alpha Q^8}{32l_p c^5} = \frac{4\pi}{10^7}, \text{ units} = \frac{\text{kg.m}}{\text{s}^2 A^2} = \frac{\text{kg}^6}{q^4 s} \quad (11)$$

3.4. Rewriting Planck length l_p in terms of Q, c, α, μ_0 ;

$$l_p = \frac{\pi^2 \alpha Q^8}{32\mu_0 c^5}, \text{ unit} = \frac{q^2 s}{\text{kg}} = m \quad (12)$$

3.5. A magnetic monopole in terms of Q, c, α, l_p ;

The ampere-meter is the SI unit for pole strength (the product of charge and velocity) in a magnet ($Am = ec$). A magnetic monopole σ_e is a hypothetical particle that is a magnet with only 1 pole [12]. I propose a magnetic monopole σ_e from α, e, c ($\sigma_e = 0.13708563 \times 10^{-6}$);

$$\sigma_e = \frac{3\alpha^2 ec}{2\pi^2}, \text{ units} = \frac{q^5 s}{\text{kg}^4} \quad (13)$$

I then use this monopole to construct an electron frequency function f_e ($f_e = 0.2389545 \times 10^{23}$);

$$f_e = \frac{\sigma_e^3}{t_p} = \frac{2^8 3^3 \alpha^3 l_p^2 c^{10}}{\pi^6 Q^9} = \frac{3^3 \alpha^5 Q^7}{4\pi^2 \mu_0^2}, \text{ units} = \frac{q^{15} s^2}{\text{kg}^{12}} \quad (14)$$

3.6. The most precisely measured of the natural constants is the Rydberg constant R_∞ (see table) and so it is important to this model. The unit for R_∞ is $1/m$. For m_e see eq(22);

$$R_\infty = \frac{m_e e^4 \mu_0^2 c^3}{8h^3} = \frac{2^5 c^5 \mu_0^3}{3^3 \pi \alpha^8 Q^{15}}, \text{ units} = \frac{1}{m} = \frac{\text{kg}^{13}}{q^{17} s^3} \quad (15)$$

This however now gives us 2 solutions for length m , see eq(1) and eq(15), if they are both valid then there must be a ratio whereby the units q, s, kg overlap and cancel;

$$m = \frac{q^2 s}{kg} \cdot \frac{q^{15} s^2}{kg^{12}} = \frac{q^{17} s^3}{kg^{13}}; \text{ thus } \frac{q^{15} s^2}{kg^{12}} = 1 \quad (16)$$

and so we can further reduce the number of units required, for example we can define s in terms of kg, q ;

$$s = \frac{kg^6}{q^{15/2}} \quad (17)$$

$$\mu_0 = \frac{kg^6}{q^4 s} = q^{7/2} \quad (18)$$

3.7. We find that this ratio is embedded in that electron function f_e (eq 14), and so f_e is a dimensionless mathematical constant whose function appears to be dictating the frequency of the Planck units;

$$f_e = \frac{\sigma_e^3}{t_p}; \text{ units} = \frac{q^{15} s^2}{kg^{12}} = 1 \quad (19)$$

Replacing q with the more familiar m gives this ratio;

$$q^2 = \frac{kg \cdot m}{s}; q^{30} = \left(\frac{kg \cdot m}{s}\right)^{15} = \frac{kg^{24}}{s^4} \quad (20)$$

$$\text{units} = \frac{kg^9 s^{11}}{m^{15}} = 1 \quad (21)$$

Electron mass as frequency of Planck mass:

$$m_e = \frac{m_p}{f_e}, \text{ unit} = kg \quad (22)$$

Electron wavelength via Planck length:

$$\lambda_e = 2\pi l_p f_e, \text{ units} = m = \frac{q^2 s}{kg} \quad (23)$$

Gravitation coupling constant:

$$\alpha_G = \left(\frac{m_e}{m_p}\right)^2 = \frac{1}{f_e^2}, \text{ units} = 1 \quad (24)$$

3.8. The Rydberg constant $R_\infty = 10973731.568508(65)$ [15] has been measured to a 12 digit precision. The known precision of Planck momentum and so Q is low, however with the solution for the Rydberg eq(15) we may re-write Q as Q^{15} in terms of; c, μ_0, R and α ;

$$Q^{15} = \frac{2^5 c^5 \mu_0^3}{3^3 \pi \alpha^8 R}, \text{ units} = \frac{kg^{12}}{s^2} = q^{15} \quad (25)$$

Using the formulas for Q^{15} eq(25) and l_p eq(12) we can re-write the least accurate dimensioned constants in terms of the

most accurate constants; R, c, μ_0, α . I first convert the constants until they include a Q^{15} term which can then be replaced by eq(25). Setting unit x as;

$$\text{unit } x = \frac{kg^{12}}{q^{15} s^2} = 1 \quad (26)$$

Elementary charge $e = 1.602\ 176\ 51130\ \text{e-19}$ (table p1)

$$e = \frac{16l_p c^2}{\alpha Q^3} = \frac{\pi^2 Q^5}{2\mu_0 c^3}, \text{ units} = \frac{q^3 s}{kg^3} \quad (27)$$

$$e^3 = \frac{\pi^6 Q^{15}}{8\mu_0^3 c^9} = \frac{4\pi^5}{3^3 c^4 \alpha^8 R}, \text{ units} = \frac{kg^3 s}{q^6} = \left(\frac{q^3 s}{kg^3}\right)^3 \cdot x \quad (28)$$

Planck constant $h = 6.626\ 069\ 134\ \text{e-34}$

$$h = 2\pi Q^2 2\pi l_p = \frac{4\pi^4 \alpha Q^{10}}{8\mu_0 c^5}, \text{ units} = \frac{q^4 s}{kg} \quad (29)$$

$$h^3 = \left(\frac{4\pi^4 \alpha Q^{10}}{8\mu_0 c^5}\right)^3 = \frac{2\pi^{10} \mu_0^3}{3^6 c^5 \alpha^{13} R^2}, \text{ units} = \frac{kg^{21}}{q^{18} s} = \left(\frac{q^4 s}{kg}\right)^3 \cdot x^2 \quad (30)$$

Boltzmann constant $k_B = 1.379\ 510\ 14752\ \text{e-23}$

$$k_B = \frac{\pi^2 \alpha Q^5}{4c^3}, \text{ units} = \frac{kg^3}{q} \quad (31)$$

$$k_B^3 = \frac{\pi^5 \mu_0^3}{3^3 2c^4 \alpha^5 R}, \text{ units} = \frac{kg^{21}}{q^{18} s^2} = \left(\frac{kg^3}{q}\right)^3 \cdot x \quad (32)$$

Gravitation constant $G = 6.672\ 497\ 19229\ \text{e-11}$

$$G = \frac{c^2 l_p}{m_p} = \frac{\pi \alpha Q^6}{64\mu_0 c^2}, \text{ units} = \frac{q^6 s}{kg^4} \quad (33)$$

$$G^5 = \frac{\pi^3 \mu_0}{2^{20} 3^6 \alpha^{11} R^2}, \text{ units} = kg^4 s = \left(\frac{q^6 s}{kg^4}\right)^5 \cdot x^2 \quad (34)$$

Planck length

$$l_p^{15} = \frac{\pi^{22} \mu_0^9}{2^{35} 3^{24} c^{35} \alpha^{49} R^8}, \text{ units} = \frac{kg^{81}}{q^{90} s} = \left(\frac{q^2 s}{kg}\right)^{15} \cdot x^8 \quad (35)$$

Planck mass

$$m_p^{15} = \frac{2^{25} \pi^{13} \mu_0^6}{3^6 c^5 \alpha^{16} R^2}, \text{ units} = kg^{15} = \frac{kg^{39}}{q^{30} s^4} \cdot \frac{1}{x^2} \quad (36)$$

Electron mass $m_e = 9.109\ 382\ 31256\ \text{e-31}$

$$m_e^3 = \frac{16\pi^{10} R \mu_0^3}{3^6 c^8 \alpha^7}, \text{ units} = kg^3 = \frac{kg^{27}}{q^{30} s^4} \cdot \frac{1}{x^2} \quad (37)$$

Ampere

$$A_Q^5 = \frac{2^{10} \pi^3 c^{10} \alpha^3 R}{\mu_0^3}, \text{ units} = \frac{q^{30} s^2}{kg^{27}} = \left(\frac{q^3}{kg^3}\right)^5 \cdot \frac{1}{x} \quad (38)$$

3.9. ($r = \sqrt{q}$)

There is a solution for an $r^2 = q$, it is the radiation density constant from the Stefan Boltzmann constant σ ;

$$\sigma = \frac{2\pi^5 k_B^4}{15h^3 c^2}, \quad r_d = \frac{4\sigma}{c}, \quad \text{units} = r \quad (39)$$

$$r_d^3 = \frac{3^3 4\pi^5 \mu_0^3 \alpha^{19} R^2}{5^3 c^{10}}, \quad \text{units} = \frac{kg^{30}}{q^{36} s^5} \cdot \frac{1}{x^2} = \frac{kg^6}{q^6 s} = r^3 \quad (40)$$

4 Geometrical units

4.1. The formula for the electron f_e incorporates dimensionful quantities but itself is dimensionless. This means that its numerical value is a mathematical constant, independent of which set of units we may use; $f_e = .23895453... \times 10^{23}$ units = 1. Setting $T = t_p$, $A = eT$, $V = 2l_p/T$;

$$f_e = \left(\frac{3\alpha^2 ATV}{2\pi^2}\right)^3 \frac{1}{T} = 0.239 \times 10^{23}, \quad \text{unit} = 1 \quad (41)$$

Because the numerical value is fixed and units = 1, we can look for other, non SI sets of ATV (AL) units which can also be used to solve f_e . In this section I describe 1 solution that constructs the base units from a geometrical component in terms of 2 fixed dimensionless mathematical constants; the fine structure constant alpha and a proposed Omega (4.5), from 2 variable unit-dependent scalars (from the list *ktlvpa*), and a unit u that replaces the SI units (kg, m, s, A, k).

$$M = (1)k, \quad \text{unit} = u^{15} \text{ (mass)} \quad (42)$$

$$T = (2\pi)t, \quad \text{unit} = u^{-30} \text{ (time)} \quad (43)$$

$$P = (\Omega)p, \quad \text{unit} = u^{16} \text{ (sqrt of momentum)} \quad (44)$$

$$V = (2\pi\Omega^2)v, \quad \text{unit} = u^{17} \text{ (velocity)} \quad (45)$$

$$L = (2\pi^2\Omega^2)l, \quad \text{unit} = u^{-13} \text{ (length)} \quad (46)$$

$$A = \left(\frac{2^6\pi^3\Omega^3}{\alpha}\right)a, \quad \text{unit} = u^3 \text{ (ampere)} \quad (47)$$

4.2 In the previous section we needed only 2 base units to define the others. In this example I use P, V to derive $MLTA$ and then the physical constants. Scaling p, v to their SI values gives $M=m_p, L=l_p, T=t_p, V=c, P=Q$

$$P = (\Omega)p, \quad \text{unit} = u^{16} \quad (48)$$

$$V = (2\pi\Omega^2)v, \quad \text{unit} = u^{17} \quad (49)$$

$$M = \frac{2\pi P^2}{V} = (1)\frac{P^2}{v}, \quad \text{unit} = u^{16 \cdot 2 - 17 = 15} \quad (50)$$

$$T^2 = (2\pi\Omega)^{15} \frac{P^9}{2\pi V^{12}} \quad (51)$$

$$T = (2\pi)\frac{P^{9/2}}{v^6}, \quad \text{unit} = u^{16 \cdot 9/2 - 17 \cdot 6 = -30} \quad (52)$$

$$L = \frac{TV}{2} = (2\pi^2\Omega^2)\frac{P^{9/2}}{v^5}, \quad \text{unit} = u^{16 \cdot 9/2 - 17 \cdot 5 = -13} \quad (53)$$

$$A = \frac{8V^3}{\alpha P^3} = \left(\frac{2^6\pi^3\Omega^3}{\alpha}\right)\frac{v^3}{P^3}, \quad \text{unit} = u^{17 \cdot 3 - 16 \cdot 3 = 3} \quad (54)$$

For convenience here I assign $r = \sqrt{p}$, unit $u^{16/2=8}$;

$$G^* = \frac{V^2 L}{M} = 2^3 \pi^4 \Omega^6 \frac{r^5}{v^2}, \quad u^{34 - 13 - 15 = 8 \cdot 5 - 17 \cdot 2 = 6} \quad (55)$$

$$h^* = 2\pi MVL = 2^3 \pi^4 \Omega^4 \frac{r^{13}}{v^5}, \quad u^{15 + 17 - 13 = 8 \cdot 13 - 17 \cdot 5 = 19} \quad (56)$$

$$T_p^* = \frac{AV}{\pi} = \frac{2^7 \pi^3 \Omega^5 v^4}{\alpha r^6}, \quad u^{3 + 17 = 17 \cdot 4 - 6 \cdot 8 = 20} \quad (57)$$

$$e = AT = \frac{2^7 \pi^4 \Omega^3 r^3}{\alpha v^3}, \quad u^{3 - 30 = 3 \cdot 8 - 17 \cdot 3 = -27} \quad (58)$$

$$k_B^* = \frac{\pi VM}{A} = \frac{\alpha r^{10}}{2^5 \pi \Omega v^3}, \quad u^{17 + 15 - 3 = 10 \cdot 8 - 17 \cdot 3 = 29} \quad (59)$$

$$\mu_0^* = \frac{\pi V^2 M}{\alpha LA^2} = \frac{\alpha}{2^{11} \pi^5 \Omega^4} r^7, \quad u^{17 \cdot 2 + 15 + 13 - 6 = 7 \cdot 8 = 56} \quad (60)$$

$$\epsilon_0^{*-1} = \frac{\alpha}{2^9 \pi^3} v^2 r^7, \quad u^{34 + 56 = 90} \quad (61)$$

$$r_\sigma^* = \left(\frac{8\pi^5 k_B^4}{15h^3 c^3}\right) = \frac{\alpha}{2^{29} 15\pi^{14} \Omega^{22}} r, \quad u^{29 \cdot 4 - 19 \cdot 3 - 17 \cdot 3 = 8} \quad (62)$$

$$R^* = \left(\frac{m_e}{4\pi l_p \alpha^2 m_p}\right) = \frac{1}{2^{23} 3^3 \pi^{11} \alpha^5 \Omega^{17}} \frac{v^5}{r^9}, \quad u^{13} \quad (63)$$

Scalars r, v were chosen as they can be determined directly from c, μ_0 (eq 49, 60);

$$v = \frac{c}{2\pi\Omega^2} \quad (64)$$

$$r^7 = \frac{2^6 \pi^6 \Omega^4}{5^7 \alpha} \quad (65)$$

In those (electron) ratios the scalar numerical values and units cancel. Using SI Planck units $M = m_p, T = t_p, L = l_p$ (4.1);

$$\frac{L^{15}}{M^9 T^{11}} = \frac{l_p^{15}}{m_p^9 t_p^{11}} = \frac{(2\pi^2\Omega^2)^{15}}{(1)^9 (2\pi)^{11}} \cdot \frac{l_p^{15}}{k^9 t_p^{11}} = 2^4 \pi^{19} \Omega^{30} \quad (66)$$

$$\frac{l_p^{15}}{k^9 t_p^{11}} = \frac{(.20322087^{-36})^{15}}{(.217672818^{-7})^9 (.171585513^{-43})^{11}} \frac{u^{-13 \cdot 15}}{u^{15 \cdot 9} u^{-30 \cdot 11}} = 1 \quad (67)$$

$$\frac{A^3 L^3}{T} = \frac{A_Q^3 l_p^3}{t_p} = \frac{(2^6 \pi^3 \Omega^3)^3 (2\pi^2 \Omega^2)^3}{(\alpha)^3 (2\pi)} \cdot \frac{a^3 l_p^3}{t} = \frac{2^{20} \pi^{14} \Omega^{15}}{\alpha^3} \quad (68)$$

$$\frac{a^3 l_p^3}{t} = \frac{(.12691859^{23})^3 (.20322087^{-36})^3}{(.171585513^{-43})} \frac{u^{3 \cdot 3} u^{-13 \cdot 3}}{u^{-30}} = 1 \quad (69)$$

4.3. The electron function f_e is both unit-less and non scalable $v^0 r^0 u^0 = 1$. It is therefore a natural (mathematical) constant.

$$\sigma_e = \frac{3\alpha^2 AL}{\pi^2} = 2^7 3\pi^3 \alpha \Omega^5 \frac{r^3}{v^2}, \quad u^{-10} \quad (70)$$

$$f_e = \frac{\sigma_e^3}{T} = 4\pi^2(2^6 3\pi^2 \alpha \Omega^5)^3, \text{ units} = \frac{u^{30}}{(u^{10})^3} = 1 \quad (71)$$

$$\sigma_{tp} = \frac{3\alpha^2 T_P}{2\pi} = 2^6 3\pi^2 \alpha \Omega^5 \frac{v^4}{r^6}, \text{ units} = u^{20} \quad (72)$$

$$f_e = t_p^2 \sigma_{tp}^3 = 4\pi^2(2^6 3\pi^2 \alpha \Omega^5)^3, \text{ units} = \frac{(u^{20})^3}{(u^{30})^2} = 1 \quad (73)$$

4.4. The Sommerfeld fine structure constant alpha is a dimensionless mathematical constant. The following use a well known formula for alpha;

$$\alpha = \frac{2h}{\mu_0 e^2 c} = 2.2\pi Q^2 2\pi l_p \cdot \frac{32l_p c^5}{\pi^2 \alpha Q^8} \cdot \frac{\alpha^2 Q^6}{256l_p^2 c^4} \cdot \frac{1}{c} = \alpha \quad (74)$$

$$\alpha = 2(8\pi^4 \Omega^4) / \left(\frac{\alpha}{2^{11} \pi^5 \Omega^4} \right) \left(\frac{128\pi^4 \Omega^3}{\alpha} \right)^2 (2\pi \Omega^2) = \alpha \quad (75)$$

$$\text{units} = \frac{u^{19}}{u^{56}(u^{-27})^2 u^{17}} = 1 \quad (76)$$

4.5. I have also premised a 2nd mathematical constant which I have denoted Omega. We can find a numerical solution using the precise c^*, μ_0^*, R^* ;

$$\Omega = 2.0071349496\dots;$$

$$\frac{(c^*)^{35}}{(\mu_0^*)^9 (R^*)^7}, \text{ units} = \frac{(u^{17})^{35}}{(u^{56})^9 (u^{13})^7} = 1 \quad (77)$$

$$\frac{(c^*)^{35}}{(\mu_0^*)^9 (R^*)^7} = (2\pi \Omega^2)^{35} / \left(\frac{\alpha}{2^{11} \pi^5 \Omega^4} \right)^9 \cdot \left(\frac{1}{2^{23} 3^3 \pi^{11} \alpha^5 \Omega^{17}} \right)^7 \quad (78)$$

$$\Omega^{225} = \frac{(c^*)^{35}}{2^{295} 3^{21} \pi^{157} (\mu_0^*)^9 (R^*)^7 \alpha^{26}}, \text{ units} = 1 \quad (79)$$

There is a close sqrt natural number solution for Ω ;

$$\Omega = \sqrt{\left(\frac{\pi^e}{e^{(e-1)}} \right)} = 2.0071 \ 349 \ 5432\dots \quad (80)$$

4.6. $\Omega^{(15)^n}$ apparently has a 'buffer' role. As constants of the series $u^{15}, u^{30}, \dots, u^{90}$ have no Ω , this suggests a base structure. For example;

$$\beta = \frac{\Omega}{a^{1/3}} = \frac{\Omega v}{r^2} = \frac{\Omega}{t^{2/15} k^{1/5}} \dots, \text{ unit} = u \quad (81)$$

Using M and β (see 4.2);

$$M = \frac{r^4}{v}, u^{15} \text{ (kg)} \quad (82)$$

$$P = M\beta = \Omega r^2, u^{16} \text{ (q)} \quad (83)$$

$$V = 2\pi M\beta^2 = 2\pi \Omega^2 v, u^{17} \text{ (m/s)} \quad (84)$$

$$T^2 = (2\pi \Omega)^{15} \frac{P^9}{2\pi V^{12}} = (2\pi \frac{r^9}{v^6})^2, u^{-60} \text{ (s}^2\text{)} \quad (85)$$

Using only β we find this unit-less ratio r^{17}/v^8 that maintains the numerical values of the dimensionful constants within a boundary, in SI units $r^{17}/v^8 = 0.813 \times 10^{-59}$;

$$A = \beta^3 \left(\frac{2^6 \pi^3}{\alpha} \right) = \frac{2^6 \pi^3 \Omega^3 v^3}{\alpha r^6}, u^3 \text{ (a)} \quad (86)$$

$$R = \beta^8 \left(\frac{r^{17}}{v^8} \right) = \Omega^8 r, u^8 \quad (87)$$

$$L^{-1} = \beta^{13} \frac{1}{(2\pi^2 \Omega^{15})} \left(\frac{r^{17}}{v^8} \right) = \frac{1}{2\pi^2 \Omega^2} \frac{v^5}{r^9}, u^{13} \text{ (1/m)} \quad (88)$$

$$M = \beta^{15} \left(\frac{1}{\Omega^{15}} \right) \left(\frac{r^{17}}{v^8} \right)^2 = \frac{r^4}{v}, u^{15} \text{ (kg)} \quad (89)$$

$$P = \beta^{16} \left(\frac{1}{\Omega^{15}} \right) \left(\frac{r^{17}}{v^8} \right)^2 = \Omega r^2, u^{16} \text{ (q)} \quad (90)$$

$$V = \beta^{17} \left(\frac{2\pi}{\Omega^{15}} \right) \left(\frac{r^{17}}{v^8} \right)^2 = 2\pi \Omega^2 v, u^{17} \text{ (m/s)} \quad (91)$$

$$T^{-1} = \beta^{30} \left(\frac{1}{2\pi \Omega^{30}} \right) \left(\frac{r^{17}}{v^8} \right)^3 = \frac{1}{2\pi} \frac{v^6}{r^9}, u^{30} \text{ (1/s)} \quad (92)$$

$$\mu_0^* = \beta^{56} \left(\frac{\alpha}{2^{11} \pi^5 \Omega^{60}} \right) \left(\frac{r^{17}}{v^8} \right)^7 = \frac{\alpha}{2^{11} \pi^5 \Omega^4} r^7, u^{56} \quad (93)$$

$$\epsilon_0^{*-1} = \beta^{90} \left(\frac{\alpha}{2^9 \pi^3 \Omega^{90}} \right) \left(\frac{r^{17}}{v^8} \right)^{11} = \frac{\alpha}{2^9 \pi^3} v^2 r^7, u^{90} \quad (94)$$

4.7. Unit = 1 combinations (examples)

$$\left(\frac{r^{17}}{v^8} = \frac{u^{8*17}}{u^{17*8}} \right) = (k^2 t = \frac{u^{15*2}}{u^{30}}), \text{ units} = 1 \quad (95)$$

$$M^2 = \frac{r^8}{v^2}, \text{ unit} = u^{8*8-17*2=30} \text{ (kg}^2\text{)} \quad (96)$$

$$T^{-1} = \frac{v^6}{2\pi r^9}, \text{ unit} = u^{17*6-8*9=30} \text{ (1/s)} \quad (97)$$

$$M^2 T = \frac{r^8}{v^2} \frac{2\pi r^9}{v^6} = 2\pi \left(\frac{r^{17}}{v^8} \right), \text{ unit} = 1 \quad (98)$$

4.8. We can numerically solve the physical constants by replacing the mathematical (c^*, μ_0^*, R^*) with the CODATA mean values for (c, μ_0, R) as in section 3.8.

$$h^* = 2^3 \pi^4 \Omega^4 \frac{r^{13}}{v^5}, u^{19} \quad (99)$$

We then find there is a combination of (c^*, μ_0^*, R^*) which reduces to h^3 .

$$(h^*)^3 = \frac{2\pi^{10} (\mu_0^*)^3}{3^6 (c^*)^5 \alpha^{13} (R^*)^2}, \text{ unit} = u^{57} \quad (100)$$

Likewise with the other dimensionful constants, we note that these equations are equivalent to section 3.8;

$$(e^*)^3 = \frac{4\pi^5}{3^3 (c^*)^4 \alpha^8 (R^*)}, \text{ unit} = u^{-81} \quad (101)$$

$$(k_B^*)^3 = \frac{\pi^5(\mu_0^*)^3}{3^3 2(c^*)^4 \alpha^5 (R^*)^2}, \text{ unit} = u^{87} \quad (102)$$

$$(G^*)^5 = \frac{\pi^3(\mu_0^*)}{2^{20} 3^6 \alpha^{11} (R^*)^2}, \text{ unit} = u^{30} \quad (103)$$

$$(m_e^*)^3 = \frac{16\pi^{10}(R^*)(\mu_0^*)^3}{3^6 (c^*)^8 \alpha^7}, \text{ unit} = u^{45} \quad (104)$$

$$(r_d)^3 = \frac{3^3 4\pi^5 (\mu_0^*)^3 \alpha^{19} (R^*)^2}{5^3 (c^*)^{10}}, \text{ unit} = u^{24} \quad (105)$$

5 Virtual universe

The electron formula f_e can be constructed from ampere-meters AL and time T and yet it is a dimensionless (mathematical) constant;

$$f_e = (AL)^3/T = 0.239 \times 10^{23}, \text{ units} = 1$$

This formula has 3 space dimensions L^3 and 1 time dimension T . We could then speculate that if the vacuum of our 3-D space is electro-magnetic in nature such that it is also a construct of ampere-meters $(AL)^3$ instead of an empty void measured in meters, then the sum universe may also be a dimensionless (mathematical) constant (aka a virtual universe);

$$f_{universe} = X(AL)^3/T, \text{ units} = 1$$

6 Notes

In 1963, Dirac noted regarding the fundamental constants; "The physics of the future, of course, cannot have the three quantities \hbar, e, c all as fundamental quantities. Only two of them can be fundamental, and the third must be derived from those two." [25]

In the article "Surprises in numerical expressions of physical constants", Amir et al write ... In science, as in life, 'surprises' can be adequately appreciated only in the presence of a null model, what we expect a priori. In physics, theories sometimes express the values of dimensionless physical constants as combinations of mathematical constants like pi or e. The inverse problem also arises, whereby the measured value of a physical constant admits a 'surprisingly' simple approximation in terms of well-known mathematical constants. Can we estimate the probability for this to be a mere coincidence? [24]

J. Barrow and J. Webb on the fundamental constants; 'Some things never change. Physicists call them the *constants of nature*. Such quantities as the velocity of light, c , Newton's constant of gravitation, G , and the mass of the electron, m_e , are assumed to be the same at all places and times in the universe. They form the scaffolding around which theories of physics are erected, and they define the fabric of our universe. Physics has progressed by making ever more accurate measurements of their values. And yet, remarkably, no

one has ever successfully predicted or explained any of the constants. Physicists have no idea why they take the special numerical values that they do. In SI units, c is 299,792,458; G is 6.673e-11; and m_e is 9.10938188e-31 -numbers that follow no discernible pattern. The only thread running through the values is that if many of them were even slightly different, complex atomic structures such as living beings would not be possible. The desire to explain the constants has been one of the driving forces behind efforts to develop a complete unified description of nature, or "theory of everything". Physicists have hoped that such a theory would show that each of the constants of nature could have only one logically possible value. It would reveal an underlying order to the seeming arbitrariness of nature.' [6].

At present, there is no candidate theory of everything that is able to calculate the mass of the electron [23].

"There are two kinds of fundamental constants of Nature: dimensionless (alpha) and dimensionful (c, h, G). To clarify the discussion I suggest to refer to the former as fundamental parameters and the latter as fundamental (or basic) units. It is necessary and sufficient to have three basic units in order to reproduce in an experimentally meaningful way the dimensions of all physical quantities. Theoretical equations describing the physical world deal with dimensionless quantities and their solutions depend on dimensionless fundamental parameters. But experiments, from which these theories are extracted and by which they could be tested, involve measurements, i.e. comparisons with standard dimensionful scales. Without standard dimensionful units and hence without certain conventions physics is unthinkable" -*Dialogue* [5].

"The fundamental constants divide into two categories, units independent and units dependent, because only the constants in the former category have values that are not determined by the human convention of units and so are true fundamental constants in the sense that they are inherent properties of our universe. In comparison, constants in the latter category are not fundamental constants in the sense that their particular values are determined by the human convention of units" -*L. and J. Hsu* [4].

A charged rotating black hole is a black hole that possesses angular momentum and charge. In particular, it rotates about one of its axes of symmetry. In physics, there is a speculative notion that if there were a black hole with the same mass and charge as an electron, it would share many of the properties of the electron including the magnetic moment and Compton wavelength. This idea is substantiated within a series of papers published by Albert Einstein between 1927 and 1949. In them, he showed that if elementary particles were treated as singularities in spacetime, it was unnecessary to postulate geodesic motion as part of general relativity [26].

The Dirac Kerr–Newman black-hole electron was introduced by Burinskii using geometrical arguments. The Dirac wave function plays the role of an order parameter that signals a broken symmetry and the electron acquires an extended space-time structure. Although speculative, this idea was corroborated by a detailed analysis and calculation [8].

Max Tegmark’s Mathematical Universe Hypothesis: Our external physical reality is a mathematical structure. That is, the physical universe is mathematics in a well-defined sense, and “in those [worlds] complex enough to contain self-aware substructures [they] will subjectively perceive themselves as existing in a physically ‘real’ world” [10].

Note: This article is an extension of an earlier article [26] which has been incorporated as section 3.

References

1. Linnebo, Øystein, “Platonism in the Philosophy of Mathematics”, *The Stanford Encyclopedia of Philosophy* (Summer 2017 Edition), Edward N. Zalta (ed.), plato.stanford.edu/archives/sum2017/entries/platonism-mathematics
2. Nick Bostrom, *Philosophical Quarterly* 53 (211):243–255 (2003)
3. <https://en.wikipedia.org/wiki/Philosophy-of-mathematics> (22, Oct 2017)
4. Leonardo Hsu, Jong-Ping Hsu; *The physical basis of natural units*; *Eur. Phys. J. Plus* (2012) 127:11
5. Michael J. Duff et al *JHEP03(2002)023* Dialogue on the number of fundamental constants
6. J. Barrow, J. Webb, *Inconstant Constants*, *Scientific American* 292, 56 - 63 (2005)
7. M. Planck, *Über irreversible Strahlungsvorgänge*. *Ann. d. Phys.* (4), (1900) 1, S. 69-122.
8. A. Burinskii, *Gravitation and Cosmology*, 2008, Vol. 14, No. 2, pp. 109–122; Pleiades Publishing, 2008. DOI: 10.1134/S0202289308020011
9. Feynman, Richard P. (1977). *The Feynman Lectures on Physics*, vol. I. Addison-Wesley. p. 22-10. ISBN 0-201-02010-6.
10. Tegmark, Max (February 2008). “The Mathematical Universe”. *Foundations of Physics*. 38 (2): 101–150.
11. planckmomentum.com/calc/ (online calculator)
12. Magnetic monopole
en.wikipedia.org/wiki/Magnetic-monopole (10/2015)
13. Fine structure constant
en.wikipedia.org/wiki/Fine-structure-constant (06/2015)
14. Burinskii, A. (2005). “The Dirac–Kerr electron”. arXiv:hep-th/0507109
15. Rydberg constant
<http://physics.nist.gov/cgi-bin/cuu/Value?ryd>
16. Planck constant
<http://physics.nist.gov/cgi-bin/cuu/Value?ha>
17. Electron mass
<http://physics.nist.gov/cgi-bin/cuu/Value?me>
18. Fine structure constant
<http://physics.nist.gov/cgi-bin/cuu/Value?alphinv>
19. Elementary charge
<http://physics.nist.gov/cgi-bin/cuu/Value?e>
20. Vacuum of permeability
<http://physics.nist.gov/cgi-bin/cuu/Value?mu0>
21. Gravitation constant
<http://physics.nist.gov/cgi-bin/cuu/Value?bg>
22. Boltzmann constant
<http://physics.nist.gov/cgi-bin/cuu/Value?k>
23. <https://en.wikipedia.org/wiki/Theory-of-everything> (02/2016)
24. Ariel Amir, Mikhail Lemesko, Tadashi Tokieda; 26/02/2016
Surprises in numerical expressions of physical constants
arXiv:1603.00299 [physics.pop-ph]
25. Dirac, Paul, *The Evolution of the Physicist’s Picture of Nature*, June 25, 2010
<https://blogs.scientificamerican.com/guest-blog/the-evolution-of-the-physicists-picture-of-nature/>
26. Macleod, Malcolm, *A virtual black-hole electron and the sqrt of Planck momentum*,
<http://vixra.org/pdf/1102.0032v9.pdf>