

# Linkage of Classical (3 Dimensional) and QM geometry( 2 Dimensional) via Hopf mapping and its implications for relic GW power production

A. Beckwith<sup>1</sup>

Chongqing University *Department of Physics: Chongqing University,*  
Chongqing, PRC , 400044

E-mail: abeckwith@uh.edu

**Abstract.** Hopf mapping from 2 dimensions (QM) to 3 dimensions (CM) are examined in terms of a formalism started by Feynman which has linkage to the (CM) equations of motion have linkage to the Serret - Frenet form (for Differential equations). We argue that in doing so we may then link QM representations of qubits to a solved version of the rotating rod problem. Furthermore since a 'generic' solid body rotation equivalent to the rotating rod problem has linkage to GW generation, as given by Lightman et. al. it is a way to tie qubits (Quantum information) to GW generation. We then make observations as to what the results mean in terms of QM initial states and the power of GW production from early universe conditions.

## 1. Introduction

We examine the results of the Hopf mapping from 2 dimensions (QM) to 3 dimensions (CM) in terms of generalizations to a rigid rod rotation [1] which could generate GW (gravitational waves) . This paper was initiated by a question by Stephen Kauffman about black holes. I.e. the reality of black hole singularities. To this end, this formalism was chosen to be the basis of finding a solution to trying to investigate black hole physics. What we are addressing is a problem created by the following confusion in physics. Namely that decoherence theorists are firm in their assertion that the decoherence of the wavefunction (of the universe?) is automatic based on the fact that nature is measuring itself all the time. This paper addresses that problematic assumption by giving a counter poise to necessary conditions to decoherence of an initial wavefunction of the universe, by specific requirements for initial two level QM system analysis prior to generation of GW in the electroweak era of cosmology. Afterwards, we may be able to give a succinct answer to if qubits of information [1] are thereby created in the prior to GW generation era. *Note that this transferal from 2 dimensions to 3 dimensions is in reality the holographic principle and its aftermath, with what is very similar to a shock wave in the time of the event horizon* [2], After this formalism is further developed, we will use the two to the three sphere mapping to ask if singularities have a presence in astrophysical problems. To start developing mathematics relevant to that future development, we solve on our own a set of equations ( in 3 dimensions) pertinent to a non-symmetric object in rotation in early universe which is a way to generate GW and from there state some caveats as to the power of GW which may ensue. The final conclusion of our document is in linking a quantum qubit form with the power created by/ during GW generation which conceivably could be identified by a suitably designed detector. The document first examines what Feynman did with respect to 2 level QM systems, their

---

<sup>1</sup> To whom any correspondence should be addressed.

generalization to Classical rigid rod rotation, and then we solve the resulting CM equation of motion. Feynman decomposed the solutions in x, y, and z in terms of the 2 level QM system [3], [4] a decomposition which we hold as still relevant and valid, and then, using the case of a uniform magnetic field ‘down’ the z axis, as a driver to the physical process leading to non-symmetric rigid body rotation. That the body is non-symmetric allows us to approximate the GW power generated as to the conventions outlined by Lightman, Press, Price, and Teukolsky [5] . We then conclude with a description of what our model says about QM generation of states relevant to GWs. In the early universe

## 2. Outlining the Feynman development of a classical system from 2 level QM system.

We look at how Feynman [1], [2], [3] linked a 2 dim quantum system to a 3 dimensional rigid rod style classical mechanics system. In doing so, Feynman worked with a quantum system given as

$$i \cdot \frac{d}{dt} \begin{pmatrix} a \\ b \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} H_z & H_x - i \cdot H_y \\ H_x + i \cdot H_y & -H_z \end{pmatrix} \cdot \begin{pmatrix} a \\ b \end{pmatrix} \quad (1)$$

In doing so via the transformation

$$\begin{aligned} x &= a \cdot b^* + b \cdot a^* \\ y &= i \cdot (a \cdot b^* - b \cdot a^*) \\ z &= a \cdot a^* - b \cdot b^* \end{aligned} \quad (2)$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} 0 & H_z & -H_y \\ -H_z & 0 & H_x \\ H_y & -H_x & 0 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (3)$$

The simplest decomposition of this problem is to set  $H_y = H_x = 0$  so then the situation is that we have

$$\begin{pmatrix} a(t) \\ b(t) \end{pmatrix} = \begin{pmatrix} a(0) \cdot \exp(i \cdot t \cdot H_z / 2) \\ b(0) \cdot \exp(-i \cdot t \cdot H_z / 2) \end{pmatrix} \quad (4)$$

And

$$\begin{pmatrix} x(t) \\ y(t) \\ z(t) \end{pmatrix} = \begin{pmatrix} x(0) \cdot \cos(H_z \cdot t) + y(0) \cdot \sin(H_z \cdot t) \\ y(0) \cdot \cos(H_z \cdot t) - x(0) \cdot \sin(H_z \cdot t) \\ z(0) \end{pmatrix} \quad (5)$$

As can be seen by Maggorie [5] and also Lightman, Press, Price, and Teukolsky [4] since the solution as given by Eq.(5) is for a circular moment of a GW there would be a GW associated with it,

We also will be looking at a more complex three dimensional example of motion which is highly complex Non-withstanding we go to a more complete version of Eq.(1) to Eq.(3) with only  $H_y = 0$  . Then we get

$$i \cdot \frac{d}{dt} \begin{pmatrix} a \\ b \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} H_z & H_x \\ H_x & -H_z \end{pmatrix} \cdot \begin{pmatrix} a \\ b \end{pmatrix} \quad (6)$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} 0 & H_z & 0 \\ -H_z & 0 & H_x \\ 0 & -H_x & 0 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (7)$$

The above two equations have the setting of what is called the Serret – Frenet form and we will solve these two DE equation systems taking the approximation that  $H_z, H_x$  are constants in lieu of the first example. The next section solves these two equations with this in mind, leading to a non-symmetric rotation in 3 dimensional space which is needed for GW production.

### 3. Solving a simplified version of Eq. (6) and Eq. (7) to come up with a non- symmetric rigid body rotation sufficient to obtain GW.

To do this we look at Eq. (7) in such a way as to have

$$\begin{aligned} x(t) &= x_0 + x_1 \cdot \sin(H_z \cdot t) \\ y(t) &= \frac{2}{H_x} \cdot \frac{dz(t)}{dt} = -2x_1 \cdot \frac{H_z}{H_x} \cos(H_z \cdot t) \\ z(t) &= z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t) \\ \frac{dy(t)}{dt} &= \frac{H_z}{2} \cdot x(t) - \frac{H_x}{2} \cdot z(t) \end{aligned} \quad (8)$$

To which we add in the reconciliation of the variables equation result from the last part of Eq.(8), namely

$$2x_1 \frac{H_z}{H_x} \cdot \sin(H_z \cdot t) = \frac{1}{2} \cdot [x_0 + x_1 \cdot \sin(H_z \cdot t)] - \frac{1}{2} \cdot \left( \frac{H_x}{H_z} \right) \cdot \left( z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t) \right) \quad (9)$$

Leading to

$$x_0 = z_0 \cdot \left( \frac{H_x}{H_z} \right) + x_1 \cdot \left( 4 \cdot \frac{H_z}{H_x} - 1 - \frac{1}{2} \cdot \frac{H_x}{H_z} \right) \cdot \sin(H_z \cdot t) \quad (10)$$

Leading to

$$\begin{aligned} x(t) &= z_0 \cdot \frac{H_x}{H_z} + 4x_1 \cdot \left[ 4 \cdot \frac{H_z}{H_x} - \frac{1}{2} \cdot \frac{H_x}{H_z} \right] \sin(H_z \cdot t) \\ y(t) &= -2x_1 \cdot \frac{H_z}{H_x} \cos(H_z \cdot t) \\ z(t) &= z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t) \end{aligned} \quad (11)$$

Combined once again with Eq.(2), and assuming that the quantity we roughly identify with the “magnetic field  $H_z$ ” is parallel to the  $z$  axis, so long as  $z_0$  as an initial starting point for the  $z$  axis is non-zero, then we have fulfilled the requirement for a non-uniform motion of a ‘rigid body’ which if related to the quantities in Eq.(2) and also Maggiore’s criteria for GW from a non-uniform non spherical generation of GW leads to the final part of the GW requirement of non-spherically symmetric motion which lends itself to GW generation. We will then make a comment as to how to link this to GW power using Lightman, Press, Price, and Teukolsky [3] to show how frequency from this example can lead to GW generation.

#### 4. Conditions permitting GW power production using the inputs from Eq. (5)

The idea is that we need to calculate the following, i.e. a moment of inertia for a system, and also the frequency. As to Eq. (5) according to the following, we can come up with a generic Eqn. of motion, namely if we do averaging and set out a general time averaging. Fortunately for us the trig identities naturally vanish.

$$I = \alpha \cdot m \cdot r^2 = \alpha \cdot m \cdot \left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{Eq.(5)} = \alpha \cdot m \cdot (x^2(0) + y^2(0) + z^2(0)) \quad (12)$$

We can, as an approximation use  $m$  above to be the net mass of the assumed geometry and set.  $\alpha \approx \frac{1}{12}$  I.e.

Then we look at the power loss according to a 'rigid rod' construction for GW power generation[4] ,[5]i.e.

$$\frac{d\varepsilon}{dt} = -\frac{32}{5} G \cdot I^2 \omega^6 \quad (13)$$

Note that we can approximate the frequency in this case as directly proportional to the input frequency of the magnetic field parallel to the z axis, i.e. looking to first approximation at  $H_z \sim \omega$  according to the conventions as given by Kholodenko [1] on page 157. This means that up to a point, if one picks representative positions as given by  $x^2(0) + y^2(0) + z^2(0)$  with each of these initial positions, squared, and a net mass  $m$ . Then we can calculate the net GW (gravitational wave) power loss of this system. We will in the end make a comment as to this Eq.(13) value, for the specified inputs into the equation and the Feynman qubit (QM) results for while comparing them to what we can infer as to Eq. (4), and its up and down 2 dimensional QM states. I.e. this problem is comparatively easy to calculate. In this case the value of Eq.(13) if we are near the cosmological origin would have a value of about

$$\left\langle \dot{\varepsilon} \sim 10^{45} - 10^{50} \text{ Joule / sec} \right\rangle_{relic-condit} \Rightarrow \left\langle \dot{\varepsilon} / A \sim 10^2 \text{ Joule / sec} \right\rangle_{Earth} \quad (14)$$

Next we will look at what happens if we assume the input geometry as given by Eq. (11).

Both of these results will be then compared to as to the simple case of Eq.(4) as due to the first set of inputs into Eq. (13) if the spatial geometry of Eq.(5) is used, and then Eq. (4) will be guessed at if we use the geometry of Eq. (11). I.e. we will guess what Eq. (11) does to Eq. (4) and compare that with what Eq. (11) does to Eq. (4)

#### 5. Conditions permitting GW power production using the inputs from Eq. (11)

This is a mess. I.e. what we have to do is to look at how to calculate the moment of inertia, and then going to Eq. (13), even if we assume the same mass which was used earlier to calculate Eq. (14) above for relic conditions.

To start this, look at, even if  $\alpha \approx \frac{1}{12}$

$$I = \alpha \cdot m \cdot r^2 = \alpha \cdot m \cdot \left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{Eq.(8)} \quad (15)$$

The problem starts immediately, in that the parenthesis of Eq. (15) above would have to be a time averaged quantity. I.e. we would be looking at  $x^2(t) + y^2(t) + z^2(t)$  **with left over terms in this set analytical expression, should they exist to be time averaged, i.e. if**

$z_0 = 0$ ,

$$x^2(t) + y^2(t) + z^2(t) = x_1^2 \cdot \left( \frac{H_z}{H_x} \right)^2 \cdot \left[ 4 + \left( 252 + 4 \cdot \left( \frac{H_x}{H_z} \right)^4 - \frac{655}{4} \cdot \left( \frac{H_x}{H_z} \right)^2 \right) \cdot \sin^2(H_z \cdot t) \right]_{Eq(11)} \quad (16)$$

The term to be time averaged would be  $\left\langle \sin^2(H_z \cdot t) \right\rangle_{Time-averaged} \sim 1/2$ . So the above would be approximated by

$$\left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{Time-averaged} = x_1^2 \cdot \left( \frac{H_z}{H_x} \right)^2 \cdot \left[ 4 + \left( \frac{252}{2} + 2 \cdot \left( \frac{H_x}{H_z} \right)^4 - \frac{655}{8} \cdot \left( \frac{H_x}{H_z} \right)^2 \right) \right]_{Eq(11)} \quad (17)$$

Using a ratio, as given of  $\left( \frac{H_x}{H_z} \right)^2 \sim 1/2$ , the above then becomes approximately

$$\left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{Time-averaged} = x_1^2 \cdot \left[ \frac{1441}{16} \right] \sim 91 \cdot x_1^2 \quad (18)$$

Then the magnitude of the GW power would be, per second about 10,000 times bigger.

$$\left\langle \dot{\epsilon} \sim 10^{49} - 10^{53} \text{ Joule / sec} \right\rangle_{\substack{Eq.(11) \\ Time-Averaged}}^{relic-condit} \Rightarrow \left\langle \dot{\epsilon} / A \sim 10^6 \text{ Joule / sec} \right\rangle_{\substack{Eq.(11) \\ Time-Averaged}}^{Earth} \quad (19)$$

## 6. Comparison of Eq. (14) and Eq. (19) results in terms upon solving Eq.(1)

The value for the simple geometry (in terms of simple qudits ) to understand working with both Eq. (1) and then Eq.(4) has , if a particle is in a constant magnetic field, then according to [4] it is a special case of working

with qubits, according to [1], [5] , the values of  $\begin{pmatrix} a \\ b \end{pmatrix}$  if only  $H_z$  is non- zero, for the below equation

become very simple. The problem of solving for the functions of an applied non zero  $H_z$  in [1][2] , [3]

$$\varphi = a \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + b \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} \doteq a \cdot |1\rangle + b \cdot |0\rangle \quad (20)$$

is much simpler than when  $H_z$  and  $H_x$  are both non zero. Is in the case of Eq. (4) with only  $H_z$  not equal to zero, then looking at the terms for **a** and **b** in the case of Eq. (4) is extremely simple, for the situation for Eq.(14) as diagrammed out above. It is the same problem for Eq. (19) and the much larger GW power situation,

but due to the ‘noisy’ values for **a** and **b**, **then**  $\begin{pmatrix} a \\ b \end{pmatrix}$  is the same as looking at highly non-linear inputs into

$\begin{pmatrix} a \\ b \end{pmatrix}$  QM values which are still then mapped into the 3 dimensional CM results. . Still the same rotating rigid

body problem approximated by a rod in spatial rotation, but the movements and more would become much more

complicated. And then we find that  $\begin{pmatrix} a \\ b \end{pmatrix}$  the input values are MUCH harder to solve.

## 7. Conclusion. CM and QM correspondence remains, but turbulence, a.k.a. Duerrer and Beckwith results for early universe GW generation makes the QM connection very hard to mathematically identify. Simple logical process, MESSY algebra ahead. With a possible answer to the question if singularities are essential in Cosmology.

Looking at Eq. (13) , simpler and harder cases, still in the case of relic GW production has large number correspondence and scaling as mentioned by Valev [7], with his H, not a Hamiltonian, but instead

$$r(\text{radius} - \text{of} - \text{universe}) \sim cH^{-1} \quad (21)$$

Also, the mass of the Universe, as given by Valev [7] is

$$M = (\text{Mass-of-universe}) \sim c^3 \cdot 2^{-1} \cdot (G \cdot H)^{-1} \quad (22)$$

There is here, a template for innate simplicity which can be based, also on keeping in mind the holographic principle, i.e. the QM is embedded and done first with a transferal to CM with the event horizon being the boundary for a phase transition between regimes of space-time, with some of the processes as outlined in [2]. When this is brought up, we should keep in mind that the qubits generated in BHs have their counterpart in early universe cosmology as stated in [8] which shows that event horizon and entropy generation of black holes has a similar analogy to early universe cosmology before the Electro-Weak era. Also there is, when we look at physics innate simplicity in the inter relationship, of the sort mentioned by Valev[7], in terms of space-time geometry. The inter relationship of CM and QM given by Eq. (1) and then Eq. (3) with the stunning interplay between  $\mathbf{x}, \mathbf{y}, \mathbf{z}$  and  $\mathbf{a}, \mathbf{b}$  given by Eq.(2) is, we believe, obscured by how complex the problem is of finding

$\begin{pmatrix} a \\ b \end{pmatrix}$ . However, there is an interplay between the qubits given in Eq.(20) and the complex systems given in

Eq.(20) and the inputs into Eq. (2). This interrelationship depends upon the complex systems as given by Eq. (11), for Classical mechanics or Eq.(5) as seen in references [9], and [10]. Finally the author suggests that the linkage of three sphere topology (CM) to two sphere topology(QM) is a very useful starting point to answer an objection by Dr. Kauffmann to the existence of singularities in General relativity. As well as the classical GR treatment of black holes. As related by Dr. Kauffmann, quantum mechanics routinely prunes out non-physical solutions with regards to the hydrogen atom and only uses solutions which are physically feasible. Dr. Kauffmann asserts no such pruning of solutions occurs in black hole physics or in the formation of cosmological singularities. The author in [10] used chaotic space-time conditions for generating GW in the electro-weak era. Furthermore, the author asserts that turbulence as exemplified by [9] is the driver of relic GW generation and GW power calculations. If so, further development of the Hopf mapping results from 2 dimensions (QM) to 3 dimensions (CM) may enable us to identify necessary conditions

for finding  $\begin{pmatrix} a \\ b \end{pmatrix}$  in initial QM states necessary for relic GW development. It is well

known that spherically symmetric geometry will generally not generate GW. This puts

a restriction upon finding  $\begin{pmatrix} a \\ b \end{pmatrix}$  which in turn would answer if an initial space-time singularity is even feasible. Note also that necessary conditions of how to

construct and finding inputs into  $\begin{pmatrix} a \\ b \end{pmatrix}$  will allow us to avoid a serious logical error by many physicists.

This error is due to confusion about how to form an initial wave-function of the universe. Namely as stated initially by the author, that decoherence theorists are firm in their assertion that the decoherence of the wave-function (of the universe?) is automatic, based on the fact that nature is measuring itself all of the time. This paper addresses that problematic assumption by giving a counter poise for necessary conditions to the decoherence of an initial wave-function of the universe, by specific requirements for forming an initial two level QM system for analysis prior to the generation of GW in the electro-weak era of cosmology. Also in finding

sufficient conditions for an adequate input into the initial wave-function called  $\begin{pmatrix} a \\ b \end{pmatrix}$ . Also we should note that

while String theory and loop quantum gravity via different mechanisms [11], [12] purport to have solutions to the initial

cosmological state, that the author views development of mandatory restrictions upon acceptable  $\begin{pmatrix} a \\ b \end{pmatrix}$  QM

quantum initial states for GW development as essential for finding and investigating rigorously the question if singularities are indeed necessary and even allowed in astrophysical problems. Qubit

analysis of the two level QM state created by  $\begin{pmatrix} a \\ b \end{pmatrix}$  [1] is essential for yet another procedure. In Classical and

Quantum Gravity (IOP) Borsten. Duff and Levay [13] give a plan of action statement as to how to tie in qubits into entropy of Black holes. But in doing so there is a serious confusion relating to the fact that specific

information about black holes, and also the universe is purely abstract unless a conscious observer or their measurement apparatus is there to observe and cognize it. If this is true for qubits relating to black holes, it is also even more true for the assumed singularity at the start of the initial configuration of the universe. As given by Muller, and Lousto [8] there is a way to give very similar treatment of entanglement entropy for both black holes and the initial universe. Then, the way qubits are measured and used will allow us to ask what entropy and information is stored in what is commonly purported to be a singularity. If it is one ( a singularity) , which will be the final question to answer. The author, also has an accepted publication in the Journal of Applied mathematics, SCIRP in part on this very question [14] with the answer that this question is not open and shut. I.e. that further work needs to be done in this area. This present Hopf mapping from QM to CM with , as argued here, CM turbulence generating GW, may be a way to determine restrictions upon initial states in QM which the author views, as a way to answer Dr. Kauffmann's question in a rigorous role in the near future . Qubit analysis, if stated appropriately along a simplified version of [12] while keeping in mind [13] and the cautionary example in [14] will either confirm or disavows the existences of cosmological singularities. Namely we need to know and confirm if qubit presentation of black hole and early universe information leading to entropy (which may lead to GW generation [10]), is commensurate with either singularities in space-time or their explicit disproof of existence, if that is possible .

## References

1. A. L. Kholodenko, *Applications of Contact Geometry and Topology in Physics*, World Scientific, 2013
2. F. Darabi, S. Jalalzadeh, "Cosmological quantum tunneling and the Holographic principle", **Theoretical and mathematical physics**, May 2013, Volume 175, Issue 2 PP 710-716; also see much the same thing in <http://arxiv.org/abs/1108.0682>
3. R. Feynman, F. Verno, and R. Hellwarth, "Geometrical representation of the Schrodinger Equation for solving Master Equation", **J. Appl. Phys** **28**. 49-52, (1957)
4. R. Feynman, *The Feynman Lectures on Physics, Volume 3*( Adsdison-Wesley, Reading , 1966)
5. A. Lightman, W. Press, R. Price, and S. Teukolsky, *Problem book in Relativity and Gravitation*, Princeton University Press, Princeton, New Jersey, 1979
6. M. Maggiore, *Gravitational Waves, Volume 1 : Theory and Experiment*, Oxford Univ. Press(2008)
7. D. Valev, "Estimation of the total mass and energy of the Universe", <http://arxiv.org/pdf/1004.1035.pdf>
8. R. Muller, and C. Lousto, "Entanglement entropy in curved spacetimes with event horizons" arXIV Gr-qc/9504049v1
9. R. Durrer, <http://arxiv.org/pdf/1303.7121.pdf>; R. Durrer, <http://arxiv.org/abs/0901.0650> , Phys.Rev.D79:063507, 2009,
10. A. Beckwith, , <http://vixra.org/abs/1308.0110>; ICISE inaugural conference accepted conference proceedings, in Quy Nohn, Vietnam
11. T. Padmanabhan , *Gravitation, Foundations and frontiers*, Cambridge University Press, Cambridge, UK, 2010
12. G. Ellis, R. Maartens, M. Mac Callum, *Relativistic Cosmology* Cambridge University Press, Cambridge, UK, 2012
13. G L. Borsten, M. Duff and P. Levay, ' The Black hole/qubit correspondence; an up-to-date review", *Class.Quantum.Grav.***29**(2012) 224008 (80 pp)
14. A. Beckwith, <http://vixra.org/abs/1304.0156> (accepted for publication in Journal for Applied mathematics, SCIRP, September, 2013)