

The accelerated frame in the curved time-space in the general relativity theory

Sang wha-Yi

Department of Math , Taejon University 300-716

ABSTRACT

In the general relativity theory, defines the accelerated frame that moves in \hat{r} -axis in the curved time-space. And calculates the curvature tensor of the accelerated frame in the curved time-space.

PACS Number:04.04.90.+e

Key words:The general relativity theory,

The tetrad,

The curved time-space,

The accelerated frame

The curvature tensor

e-mail address:sangwha1@nate.com

Tel:051-624-3953

I.Introduction

This theory's object is that defines the accelerated frame that moves in \hat{r} -axis in the curved time-space.

The Schwarzschild solution is

$$d\tau^2 = \left(1 - \frac{2GM}{rc^2}\right)dt^2 - \frac{1}{c^2} \left[\frac{dr^2}{1 - \frac{2GM}{rc^2}} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right] \quad (1)$$

In this time, a moving matter's acceleration is a in the Schwarzschild time-space.

$$a = \frac{d}{dt} \left(\frac{u}{\sqrt{1 - \frac{2GM}{rc^2} - \frac{u^2}{c^2}}} \right), u = \frac{1}{\sqrt{1 - \frac{2GM}{rc^2}}} \frac{dr}{dt} \quad (2)$$

If $a_0 = a / \sqrt{1 - \frac{2GM}{rc^2}}$ is,

$$a_0 = \frac{1}{\sqrt{1 - \frac{2GM}{rc^2}}} \frac{d}{dt} \left(\frac{u}{\sqrt{1 - \frac{2GM}{rc^2} - \frac{u^2}{c^2}}} \right) = \frac{d}{d\hat{t}} \left(\frac{V}{\sqrt{1 - \frac{V^2}{c^2}}} \right),$$

$$V = \frac{d\hat{r}}{d\hat{t}} = \frac{dr}{dt} \frac{1}{\left(1 - \frac{2GM}{rc^2}\right)}, \quad d\hat{t} = dt \sqrt{1 - \frac{2GM}{rc^2}}, \quad d\hat{r} = \frac{dr}{\sqrt{1 - \frac{2GM}{rc^2}}}$$

$$a_0 \hat{t} = \frac{V}{\sqrt{1 - \frac{V^2}{c^2}}}, \quad V = \frac{a_0 \hat{t}}{\sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}}}, \quad V \text{ is the } \hat{r}\text{-axis's velocity} \quad (3)$$

If $\frac{d\theta}{dt} = \frac{d\phi}{dt} = 0$, the solution is

$$d\tau^2 = \left(1 - \frac{2GM}{rc^2}\right)dt^2 - \frac{1}{c^2} \frac{dr^2}{1 - \frac{2GM}{rc^2}} = d\hat{t}^2 - \frac{1}{c^2} d\hat{r}^2 = d\hat{t}^2 \left(1 - \frac{V^2}{c^2}\right)$$

$$= \frac{d\hat{t}^2}{\sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}}} \quad (4)$$

In this time,

$$\begin{aligned}
\tau &= \int d\tau = \int \frac{d\hat{t}}{\sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}}} = \frac{c}{a_0} \sinh^{-1}\left(\frac{a_0}{c} \hat{t}\right), \\
\hat{t} &= \frac{c}{a_0} \sinh\left(\frac{a_0 \tau}{c}\right), \quad \hat{r} = \int V d\hat{t} = \int \frac{a_0 \hat{t} d\hat{t}}{\sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}}} = \frac{c^2}{a_0} \sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}} \\
&\quad = \frac{c^2}{a_0} \cosh\left(\frac{a_0 \tau}{c}\right) \\
\frac{d\hat{t}}{d\tau} &= \cosh\left(\frac{a_0}{c} \tau\right), \quad \frac{1}{c} \frac{d\hat{r}}{d\tau} = \sinh\left(\frac{a_0}{c} \tau\right)
\end{aligned} \tag{5}$$

II. The tetrad in the curved time-space

The tetrad $\epsilon^{\hat{\alpha}}_{\hat{\mu}}$ is the unit vector defined by the following formula.

$$\eta_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\alpha}}_{\hat{\mu}} \epsilon^{\hat{\beta}}_{\hat{\nu}} = g_{\hat{\mu}\hat{\nu}} \tag{6}$$

In this time, if a matter moves in \hat{r} -axis in the curved time-space,

$$\eta_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\alpha}}_{\hat{\mu}}(\tau) \epsilon^{\hat{\beta}}_{\hat{\nu}}(\tau) = g_{\hat{\mu}\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}}, \quad g_{\hat{\mu}\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}} \tag{7}$$

Hence, Eq(6),Eq(7) is

$$\eta_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\alpha}}_{\hat{0}}(\tau) \epsilon^{\hat{\beta}}_{\hat{0}}(\tau) = \eta_{\hat{0}\hat{0}} = -1 \tag{8}$$

$$\begin{aligned}
d\tau^2 &= -\frac{1}{c^2} \eta_{\hat{\alpha}\hat{\beta}} d\hat{x}^\alpha d\hat{x}^\beta \\
\rightarrow -1 &= \eta_{\hat{\alpha}\hat{\beta}} \left(\frac{1}{c} \frac{d\hat{x}^\alpha}{d\tau}\right) \left(\frac{1}{c} \frac{d\hat{x}^\beta}{d\tau}\right) = \eta_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\alpha}}_{\hat{0}}(\tau) \epsilon^{\hat{\beta}}_{\hat{0}}(\tau) \\
\hat{x}^\alpha &= (c\hat{t}, \hat{r}, \hat{\theta}, \hat{\phi})
\end{aligned} \tag{9}$$

According to Eq(5),Eq(9)

$$\epsilon^{\hat{\alpha}}_{\hat{0}}(\tau) = \frac{1}{c} \frac{d\hat{x}^\alpha}{d\tau} = (\cosh\left(\frac{a_0 \tau}{c}\right), \sinh\left(\frac{a_0 \tau}{c}\right), 0, 0) \tag{10}$$

About $\hat{\theta}$ -axis's and $\hat{\phi}$ -axis's orientation

$$\eta_{\hat{\alpha}\hat{\beta}} \epsilon^{\hat{\alpha}}_{\hat{0}}(\tau) \epsilon^{\hat{\beta}}_{\hat{0}}(\tau) = \eta_{\hat{0}\hat{0}} = 1, \quad \epsilon^{\hat{\alpha}}_{\hat{0}}(\tau) = (0, 0, 1, 0)$$

$$\eta_{\hat{3}\hat{3}} e^{\hat{3}\hat{3}}(\tau) e^{\hat{3}\hat{3}}(\tau) = \eta_{\hat{3}\hat{3}} = 1, \quad e^{\hat{3}\hat{3}}(\tau) = (0,0,0,1) \quad (11)$$

And the other vector $e^{\hat{3}\hat{1}}(\tau)$ has to satisfy the tetrad condition, Eq (6), Eq(7)

$$e^{\hat{3}\hat{1}}(\tau) = (\sinh(\frac{a_0\tau}{c}), \cosh(\frac{a_0\tau}{c}), 0, 0) \quad (12)$$

In this time,

$$\bar{e}_{\hat{t}}^{\rho} = (1/\sqrt{1 - \frac{2GM}{rc^2}}, 0, 0, 0), \quad \bar{e}_{\hat{r}}^{\rho} = (0, \sqrt{1 - \frac{2GM}{rc^2}}, 0, 0)$$

$$\bar{e}_{\hat{\theta}}^{\rho} = (0, 0, 1/r, 0), \quad \bar{e}_{\hat{\phi}}^{\rho} = (0, 0, 0, 1/r \sin\theta)$$

$$g_{\rho\sigma} \bar{e}_{\hat{\alpha}}^{\rho} \bar{e}_{\hat{\beta}}^{\sigma} = \eta_{\hat{\alpha}\hat{\beta}} \quad (13)$$

$$\frac{a_0}{c} \hat{t} = \sinh(\frac{a_0}{c} \tau) = \frac{v/c}{\sqrt{1-v^2/c^2}}, \sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}} = \cosh(\frac{a_0}{c} \tau) = \frac{1}{\sqrt{1-v^2/c^2}} \quad (14)$$

Therefore, the Lorentz transformation $B^{\hat{\alpha}\hat{\mu}}(v)$ is

$$B^{\hat{\alpha}\hat{\mu}}(v) = \begin{pmatrix} 1 & v/c & 0 & 0 \\ \frac{v/c}{\sqrt{1-v^2/c^2}} & \frac{1}{\sqrt{1-v^2/c^2}} & 0 & 0 \\ \frac{v/c}{\sqrt{1-v^2/c^2}} & \frac{1}{\sqrt{1-v^2/c^2}} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$= e^{\hat{\alpha}\hat{\mu}}(\tau) = \begin{pmatrix} \cosh(\frac{a_0}{c} \tau) & \sinh(\frac{a_0}{c} \tau) & 0 & 0 \\ \sinh(\frac{a_0}{c} \tau) & \cosh(\frac{a_0}{c} \tau) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (15)$$

$$\bar{e}_{\hat{\mu}}^{\rho} = B^{\hat{\alpha}\hat{\mu}}(v) \bar{e}_{\hat{\alpha}}^{\rho} = e^{\hat{\alpha}\hat{\mu}}(\tau) \bar{e}_{\hat{\alpha}}^{\rho} \quad (16)$$

Hence,

$$g_{\rho\sigma} \bar{e}_{\hat{\alpha}}^{\rho} \bar{e}_{\hat{\beta}}^{\sigma} = \eta_{\hat{\alpha}\hat{\beta}}$$

$$g_{\rho\sigma} B^{\hat{\alpha}\hat{\mu}}(v) \bar{e}_{\hat{\alpha}}^{\rho} B^{\hat{\beta}\hat{\nu}}(v) \bar{e}_{\hat{\beta}}^{\sigma} = g_{\rho\sigma} \bar{e}_{\hat{\mu}}^{\rho} \bar{e}_{\hat{\nu}}^{\sigma} = \eta_{\hat{\alpha}\hat{\beta}} e^{\hat{\alpha}\hat{\mu}}(\tau) e^{\hat{\beta}\hat{\nu}}(\tau) = \eta_{\hat{\mu}\hat{\nu}} \quad (17)$$

III. The accelerated frame in the curved time-space

About the accelerated frame $\hat{\xi}$ in the curved time-space,

$$\begin{aligned}
d\tau^2 &= -\frac{1}{c^2} \eta_{\hat{\alpha}\hat{\beta}} d\hat{x}^\alpha d\hat{x}^\beta = -\frac{1}{c^2} \eta_{\hat{\alpha}\hat{\beta}} \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^\mu} \frac{\partial \hat{x}^\beta}{\partial \hat{\xi}^\nu} d\hat{\xi}^\mu d\hat{\xi}^\nu \\
&= -\frac{1}{c^2} \eta_{\hat{\alpha}\hat{\beta}} e^{\hat{\alpha}}{}_{\hat{\mu}} e^{\hat{\beta}}{}_{\hat{\nu}} d\hat{\xi}^\mu d\hat{\xi}^\nu \\
&= -\frac{1}{c^2} g_{\hat{\mu}\hat{\nu}} d\hat{\xi}^\mu d\hat{\xi}^\nu
\end{aligned} \tag{18}$$

$$e^{\hat{\alpha}}{}_{\hat{\mu}} = \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^\mu}, \quad \frac{\partial e^{\hat{\alpha}}{}_{\hat{0}}}{\partial \hat{\xi}^1} = \frac{\partial^2 \hat{x}^\alpha}{c \partial \hat{\xi}^0 \partial \hat{\xi}^1} = \frac{\partial e^{\hat{\alpha}}{}_{\hat{1}}}{c \partial \hat{\xi}^0} \tag{19}$$

In this time, in Eq(10),Eq(11),Eq(12), if uses $\hat{\xi}^0$ instead of τ ,

$$e^{\hat{\alpha}}{}_{\hat{0}}(\hat{\xi}^0) = \frac{1}{c} \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^0} = ((1 + \frac{a_0 \hat{\xi}^1}{c^2}) \cosh(\frac{a_0 \hat{\xi}^0}{c}), (1 + \frac{a_0 \hat{\xi}^1}{c^2}) \sinh(\frac{a_0 \hat{\xi}^0}{c}), 0, 0) \tag{20}$$

$$e^{\hat{\alpha}}{}_{\hat{1}}(\hat{\xi}^0) = \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^1} = (\sinh(\frac{a_0 \hat{\xi}^0}{c}), \cosh(\frac{a_0 \hat{\xi}^0}{c}), 0, 0) \tag{21}$$

$$e^{\hat{\alpha}}{}_{\hat{2}}(\hat{\xi}^0) = \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^2} = (0, 0, 1, 0), \quad e^{\hat{\alpha}}{}_{\hat{3}}(\hat{\xi}^0) = \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^3} = (0, 0, 1, 0) \tag{22}$$

$$d\hat{x}^\alpha = \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^\mu} d\hat{\xi}^\mu = e^{\hat{\alpha}}{}_{\hat{0}}(\hat{\xi}^0) c d\hat{\xi}^0 + e^{\hat{\alpha}}{}_{\hat{1}}(\hat{\xi}^0) d\hat{\xi}^1 + e^{\hat{\alpha}}{}_{\hat{2}}(\hat{\xi}^0) d\hat{\xi}^2 + e^{\hat{\alpha}}{}_{\hat{3}}(\hat{\xi}^0) d\hat{\xi}^3 \tag{23}$$

Hence,

$$cd\hat{t} = cd\hat{t} \sqrt{1 - \frac{2GM}{rc^2}} = (1 + \frac{a_0 \hat{\xi}^1}{c^2}) \cosh(\frac{a_0 \hat{\xi}^0}{c}) c d\hat{\xi}^0 + \sinh(\frac{a_0 \hat{\xi}^0}{c}) d\hat{\xi}^1$$

$$d\hat{r} = \frac{dr}{\sqrt{1 - \frac{2GM}{rc^2}}} = (1 + \frac{a_0 \hat{\xi}^1}{c^2}) \sinh(\frac{a_0 \hat{\xi}^0}{c}) c d\hat{\xi}^0 + \cosh(\frac{a_0 \hat{\xi}^0}{c}) d\hat{\xi}^1$$

$$d\hat{\theta} = d\hat{\xi}^2, \quad d\hat{\phi} = d\hat{\xi}^3 \tag{24}$$

$$\begin{aligned}
d\tau^2 &= \left(1 - \frac{2GM}{rc^2}\right)dt^2 - \frac{1}{c^2} \left[\frac{dr^2}{1 - \frac{2GM}{rc^2}} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right] \\
&= d\hat{t}^2 - \frac{1}{c^2} [d\hat{r}^2 + d\hat{\theta}^2 + d\hat{\phi}^2] \\
&= (1 + \frac{a_0 \hat{\xi}^1}{c^2})^2 (d\xi^0)^2 - \frac{1}{c^2} [(d\xi^1)^2 + (d\xi^2)^2 + (d\xi^3)^2]
\end{aligned} \tag{25}$$

The coordinate transformation is

$$c\hat{t} = \left(\frac{c^2}{a_0} + \hat{\xi}^1\right) \sinh\left(\frac{a_0 \hat{\xi}^0}{c}\right), \hat{r} = \left(\frac{c^2}{a_0} + \hat{\xi}^1\right) \cosh\left(\frac{a_0 \hat{\xi}^0}{c}\right) - \frac{c^2}{a_0}$$

$$\hat{\theta} = \hat{\xi}^2, \hat{\phi} = \hat{\xi}^3 \tag{26}$$

The inverse-transformation is

$$\hat{\xi}^0 = \frac{c}{a_0} \tanh^{-1} \left(\frac{c\hat{t}}{\hat{r} + \frac{c^2}{a_0}} \right), \hat{\xi}^1 = \sqrt{(\hat{r} + \frac{c^2}{a_0})^2 - c^2 \hat{t}^2} - \frac{c^2}{a_0}$$

$$\hat{\xi}^2 = \hat{\theta}, \hat{\xi}^3 = \hat{\phi} \tag{27}$$

If calculates the curvature tensor $R_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\lambda}}(\hat{\xi})$,

$$\begin{aligned}
R_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\lambda}}(\hat{\xi}) &= \frac{\partial \hat{x}^\alpha}{\partial \hat{\xi}^\mu} \frac{\partial \hat{x}^\beta}{\partial \hat{\xi}^\nu} \frac{\partial \hat{x}^\gamma}{\partial \hat{\xi}^\rho} \frac{\partial \hat{x}^\delta}{\partial \hat{\xi}^\lambda} R_{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}(\hat{x}) \\
&= e^{\hat{\alpha}}{}_{\hat{\mu}}(\hat{\xi}^0) e^{\hat{\beta}}{}_{\hat{\nu}}(\hat{\xi}^0) e^{\hat{\gamma}}{}_{\hat{\rho}}(\hat{\xi}^0) e^{\hat{\delta}}{}_{\hat{\lambda}}(\hat{\xi}^0) R_{\hat{\alpha}\hat{\beta}\hat{\gamma}\hat{\delta}}(\hat{x}) \\
R_{\hat{t}\hat{t}\hat{t}\hat{t}} &= -R_{\hat{t}\hat{t}\hat{t}\hat{t}} = R_{\hat{t}\hat{t}\hat{t}\hat{t}} = -R_{\hat{t}\hat{t}\hat{t}\hat{t}} = \frac{2GM}{r^3 c^2}, \\
R_{\hat{t}\hat{\theta}\hat{t}\hat{\theta}} &= -R_{\hat{t}\hat{\theta}\hat{t}\hat{\theta}} = R_{\hat{\theta}\hat{t}\hat{\theta}\hat{t}} = -R_{\hat{\theta}\hat{t}\hat{\theta}\hat{t}} = -\frac{GM}{r^3 c^2} = R_{\hat{t}\hat{\phi}\hat{t}\hat{\phi}} = -R_{\hat{t}\hat{\phi}\hat{t}\hat{\phi}} = R_{\hat{\phi}\hat{t}\hat{\phi}\hat{t}} = -R_{\hat{\phi}\hat{t}\hat{\phi}\hat{t}} \\
R_{\hat{\theta}\hat{\phi}\hat{\theta}\hat{\phi}} &= -R_{\hat{\theta}\hat{\phi}\hat{\theta}\hat{\phi}} = R_{\hat{\theta}\hat{\phi}\hat{\theta}\hat{\phi}} = -R_{\hat{\theta}\hat{\phi}\hat{\theta}\hat{\phi}} = -\frac{2GM}{r^3 c^2} \\
R_{\hat{r}\hat{\theta}\hat{r}\hat{\theta}} &= -R_{\hat{r}\hat{\theta}\hat{r}\hat{\theta}} = R_{\hat{\theta}\hat{r}\hat{r}\hat{\theta}} = -R_{\hat{\theta}\hat{r}\hat{r}\hat{\theta}} = \frac{GM}{r^3 c^2} = R_{\hat{r}\hat{\phi}\hat{r}\hat{\phi}} = -R_{\hat{r}\hat{\phi}\hat{r}\hat{\phi}} = R_{\hat{\phi}\hat{r}\hat{r}\hat{\phi}} = -R_{\hat{\phi}\hat{r}\hat{r}\hat{\phi}}
\end{aligned} \tag{29}$$

Therefore,

$$\begin{aligned}
e^{\hat{\alpha}_0}(\hat{\xi}^0) &= ((1 + \frac{a_0 \hat{\xi}^1}{c^2}) \cosh(\frac{a_0 \hat{\xi}^0}{c}), (1 + \frac{a_0 \hat{\xi}^1}{c^2}) \sinh(\frac{a_0 \hat{\xi}^0}{c}), 0, 0) \\
e^{\hat{\alpha}_1}(\hat{\xi}^0) &= (\sinh(\frac{a_0 \hat{\xi}^0}{c}), \cosh(\frac{a_0 \hat{\xi}^0}{c}), 0, 0) \\
e^{\hat{\alpha}_2}(\hat{\xi}^0) &= (0, 0, 1, 0), e^{\hat{\alpha}_3}(\hat{\xi}^0) = (0, 0, 1, 0)
\end{aligned} \tag{30}$$

$$\begin{aligned}
R_{\hat{0}\hat{1}\hat{0}\hat{1}}(\hat{\xi}) &= \frac{2GM}{r^3 c^2} (1 + \frac{a_0 \hat{\xi}^1}{c^2})^2, \quad R_{\hat{0}\hat{2}\hat{0}\hat{2}}(\hat{\xi}) = R_{\hat{0}\hat{3}\hat{0}\hat{3}}(\hat{\xi}) = -\frac{GM}{r^3 c^2} (1 + \frac{a_0 \hat{\xi}^1}{c^2})^2 \\
R_{\hat{2}\hat{3}\hat{2}\hat{3}}(\hat{\xi}) &= -\frac{2GM}{r^3 c^2}, \quad R_{\hat{1}\hat{2}\hat{1}\hat{2}}(\hat{\xi}) = R_{\hat{1}\hat{3}\hat{1}\hat{3}}(\hat{\xi}) = \frac{GM}{r^3 c^2}
\end{aligned} \tag{31}$$

Specially, if $t = 0$,

$$U = \frac{1}{\sqrt{1 - \frac{2GM}{rc^2}}} \frac{dr}{dt} = 0 \rightarrow V = \frac{d\hat{t}}{d\hat{t}} = -\frac{a_0 \hat{t}}{\sqrt{1 + \frac{a_0^2 \hat{t}^2}{c^2}}} = \frac{dr}{dt} \frac{1}{(1 - \frac{2GM}{rc^2})} = 0 \tag{32}$$

Therefore, if $t = \hat{t} = \hat{\xi}^0 = 0$, the theory treats the real situation.

$$\begin{aligned}
\hat{\xi}^1 &= \sqrt{(\hat{r} + \frac{c^2}{a_0})^2 - c^2 \hat{t}^2} - \frac{c^2}{a_0} = \hat{r} \\
d\hat{r} &= \frac{dr}{\sqrt{1 - \frac{2GM}{rc^2}}} \rightarrow \hat{r} = \sqrt{r} \sqrt{r - \frac{2GM}{c^2}} + \frac{2GM}{c^2} \ln |\sqrt{r} + \sqrt{r - \frac{2GM}{c^2}}| \\
&\quad - \sqrt{r_0} \sqrt{r_0 - \frac{2GM}{c^2}} - \frac{2GM}{c^2} \ln |\sqrt{r_0} + \sqrt{r_0 - \frac{2GM}{c^2}}| \\
a_0 &= \frac{1}{\sqrt{1 - \frac{2GM}{rc^2}}} \frac{d}{dt} \left(\frac{u}{\sqrt{1 - \frac{2GM}{rc^2} - \frac{u^2}{c^2}}} \right) = \frac{a}{\sqrt{1 - \frac{2GM}{rc^2}}} \\
a &= \frac{d}{dt} \left(\frac{u}{\sqrt{1 - \frac{2GM}{rc^2} - \frac{u^2}{c^2}}} \right) \\
r_0 \text{ is the location of the stationary accelerated frame} &
\end{aligned} \tag{33}$$

In this time, in the curved time-space, the curvature tensor $R_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\lambda}}(\hat{\xi})$ of the stationary accelerated frame is

$$\begin{aligned}
R_{\hat{0}\hat{1}\hat{0}\hat{1}}(\hat{\xi}) &= \frac{2GM}{r^3c^2} \left(1 + \frac{a_0\hat{\xi}^1}{c^2}\right)^2 = \frac{2GM}{r^3c^2} \left(1 + \frac{a_0\hat{r}}{c^2}\right)^2 \\
&= \frac{2GM}{r^3c^2} \left[1 + \frac{a_0}{c^2} \left\{ \sqrt{r} \sqrt{r - \frac{2GM}{c^2}} + \frac{2GM}{c^2} \ln |\sqrt{r} + \sqrt{r - \frac{2GM}{c^2}}| \right.\right. \\
&\quad \left.\left. - \sqrt{r_0} \sqrt{r_0 - \frac{2GM}{c^2}} - \frac{2GM}{c^2} \ln |\sqrt{r_0} + \sqrt{r_0 - \frac{2GM}{c^2}}| \right\} \right]^2 \\
&= \frac{2GM}{r^3c^2} \left[1 + \frac{1}{c^2} \frac{a}{\sqrt{1 - \frac{2GM}{rc^2}}} \left\{ \sqrt{r} \sqrt{r - \frac{2GM}{c^2}} + \frac{2GM}{c^2} \ln |\sqrt{r} + \sqrt{r - \frac{2GM}{c^2}}| \right.\right. \\
&\quad \left.\left. - \sqrt{r_0} \sqrt{r_0 - \frac{2GM}{c^2}} - \frac{2GM}{c^2} \ln |\sqrt{r_0} + \sqrt{r_0 - \frac{2GM}{c^2}}| \right\} \right]^2 \\
R_{\hat{0}\hat{2}\hat{0}\hat{2}}(\hat{\xi}) &= R_{\hat{0}\hat{3}\hat{0}\hat{3}}(\hat{\xi}) = -\frac{GM}{r^3c^2} \left(1 + \frac{a_0\hat{\xi}^1}{c^2}\right)^2 = -\frac{GM}{r^3c^2} \left(1 + \frac{a_0\hat{r}}{c^2}\right)^2 \\
&= -\frac{GM}{r^3c^2} \left[1 + \frac{a_0}{c^2} \left\{ \sqrt{r} \sqrt{r - \frac{2GM}{c^2}} + \frac{2GM}{c^2} \ln |\sqrt{r} + \sqrt{r - \frac{2GM}{c^2}}| \right.\right. \\
&\quad \left.\left. - \sqrt{r_0} \sqrt{r_0 - \frac{2GM}{c^2}} - \frac{2GM}{c^2} \ln |\sqrt{r_0} + \sqrt{r_0 - \frac{2GM}{c^2}}| \right\} \right]^2 \\
&= -\frac{GM}{r^3c^2} \left[1 + \frac{1}{c^2} \frac{a}{\sqrt{1 - \frac{2GM}{rc^2}}} \left\{ \sqrt{r} \sqrt{r - \frac{2GM}{c^2}} + \frac{2GM}{c^2} \ln |\sqrt{r} + \sqrt{r - \frac{2GM}{c^2}}| \right.\right. \\
&\quad \left.\left. - \sqrt{r_0} \sqrt{r_0 - \frac{2GM}{c^2}} - \frac{2GM}{c^2} \ln |\sqrt{r_0} + \sqrt{r_0 - \frac{2GM}{c^2}}| \right\} \right]^2 \\
R_{\hat{2}\hat{3}\hat{2}\hat{3}}(\hat{\xi}) &= -\frac{2GM}{r^3c^2}, \quad R_{\hat{1}\hat{2}\hat{1}\hat{2}}(\hat{\xi}) = R_{\hat{1}\hat{3}\hat{1}\hat{3}}(\hat{\xi}) = \frac{GM}{r^3c^2}
\end{aligned}$$

r_0 is the location of the stationary accelerated frame (34)

IV. Conclusion

In the general relativity theory, defines the accelerated frame that moves in \hat{r} -axis in the curved time-space. Specially, if $t = \hat{t} = \hat{\xi}^0 = 0$, this theory treats the curvature tensor of the stationary accelerated frame in the curved time-space.

Reference

- [1]S.Weinberg, Gravitation and Cosmology (John Wiley & Sons, Inc., 1972)
- [2]P.Bergman, Introduction to the Theory of Relativity (Dover Pub. Co., Inc., New York, 1976), Chapter V
- [3]C.Misner, K.Thorne and J. Wheeler, Gravitation (W.H.Freeman & Co., 1973)
- [4]S.Hawking and G.Ellis, The Large Scale Structure of Space-Time (Cambridge University Press, 1973)
- [5]R.Adler, M.Bazin and M.Schiffer, Introduction to General Relativity (McGraw-Hill, Inc., 1965)
- [6]M.Schwarzschild, Structure and Evolution of the Stars (Princeton University Press, 1958; reprint, Dover, N.Y. 1965), chapter II
- [7]S.Chandrasekhar, Mon.Not.Roy.Astron.Soc.95.207(1935)
- [8]C.Rhoades, "Investigations in the Physics of Neutron Stars", doctoral dissertation, Princeton University
- [9]J.Oppenheimer and H.Snyder, phys.Rev.56,455(1939)