GENERALIZATION OF AN ER'S MATRIX METHOD FOR COMPUTING

Florentin Smarandache, Ph D
Associate Professor
Chair of Department of Math & Sciences
University of New Mexico
200 College Road
Gallup, NM 87301, USA
E-mail:smarand@unm.edu

Er's matrix method for computing Fibonacci numbers and their sums can be extended to the s-additive sequence:

$$g_{-s+1} = g_{-s+2} = \dots = g_{-1} = 0, g_0 = 1,$$

and

$$g_n = \sum_{i=1}^s g_{n-i}$$
 for $n > 0$.

For example, if we note $S_n = \sum_{j=1}^{n-1} g_j$, we define two $(s+1) \times (s+1)$ matrixes such

that:

$$B_{n} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ S_{n} & g_{n} & g_{n-1} & \dots & g_{n-s+2} & g_{n-s+1} \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ S_{n-s+1} & g_{n-s+1} & g_{n-s} & \dots & g_{n-2s+3} & g_{n-2s+2} \end{bmatrix},$$

 $n \ge 1$, and

$$M = \begin{bmatrix} 1 & 0 & 0 \dots & 0 \\ 1 & 1 & 1 \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & 1 & 0 & \dots & 1 \\ 1 & 1 & 0 & \dots & 0 \end{bmatrix},$$

thus, we have analogously:

$$B_{n+1} = M^{n+1}, M^{r+c} = M^r \cdot M^c,$$

whence

$$S_{r+c} = S_r + g_r S_c + g_{r-1} S_{c-1} + \dots + g_{r-s+1} S_{c-s+1},$$

$$g_{r+c} = g_r g_c + g_{r-1} g_{c-1} + \dots + g_{r-s+1} g_{c-s+1},$$

and for
$$r = c = n$$
 it results:

$$S_{2n} = S_n + g_n S_n + g_{n-1} S_{n-1} + ... + g_{n-s+1} S_{n-s+1},$$

$$g_{2n} = g_n^2 + g_{n-1}^2 + ... + g_{n-s+1}^2;$$
for $r = n$, $c = n - 1$, we find:

$$g_{2n-1} = g_n g_{n-1} + g_{n-1} g_{n-2} + ... + g_{n-s+1} g_{n-s}, \text{ etc.}$$

$$S_{2n-1} = S_n + g_n S_{n-1} + g_{n-1} S_{n-2} + ... + g_{n-s+1} S_{n-s}$$

Whence we can construct a similar algorithm as M. C. Er for computing s-additive numbers and their sums.

REFERENCE:

M. C. Er, Fast Computation of Fibonacci Numbers and Their Sums, J. Inf. Optimization Sci. (Delhi), Vol. 6 (1985), No. 1. pp. 41-47.

[Published in "GAMMA", Braşov, Anul X, No.. 1-2, October 1987, p. 8]