

Inadequacy of Hubble-Friedmann Cosmology and the Basics of Stoney Scale Black Hole Cosmology

U. V. S. Seshavatharam^{*1,2} and S. Lakshminarayana⁺

^{*1}Honorary faculty, I-SERVE, Alakapuri, Hyderabad-35, India. Email: seshavatharam.uvs@gmail.com

^{*2}Sr. QA Engineer, DIP Division, Lanco Industries Ltd, Rachagunneri, Srikalahasti-517644, AP, India.

⁺Dept. of Nuclear Physics, Andhra University, Visakhapatnam-03, India. Email: Insrirama@yahoo.com

Abstract: Throughout the cosmic evolution, currently believed cosmic ‘critical density’ can be shown to be a default result of the ‘positively curved’ light speed rotating black hole universe ‘volume density’. As there is no observational or experimental evidence to Friedmann’s second assumption, the density classification scheme of Friedmann cosmology must be reviewed at fundamental level and possibly can be relinquished. The observed cosmic redshift can be reinterpreted as an index of ‘cosmological’ thermodynamic light emission mechanism. Clearly speaking during cosmic evolution, at any time in the past, in hydrogen atom- emitted photon energy was always inversely proportional to the cosmic temperature. Thus past light emitted from older galaxy’s excited hydrogen atom will show redshift with reference to the current laboratory data. Note that there will be no change in the energy of the emitted photon during its journey from the distant galaxy to the observer. By considering the ‘Stoney mass’ as the initial mass of the baby cosmic black hole, past and current physical and thermal parameters (like angular velocity, growth rate, age, redshift, thermal energy density and matter density) of the cosmic black hole can be understood. For a cosmic temperature of 3000 K, obtained redshift is 1100. From now onwards, CMBR temperature can be called as ‘Comic Black Hole’s Thermal Radiation’ temperature and can be expressed as ‘CBHTR’ temperature. Uncertainty relation and all other microscopic physical constants play a crucial role in understanding the halt of the present cosmic expansion. In view of the confirmed zero rate of change in inverse of the Fine structure ratio (from the ground based laboratory experimental results), zero rate of change in the current CMBR temperature (from satellite data) and zero rate of change in the current Hubble’s constant (from satellite data), it can be suggested that, current cosmic expansion is almost all saturated and at present there is no significant cosmic acceleration.

Keywords: Mach’s principle, Stoney mass, Black Hole Cosmology, Cosmic growth index, Cosmic growth rate, Hubble Potential, Cosmic redshift, Cosmic age, Halting of Cosmic Expansion, Final Unification.

1. Introduction

Authors published their concepts on black hole cosmology in many online journals [1-13]. In this paper by highlighting the basic short comings of Friedmann cosmology [14] an attempt is made to review the model of black hole cosmology [15-28] in terms of cosmic redshift, CMBR redshift, cosmic growth index, cosmic growth rate and cosmic age. The basic shortcomings of modern cosmology can be expressed as follows. For more information one may see the appendix [1].

- 1) No direct observational evidence to Friedmann’s second assumption [29].
- 2) No theoretical base in Friedmann’s ‘critical density’ concept and the ‘matter density’ classification scheme.
- 3) If light is coming from the atoms of the gigantic galaxy, then redshift can also be interpreted as an index of the galactic cosmological atomic ‘light emission mechanism’. In no way it seems to be connected with ‘galaxy receding’.
- 4) No theoretical base in the currently believed wave length based redshift definition [30,31]. In terms of ‘quantum of energy’, redshift can also be interpreted as an index of cosmological thermodynamic light emission mechanism in hydrogen atom.
- 5) Merely by estimating galaxy distance and without measuring galaxy receding speed, one cannot verify its receding speed or acceleration. (Clearly speaking: two mistakes are possible here. i) Assumed galaxy receding speed is not being measured and not being confirmed. ii) Without measuring and confirming the galaxy receding speed, how can one say and confirm that it (galaxy) is accelerating).
- 6) No theoretical base in considering the Hubble’s constant merely as the cosmic expansion parameter. With reference to angular velocity it is having deep inner meaning.
- 7) No direct observational evidence for the current cosmic acceleration and the dark energy [32,33].
- 8) By substituting the geometric mean mass of $(c^3/2GH_0)$ and $\sqrt{hc/2\pi G}$ in the famous Hawking’s black hole temperature formula automatically the observed 2.725 K can be fitted very accurately.

Requesting your kind and valuable (unbiased) review comments...

- 52 9) When Friedmann's cosmology was taking its final shape, black hole physics was in its beginning stage.
53 10) No comparative and relational study in between Friedmann cosmology and microscopic physical phenomena.

54
55 Friedmann made two simple assumptions about the universe. They can be stated in the following way.

- 56
57 1. When viewed at large enough scales, universe appears the same in every direction.
58 2. When viewed at large enough scales, universe appears the same from every location.

59
60 In this regard Hawking says : "There is no scientific evidence for the Friedmann's second assumption. We believe it only
61 on grounds of modesty: it would be most remarkable if the universe looked the same in every direction around us, but not
62 around other points in the universe". This is one key point to be noted here. The term 'critical density' is the back bone of
63 modern cosmology. At any time in the past, it is generally expressed in the following way.

64
65
$$(\rho_c)_t \cong \frac{3H_t^2}{8\pi G} \quad (1)$$

66 Its current expression is as follows.

67
$$(\rho_c)_0 \cong \frac{3H_0^2}{8\pi G} \quad (2)$$

68 According to standard Friedmann cosmology,

- 69
70 1. If matter density is greater than the critical density, universe will have a positive curvature.
71 2. If matter density equals the critical density, universe will be flat.
72 3. If matter density is less than the critical density, universe will have a negative curvature.

73
74 But by considering 'black hole geometry' as the 'eternal cosmic geometry' and by assuming 'constant light speed
75 rotation' throughout the cosmic evolution, at any time the currently believed cosmic 'critical density' can be shown to be
76 the cosmic black hole's eternal 'volume density'. If mass of the black hole universe is M_t , $\left(\frac{c}{H_t}\right)$ is the radius of the
77 black hole universe that rotates at light speed and angular velocity H_t , at any time in the past,

78
79
$$\frac{2GM_t}{c^2} \cong \frac{c}{H_t} \quad \text{and} \quad M_t \cong \frac{c^3}{2GH_t}. \quad (3)$$

80
$$(\rho_v)_t \cong (M_t) \left[\frac{4\pi \left(\frac{c}{H_t}\right)^3}{3} \right]^{-1} \cong \left(\frac{c^3}{2GH_t} \right) \left[\frac{3 \left(\frac{H_t}{c}\right)^3}{4\pi} \right] \cong \frac{3H_t^2}{8\pi G} \quad (4)$$

81 At present,

82
$$(\rho_v)_0 \cong (M_0) \left[\frac{4\pi \left(\frac{c}{H_0}\right)^3}{3} \right]^{-1} \cong \left(\frac{c^3}{2GH_0} \right) \left[\frac{3 \left(\frac{H_0}{c}\right)^3}{4\pi} \right] \cong \frac{3H_0^2}{8\pi G} \quad (5)$$

83 Based on this coincidence and as there is no observational or experimental evidence to Friedmann's second assumption, the
84 density classification scheme of Friedmann cosmology must be reviewed at fundamental level.

85 **2. Possible Assumptions and Possible Explanation**

86 Possible assumptions in unified cosmic physics can be expressed in the following way.

87
88 **Assumption-1: With reference to the elementary charge and with mass similar to the Planck mass, a new mass unit**
89 **can be constructed in the following way. It can be called as the Stoney mass.**

90
91
$$(M_S)^\pm \cong \sqrt{\frac{e^2}{4\pi\epsilon_0 G}} \cong 1.859272 \times 10^{-9} \text{ Kg} \cong 1.042975 \times 10^{18} \text{ GeV}/c^2 \quad (6)$$

Requesting your kind and valuable (unbiased) review comments...

92 **Assumption-2: At any time Hubble length (c / H_t) can be considered as the gravitational or electromagnetic**
 93 **interaction range.**

94
 95 **Assumption-3: At any time, H_t being the angular velocity, universe can be considered as a growing and light speed**
 96 **rotating primordial black hole.** Thus at any given cosmic time,

97
$$R_t \cong \frac{2GM_t}{c^2} \cong \frac{c}{H_t} \text{ and } M_t \cong \frac{c^3}{2GH_t} \quad (7)$$

98
$$\text{when } M_t \rightarrow M_s, R_s \cong \frac{2GM_s}{c^2} \text{ and } H_s \cong \frac{c}{R_s} \cong \frac{c^3}{2GM_s} \quad (8)$$

99
 100 can be considered as the characteristic initial physical measurements of the universe. Here the subscript S refers to the
 101 initial conditions of the universe and can be called as the Stoney scale. Similarly
 102

103
$$R_0 \cong \frac{2GM_0}{c^2} \cong \frac{c}{H_0}, M_0 \cong \frac{c^3}{2GH_0} \text{ and } H_0 \cong \frac{c^3}{2GM_0} \quad (9)$$

104
 105 can be considered as the characteristic current physical measurements of the universe.
 106

107 **Assumption-4: During cosmic evolution, at any time the past, in hydrogen atom emitted photon energy was always**
 108 **inversely proportional to the cosmic temperature. Thus past light emitted from older galaxy's hydrogen atom will**
 109 **show redshift with reference to the current laboratory data. There will be no change in the energy of the emitted**
 110 **photon during its journey from the distant galaxy to the observer.**

111
 112
$$\frac{E_t}{E_0} \cong \frac{\lambda_0}{\lambda_t} \cong \frac{T_t}{T_0} \quad (10)$$

113 Here, E_t is the energy of emitted photon from the galactic hydrogen atom and E_0 is the corresponding energy in the
 114 laboratory. λ_t is the wave length of emitted and received photon from the galactic hydrogen atom and λ_0 is the
 115 corresponding wave length in the laboratory. T_t is the cosmic temperature at the time when the photon was emitted and is
 116 T_0 the current cosmic temperature.

117
 118 **Assumption-5: At any given time, ratio of volume energy density and thermal energy density can be called as the**
 119 **cosmic growth index and can be expressed as follows.**

120
 121
$$\frac{3H_t^2 c^2}{8\pi G a T_t^4} \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^2 \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^2 \quad (11)$$

$$\cong \text{Cosmic Growth index}$$

122 Thus at the Stoney scale,

123
$$\frac{3H_s^2 c^2}{8\pi G a T_s^4} \cong \left[1 + \ln \left(\frac{M_s}{M_s} \right) \right]^2 \cong \left[1 + \ln \left(\frac{H_s}{H_s} \right) \right]^2 \cong 1 \quad (12)$$

124 **Assumption-6: At any given time, cosmic black hole's growth rate can be expressed as $g_t \cong \left(\frac{3H_t^2 c^2}{8\pi G a T_t^4} \right)^{-1} c$. With this**
 125 **idea and by considering the average growth rate cosmic age can be estimated.**

126
 127
$$g_t \cong \text{Cosmic growth rate} \cong \frac{c}{\text{cosmic growth index}} \quad (13)$$

$$\cong \left(\frac{3H_t^2 c^2}{8\pi G a T_t^4} \right)^{-1} c \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^{-2} c \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-2} c$$

Requesting your kind and valuable (unbiased) review comments...

128 At the Stoney scale,

$$129 \quad g_s \cong \left(\frac{3H_s^2 c^2}{8\pi G a T_s^4} \right)^{-1} \quad c \cong \left[1 + \ln \left(\frac{M_s}{M_s} \right) \right]^{-2} \quad c \cong \left[1 + \ln \left(\frac{H_s}{H_s} \right) \right]^{-2} \quad c \cong c \quad (14)$$

130

131

2.1 Possible Explanation for the proposed Assumptions

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

To have some clarity and to have some quantitative measurements and fittings of initial and current states of the black hole universe - instead of considering 'star - black hole explosions' and 'higher dimensions', the authors of this paper focused their attention only on the old and famous Mach's principle [34], 'Hubble volume' and 'primordial evolving black holes'. Some cosmologists use the term 'Hubble volume' to refer to the volume of the observable universe. There is no perfect theory that defines the lower and upper limits of a massive black hole. Most of the theoretical models assume a lower mass limit close to the 'Planck mass'. Astronomers believe that black holes that are as large as a billion solar masses can be found at the centre of most of the galaxies. Here the fundamental questions to be answered are: If the galactic central black hole mass is 10 billion solar masses and density is less than 1 kg/m^3 - with such a small density and large mass, without collapsing - how it is able to hold a gigantic galaxy? What force makes the black hole stable? Recent observations confirm that, instead of collapsing, galactic central black holes are growing faster and spinning with light speed. Even though mass is too high and density is too low, light speed rotation certainly helps in maintaining black hole's stability from collapsing with maximum possible outward radial force of the magnitude close to (c^4/G) . Based on these points the authors propose the following picture of Black hole cosmology. Forever rotating at light speed, high temperature and high angular velocity small sized primordial cosmic black hole of mass $M_s \cong \sqrt{e^2/4\pi\epsilon_0 G}$ gradually transforms into a low temperature and low angular velocity large sized massive primordial cosmic black hole. At any given cosmic time, for the primordial growing black hole universe, its 'Schwarzschild radius' can be considered as its characteristic possible minimum radius and 'constant light speed rotation' will give the maximum possible stability from collapsing. Here $M_s \cong \sqrt{e^2/4\pi\epsilon_0 G}$ can be called as the mass of the primordial baby black hole universe. Here 4 important points can be stated as follows.

1. It is well known that e, c, G play a vital role in fundamental physics. With these 3 constants space-time curvature concepts at a charged particle surface can be studied. Note that the basic concept of unification is to understand the origin of 'mass' of any particle. Mass is the basic property in 'gravitation' and charge is the basic property in 'atomicity'. So far no model established a cohesive relation in between 'electric charge' and 'mass' of any 'elementary particle' or 'cosmic dust'. From physics point of view, the fundamental questions to be answered are: 1) Without charge, is there any independent existence to "mass"? 2) Without mass, is there any independent existence to "charge"? From cosmology point of view the fundamental questions to be answered are: 1) What is 'cosmic dust'? 2) Without charge, is there any independent existence to "cosmic dust"? From astrophysics point of view the fundamental questions to be answered are: 1) Without charge, is there any independent existence to 'mass' of any star? 2) Is black hole - a neutral body or electrically a neutralized body? To understand these questions the authors made an attempt to construct the above unified mass unit. It is having a long history. It was first introduced by the physicist George Johnstone Stoney [35]. He is most famous for introducing the term 'electron' as the 'fundamental unit quantity of electricity'. With this mass unit in unification program with a suitable proportionality it may be possible to represent the characteristic mass of elementary charge. It can be considered as the seed of galactic matter or galactic central black hole. It can also be considered as the seed of any cosmic structure. If 2 such oppositely charged particles annihilates, a large amount of energy can be released. If so under certain extreme conditions at the vicinity of massive stars or black holes, a very high energy radiation can be seen to be emitted by the pair annihilation of M_s . With this mass unit, proton-electron mass ratio and proton and electron rest masses can be fitted. Thus with reference to the elementary charge and electron & proton rest masses, magnitude of the gravitational constant can be fitted [1,2].
2. In theoretical physics, particularly in discussions of gravitation theories, Mach's principle is the name given by Einstein to an interesting hypothesis often credited to the physicist and philosopher Ernst Mach. The idea is that the local motion of a rotating reference frame is determined by the large scale distribution of matter. With reference to the Mach's principle and the Hubble volume, at any cosmic time, if 'Hubble mass' is the product of cosmic 'critical density' and the 'Hubble volume', then it can be suggested that, i) Each and every point in the free space is influenced by the Hubble mass, ii) Hubble volume and Hubble mass play a vital role in understanding the properties of electromagnetic and nuclear interactions and iii) Hubble volume and Hubble mass play a key role in understanding the geometry of the universe. With reference to the famous Mach's principle, 'Hubble volume' and 'Hubble mass' both can

Requesting your kind and valuable (unbiased) review comments...

- 181 be considered as quantitative measurements of the ‘distance cosmic back ground’. As a first attempt, in this paper
182 authors proposed a semi empirical relation that connects the CMBR energy density, Hubble’s constant and
183 $\sqrt{e^2/4\pi\epsilon_0 G}$.
- 184 3. Starting from an electron to any gigantic galaxy, rotation is a common phenomenon in atomic experiments and
185 astronomical observations. From Newton’s laws of motion and based on the Mach’s principle, sitting inside a closed
186 universe, one cannot comment whether the universe is rotating or not. We have to search for alternative means for
187 confirming the cosmic rotation. Recent findings from the University of Michigan [36] suggest that the shape of the Big
188 Bang might be more complicated than previously thought, and that the early universe spun on an axis. A left-handed
189 and right-handed imprint on the sky as reportedly revealed by galaxy rotation would imply the universe was rotating
190 from the very beginning and retained an overwhelmingly strong angular momentum. An anonymous referee who
191 reviewed the paper for Physics Letters said, “In the paper the author claims that there is a preferred handedness of
192 spiral galaxies indicating a preferred direction in the universe. Such a claim, if proven true, would have a profound
193 impact on cosmology and would very likely result in a “Nobel prize”. The consequences of a spinning universe [36-49]
194 seem to be profound and natural. Not only that, with ‘constant rotation speed’ ‘cosmic collapse’ can be prevented and
195 can be considered as an alternative to the famous ‘repulsive gravity’ concept. If so, at any time to have maximum
196 possible stability from collapsing ‘constant light speed rotation’ can be considered as a constructive and workable
197 concept.
- 198 4. Recent observations confirm black hole’s light speed rotation. In 2013 February, using NASA’s newly launched NuStar
199 telescope and the European Space Agency’s workhorse XMM-Newton, an international team observed high-energy X-
200 rays released by a super massive black hole in the middle of a nearby galaxy. They calculated its spin at close to the
201 speed of light: 670 million mph [50,51]. Please note that, for any black hole even though its mass is too high and
202 density is too low, light speed rotation certainly helps in maintaining its stability from collapsing with maximum
203 possible outward radial force of magnitude (c^4/G) . At the beginning of comic evolution if rotation speed was zero and
204 there was no big bang - definitely it will cast a doubt on the stability, existence and angular velocity of the assumed
205 initial primordial cosmic baby black hole. Hence at the beginning also, to guess or define the angular velocity and to
206 have maximum possible stability it is better to assume light speed rotation for the cosmic baby black hole. At present if
207 rate of cosmic expansion is very slow, then rate of decrease in angular velocity will be very small and practically can
208 be considered as zero. Along with (practically) constant angular velocity, at present if constant light speed rotation is
209 assumed to be maintained then cosmic stability will be maximum and rate of change in cosmic size will be practically
210 zero and hence this idea helps us to believe in present Hubble length along with the observed ordered galactic
211 structures and uniform thermal energy density.

212 **2.2 To reinterpret the Hubble’s constant**

213 With a simple derivation it is possible to show that, Hubble’s constant H_i represents the cosmological angular velocity.
214 Authors presented this derivation in their published papers. Basic idea of this derivation is to express the angular velocity
215 of any rotating celestial body in terms of its mass, radius, mass density and surface escape velocity. Assume that, a planet
216 of mass M and radius R rotates with angular velocity ω_e and linear velocity v_e in such a way that, free or loosely bound
217 particle of mass m lying on its equator gains a kinetic energy equal to potential energy as,

$$218 \quad \frac{1}{2}mv_e^2 = \frac{GMm}{R} \quad (15)$$

$$219 \quad R\omega_e = v_e = \sqrt{\frac{2GM}{R}} \quad \text{and} \quad \omega_e = \frac{v_e}{R} = \sqrt{\frac{2GM}{R^3}} \quad (16)$$

220 i.e Linear velocity of planet’s rotation is equal to free particle’s escape velocity. Without any external power or energy, test
221 particle gains escape velocity by virtue of planet’s rotation. Note that if Earth completes one rotation in one hour then free
222 particles lying on the equator will get escape velocity. Now writing $M = \frac{4\pi}{3}R^3\rho_e$,

$$223 \quad \omega_e = \frac{v_e}{R} = \sqrt{\frac{8\pi G\rho_e}{3}} \quad \text{Or} \quad \omega_e^2 = \frac{8\pi G\rho_e}{3} \quad (17)$$

$$224 \quad \text{Density, } \rho_e = \frac{3\omega_e^2}{8\pi G} \quad (18)$$

Requesting your kind and valuable (unbiased) review comments...

225 In real time, this obtained density may or may not be equal to the actual density. But the ratio $\frac{8\pi G \rho_{real}}{3\omega_{real}^2}$ may have some
226 physical significance. The most important point to be noted here, is that, as far as dimensions and units are considered,
227 from equation (18), it is very clear that, proportionality constant being $\frac{3}{8\pi G}$,
228

$$229 \text{ density} \propto (\text{angular velocity})^2 \quad (19)$$

230 Equation (18) is similar to “flat model concept” of cosmic “critical density”
231

$$232 \rho_c = \frac{3H_t^2}{8\pi G} \quad (20)$$

233 Comparing equations (18) and (20) dimensionally and conceptually, i.e.
234

$$235 \rho_e = \frac{3\omega_e^2}{8\pi G} \text{ with } \rho_c = \frac{3H_t^2}{8\pi G} \quad (21)$$

$$236 H_t^2 \rightarrow \omega_e^2 \text{ and } H_t \rightarrow \omega_e \quad (22)$$

237 It is very clear that, dimensions of ‘Hubble’s constant’ must be ‘radian/second’. In any physical system under study, for
238 any one ‘simple physical parameter’ there will not be two different units and there will not be two different physical
239 meanings. This is a simple clue and brings ‘cosmic rotation’ into picture. This is possible in a closed universe only. Cosmic
240 models that depend on this “critical density” may consider ‘angular velocity of the universe’ in the place of ‘Hubble’s
241 constant’. In the sense, with a great confidence ‘cosmic rotation’ can be included in the existing models of cosmology. Then
242 the term ‘critical density’ appears to be the ‘volume density’ of the closed and expanding universe. Thinking in this way,
243 considering ‘black hole geometry’ as the ‘eternal cosmic geometry’ and by assuming ‘constant light speed rotation’
244 throughout the cosmic evolution, at any time the currently believed cosmic ‘critical density’ can be shown to be the cosmic
245 black hole’s eternal ‘volume density’. Thus based on the Mach’s principle, ‘distance cosmic back ground’ can be quantified
246 in terms of ‘Hubble volume’ and ‘Hubble mass’.

247 **2.3 To reinterpret the Cosmic redshift**

248 Hubble initially interpreted red shifts [30] as a Doppler effect, due to the motion of the galaxies as they receded for our
249 location in the Universe[52]. He called it a ‘Doppler effect’ as though the galaxies were moving ‘through space’; that is
250 how some astronomers initially perceived it. This is different to what has now become accepted but observations alone
251 could not distinguish between the two concepts. In 1947 he [31] stated that: “The red shifts are more easily interpreted as
252 evidence of motion in the line of sight away from the earth – as evidence that the nebulae in all directions are rushing away
253 from us and that the farther away they are, the faster they are receding. This interpretation lends itself directly to theories of
254 expanding universe. The interpretation is not universally accepted, but even the most cautious of us admit that red shifts are
255 evidence of either an expanding universe or of some hitherto unknown principle of nature”. “Attempts have been made to
256 attain the necessary precision with the 100 inch, and the results appear to be significant. If they are valid, it seems likely
257 that the red-shifts may not be due to an expanding universe, and much of the current speculation on the structure of the
258 universe may require re-examination. The significant data, however, were necessarily obtained at the very limit of a single
259 instrument, and there were no possible means of checking the results by independent evidence. Therefore the results must
260 be accepted for the present as suggestive rather than definitive”. “We may predict with confidence that the 200 inch will
261 tell us whether the red shifts must be accepted as evidence of a rapidly expanding universe, or attributed to some new
262 principle in nature. Whatever may be the answer, the result may be welcomed as another major contribution to the
263 exploration of the universe.”

264 In this regard if one is willing to consider the proposed assumptions, in hydrogen atom emitted photon energy can be under
265 stood as follows.

- 266 1. As the cosmic time increases cosmic angular velocity and hence cosmic temperature both decrease. As a result, during
267 cosmic evolution, in hydrogen atom, binding energy increases in between proton and electron.

Requesting your kind and valuable (unbiased) review comments...

- 268 2. As cosmic temperature decreases, it requires more excitation energy to break the bond between electron and the proton.
269 In this way, during cosmic evolution, whenever it is excited, hydrogen atom emits photons with increased quantum of
270 energy.
271 3. Thus past light quanta emitted from old galaxy's excited hydrogen atom will have less energy and show a red shift with
272 reference to the current laboratory magnitude.
273 4. During journey light quanta will not lose energy and there will be no change in light wavelength.
274 5. Galactic photon energy in hydrogen atom when it was emitted can be estimated as follows.
275

276
$$E_t \cong \frac{hc}{\lambda_t} \cong \left(\frac{T_0}{T_t}\right) \left(\frac{hc}{\lambda_0}\right) \cong \left(\frac{T_0}{T_t}\right) E_0 \quad (23)$$

277 Here, λ_0 is the wavelength of photon in the laboratory.

278 E_t is the energy of received photon when it was emitted in the distant galaxy.

279 E_0 is the corresponding energy of photon in the current laboratory methods.

280 λ_t is the wavelength of emitted and received photon when it was emitted in the distant galaxy.

281 T_t is the cosmic temperature at the time when the photon was emitted and T_0 is the current cosmic temperature.

282 In subsection 2.5 an attempt is made to understand the cosmological thermodynamic light emission mechanism in hydrogen
283 atom in a unified approach.

284 **2.4 To reinterpret the Hubble's Law**

285 Based on the assumptions it is possible to say that, during cosmic evolution, at any time, any galaxy will have revolution
286 speed as well as receding speed simultaneously and both can be expressed in the following way.
287

288
$$(V_G)_{revolution} \cong \left(\frac{r}{R_t}\right) c \cong rH_t \quad \text{where } r \leq \left(R_t \cong \frac{c}{H_t}\right) \quad (24)$$

289

290 r is the distance between galaxy and the cosmic center and R_t is the cosmic radius at time t .

291

292
$$(V_G)_{receding} \cong \left(\frac{r}{R_t}\right) g_t \cong \left(\frac{r}{R_t}\right) \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} c \quad (25)$$
$$\cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} rH_t \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} (V_G)_{revolution}$$

293

294
$$\frac{(V_G)_{revolution}}{(V_G)_{receding}} \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^2 \quad (26)$$

295

296 Please note that both the relations are independent of the observed redshift. This is for further study.
297

298 **2.5. To understand the cosmological thermodynamic light emission mechanism**

299

300 It is very tempting to make an analogy between the status of the cosmological 'Standard Model' and that of particle physics
301 [53]. In cosmology there are about 10 free parameters, each of which is becoming well determined, and with a great deal of
302 consistency between different measurements. However, none of these parameters can be calculated from a fundamental
303 theory, and so hints of the bigger picture, 'physics beyond the Standard Model,' are being searched for with ever more
304 ambitious experiments. Despite this analogy, there are some basic differences. For one thing, many of the cosmological
305 parameters change with cosmic epoch, and so the measured values are simply the ones determined today, and hence they
306 are not 'constants,' like particle masses for example (although they are deterministic, so that if one knows their values at
307 one epoch, they can be calculated at another). Moreover, the parameter set is not as well defined as it is in the particle
308 physics Standard Model; different researchers will not necessarily agree on which parameters should be considered as free,

Requesting your kind and valuable (unbiased) review comments...

309 and the set can be extended as the quality of the data improves. In a more general sense, the cosmological ‘Standard Model’
 310 is much further from the underlying ‘fundamental theory,’ which will ultimately provide the values of the parameters from
 311 first principles. Nevertheless, any genuinely complete ‘theory of everything’ must include an explanation for the values of
 312 these cosmological parameters as well as the parameters of the Standard Model of particle physics. Current magnitude of
 313 Hubble constant [53-57] is (67.80 ± 0.77) km/sec/Mpc, (68.1 ± 1.2) km/sec/Mpc,
 314 (67.3 ± 1.2) km/sec/Mpc, (69.7 ± 2.0) km/sec/Mpc, (70.0 ± 2.2) km/sec/Mpc, (70.6 ± 3.3) km/sec/Mpc,
 315 (73.8 ± 2.4) km/sec/Mpc, and (72.5 ± 2.5) km/sec/Mpc. In a cosmological approach with various trial-error methods, at
 316 present in hydrogen atom, if $H_0 \cong 71$ km/sec/Mpc, Bohr radius [58] can be fitted as follows.
 317

$$\begin{aligned}
 (a_B)_0 &\cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{GM_0}{c^2} \right) \cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{2H_0} \right) \\
 &\cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{2H_0} \right) \cong \frac{1}{2} \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{H_0} \right) \\
 &\cong 5.27225 \times 10^{-11} \text{ m.}
 \end{aligned} \tag{27}$$

319 $\left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right)$ is the electromagnetic and gravitational force ratio of proton. This relation seems to be very simple and needs
 320 no further derivation. Factor 2 may be connected with half of the current Hubble length $\left(\frac{1}{2} \frac{c}{H_0} \right)$. For any physicist or
 321 cosmologist it will be a very big surprise. Note that, this relation is free from the famous reduced Planck’s constant,
 322 electron rest mass and other arbitrary numbers or coefficients. After simplification and considering the ground state, it is
 323 possible to express the ground state potential energy of electron in the following way.
 324

$$\begin{aligned}
 (E_{\text{pot}})_0 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{4\pi\epsilon_0 G M_0} \right) \cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{1}{2} \frac{c}{H_0} \right)^{-1} \\
 &\cong -2 \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned} \tag{28}$$

326 Here $\left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)$ can be called as the current Hubble potential. Characteristic ground state kinetic energy of electron can be
 327 expressed in the following way.

$$\begin{aligned}
 (E_{\text{kin}})_0 &\cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{8\pi\epsilon_0 G M_0} \right) \cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{c^2}{2GM_0} \right) \\
 &\cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned} \tag{29}$$

329 Characteristic ground state total energy of electron can be expressed in the following way.
 330

$$\begin{aligned}
 (E_{\text{tot}})_0 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{8\pi\epsilon_0 G M_0} \right) \cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{c^2}{2GM_0} \right) \\
 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned} \tag{30}$$

332 If $H_0 \cong 71$ km/sec/Mpc, $(E_{\text{tot}})_0 \cong -13.66$ eV. Based on this coincidence, this proposed new concept can be given some
 333 consideration and it can be suggested that the best value of H_0 lies in between 70 and 71 km/sec/Mpc. Unfortunately
 334 these relations seem to be independent of the reduced Planck’s constant [59,60]. If one is willing to linkup these relations
 335 with the observed ‘discrete’ energy spectrum of the hydrogen atom, then the desired cosmological light emission
 336 mechanism can be developed in a unified picture. Considering the concept of stationary orbits and jumping nature of
 337

Requesting your kind and valuable (unbiased) review comments...

338 electron, emitted photon energy can be expressed in the following way.

339
$$(E_{\text{photon}})_0 \cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \quad (31)$$

340 where $n_1 = n_2 \cong 1, 2, 3, \dots$ and $n_2 > n_1$. The best fit of H_0 can be obtained in the following way.

341
$$\left. \begin{aligned} & \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \cong \frac{e^4 m_e}{32\pi^2 \epsilon_0^2 \hbar^2} \\ & \text{and } H_0 \cong \frac{G m_p^2 m_e c}{2\hbar^2} \cong 70.738 \text{ km/sec/Mpc} \end{aligned} \right\} \quad (32)$$

343 At any time in the past - in support of the proposed cosmological red shift interpretation, above relations can be re-
344 expressed as follows.
345

346
$$(E_{\text{pot}})_t \cong - \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{4\pi\epsilon_0 G M_0} \right) \cong -2 \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (33)$$

347
$$(E_{\text{kin}})_t \cong \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (34)$$

348
$$(E_{\text{tot}})_t \cong - \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (35)$$

351 This can be considered as the base for the ‘cosmological thermodynamic light emission mechanism’. At any time in the
352 past, at any galaxy, emitted photon energy can be expressed as follows.

353
$$(E_{\text{photon}})_t \cong \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]_t \quad (36)$$

354 This issue is for further study. In a unified picture, with reference to the current cosmic temperature, electron’s current
355 quantum of angular momentum can be expressed as follows.

356
$$\hbar \cong m_p \sqrt{\frac{G m_e c}{2 H_0}} \cong \frac{G m_p \sqrt{m_e M_0}}{c} \cong \hbar_0 \quad (37)$$

357 If atomic nuclear mass increases in integral multiples of the proton mass, then the observed discreteness of the reduced
358 Planck’s constant can be expressed as follows.
359

360
$$n\hbar \cong \frac{G(n.m_p) \sqrt{m_e M_0}}{c} \cong n\hbar_0 \quad (38)$$

361 where $n = 1, 2, 3, \dots$. This issue is also for further study. At any time in the past, hypothetically, in terms of the current and
362 past ‘primordial’ cosmic temperatures, it is possible to express the cosmological ‘variable quantum of angular momentum’
363 of electron in the following way. Whether it is virtual or real or speculative - to be confirmed from further study.

364
$$\hbar_t \cong \sqrt{\frac{T_t}{T_0}} \cdot \hbar_0 \cong \sqrt{\frac{\lambda_t}{\lambda_0}} \cdot \hbar_0 \quad (39)$$

365 It may be noted that, throughout the cosmic evolution, Planck’s constant and the Uncertainty constant both can be conside-
366 red as ‘constants’. Now the fundamental questions to be answered are –
367

Requesting your kind and valuable (unbiased) review comments...

- 368 1) Is reduced Planck's constant – an output of the atomic system?
369 2) Is the reduced Planck's constant – a cosmological variable?
370 3) Is the Planck's constant – a cosmological constant?
371 4) How to understand and how to consider the constancy of the Planck's constant along with the variable reduced
372 Planck's constant?
373 5) Is the condition, $\hbar \rightarrow (h/2\pi)$ an indication of saturation or halt of cosmological expansion?

374 **3. Connecting Cosmic Thermal and Physical Parameters** 375

376 **3.1 Cosmic Thermal energy density and Matter energy density**

377 It may be noted that connecting CMBR energy density with Hubble's constant is really a very big task and mostly preferred
378 in cosmology. At any given cosmic time, thermal energy density can be expressed with the following semi empirical
379 relation.

$$380 \quad aT_i^4 \cong \left[1 + \ln \left(\frac{M_i}{M_s} \right) \right]^{-2} \left(\frac{3H_i^2 c^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{H_s}{H_i} \right) \right]^{-2} \left(\frac{3H_i^2 c^2}{8\pi G} \right) \quad (40)$$

$$381 \quad T_i \cong \left[1 + \ln \left(\frac{H_s}{H_i} \right) \right]^{-\frac{1}{2}} \left(\frac{3H_i^2 c^2}{8\pi G a} \right)^{\frac{1}{4}} \quad (41)$$

382 With a suitable derivation if above expression is obtained, then certainly the subject of black hole cosmology is put into
383 main stream physics. Thus at present, if H_0 is close to 71 km/sec/Mpc, obtained CMBR temperature is 2.723 K [53-57].
384 For the time being this can be considered as a remarkable discovery and an accurate fit.
385

$$386 \quad aT_0^4 \cong \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-2} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{M_0}{M_s} \right) \right]^{-2} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \quad (42)$$

$$387 \quad T_0 \cong \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-\frac{1}{2}} \left(\frac{3H_0^2 c^2}{8\pi G a} \right)^{\frac{1}{4}} \quad (43)$$

388 With reference to the current cosmic temperature, at any time in the past,

$$389 \quad \frac{T_i}{T_0} \cong \frac{\left[\frac{1 + \ln \left(\frac{H_s}{H_0} \right)}{1 + \ln \left(\frac{H_s}{H_i} \right)} \right]^{\frac{1}{2}} H_i}{H_0} \quad (44)$$

390 Using this relation, cosmic redshift data can be fitted. When the assumed CMBR temperature is 2999 K, estimated redshift
391 is 1099 and is in very good agreement with the standard model of cosmology.
392

393 Mostly at the ending stage of expansion, rate of change in H_i will be practically zero and can be considered as practically
394 constant. Thus at its ending stage of expansion, for the whole cosmic black hole as H_i practically remains constant, its
395 corresponding thermal energy density will be 'the same' throughout its volume. This 'sameness' may be the reason for the
396 observed 'isotropic' nature of the current CMB radiation. With this coincidence it can be suggested that, at the beginning of
397 cosmic evolution,
398

$$399 \quad aT_s^4 \cong \left(\frac{3H_s^2 c^2}{8\pi G} \right) \quad (45)$$

400 Matter-energy density can be considered as the geometric mean density of volume energy density and the thermal energy

Requesting your kind and valuable (unbiased) review comments...

401 density and it can be expressed with the following semi empirical relation.

$$402 \quad (\rho_m)_t c^2 \cong \sqrt{\left(\frac{3H_t^2 c^2}{8\pi G}\right)} (aT_t^4) \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-1} \left(\frac{3H_t^2 c^2}{8\pi G}\right) \cong \left[1 + \ln\left(\frac{M_t}{M_s}\right)\right]^{-1} \left(\frac{3H_0^2 c^2}{8\pi G}\right) \quad (46)$$

403 Here one important observation to be noted is that, at any time

$$404 \quad \frac{3H_t^2}{8\pi G (\rho_m)_t} \cong \left[1 + \ln\left(\frac{M_t}{M_s}\right)\right] \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right] \quad (47)$$

405 Thus at present,

$$406 \quad (\rho_m)_0 \cong \frac{1}{c^2} \sqrt{\left(\frac{3H_0^2 c^2}{8\pi G}\right)} (aT_0^4) \cong \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-1} \left(\frac{3H_0^2}{8\pi G}\right) \cong \left[1 + \ln\left(\frac{M_0}{M_s}\right)\right]^{-1} \left(\frac{3H_0^2}{8\pi G}\right) \quad (48)$$

$$\cong 6.6 \times 10^{-32} \text{ gram / cm}^3$$

407 Based on the average mass-to-light ratio for any galaxy present matter density can be expressed with the following relation
408 [61].

$$410 \quad (\rho_m)_0 \cong 1.5 \times 10^{-32} \eta h_0 \text{ gram/cm}^3 \quad (49)$$

411 Here $\eta \cong \left\langle \frac{M}{L} \right\rangle_{\text{galaxy}} / \left\langle \frac{M}{L} \right\rangle_{\text{sun}}$, $h_0 \cong H_0 / 100 \text{ Km/sec/Mpc} \cong 0.71$ Note that elliptical galaxies probably comprise about

412 60% of the galaxies in the universe and spiral galaxies thought to make up about 20% percent of the galaxies in the
413 universe. Almost 80% of the galaxies are in the form of elliptical and spiral galaxies. For spiral galaxies, $\eta h_0^{-1} \cong 9 \pm 1$ and
414 for elliptical galaxies, $\eta h_0^{-1} \cong 10 \pm 2$ For our galaxy inner part, $\eta h_0^{-1} \cong 6 \pm 2$. Thus the average ηh_0^{-1} is very close to 8 to 9
415 and its corresponding matter density is close to $(6.0 \text{ to } 6.7) \times 10^{-32} \text{ gram/cm}^3$ and can be compared with the above proposed
416 magnitude of $6.6 \times 10^{-32} \text{ gram/cm}^3$.

418 3.2 Age of the Growing Cosmic black hole

419 Age of the growing cosmic black hole can be assumed as the time taken to grow from the assumed Stoney scale to the
420 current scale. At present,

$$422 \quad g_0 \cong \left(\frac{8\pi G a T_0^4}{3H_0^2 c^2}\right) c \cong \left[1 + \ln\left(\frac{M_0}{M_s}\right)\right]^{-2} c \cong \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2} c \cong 14.66 \text{ km/sec} \quad (50)$$

423 Clearly speaking, at present, Hubble volume is growing at 14.66 km/sec in a decelerating trend. Starting from the Stoney
424 scale, if the assumed growth rate is gradually decreasing, at any time average growth rate can be expressed as follows.

$$427 \quad \frac{g_s + g_t}{2} \cong \frac{1}{2} \left\{1 + \left[1 + \ln\left(\frac{M_t}{M_s}\right)\right]^{-2}\right\} c \cong \frac{1}{2} \left\{1 + \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2}\right\} c \quad (51)$$

428 For the current scale, average growth rate can be expressed as follows.

$$430 \quad \frac{g_s + g_0}{2} \cong \frac{1}{2} \left\{1 + \left[1 + \ln\left(\frac{M_0}{M_s}\right)\right]^{-2}\right\} c \cong \frac{1}{2} \left\{1 + \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2}\right\} c \quad (52)$$

431 Time taken to reach from the Stoney scale to any assumed scale can be expressed as follows.

$$434 \quad \left(\frac{g_s + g_t}{2}\right) t \cong (R_t - R_s) \cong R_t \quad (53)$$

435 where, $R_t \gg R_s$ and $R_s \approx 0$. Hence for the current scale,

436

Requesting your kind and valuable (unbiased) review comments...

437
$$\left(\frac{g_s + g_0}{2}\right) t_0 \cong (R_0 - R_s) \cong R_0 \cong \frac{c}{H_0} \quad (54)$$

438
$$t_0 \cong \left(\frac{g_s + g_0}{2}\right)^{-1} \frac{c}{H_0} \cong \left\{1 + \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2}\right\}^{-1} \frac{2}{H_0} \cong 27.496 \text{ Gyr.} \quad (55)$$

439 where $\left\{1 + \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2}\right\}^{-1} \cong 0.99995$. This proposal is for further study. Based on this proposal, after one second from

440 the Stoney scale, cosmic angular velocity is 2 rad/sec, growth rate is 29 km/sec and cosmic temperature is 3×10^9 K.

441
442 With reference to the current and past cosmic temperatures, at any time in the past, at any galaxy, for any hydrogen atom,
443

444
$$\frac{E_0}{E_t} \cong \frac{\lambda_t}{\lambda_0} \cong \frac{T_t}{T_0} \cong \frac{\left[\frac{\left[1 + \ln\left(\frac{H_s}{H_0}\right)\right] H_t}{\left[1 + \ln\left(\frac{H_s}{H_t}\right)\right] H_0}\right]^{\frac{1}{2}}}{\left[\frac{\left[1 + \ln\left(\frac{R_0}{R_s}\right)\right] R_0}{\left[1 + \ln\left(\frac{R_t}{R_s}\right)\right] R_t}\right]^{\frac{1}{2}}} \quad (56)$$

445 By guessing H_t , $(z_0 + 1)$ can be estimated. It seems to be a full and absolute definition for the cosmic redshift. Thus at any
446 time in the past,

447
$$\left(\frac{E_0}{E_t} - 1\right) \cong \left(\frac{\lambda_t}{\lambda_0} - 1\right) \cong \left(\frac{T_t}{T_0} - 1\right) \cong \frac{\left[\frac{\left[1 + \ln\left(\frac{H_s}{H_0}\right)\right] H_t}{\left[1 + \ln\left(\frac{H_s}{H_t}\right)\right] H_0}\right]^{\frac{1}{2}}}{\left[\frac{\left[1 + \ln\left(\frac{R_0}{R_s}\right)\right] R_0}{\left[1 + \ln\left(\frac{R_t}{R_s}\right)\right] R_t}\right]^{\frac{1}{2}}} - 1 \cong z_0 \quad (57)$$

448 Please see the following table-1 for the cosmic physical and thermal parameters. This table prepared with C++ program
449 with reference to the observed 2.725 K. In this table:

- 450
451 Column-1 = Assumed cosmic angular velocity.
452 Column-2 = Estimated cosmic radius, from relation (7).
453 Column-3 = Estimated cosmic mass, from relation (7).
454 Column-4 = Estimated cosmic growth index, from relation (10).
455 Column-5 = Estimated cosmic growth rate, from relation (12).
456 Column-6 = Estimated cosmic time, from relation (53).
457 Column-7 = Estimated cosmic temperature, from relation (41)
458 Column-8 = Estimated cosmic redshift, from relation (57)

Table-1: Assumed Cosmic angular velocity and estimated other cosmic physical and thermal parameters

Assumed Cosmic Angular velocity	Estimated Cosmic radius	Estimated Cosmic mass	Cosmic Growth index $\cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^2$	Estimated Cosmic Growth rate	Estimated Cosmic time	Estimated Cosmic temperature	Estimated Cosmic Redshift z_0
(rad/sec)	(meter)	(kg)	(number)	(km/sec)	(sec)	(K)	(number)
1.086E+44	2.761E-36	1.859E-09	1	299792	0.000E+00	2.237E+32	8.207E+31
2.305E+43	1.301E-35	8.759E-09	6.50173	46109.6	5.924E-44	6.455E+31	2.368E+31
2.305E+42	1.301E-34	8.759E-08	23.5461	12732.1	8.148E-43	1.480E+31	5.428E+30
2.305E+41	1.301E-33	8.759E-07	51.1943	5855.97	8.493E-42	3.853E+30	1.414E+30
2.305E+40	1.301E-32	8.759E-06	89.4463	3351.65	8.580E-41	1.060E+30	3.888E+29

Requesting your kind and valuable (unbiased) review comments...

2.305E+39	1.301E-31	8.759E-05	138.302	2167.66	8.615E-40	3.006E+29	1.103E+29
2.305E+38	1.301E-30	8.759E-04	197.762	1515.93	8.634E-39	8.692E+28	3.189E+28
2.305E+37	1.301E-29	8.759E-03	267.825	1119.36	8.645E-38	2.548E+28	9.347E+27
2.305E+36	1.301E-28	8.759E-02	348.492	860.256	8.653E-37	7.544E+27	2.768E+27
2.305E+35	1.301E-27	8.759E-01	439.763	681.714	8.658E-36	2.251E+27	8.258E+26
2.305E+34	1.301E-26	8.759E+00	541.638	553.492	8.662E-35	6.756E+26	2.479E+26
2.305E+33	1.301E-25	8.759E+01	654.116	458.317	8.665E-34	2.038E+26	7.477E+25
2.305E+32	1.301E-24	8.759E+02	777.199	385.735	8.667E-33	6.173E+25	2.265E+25
2.305E+31	1.301E-23	8.759E+03	910.885	329.122	8.668E-32	1.876E+25	6.883E+24
2.305E+30	1.301E-22	8.759E+04	1055.17	284.116	8.670E-31	5.719E+24	2.098E+24
2.305E+29	1.301E-21	8.759E+05	1210.07	247.748	8.671E-30	1.748E+24	6.411E+23
2.305E+28	1.301E-20	8.759E+06	1375.57	217.941	8.671E-29	5.352E+23	1.964E+23
2.305E+27	1.301E-19	8.759E+07	1551.67	193.207	8.672E-28	1.642E+23	6.025E+22
2.305E+26	1.301E-18	8.759E+08	1738.37	172.456	8.673E-27	5.048E+22	1.852E+22
2.305E+25	1.301E-17	8.759E+09	1935.68	154.877	8.673E-26	1.554E+22	5.701E+21
2.305E+24	1.301E-16	8.759E+10	2143.59	139.855	8.674E-25	4.790E+21	1.757E+21
2.305E+23	1.301E-15	8.759E+11	2362.11	126.917	8.674E-24	1.478E+21	5.424E+20
2.305E+22	1.301E-14	8.759E+12	2591.23	115.695	8.674E-23	4.568E+20	1.676E+20
2.305E+21	1.301E-13	8.759E+13	2830.96	105.898	8.675E-22	1.413E+20	5.184E+19
2.305E+20	1.301E-12	8.759E+14	3081.28	97.2947	8.675E-21	4.375E+19	1.605E+19
2.305E+19	1.301E-11	8.759E+15	3342.21	89.6987	8.675E-20	1.356E+19	4.973E+18
2.305E+18	1.301E-10	8.759E+16	3613.75	82.9588	8.675E-19	4.204E+18	1.542E+18
2.305E+17	1.301E-09	8.759E+17	3895.89	76.951	8.676E-18	1.305E+18	4.786E+17
2.305E+16	1.301E-08	8.759E+18	4188.63	71.5729	8.676E-17	4.052E+17	1.486E+17
2.305E+15	1.301E-07	8.759E+19	4491.98	66.7395	8.676E-16	1.259E+17	4.619E+16
2.305E+14	1.301E-06	8.759E+20	4805.93	62.3797	8.676E-15	3.915E+16	1.436E+16
2.305E+13	1.301E-05	8.759E+21	5130.48	58.4336	8.676E-14	1.218E+16	4.468E+15
2.305E+12	1.301E-04	8.759E+22	5465.64	54.8504	8.676E-13	3.791E+15	1.391E+15
2.305E+11	1.301E-03	8.759E+23	5811.41	51.5869	8.676E-12	1.180E+15	4.331E+14
2.305E+10	1.301E-02	8.759E+24	6167.77	48.6063	8.676E-11	3.678E+14	1.349E+14
2.305E+09	1.301E-01	8.759E+25	6534.74	45.8767	8.676E-10	1.146E+14	4.206E+13
2.305E+08	1.301E+00	8.759E+26	6912.31	43.3708	8.677E-09	3.575E+13	1.311E+13
2.305E+07	1.301E+01	8.759E+27	7300.49	41.0647	8.677E-08	1.115E+13	4.091E+12
2.305E+06	1.301E+02	8.759E+28	7699.27	38.9378	8.677E-07	3.480E+12	1.277E+12
2.305E+05	1.301E+03	8.759E+29	8108.66	36.9719	8.677E-06	1.086E+12	3.985E+11
2.305E+04	1.301E+04	8.759E+30	8528.65	35.1512	8.677E-05	3.392E+11	1.244E+11
2.305E+03	1.301E+05	8.759E+31	8959.24	33.4618	8.677E-04	1.059E+11	3.887E+10
2.305E+02	1.301E+06	8.759E+32	9400.43	31.8913	8.677E-03	3.310E+10	1.214E+10
2.305E+01	1.301E+07	8.759E+33	9852.23	30.4289	8.677E-02	1.035E+10	3.796E+09
2.305E+00	1.301E+08	8.759E+34	10314.6	29.0648	8.677E-01	3.234E+09	1.187E+09
2.305E-01	1.301E+09	8.759E+35	10787.6	27.7904	8.677E+00	1.011E+09	3.710E+08
2.305E-02	1.301E+10	8.759E+36	11271.3	26.598	8.677E+01	3.163E+08	1.161E+08
2.305E-03	1.301E+11	8.759E+37	11765.5	25.4807	8.677E+02	9.897E+07	3.631E+07

Requesting your kind and valuable (unbiased) review comments...


2.305E-04	1.301E+12	8.759E+38	12270.3	24.4324	8.677E+03	3.097E+07	1.136E+07
2.305E-05	1.301E+13	8.759E+39	12785.7	23.4475	8.677E+04	9.693E+06	3.556E+06
2.305E-06	1.301E+14	8.759E+40	13311.7	22.5209	8.677E+05	3.034E+06	1.113E+06
2.305E-07	1.301E+15	8.759E+41	13848.4	21.6482	8.677E+06	9.501E+05	3.486E+05
2.305E-08	1.301E+16	8.759E+42	14395.6	20.8253	8.677E+07	2.976E+05	1.092E+05
2.305E-09	1.301E+17	8.759E+43	14953.4	20.0484	8.677E+08	9.321E+04	3.419E+04
2.305E-10	1.301E+18	8.759E+44	15521.9	19.3142	8.677E+09	2.920E+04	1.071E+04
2.305E-11	1.301E+19	8.759E+45	16100.9	18.6196	8.677E+10	9.150E+03	3.356E+03
2.52E-12	1.19E+20	8.01E+46	16667.6	17.9865	7.94E+11	2998.85	1099.21
2.305E-12	1.301E+20	8.759E+46	16690.6	17.9618	8.677E+11	2.868E+03	1.051E+03
2.305E-13	1.301E+21	8.759E+47	17290.8	17.3382	8.677E+12	8.988E+02	3.288E+02
2.305E-14	1.301E+22	8.759E+48	17901.7	16.7466	8.677E+13	2.818E+02	1.024E+02
2.305E-15	1.301E+23	8.759E+49	18523.2	16.1847	8.677E+14	8.835E+01	3.141E+01
2.305E-16	1.301E+24	8.759E+50	19155.2	15.6507	8.677E+15	2.771E+01	9.164E+00
2.305E-17	1.301E+25	8.759E+51	19797.9	15.1427	8.677E+16	8.689E+00	2.188E+00
2.305E-18	1.301E+26	8.759E+52	20451.2	14.6589	8.677E+17	2.726E+00	0.000E+00

462
463
464

Please see the below graph for the cosmic growth index for ~ 61 values starting from 1 to 20451.2 of Column-4 in table-1.

465

Cosmic Growth Index



466
467

3.3. Direct fitting of the two current CMBR wavelengths

468
469
470
471
472
473
474

Note that the spectrum from Planck's law of black body radiation takes a different shape in the frequency domain from that of the wavelength domain, the frequency location of the peak emission does not correspond to the peak wavelength using the simple relationship between frequency, wavelength, and the speed of light. In other words, the peak wavelength and the peak frequency do not correspond. The frequency form of Wien's displacement law is derived using similar methods, but starting with Planck's law in terms of frequency instead of wavelength. The effective result is to substitute 3 for 5 in the

Requesting your kind and valuable (unbiased) review comments...

475 equation for the peak wavelength. Thus it is possible to say that [62],
476

$$477 \quad \sqrt{\frac{c}{\lambda_m f_m}} \cong \sqrt{1.75978} \cong 1.326567 \cong \frac{4}{3} \quad (58)$$

478 where λ_m and f_m are the peak wavelength in wavelength domain and peak frequency in frequency domain respectively.

479 Let λ_f is the wavelength corresponding to $\frac{dE_\nu}{d\nu}$ and E_ν is the total energy at all frequencies up to and including ν , at any
480 given cosmic time. λ_m is the wavelength corresponding to $\frac{dE_\lambda}{d\lambda}$ and E_λ is the total energy at all wavelengths up to and
481 including λ . Considering the observed CMBR wavelengths, it is possible to express both the wavelengths in the following
482 way.

$$483 \quad [(\lambda_m)_t \text{ and } (\lambda_f)_t] \propto \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \quad (59)$$

$$484 \quad [(\lambda_m)_t \text{ and } (\lambda_f)_t] \propto \sqrt{\left(\frac{4\pi G M_t}{c^2}\right) \cdot \left(\frac{4\pi G M_S}{c^2}\right)} \quad (60)$$

486 Guessing in this way it is noticed that,
487
488

$$489 \quad (\lambda_f)_t \cong \left(\frac{4}{3}\right) \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2} \quad (61)$$

$$\cong \left(\frac{4}{3}\right) \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2}$$

$$490 \quad (\lambda_m)_t \cong \left(\frac{3}{4}\right) \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2} \quad (62)$$

$$\cong \left(\frac{3}{4}\right) \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2}$$

491 Thus it is possible to express both the wavelength relations in the following way.
492

$$493 \quad (\lambda_f, \lambda_m)_t \cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2} \quad (63)$$

$$\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{H_S}{H_t}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_t}} \cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{2\pi c}{\sqrt{H_S H_t}}$$

494 Alternatively geometric mean of $(\lambda_f, \lambda_m)_t$ can be expressed as follows.
495
496

$$\sqrt{(\lambda_m)_t (\lambda_f)_t} \cong \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2} \quad (64)$$

$$\cong \sqrt{1 + \ln\left(\frac{H_S}{H_t}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_t}} \cong \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{2\pi c}{\sqrt{H_S H_t}}$$

Requesting your kind and valuable (unbiased) review comments...

497 At present, if H_0 is close to 71 km/sec/Mpc,
498

$$\begin{aligned}
 (\lambda_f, \lambda_m)_0 &\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{M_0}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_0 M_S}}{c^2} \\
 &\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{H_S}{H_0}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_0}} \cong (1.90 \text{ mm}, 1.069 \text{ mm})
 \end{aligned}
 \tag{65}$$

500 With reference to $(\lambda_m)_t$ and Wien's displacement constant, from above relations $k_B T_t$ can be expressed as follows.
501

$$\begin{aligned}
 T_t &\cong \frac{2.898 \times 10^{-3}}{(\lambda_m)_t} \cong \left(\frac{hc}{4.965114 k_B}\right) \left(\frac{1}{(\lambda_m)_t}\right) \text{ and} \\
 k_B T_t &\cong \left(\frac{4}{3x}\right) \sqrt{\left(1 + \ln\left(\frac{M_t}{M_S}\right)\right)^{-1} \left(\frac{M_t}{M_S}\right)} \cdot \left(\frac{hc^3}{4\pi G M_t}\right)
 \end{aligned}
 \tag{66}$$

503 where $x \cong 4.965114$.

$$k_B T_t \propto \left(\frac{hc^3}{4\pi G M_t}\right) \cong \frac{h H_t}{2\pi} \cong h \left(\frac{H_t}{2\pi}\right)
 \tag{67}$$

505 This relation may not be identical but similar to the famous Hawking's black hole temperature formula [63].
506

$$k_B T_t \propto \sqrt{\left(1 + \ln\left(\frac{M_t}{M_S}\right)\right)^{-1} \left(\frac{M_t}{M_S}\right)}
 \tag{68}$$

508 In this way in a very simple approach observed CMBR and the proposed Black hole universe concepts can be put into
509 single frame of reference. Here the very interesting and strange observation is that, at present
510

$$\left(1 + \ln\left(\frac{M_0}{M_S}\right)\right)^{-1} \left(\frac{M_0}{M_S}\right) \cong \exp\left(\frac{1}{\alpha}\right)
 \tag{69}$$

512 where $\left(\frac{1}{\alpha}\right)$ is the inverse of the fine structure ratio. For any mathematician this seems be a fun. For a cosmologist it
513 may be an accidental coincidence. For any physicist it is an astounding and exciting coincidence. Even though it depends
514 upon one's own choice of scientific interest, from unification point of view, assuming it to be a cosmological variable it is
515 possible to express $\left(\frac{1}{\alpha}\right)$ in the following way.

$$\left(\frac{1}{\alpha}\right)_0 \cong \ln \left[\left(1 + \ln\left(\frac{M_0}{M_S}\right)\right)^{-1} \left(\frac{M_0}{M_S}\right) \right] \cong 137.047
 \tag{70}$$

517 Here $\left(\frac{1}{\alpha}\right)_0$ may be considered as the current magnitude of 'inverse of the fine structure ratio. In atomic and nuclear physics,
518 the fine-structure ratio (α) is a fundamental physical constant namely the coupling constant characterizing the strength
519 [64-66] of the electromagnetic interaction. Being a dimensionless quantity, it has a constant numerical value in all systems
520 of units. Note that, from unification point of view, till today role of dark energy or dark matter is unclear and undecided.
521 Their laboratory or physical existence is also not yet confirmed. In this critical situation this application or coincidence
522 can be considered as a key tool in particle cosmology. Based on the above heuristic observation and for the assumed initial
523 conditions of the universe, if $M_t \rightarrow M_S$, $\left(\frac{1}{\alpha}\right)_S \rightarrow 0$. Based on the relation (70), if one is willing to consider the cos-

Requesting your kind and valuable (unbiased) review comments...

524 mological variable nature of $\left(\frac{1}{\alpha}\right)$, relation (66) can be expressed as follows.

525

526
$$T_t \cong \sqrt{\left(\frac{1}{e^\alpha}\right)_t} \cdot \left(\frac{bc^2}{3\pi GM_t}\right) \quad (71)$$

527 At the beginning of cosmic evolution for the Stoney scale,

528
$$T_s \cong \left(\frac{bc^2}{3\pi GM_s}\right) \quad (72)$$

529 From now onwards, CMBR temperature can be called as '**Comic Black Hole's Thermal Radiation**' temperature and can
530 be expressed as '**CBHTR**' temperature. From ground based laboratory experiments, it is possible to measure the rate of

531 change in $\frac{d}{dt}\left(\frac{1}{\alpha_t}\right)$. Hence the absolute cosmic rate of expansion can be measured. Thus at any time based on

532 $\left[\frac{d}{dt}[(\lambda_m)_t \text{ and } (\lambda_f)_t], \frac{d}{dt}(T_t) \text{ and } \frac{d}{dt}(H_t)\right]$, the absolute cosmic rate of expansion can be confirmed. At present with

533 reference to $\left[\frac{d}{dt}[(\lambda_m)_0 \text{ and } (\lambda_f)_0], \frac{d}{dt}(T_0) \text{ and } \frac{d}{dt}(H_0)\right]$ current 'true' cosmic rate of expansion can be understood.

534 Drop in current 'cosmic temperature' can be considered as a measure of the current cosmic expansion and 'rate of decrease
535 in current cosmic temperature' can be considered as a measure of the current cosmic 'rate of expansion'. But if rate of
536 decrease in temperature is very small and is beyond the scope of current experimental verification, then the two possible
537 states are: a) cosmic temperature is decreasing at a very slow rate and universe is expanding at a very slow rate and b) there
538 is no 'observable' thermal expansion and there is no 'observable' cosmic expansion. If observed CMBR temperature is
539 2.725 K and is very low in magnitude and is very close to absolute zero, then thinking about and confirming the 'cosmic
540 acceleration' may not be reasonable. Similarly 'rate of decrease in current 'Hubble's constant' can be considered as a
541 measure of current cosmic 'rate of expansion'. If rate of decrease in current 'Hubble's constant is very small and is beyond
542 the scope of current experimental verification, then the two possible states are: a) current 'Hubble's constant is decreasing
543 at a very slow rate and current universe is expanding at a very slow rate and b) at present there is no 'observable' cosmic
544 expansion. Fortunately as per the Cobe/Planck satellite data current CMBR temperature is very smooth and isotropic, and
545 there is no data that refers to the rate of change in the current Hubble's constant. Hence it can be suggested that at present
546 there is no significant cosmic expansion. Even though this suggestion is completely against to the current notion of cosmic
547 acceleration [32,33], based on the proposed arguments, relations and observed data authors request the science
548 community to review the standard cosmology. If observed CMB radiation temperature is 2.725 K and is very low in
549 magnitude and is very close to absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be
550 reasonable.
551

552 **4. To understand the physical significance of large numbers in cosmology**

553

554 Great cosmologists proposed many interesting large numbers in cosmology [67-74]. Ultimately the essence of any
555 cosmological number or ratio is to connect the microscopic and macroscopic physical constants with a possible physical
556 meaning with in the 'evolving universe'. Clearly speaking large dimensionless constants and compound physical constants
557 must reflect an 'observable' intrinsic property of any natural physical phenomenon. Then only the real meaning of any
558 cosmological number can be explored. In this regard authors proposed many interesting relations in the previous sections of
559 this paper. Authors noticed that uncertainty relation or Planck's constant or reduced Planck's constant or inverse of the Fine
560 structure ratio or characteristic nuclear potential radius or rms radius of proton or classical radius of electron - play a
561 crucial role in the understanding the halt of cosmic expansion. The basic questions to be answered are: 1) The general idea
562 of large number coincidence is interesting, yet is there any observational proves? and 2) How Einstein's general theory of
563 relativity is fitted in the theory of the large cosmological numbers? In this regard the characteristic and key relation can be
564 expressed in the following way.

Requesting your kind and valuable (unbiased) review comments...

565

$$\frac{c^3}{2GM_0} \cong H_0 \quad \text{Or} \quad \frac{c^3}{2GH_0} \cong M_0 \quad (73)$$

566

Here (M_0, H_0) can be considered as the current mass and current angular velocity of the black hole universe respectively.

567

By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.

568

569

$$\frac{c^3}{2GM_{sat}} \cong H_{sat} \quad \text{Or} \quad \frac{c^3}{2GH_{sat}} \cong M_{sat} \quad (74)$$

570

Here (M_{sat}, H_{sat}) can be considered as the saturated mass and saturated angular velocity of the black hole universe at its

571

ending stage of expansion. Fortunately it is noticed that, $M_{sat} \cong M_0$ and $H_{sat} \cong H_0$. Authors strongly believe that the

572

following relations certainly help in understanding the mystery of the halting of the present cosmic expansion.

573

574

4.1 Role of the Uncertainty relation

575

It is noticed that,

576

577

$$\frac{Gm_p m_e}{R_p H_0} \cong \frac{h}{4\pi} \quad (75)$$

579

Here $R_p \cong (0.84184 \text{ to } 0.87680) \text{ fm}$ is the rms radius of proton [75,76]. After re-arranging, it can be expressed in the

580

following way.

581

582

$$\left(\frac{2Gm_p}{c^2 R_p} \right) \frac{m_e c^2}{H_0} \cong \left(\frac{2Gm_p}{c^2 R_p} \right) \left[m_e c \left(\frac{2\pi c}{H_0} \right) \right] \cong h \quad (76)$$

583

By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.

584

585

586

$$H_{sat} \Rightarrow \frac{4\pi Gm_p m_e}{hR_p} \cong \frac{Gm_p m_e}{(h/4\pi)R_p} \quad (77)$$
$$\Rightarrow H_{sat} \cong (67.87 \text{ to } 70.69) \text{ km/sec/Mpc}$$

587

This is a remarkable fit and needs further study.

588

589

4.2 Role of the classical radius of electron

590

It is noticed that,

591

592

593

$$\sqrt{\left(\frac{2G\sqrt{m_p m_e}}{c^2} \right) \left(\frac{c}{H_0} \right)} \cong \sqrt{\left(\frac{2G\sqrt{m_p m_e}}{c^2} \right) \left(\frac{2GM_0}{c^2} \right)} \cong \left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right) \quad (78)$$

594

$\left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right)$ is nothing but the presently believed classical radius of electron. In a broad picture or considering the

595

interaction in between proton and electron it is a very general idea to consider the geometric mean mass of proton and

596

electron. By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as

597

follows.

598

$$\left(\frac{c}{H_{sat}} \right) \Rightarrow \left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right)^2 \left(\frac{c^2}{2G\sqrt{m_p m_e}} \right) \quad (79)$$

Requesting your kind and valuable (unbiased) review comments...

599

$$H_{sat} \Rightarrow \frac{2G\sqrt{m_p m_e}}{c} \left(\frac{4\pi\epsilon_0 m_e c^2}{e^2} \right)^2 \cong 67.533 \text{ km/sec/Mpc} \quad (80)$$

600
601

This is also a remarkable fit and needs further study.

4.3 Role of the characteristic nuclear potential radius

602
603
604

It is noticed that,

605

$$\frac{G\sqrt{M_0\sqrt{m_p m_e}}}{c^2} \cong \sqrt{\left(\frac{GM_0}{c^2}\right)\left(\frac{G\sqrt{m_p m_e}}{c^2}\right)} \cong 1.4 \times 10^{-15} \text{ m} \cong R_n \quad (81)$$

606
607
608
609

R_n is nothing but the presently believed characteristic nuclear potential radius [77] or the nuclear strong interaction range as proposed by Yukawa [78]. By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows [79-81].

610

$$\frac{G\sqrt{M_{sat}\sqrt{m_p m_e}}}{c^2} \Rightarrow R_n \quad (82)$$

611

$$H_{sat} \Rightarrow \frac{G\sqrt{m_p m_e}}{2cR_n^2} \quad (83)$$

612
613
614
615

This is also a remarkable coincidence and accuracy mainly depends upon the magnitude of the characteristic nuclear potential radius. Further study may reveal the mystery.

4.4 Role of the ‘inverse’ of the Fine structure ratio

616
617
618
619

Total thermal energy in the present Hubble volume can be expressed as follows.

620

$$(E_T)_0 \cong aT_0^4 \cdot \frac{4\pi}{3} \left(\frac{c}{H_0} \right)^3 \quad (84)$$

621
622

Thermal energy present in half of the current Hubble volume can be expressed as follows.

623

$$\frac{(E_T)_0}{2} \cong \frac{1}{2} \left[aT_0^4 \cdot \frac{4\pi}{3} \left(\frac{c}{H_0} \right)^3 \right] \quad (85)$$

624
625

If (c/H_0) is the present electromagnetic interaction range, then present characteristic Hubble potential can be expressed as

626

$$(E_e)_0 \cong \frac{e^2}{4\pi\epsilon_0 (c/H_0)} \cong \frac{e^2 H_0}{4\pi\epsilon_0 c} \quad (86)$$

627
628
629

If H_0 is close to 71 km/sec/Mpc and $T_0 \cong 2.725 \text{ K}$, it is noticed that,

630

$$\ln \sqrt{\frac{[(E_T)_0/2]}{(E_e)_0}} \cong 137.05 \quad (87)$$

631
632
633

By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.

Requesting your kind and valuable (unbiased) review comments...

$$\ln \sqrt{\frac{[(E_T)_0/2]}{(E_e)_0}} \cong \ln \sqrt{\frac{[(E_T)_{sat}/2]}{(E_e)_{sat}}} \Rightarrow \left(\frac{1}{\alpha}\right) \quad (88)$$

$(E_T)_{sat}$ can be considered as the total thermal energy in the Hubble volume at the end of cosmic expansion.

$(E_e)_{sat}$ can be considered as the Hubble potential at the end of cosmic expansion.

5. To fit the nuclear charge radius and the Planck's constant

The subject of final unification is having a long history. After the nucleus was discovered [77] in 1908, it was clear that a new force was needed to overcome the electrostatic repulsion of the positively charged protons. Otherwise the nucleus could not exist. Moreover, the force had to be strong enough to squeeze the protons into a volume of size 10^{-15} meter. In general the word 'strong' is used since the strong interaction is the "strongest" of the four fundamental forces. Its observed strength is around 10^2 times that of the electromagnetic force, some 10^5 times as great as that of the weak force, and about 10^{39} times that of gravitation.

The aim of unification is to understand the relation that connects 'gravity', 'mass', 'charge' and the 'microscopic space-time curvature'. Many scientists addressed this problem in different ways [79-81]. The authors also made many attempts in their previously published papers [82-85]. Experimentally observed nuclear charge radius R_{ch} can be fitted with the following strange and simple unified relation.

$$R_{ch} \cong \sqrt{\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{2GM_s}{c^2}\right)} \cong 1.252 \text{ fermi} \quad (89)$$

Considering the rest energy of proton and 1.25 fermi, semi empirical mass formula energy coefficients can be fitted very easily.

$$\frac{R_{ch} c^2}{2GM_s} \cong \sqrt{\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right)} \quad (90)$$

Whether the expression $\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cong 90.62$ playing a 'key unified role' or 'only a fitting role' to be confirmed.

With a great accuracy the famous Planck's constant can be fitted with the following relation.

$$\begin{aligned} h &\cong \frac{1}{2} \ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot (\sqrt{m_p m_e} \cdot c \cdot R_{ch}) \\ &\cong \ln\sqrt{\frac{e^2}{4\pi\epsilon_0 G m_p m_e}} \cdot (\sqrt{m_p m_e} \cdot c \cdot R_{ch}) \cong 6.63862 \times 10^{-34} \text{ J.sec} \end{aligned} \quad (91)$$

Recommended value of h is $6.6260695729 \times 10^{-34}$ J.sec and the error is 0.189%. Now above relation can be simplified into the following form [75].

$$h \cong \left[\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \right]^{3/2} \left(\frac{e^2}{4\pi\epsilon_0 c}\right) \quad (92)$$

Connecting quantum constants and gravity is really a very big task. At this juncture this relation can be given a chance. It casts a doubt on the independent existence of quantum mechanics. With this relation, obtained magnitude of the gravitational constant is, $G \cong 7.48183566 \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{sec}^{-2}$. Independent of 'length', 'force' and other physical considerations, with this relation order of magnitude of G can be confirmed from atomic physical constants. To proceed further - at first the hierarchy of physical constants must be established and it needs further study and analysis.

670 **6. Conclusions**

671 **6.1 Need of the mass unit $M_S \cong \sqrt{e^2/4\pi\epsilon_0 G}$ in unification**

672 The basic idea of unification is – 1) To minimize the number of physical constants and to merge a group of different
673 fundamental constants into one compound physical constant with appropriate unified interpretation and 2) To merge and
674 minimize various branches of physics. In this journey, the first step is to see the numerical coincidences, second step is to
675 interpret the numerical coincidences and the third step is to synchronize the current interpretations and new interpretations.
676 When the new interpretation disagrees with the current interpretation, generally with the help of emerging science and
677 technology, discrepancies can be resolved with future observations, experiments and analysis. The first step in unification
678 is to understand the origin of the rest mass of a charged elementary particle. Second step is to understand the combined
679 effects of its electromagnetic (or charged) and gravitational interactions. Third step is to understand its behavior with
680 surroundings when it is created. Fourth step is to understand its behavior with cosmic space-time or other particles. Right
681 from its birth to death, in all these steps the underlying fact is that whether it is a strongly interacting particle or weakly
682 interacting particle, it is having some rest mass. To understand the first two steps somehow one can implement the
683 gravitational constant in sub atomic physics. In this regard $M_S \cong \sqrt{e^2/4\pi\epsilon_0 G}$ can be considered as the nature's given true
684 unified mass unit [35].
685
686

687 **6.2 To consider the universe as a growing and light speed rotating primordial black hole**

688 If 'black hole geometry' is more intrinsic compared to the black hole 'mass' and 'density' parameters, if universe
689 constitutes so many galaxies and if each galaxy constitutes a central growing and fast spinning black hole then considering
690 universe as a 'growing and light speed rotating primordial black hole' may not be far away from reality. If universe is
691 having no black hole geometry - any massive body (which is bound to the universe) may not show a black hole structure.
692 That is black hole structure or geometry may be a subset of the cosmic geometry. At this juncture considering or rejecting
693 this proposal completely depends on the observed cosmic redshift. Based on the relations proposed in sections 2 and 4
694 observed cosmic redshift can be considered as a result of cosmological light emission mechanism. Authors are working on
695 the assumed Hubble volume and Hubble mass in different directions with different applications [1-13] that connect micro
696 physics and macro physics. Based on the proposed applications and short comings of the standard model of cosmology -
697 concepts of black hole cosmology may be given at least 99% priority.
698

699 **6.3 About the current cosmic black hole's deceleration**

700 In view of the applications proposed in sections (2) to (4) and with reference to the zero rate of change in inverse of the
701 fine structure ratio (from ground based experiments), zero rate of change in the 'current CMBR temperature' (from
702 Cobe/Planck satellite data) and zero rate of change in the 'current Hubble's constant' (from Cobe/Planck satellite data) it
703 can be suggested that, current cosmic expansion is almost all saturated and at present there is no significant cosmic
704 acceleration [47,48]. Clearly speaking, Stoney scale cosmic black hole's growth rate is equal to the speed of light and
705 current cosmic black hole is growing at 14.66 km/sec in a decelerating trend. It can be also be possible to suggest that
706 currently believed 'dark energy' is a pure, 'mathematical concept' and there exists no physical base behind its confirmation.
707 Now the key leftover things are nucleosynthesis and structure formation. Authors are working in this direction. As nuclear
708 binding energy was zero at the beginning of cosmic evolution, by considering the time dependent variable nature of
709 magnitudes of the semi empirical mass formula energy coefficients it is possible to show that, at the beginning of formation
710 of nucleons, nuclear stability is maximum for light atoms only. If so it can be suggested that, from the beginning of
711 formation of nucleons, in any galaxy, maximum scope is being possible only for the survival of light atoms and this may be
712 the reason for the accumulation and abundance of light atoms in large proportion.
713

714 **Acknowledgements**

715 The first author is indebted to professor K. V. Krishna Murthy, Chairman, Institute of Scientific Research on Vedas
716 (I-SERVE), Hyderabad, India and Shri K. V. R. S. Murthy, former scientist IICT (CSIR) Govt. of India, Director, Research
717 and Development, I-SERVE, for their valuable guidance and great support in developing this subject. Both the authors are
718 very much thankful to the anonymous referees for their valuable comments and kind suggestions in improving and bringing
719 this subject into current main stream physics research.
720

References

- 721
722
723 [1] U. V. S. Seshavatharam, S. Lakshminarayana. Basics of black hole cosmology – first critical scientific review.
724 Physical Science International journal, Vol-4, Issue-6, p.842-879. (2014)
725 [2] Seshavatharam, U. V. S. & Lakshminarayana, S., On the Role of Hubble Volume in Black Hole Cosmology & Final
726 Unification. Prespacetime Journal, February 2014, Volume 5, Issue 2, pp. 148-173.
727 [3] U. V. S. Seshavatharam. Physics of rotating and expanding black hole universe. Progress in Physics. April, p 7-14,
728 (2010).
729 [4] U.V.S. Seshavatharam. The Primordial Cosmic Black Hole and the Cosmic Axis of Evil. International Journal of
730 Astronomy, 1(2): 20-37, (2012).
731 [5] Seshavatharam, U. V. S, Lakshminarayana, S. Applications of Hubble Volume in Atomic Physics, Nuclear Physics,
732 Particle Physics, Quantum Physics and Cosmic Physics. Journal of Nuclear Physics, Material Sciences, Radiation and
733 Applications Vol. 1, No. 1, August 2013 pp. 45–60.
734 [6] Seshavatharam U.V. S, Lakshminarayana, S, To confirm the existence of Black hole cosmology. International Journal
735 of Advanced Astronomy, 2 (1), 21-36, 2013
736 [7] U. V. S. Seshavatharam, S. Lakshminarayana, Hubble Volume and the Fundamental Interactions, International Journal
737 of Astronomy, Vol. 1 No. 5, 2012, pp. 87-100.
738 [8] U. V. S. Seshavatharam, S. Lakshminarayana, B.V.S.T. Sai. Is red shift an index of galactic‘atomic light emission’
739 mechanism? International Journal of Physics, Vol. 1, No.3, 49-64, (2013).
740 [9] U. V. S. Seshavatharam, S. Lakshminarayana Microscopic Physical Phenomena in Black Hole Cosmos Rotating at
741 Light Speed. Prespacetime Journal. October 2013, Volume 4, Issue 9, pp. 884-922.
742 [10] U. V. S. Seshavatharam, S. Lakshminarayana. Black Hole Cosmology: A Biological Boom. Journal of Astrobiology
743 and Outreach, 2014, 2:1. <http://dx.doi.org/10.4172/2332-2519.1000108>.
744 [11] U. V. S. Seshavatharam, S. Lakshminarayana. Basic Interactions in Black Hole Cosmology. American Journal of
745 Astronomy and Astrophysics. Vol. 2, No. 1, 2013, pp. 6-17. doi: 10.11648/j.ajaa.20140201.12
746 [12] U. V. S. Seshavatharam, S. Lakshminarayana. Friedman Cosmology: Reconsideration and New Results, International
747 Journal of Astronomy, Astrophysics and Space Science. Vol. 1, No. 2, 2014, pp. 16-26.
748 [13] Seshavatharam, U. V. S. and Lakshminarayana, S. The Reduced Planck's Constant, Mach's Principle, Cosmic
749 Acceleration and the Black Hole Universe. Journal of Physical Science and Application. Vol.2 (10) 441-447. (2012)
750 [14] Friedmann. A. On the Curvature of Space. General Relativity and Gravitation 31 (12): 1991-2000. 1999
751 [15] Pathria, R. K. The Universe as a Black Hole. Nature 240 (5379):298-299.doi:10.1038/240298a0 (1972).
752 [16] Good, I. J. Chinese universes. Physics Today 25 (7): 15. July. doi:10.1063/1.3070923 (1972).
753 [17] Joel Smoller and Blake Temple. Shock-wave cosmology inside a black hole. Proc Natl Acad Sci U S A. September 30;
754 100(20): 1121611218. (2003).
755 [18] Chul-Moon Yoo et al. Black Hole Universe. Time evolution. Phys. Rev. Lett. 111, 161102 (2013).
756 [19] Michael E. McCulloch. A Toy Cosmology Using a Hubble-Scale Casimir Effect. Galaxies 2014, 2, 81-88.
757 [20] T.X. Zhang and C. Frederic. Acceleration of black hole universe. Astrophysics and Space Science, Volume 349, Issue
758 1, pp 567-573. (2013).
759 [21] Zhang, Tianxi. Cosmic microwave background radiation of black hole universe. Astrophysics and Space Science,
760 Volume 330, Issue 1, pp 157-165. (2010).
761 [22] Zhang, Tianxi. Quasar Formation and Energy Emission in Black hole universe
762 <http://downloads.hindawi.com/journals/aa/2012/625126.pdf>. Progress in Physics, 3: 48-53, (2012).
763 [23] Poplawski, N. J. Radial motion into an Einstein-Rosen bridge. Physics Letters B 687 (23): 110-113. (2010).
764 [24] Poplawski, N. J. Big bounce from spin and torsion. General Relativity and Gravitation Vol. 44, No. 4 (2012) pp.
765 1007–1014.
766 [25] Poplawski, N. J. Energy and momentum of the Universe. Class. Quantum Grav. 31, 065005 (2014).
767 [26] Pourhasan R, Afshordi N and Mann R.B. Did a hyper black hole spawn the universe? Nature - International weekly
768 journal of science. 13 September 2013, doi:10.1038/nature.2013.13743, arXiv: 1309. 1487v2.
769 [27] Andy Gardner, Joseph P. Conlon. Cosmological natural selection and the purpose of the universe. Complexity.
770 Vol.18, Issue 5, pp48-56. 2013
771 [28] Smolin, L. Cosmological natural selection as the explanation for the complexity of the universe. Physica A 340, 705-
772 713. 2004.
773 [29] Hawking S.W. A Brief History of Time. Bantam Dell Publishing Group. 1988
774 [30] Hubble E. P, A relation between distance and radial velocity among extra-galactic nebulae, PNAS, 1929, vol. 15, 1929,
775 pp.168-173.
776 [31] Hubble, E.P, The 200-inch telescope and some problems it may solve. PASP, 59, pp153-167, 1947.
777 [32] J. A. Frieman et al. Dark energy and the accelerating universe. Ann.Rev.Astron.Astrophys.46, 2008, p 385.

Requesting your kind and valuable (unbiased) review comments...

- 778 [33] The Accelerating Universe. The Royal Swedish Academy of sciences. 2011 Nobel prize in physics.
779 www.nobelprize.org/nobel_prizes/physics/laureates/2011/advanced-physicsprize2011.pdf
- 780 [34] Hawking, S.W.; Ellis, G.F.R. (1973). *The Large-Scale Structure of Space-Time*. Cambridge University Press.
- 781 [35] G.J. Stoney, On the Physical Units of Nature. *Phil.Mag.* 11 (1881) 381-91.
- 782 [36] Michael J. Longo, Detection of a Dipole in the Handedness of Spiral Galaxies with Redshifts $z < 0.04$, *Phys. Lett. B* 699,
783 224-229 2011.
- 784 [37] S.-C. Su and M.-C. Chu. Is the universe rotating? *Astrophysical Journal*, 703 354. 2009.
- 785 [38] J. D. McEwen et al. Bayesian analysis of anisotropic cosmologies: Bianchi VIII and WMAP. *Mon. Not. R. Astron.*
786 *Soc.* 000, 1–15 (2013). arXiv:1303.3409v1.
- 787 [39] L. M. Chechin. On the Modern Status of the Universe Rotation Problem. *Journal of Modern Physics*, 2013, 4, 126-132.
- 788 [40] C Sivaram and Kenath Arun, Primordial Rotation of the Universe, Hydrodynamics, Vortices and Angular Momenta of
789 Celestial Objects. *The Open Astronomy Journal*, 2012, 5, 7-11
- 790 [41] Sidharth, B.G. Is the Universe Rotating? *Prespacetime Journal*. October 2010, Vol. 1, Issue 7, pp. 1168-1173.
- 791 [42] Marcelo Samuel Berman, Fernando de Mello Gomide. Local and Global Stability of the Universe. *Journal of Modern*
792 *Physics*, 2013, 4, 7-9
- 793 [43] Robert V Gentry. New Cosmic Center Universe Model Matches Eight of Big Bang's Major Predictions Without The
794 F-L Paradigm. CERN preprint, EXT-2003-022, 14 Apr 2003.
- 795 [44] G. Chapline et al. Tommy Gold Revisited: Why Does Not The Universe Rotate? *AIP Conf.Proc.*822:160-165, 2006.
796 <http://arxiv.org/abs/astro-ph/0509230>.
- 797 [45] Dmitri Rabounski. On the Speed of Rotation of Isotropic Space: Insight into the Redshift Problem. *The Abraham*
798 *Zelmanov Journal*, Vol. 2, 2009, 208-223.
- 799 [46] Kurt Godel. Rotating Universes in General Relativity Theory. *Proceedings of the international Congress of*
800 *Mathematicians in Cambridge*, 1: 175-81, 1950.
- 801 [47] S.W. Hawking. On the rotation of the universe. *Mon. Not. Royal. Astr. Soc.* 142, 129-141.1969.
- 802 [48] M. Novello and M. J. Reboucas. Rotating universe with successive causal and noncausal regions. *Phys. Rev. D* 19,
803 2850-2852 (1979)
- 804 [49] Barrow J D, Juszkiewicz R, Sonoda DH. Universal rotation - How large can it be? *Mon. Not. R. Astron. Soc.* 1985;
805 213: 917.
- 806 [50] Christopher S. Reynolds. Astrophysics: Black holes in a spin. *Nature*. 494, 432–433 (28 February 2013)
- 807 [51] U. V. S. Seshavatharam. Light speed rotating black holes: The special holes *International Journal of Advanced*
808 *Astronomy*, 1 (1) (2013) 13-20.
- 809 [52] Louis Marmet. On the Interpretation of Red-Shifts: A Quantitative Comparison of Red-Shift Mechanisms.
810 www.marmet.org/louis/index.html
- 811 [53] J. Beringer et al. Particle Data Group. *Phys. Rev. D*86, 010001 (2012)
- 812 [54] C. L. Bennett et al, Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and
813 Results. Submitted to *Astrophysical Journal Supplement Series*. <http://arxiv.org/abs/1212.5225v1>
- 814 [55] David N. Spergel et al. Planck Data Reconsidered. <http://arxiv.org/pdf/1312.3313.pdf>
- 815 [56] W. L. Freedman et al. Final Results from the Hubble Space Telescope Key Project to Measure the Hubble Constant.
816 *The Astrophysical Journal* 553 (1): 47-72. 2001.
- 817 [57] P.J. Mohr, B.N. Taylor, and D.B. Newell in arXiv:1203.5425 and *Rev. Mod. Phys.* (to be published).
818 <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-phys-constants.pdf>
- 819 [58] N. Bohr. On the Constitution of Atoms and Molecules. (Part-1) *Philos. Mag.* 26, 1913, p 1.
- 820 [59] Abdus Salam. Einstein's Last Dream: The Space -Time Unification of Fundamental Forces, *Physics News*, 12(2):36,
821 June 1981.
- 822 [60] David Gross, Einstein and the search for Unification. *Current science*, Vol. 89, No. 2005, p 12.
- 823 [61] J. V. Narlikar. *Introduction to cosmology*. Cambridge Univ Press, 2002.
- 824 [62] Lianxi Ma et al. Two forms of Wien's displacement law. *Lat. Am. J. Phys. Educ.* Vol. 3, No. 3, Sept. 2009
- 825 [63] Hawking S.W. Particle creation by black holes. *Commun. Math. Phys.*,1975, v.43, 199–220.
- 826 [64] Gaurab Ganguly et al. SeD Radical: A probe for measurement of time variation of Fine Structure Constant (α) and
827 Proton to Electron Mass Ratio (μ). <http://arxiv.org/pdf/1403.4061v2.pdf>.
- 828 [65] J.K. Webb et al. Indications of a spatial variation of the fine structure constant. *Physical Review letters*, 107 (19) 2011
- 829 [66] Srianand R. et al., Time Variation of the Fine Structure Constant. *The Messenger*. No.116, 25-28 (2004)
- 830 [67] P. A. M. Dirac. The cosmological constants. *Nature*, 139, 1937, p 323.
- 831 [68] P. A. M. Dirac. A new basis for cosmology. *Proc. Roy. Soc. A* 165, 1938, p 199.
- 832 [69] Brandon Carter. Large number coincidences and the anthropic principle in cosmology. *General Relativity and*
833 *Gravitation.*, Volume 43, Issue 11, pp 3225-3233, (2011)
- 834 [70] Ross A. McPherson. The Numbers Universe: An Outline of the Dirac/Eddington Numbers as Scaling Factors for
835 Fractal, Black Hole Universes. *EJTP* 5, No. 18 (2008) 81–94;
- 836 [71] Scott Funkhouser. A new large-number coincidence and a scaling law for the cosmological constant. *Proc. R. Soc. A* 8
837 May 2008 vol. 464 no. 20931345-1353;

Requesting your kind and valuable (unbiased) review comments...

- 838 [72] Barrow. J.D. The Constants of Nature From Alpha to Omega-The Numbers that Encode the Deepest Secrets of the
839 Universe. Pantheon Books. 2002:
- 840 [73] Gamov G. Numerology for the constants of nature. Proceedings of the National Academy of Science U.S.A., 1968, v.
841 59(2). 313–318:
- 842 [74] Saibal Rav. Utpal Mukhopadhyay and Partha Pratim Ghosh. Large Number Hypothesis : A Review.
843 <http://arxiv.org/pdf/0705.1836.pdf>
- 844 [75] P.J. Mohr, B.N. Taylor, and D.B. Newell in arXiv:1203.5425 and Rev. Mod. Phys.
845 <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-phys-constants.pdf>
- 846 [76] Michael O. Distler et al. The RMS Charge Radius of the Proton and Zemach Moments. Phys. Lett.B. 696: 343-
847 347,2011
- 848 [77] Geiger H and Marsden E. On a diffuse reaction of the particles. Proc. Roy. Soc., Ser. A 82: 495-500, 1909.
- 849 [78] H. Yukawa. On the Interaction of Elementary Particles. Proc. Phys. Math. Soc. Jap. 17 (48). 1935
- 850 [79] Recami E. Elementary Particles as Micro-Universes, and “Strong Black-holes”: A Bi-Scale Approach to Gravitational
851 and Strong Interactions. Preprint NSF-ITP-02-94. posted in the arXives as the e-print physics/0505149, and references
852 therein.
- 853 [80] Salam A. and Sivaram C. Strong Gravity Approach to QCD and Confinement. Mod. Phys. Lett., 1993, v. A8(4), 321-
854 326.
- 855 [81] Abdus Salam. Strong Interactions, Gravitation and Cosmology. Publ. in: NATO Advanced Study Institute, Erice,
856 June16-July 6, 1972 ; in: High Energy Astrophysics and its Relation to Elementary Particle Physics, 441-452 MIT
857 Press, Cambridge (1974).
- 858 [82] U. V. S. Seshavatharam and S. Lakshminarayana, Strong nuclear gravitational constant and the origin of nuclear
859 planck scale. Progress in Physics, vol. 3, July, 2010, p. 31-38.
- 860 [83] U. V. S. Seshavatharam and S. Lakshminarayana, Role of Avogadro number in grand unification. Hadronic Journal.
861 Vol-33, No 5, 2010 October. p 513.
- 862 [84] U. V. S. Seshavatharam and S. Lakshminarayana. Accelerating universe and the expanding atom. Hadronic journal,
863 35(3): 271, 2012.
- 864 [85] U. V. S. Seshavatharam and S. Lakshminarayana. Nucleus in Strong nuclear gravity. Proceedings of the DAE Symp.
865 On Nucl. Phys. 56: 302, 2011

866

APPENDIX: Major shortcomings of modern big bang cosmology

- 868 1) It may be noted that, increased redshifts and increased distances forced Edwin Hubble to propose the Hubble’s law . In
869 fact there is no chance or scope or place for ‘galaxy receding’. It is only our belief in its 'given' (Doppler shift based)
870 interpretation. Even then, merely by estimating galaxy distance and without measuring galaxy receding speed, one
871 cannot verify its acceleration. Clearly speaking: two mistakes are possible here. i) Assumed galaxy receding speed is
872 not being measured and not being confirmed. ii) Without measuring and confirming the galaxy receding speed, how
873 can one say and confirm that it (galaxy) is accelerating. It is really speculative.
- 874 2) If light is coming from the atoms of the gigantic galaxy, then redshift can also be interpreted as an index of the galactic
875 cosmological atomic ‘light emission mechanism’. In no way it seems to be connected with ‘galaxy receding’.
- 876 3) According to the modern cosmological approach, bound systems like ‘atoms’ which are found to be the major
877 constituents of galactic matter - will not change with cosmic expansion/acceleration. As per the present observational
878 data this may be true. But it might be the result of ending stage of cosmic expansion. As the issue is directly related
879 with unification it requires lot of research in basic physics to confirm. In this regard, without considering and without
880 analysing the past data, one can not come to a conclusion. If one is willing to think in this direction observed galactic
881 redshift data can be considered for this type of new analysis.
- 882 4) ‘Rate of decrease in current ‘Hubble’s constant’ can be considered as a measure of current cosmic ‘rate of expansion’.
883 If rate of decrease in current ‘Hubble’s constant is very small and is beyond the scope of current experimental
884 verification, then the two possible states are: a) current ‘Hubble’s constant is decreasing at a very slow rate and current
885 universe is expanding at a very slow rate and b) at present there is no ‘observable’ cosmic expansion. Without a proper
886 confirmation procedure for the absolute cosmic expansion and guessing that current universe is expanding -
887 cosmologists proposed and confirmed the existence of dark energy indirectly. It may not be reasonable. Quantitatively
888 or at least qualitatively standard model of cosmology does not throw light on the generation and (normal) physical
889 properties of ‘dark energy’.
- 890 5) The standard Big Bang model tells us that the Universe exploded out of an infinitely dense point. But nobody knows
891 what would have triggered this outburst: the known laws of physics cannot tell us what happened at that moment.
- 892 6) Really if there was a ‘big bang’ in the past, with reference to formation of the big bang as predicted by general theory
893 of relativity and with reference to the cosmic expansion that takes place simultaneously in all directions at a uniform

Requesting your kind and valuable (unbiased) review comments...

- 894 rate at that time about the point of big bang - 'point' of big bang can be considered as the centre or characteristic
895 reference point of cosmic expansion in all directions. In this case, saying that there is no preferred direction in the
896 expanding universe - may not be correct.
- 897 7) Either in the big bang or in the inflation, quantification of the initial assumed conditions seem to be poor, unclear and
898 not linked with fundamental constants. The earliest phases of the Big Bang are subject to much speculation and
899 inflation requires 'fine tuning'.
 - 900 8) Standard cosmology does not give information on the origin of 'inflation'. Inflation is often called a period
901 of accelerated expansion. With respect to 'no hair theorem' some similarities are there for cosmic inflation and black
902 holes. Conceptually 'inflation' can be accommodated in any model of cosmology like open model or closed model.
 - 903 9) A key requirement is that inflation must continue 'long enough' to produce the present observable universe from a
904 single, small inflationary Hubble volume. Assuming a rapid rate of cosmic expansion and steady rate of time may not
905 be reasonable. If space-time is interrelated then 'space' and 'time' both should simultaneously follow the momentary
906 rapid exponential expansion. For example if space expands by a factor 10^{26} in size within a very 'short span', cosmic
907 time should also increase in the same proportion. 'Time' seems to be a silent observer in the presently believed
908 'cosmic inflation'. It may not be reasonable.
 - 909 10) There is no scientific evidence for the Friedmann's second assumption. We believe it only on the grounds of modesty.
 - 910 11) Dimensionally it is perfectly possible to show that, the dimensions of Hubble's constant and angular velocity are same.
911 If so considering Hubble's constant merely as an expansion parameter may not be correct.
 - 912 12) Even though it was having strong footing, Mach's principle was not implemented successfully in standard cosmology.
913 Clearly speaking the term "distance cosmic back ground" is not being defined and not being quantified in a physical
914 approach.
 - 915 13) At any given cosmic time, the product of 'critical density' and 'Hubble volume' gives a characteristic cosmic mass
916 and it can be called as the 'Hubble mass'. Interesting thing is that, Schwarzschild radius of the 'Hubble mass' again
917 matches with the 'Hubble length'. Most of the cosmologists believe that this is merely a coincidence. Here the
918 researchers emphasize the fact that this coincidence is having deep connection with cosmic geometry and the
919 cosmological physical phenomena.
 - 920 14) Somehow and by any reason, magnitude of the current Hubble mass being the same, hypothetically if volume density
921 approaches the current matter density, then Hubble length increases by a factor ~ 5 . Similarly if volume density
922 approaches the current thermal energy density, then Hubble length increases by a factor ~ 27 . These two numbers
923 can be compared with the presently believed first two of the three cosmological numbers 4.9%, 26.8% and 68.3%.
924 Based on this coincidence and as the currently believed third number $\sim 68\%$ is obtained from the relation $(100 - (4.9 + 26.8))\%$, its proposed existence seems to be ad-hoc.
 - 925 15) If 'Planck mass' is the characteristic beginning 'mass scale' of the universe, then by substituting the geometric mean
926 mass of the present Hubble mass and the Planck mass in the famous Hawking's black hole temperature formula
927 automatically the observed 2.725 K can be fitted very accurately. Standard cosmology is not throwing any light on this
928 surprising coincidence.
 - 929 16) If cosmic expansion is continuous and accelerating and redshift is a measure of cosmic expansion, then 'rate of
930 increase in redshift' can be considered as a measure of cosmic 'rate of expansion'. Then there is no possibility to
931 observe a 'constant' red shift. More over the current definition of red shift seems to be ad-hoc and not absolute. Hence
932 one may not be able to understand or confirm the actual cosmic rate of expansion.
 - 933 17) Even though the whole physics strictly follows the 'constancy of speed of light', cosmic acceleration seems to violate
934 it. This is really doubtful.
 - 935 18) Drop in current 'cosmic temperature' can be considered as a measure of the current cosmic expansion and 'rate of
936 decrease in current cosmic temperature' can be considered as a measure of the current cosmic 'rate of expansion'. But
937 if rate of decrease in temperature is very small and is beyond the scope of current experimental verification, then the
938 two possible states are: a) current cosmic temperature is decreasing at a very slow rate and current universe is
939 expanding at a very slow rate and b) at present there is no 'observable' thermal expansion and there is no 'observable'
940 cosmic expansion. If observed CMBR temperature is 2.725 K and is very low in magnitude and is very close to
941 absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be reasonable.
 - 942 19) If observed cosmic microwave back ground radiation temperature is 2.725 K and is very low in magnitude and is very
943 close to absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be reasonable.
 - 944 20) In the standard model of cosmology, there is no clear cut information about the 'uniqueness' of the assumed 'dark
945 energy'. If its identification is not unique in nature, then different cosmology models can be developed with different
946 forms of 'dark energy'. If so understanding the absolute cosmic expansion rate with dark energy seems to be doubtful.
 - 947 21) So far no ground based experiment confirmed the existence of dark energy. There is no single clue or evidence to any
948 of the natural physical properties of (the assumed) dark energy.
949

Requesting your kind and valuable (unbiased) review comments...

- 950 22) If 'Dark energy' is the major outcome of the 'accelerating universe', it is very important to note that - in understanding
951 the basic concepts of unification or other fundamental areas of physics, role of dark energy is very insignificant.
- 952 23) If existence of dark energy is true and dark energy is supposed to have a key role in the past and current cosmic
953 expansion, then it must have also played a key role in the beginning of cosmic evolution. In this regard no
954 information is available in standard cosmology.
- 955 24) Standard model of cosmology does not throw light on the generation and existence of atomic physical constants like
956 Planck's constant, reduced Planck's constant, inverse of fine structure ratio and nuclear charge radius etc. Clearly
957 speaking synthesis of elementary physical constants seem to be more important than the cosmological nucleosynthesis.
- 958 25) General theory of relativity does not throw any light on the 'mass generation' of charged particles. It only suggests
959 that space-time is curved near the massive celestial objects. More over it couples the cosmic (dust) matter with
960 geometry. But how matter/dust is created? Why and how elementary particle possesses both charge and mass? Such
961 types of questions are not being discussed in the frame work of general relativity.
- 962 26) Standard model of cosmology does not throw light on the charge-mass unification scheme of atomic particles. The
963 main object of unification is to understand the origin of elementary particles rest mass, magnetic moments and their
964 forces. Right now and till today 'string theory' with 4 + 6 extra dimensions is not in a position to explain the
965 unification of gravitational and non-gravitational forces. More clearly speaking it is not in a position to merge the
966 Planck scale and cosmic scale with the characteristic nuclear scale.
- 967 27) Either general theory of relativity or standard cosmology does not give any information on the applications of the
968 classical force limit (c^4/G) and the classical power limit (c^5/G). Compared to the hypothetical 'dark energy', with a
969 coefficient of unity, (c^4/G) can be considered as the cosmic vacuum force and (c^5/G) can be considered as the cosmic
970 vacuum power.
- 971 28) In Big bang model, confirmation of all the observations directly depend on the large scale galactic distances that are
972 beyond human reach and raise ambiguity in all respects. The subject of modern black hole physics is absolutely
973 theoretical. Advantage of Black hole cosmology lies in confirming its validity through the ground based atomic and
974 nuclear experimental results.

976 If one is willing to think in this new direction, certainly other hidden short comings can also be surfaced out. Most of the
977 modern cosmologists are enforced with 85 years old Hubble's interpretation. This is the time to re-interpret the Hubble's
978 law and to revise the basics of modern cosmology. Based on the proposed short comings the concepts of 'big bang
979 cosmology' can be relinquished and Black hole cosmology can be invoked for in-depth discussion.