

Inadequacy of Hubble-Friedmann Cosmology and the Basics of Stoney Scale Black Hole Cosmology

U. V. S. Seshavatharam^{*1,2} and S. Lakshminarayana⁺

^{*1}Honorary faculty, I-SERVE, Alakapuri, Hyderabad-35, India. Email: seshavatharam.uvs@gmail.com

^{*2}Sr. QA Engineer, DIP Division, Lanco Industries Ltd, Rachagunneri, Srikalahasti-517644, AP, India.

⁺Dept. of Nuclear Physics, Andhra University, Visakhapatnam-03, India. Email: Insrirama@yahoo.com

Abstract: Throughout the cosmic evolution, currently believed cosmic ‘critical density’ can be shown to be a default result of the ‘positively curved’ light speed rotating black hole universe ‘volume density’. As there is no observational or experimental evidence to Friedmann’s second assumption, the density classification scheme of Friedmann cosmology must be reviewed at fundamental level and possibly can be relinquished. The observed cosmic redshift can be reinterpreted as an index of ‘cosmological’ thermodynamic light emission mechanism. Clearly speaking during cosmic evolution, at any time in the past, in hydrogen atom- emitted photon energy was always inversely proportional to the cosmic temperature. Thus past light emitted from older galaxy’s excited hydrogen atom will show redshift with reference to the current laboratory data. Note that there will be no change in the energy of the emitted photon during its journey from the distant galaxy to the observer. By considering the ‘Stoney mass’ as the initial mass of the baby cosmic black hole, past and current physical and thermal parameters (like angular velocity, growth rate, age, redshift, thermal energy density and matter density) of the cosmic black hole can be understood. For a cosmic temperature of 3000 K, obtained redshift is 1100. From now onwards, CMBR temperature can be called as ‘Comic Black Hole’s Thermal Radiation’ temperature and can be expressed as ‘CBHTR’ temperature. Current cosmic black hole is growing at a rate of 14.66 km/sec in a decelerating mode. Uncertainty relation and all other microscopic physical constants play a crucial role in understanding the halt of the present cosmic expansion. In view of the confirmed zero rate of change in inverse of the Fine structure ratio (from the ground based laboratory experimental results), zero rate of change in the current CMBR temperature (from satellite data) and zero rate of change in the current Hubble’s constant (from satellite data), it can be suggested that, current cosmic expansion is almost all saturated and at present there is no significant cosmic acceleration.

Keywords: Mach’s principle, Stoney mass, Black Hole Cosmology, Cosmic growth index, Cosmic growth rate, Hubble Potential, Cosmic redshift, Cosmic age, Halting of Cosmic Expansion, Final Unification.

1. Introduction

Authors published their concepts on black hole cosmology in many online journals [1-13]. In this paper by highlighting the basic short comings of Friedmann cosmology [14] an attempt is made to review the model of black hole cosmology [15-28] in terms of cosmic redshift, CMBR redshift, cosmic growth index, cosmic growth rate and cosmic age. The basic shortcomings of modern cosmology can be expressed as follows. For more information one may see the appendix [1].

- 1) No direct observational evidence to Friedmann’s second assumption [29].
- 2) No theoretical base in Friedmann’s ‘critical density’ concept and the ‘matter density’ classification scheme.
- 3) If light is coming from the atoms of the gigantic galaxy, then redshift can also be interpreted as an index of the galactic cosmological atomic ‘light emission mechanism’. In no way it seems to be connected with ‘galaxy receding’.
- 4) No theoretical base in the currently believed wave length based redshift definition [30,31]. In terms of ‘quantum of energy’, redshift can also be interpreted as an index of cosmological thermodynamic light emission mechanism in hydrogen atom.
- 5) Merely by estimating galaxy distance and without measuring galaxy receding speed, one cannot verify its receding speed or acceleration. (Clearly speaking: two mistakes are possible here. i) Assumed galaxy receding speed is not being measured and not being confirmed. ii) Without measuring and confirming the galaxy receding speed, how can one say and confirm that it (galaxy) is accelerating).
- 6) No theoretical base in considering the Hubble’s constant merely as the cosmic expansion parameter. With reference to angular velocity it is having deep inner meaning.
- 7) No direct observational evidence for the current cosmic acceleration and the dark energy [32,33].

Requesting your kind and valuable (unbiased) review comments...

- 51 8) By substituting the geometric mean mass of $(c^3/2GH_0)$ and $\sqrt{hc/2\pi G}$ in the famous Hawking's black hole
52 temperature formula automatically the observed 2.725 K can be fitted very accurately.
53 9) When Friedmann's cosmology was taking its final shape, black hole physics was in its beginning stage.
54 10) No comparative and relational study in between Friedmann cosmology and microscopic physical phenomena.
55

56 Friedmann made two simple assumptions about the universe. They can be stated in the following way.

- 57 1. When viewed at large enough scales, universe appears the same in every direction.
- 58 2. When viewed at large enough scales, universe appears the same from every location.

60
61 In this regard Hawking says : "There is no scientific evidence for the Friedmann's second assumption. We believe it only
62 on grounds of modesty: it would be most remarkable if the universe looked the same in every direction around us, but not
63 around other points in the universe". This is one key point to be noted here. The term 'critical density' is the back bone of
64 modern cosmology. At any time in the past, it is generally expressed in the following way.
65

$$66 \quad (\rho_c)_t \cong \frac{3H_t^2}{8\pi G} \quad (1)$$

67 Its current expression is as follows.

$$68 \quad (\rho_c)_0 \cong \frac{3H_0^2}{8\pi G} \quad (2)$$

69 According to standard Friedmann cosmology,

- 70 1. If matter density is greater than the critical density, universe will have a positive curvature.
- 71 2. If matter density equals the critical density, universe will be flat.
- 72 3. If matter density is less than the critical density, universe will have a negative curvature.

73
74
75 But by considering 'black hole geometry' as the 'eternal cosmic geometry' and by assuming 'constant light speed
76 rotation' throughout the cosmic evolution, at any time the currently believed cosmic 'critical density' can be shown to be
77 the cosmic black hole's eternal 'volume density'. If mass of the black hole universe is M_t , $\left(\frac{c}{H_t}\right)$ is the radius of the
78 black hole universe that rotates at light speed and angular velocity H_t , at any time in the past,
79

$$80 \quad \frac{2GM_t}{c^2} \cong \frac{c}{H_t} \quad \text{and} \quad M_t \cong \frac{c^3}{2GH_t}. \quad (3)$$

$$81 \quad (\rho_v)_t \cong (M_t) \left[\frac{4\pi \left(\frac{c}{H_t}\right)^3}{3} \right]^{-1} \cong \left(\frac{c^3}{2GH_t} \right) \left[\frac{3 \left(\frac{H_t}{c}\right)^3}{4\pi \left(\frac{c}{H_t}\right)} \right] \cong \frac{3H_t^2}{8\pi G} \quad (4)$$

82 At present,

$$83 \quad (\rho_v)_0 \cong (M_0) \left[\frac{4\pi \left(\frac{c}{H_0}\right)^3}{3} \right]^{-1} \cong \left(\frac{c^3}{2GH_0} \right) \left[\frac{3 \left(\frac{H_0}{c}\right)^3}{4\pi \left(\frac{c}{H_0}\right)} \right] \cong \frac{3H_0^2}{8\pi G} \quad (5)$$

84 Based on this coincidence and as there is no observational or experimental evidence to Friedmann's second assumption, the
85 density classification scheme of Friedmann cosmology must be reviewed at fundamental level.

86 **2. Possible Assumptions and Possible Explanation**

87 Possible assumptions in unified cosmic physics can be expressed in the following way.

88
89 **Assumption-1: With reference to the elementary charge and with mass similar to the Planck mass, a new mass unit
90 can be constructed in the following way. It can be called as the Stoney mass.**
91

Requesting your kind and valuable (unbiased) review comments...

92
$$(M_S)^\pm \cong \sqrt{\frac{e^2}{4\pi\epsilon_0 G}} \cong 1.859272 \times 10^{-9} \text{ Kg} \cong 1.042975 \times 10^{18} \text{ GeV}/c^2 \quad (6)$$

93 **Assumption-2: At any time Hubble length (c/H_t) can be considered as the gravitational or electromagnetic**
 94 **interaction range.**

95
 96 **Assumption-3: At any time, H_t being the angular velocity, universe can be considered as a growing and light speed**
 97 **rotating primordial black hole.** Thus at any given cosmic time,

98
$$R_t \cong \frac{2GM_t}{c^2} \cong \frac{c}{H_t} \text{ and } M_t \cong \frac{c^3}{2GH_t} \quad (7)$$

99 when $M_t \rightarrow M_S$, $R_S \cong \frac{2GM_S}{c^2}$ and $H_S \cong \frac{c}{R_S} \cong \frac{c^3}{2GM_S}$ (8)

100
 101 can be considered as the characteristic initial physical measurements of the universe. Here the subscript S refers to the
 102 initial conditions of the universe and can be called as the Stoney scale. Similarly
 103

104
$$R_0 \cong \frac{2GM_0}{c^2} \cong \frac{c}{H_0}, \quad M_0 \cong \frac{c^3}{2GH_0} \text{ and } H_0 \cong \frac{c^3}{2GM_0} \quad (9)$$

105
 106 can be considered as the characteristic current physical measurements of the universe.

107
 108 **Assumption-4: During cosmic evolution, at any time the past, in hydrogen atom emitted photon energy was always**
 109 **inversely proportional to the cosmic temperature. Thus past light emitted from older galaxy's hydrogen atom will**
 110 **show redshift with reference to the current laboratory data. There will be no change in the energy of the emitted**
 111 **photon during its journey from the distant galaxy to the observer.**

112
 113
$$\frac{E_t}{E_0} \cong \frac{\lambda_0}{\lambda_t} \cong \frac{T_t}{T_0} \quad (10)$$

114 Here, E_t is the energy of emitted photon from the galactic hydrogen atom and E_0 is the corresponding energy in the
 115 laboratory. λ_t is the wave length of emitted and received photon from the galactic hydrogen atom and λ_0 is the
 116 corresponding wave length in the laboratory. T_t is the cosmic temperature at the time when the photon was emitted and is
 117 T_0 the current cosmic temperature.

118
 119 **Assumption-5: At any given time, ratio of volume energy density and thermal energy density can be called as the**
 120 **cosmic growth index and can be expressed as follows.**

121
 122
$$\frac{3H_t^2 c^2}{8\pi G a T_t^4} \cong \left[1 + \ln \left(\frac{M_t}{M_S} \right) \right]^2 \cong \left[1 + \ln \left(\frac{H_S}{H_t} \right) \right]^2 \quad (11)$$

$$\cong \text{Cosmic Growth index}$$

123 Thus at the Stoney scale,

124
$$\frac{3H_S^2 c^2}{8\pi G a T_S^4} \cong \left[1 + \ln \left(\frac{M_S}{M_S} \right) \right]^2 \cong \left[1 + \ln \left(\frac{H_S}{H_S} \right) \right]^2 \cong 1 \quad (12)$$

125 **Assumption-6: At any given time, cosmic black hole's growth rate can be expressed as $g_t \cong \left(\frac{3H_t^2 c^2}{8\pi G a T_t^4} \right)^{-1} c$. With this**
 126 **idea and by considering the average growth rate cosmic age can be estimated.**

127

Requesting your kind and valuable (unbiased) review comments...

$$g_t \cong \text{Cosmic growth rate} \cong \frac{c}{\text{cosmic growth index}} \quad (13)$$
$$\cong \left(\frac{3H_t^2 c^2}{8\pi G a T_t^4} \right)^{-1} c \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^{-2} c \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-2} c$$

At the Stoney scale,

$$g_s \cong \left(\frac{3H_s^2 c^2}{8\pi G a T_s^4} \right)^{-1} c \cong \left[1 + \ln \left(\frac{M_s}{M_s} \right) \right]^{-2} c \cong \left[1 + \ln \left(\frac{H_s}{H_s} \right) \right]^{-2} c \cong c \quad (14)$$

2.1 Possible Explanation for the proposed Assumptions

To have some clarity and to have some quantitative measurements and fittings of initial and current states of the black hole universe - instead of considering 'star - black hole explosions' and 'higher dimensions', the authors of this paper focused their attention only on the old and famous Mach's principle [34], 'Hubble volume' and 'primordial evolving black holes'. Some cosmologists use the term 'Hubble volume' to refer to the volume of the observable universe. There is no perfect theory that defines the lower and upper limits of a massive black hole. Most of the theoretical models assume a lower mass limit close to the 'Planck mass'. Astronomers believe that black holes that are as large as a billion solar masses can be found at the centre of most of the galaxies. Here the fundamental questions to be answered are: If the galactic central black hole mass is 10 billion solar masses and density is less than 1 kg/m^3 - with such a small density and large mass, without collapsing - how it is able to hold a gigantic galaxy? What force makes the black hole stable? Recent observations confirm that, instead of collapsing, galactic central black holes are growing faster and spinning with light speed. Even though mass is too high and density is too low, light speed rotation certainly helps in maintaining black hole's stability from collapsing with maximum possible outward radial force of the magnitude close to (c^4/G) . Based on these points the authors propose the following picture of Black hole cosmology. Forever rotating at light speed, high temperature and high angular velocity small sized primordial cosmic black hole of mass $M_s \cong \sqrt{e^2/4\pi\epsilon_0 G}$ gradually transforms into a low temperature and low angular velocity large sized massive primordial cosmic black hole. At any given cosmic time, for the primordial growing black hole universe, its 'Schwarzschild radius' can be considered as its characteristic possible minimum radius and 'constant light speed rotation' will give the maximum possible stability from collapsing. Here $M_s \cong \sqrt{e^2/4\pi\epsilon_0 G}$ can be called as the mass of the primordial baby black hole universe. Here 4 important points can be stated as follows.

1. It is well known that e, c, G play a vital role in fundamental physics. With these 3 constants space-time curvature concepts at a charged particle surface can be studied. Note that the basic concept of unification is to understand the origin of 'mass' of any particle. Mass is the basic property in 'gravitation' and charge is the basic property in 'atomicity'. So far no model established a cohesive relation in between 'electric charge' and 'mass' of any 'elementary particle' or 'cosmic dust'. From physics point of view, the fundamental questions to be answered are: 1) Without charge, is there any independent existence to "mass"? 2) Without mass, is there any independent existence to "charge"? From cosmology point of view the fundamental questions to be answered are: 1) What is 'cosmic dust'? 2) Without charge, is there any independent existence to "cosmic dust"? From astrophysics point of view the fundamental questions to be answered are: 1) Without charge, is there any independent existence to 'mass' of any star? 2) Is black hole - a neutral body or electrically a neutralized body? To understand these questions the authors made an attempt to construct the above unified mass unit. It is having a long history. It was first introduced by the physicist George Johnstone Stoney [35]. He is most famous for introducing the term 'electron' as the 'fundamental unit quantity of electricity'. With this mass unit in unification program with a suitable proportionality it may be possible to represent the characteristic mass of elementary charge. It can be considered as the seed of galactic matter or galactic central black hole. It can also be considered as the seed of any cosmic structure. If 2 such oppositely charged particles annihilates, a large amount of energy can be released. If so under certain extreme conditions at the vicinity of massive stars or black holes, a very high energy radiation can be seen to be emitted by the pair annihilation of M_s . With this mass unit, proton-electron mass ratio and proton and electron rest masses can be fitted. Thus with reference to the elementary charge and electron & proton rest masses, magnitude of the gravitational constant can be fitted [1,2].
2. In theoretical physics, particularly in discussions of gravitation theories, Mach's principle is the name given by Einstein to an interesting hypothesis often credited to the physicist and philosopher Ernst Mach. The idea is that the

Requesting your kind and valuable (unbiased) review comments...

176 local motion of a rotating reference frame is determined by the large scale distribution of matter. With reference to the
177 Mach's principle and the Hubble volume, at any cosmic time, if 'Hubble mass' is the product of cosmic 'critical
178 density' and the 'Hubble volume', then it can be suggested that, i) Each and every point in the free space is influenced
179 by the Hubble mass, ii) Hubble volume and Hubble mass play a vital role in understanding the properties of
180 electromagnetic and nuclear interactions and iii) Hubble volume and Hubble mass play a key role in understanding the
181 geometry of the universe. With reference to the famous Mach's principle, 'Hubble volume' and 'Hubble mass' both can
182 be considered as quantitative measurements of the 'distance cosmic back ground'. As a first attempt, in this paper
183 authors proposed a semi empirical relation that connects the CMBR energy density, Hubble's constant and
184 $\sqrt{e^2/4\pi\epsilon_0 G}$.

- 185 3. Starting from an electron to any gigantic galaxy, rotation is a common phenomenon in atomic experiments and
186 astronomical observations. From Newton's laws of motion and based on the Mach's principle, sitting inside a closed
187 universe, one cannot comment whether the universe is rotating or not. We have to search for alternative means for
188 confirming the cosmic rotation. Recent findings from the University of Michigan [36] suggest that the shape of the Big
189 Bang might be more complicated than previously thought, and that the early universe spun on an axis. A left-handed
190 and right-handed imprint on the sky as reportedly revealed by galaxy rotation would imply the universe was rotating
191 from the very beginning and retained an overwhelmingly strong angular momentum. An anonymous referee who
192 reviewed the paper for Physics Letters said, "In the paper the author claims that there is a preferred handedness of
193 spiral galaxies indicating a preferred direction in the universe. Such a claim, if proven true, would have a profound
194 impact on cosmology and would very likely result in a "Nobel prize". The consequences of a spinning universe [36-49]
195 seem to be profound and natural. Not only that, with 'constant rotation speed' 'cosmic collapse' can be prevented and
196 can be considered as an alternative to the famous 'repulsive gravity' concept. If so, at any time to have maximum
197 possible stability from collapsing 'constant light speed rotation' can be considered as a constructive and workable
198 concept.
- 199 4. Recent observations confirm black hole's light speed rotation. In 2013 February, using NASA's newly launched NuStar
200 telescope and the European Space Agency's workhorse XMM-Newton, an international team observed high-energy X-
201 rays released by a super massive black hole in the middle of a nearby galaxy. They calculated its spin at close to the
202 speed of light: 670 million mph [50,51]. Please note that, for any black hole even though its mass is too high and
203 density is too low, light speed rotation certainly helps in maintaining its stability from collapsing with maximum
204 possible outward radial force of magnitude (c^4/G) . At the beginning of comic evolution if rotation speed was zero and
205 there was no big bang - definitely it will cast a doubt on the stability, existence and angular velocity of the assumed
206 initial primordial cosmic baby black hole. Hence at the beginning also, to guess or define the angular velocity and to
207 have maximum possible stability it is better to assume light speed rotation for the cosmic baby black hole. At present if
208 rate of cosmic expansion is very slow, then rate of decrease in angular velocity will be very small and practically can
209 be considered as zero. Along with (practically) constant angular velocity, at present if constant light speed rotation is
210 assumed to be maintained then cosmic stability will be maximum and rate of change in cosmic size will be practically
211 zero and hence this idea helps us to believe in present Hubble length along with the observed ordered galactic
212 structures and uniform thermal energy density.

213 **2.2 To reinterpret the Hubble's constant**

214 With a simple derivation it is possible to show that, Hubble's constant H_t represents the cosmological angular velocity.
215 Authors presented this derivation in their published papers. Basic idea of this derivation is to express the angular velocity
216 of any rotating celestial body in terms of its mass, radius, mass density and surface escape velocity. Assume that, a planet
217 of mass M and radius R rotates with angular velocity ω_e and linear velocity v_e in such a way that, free or loosely bound
218 particle of mass m lying on its equator gains a kinetic energy equal to potential energy as,

$$219 \quad \frac{1}{2}mv_e^2 = \frac{GMm}{R} \quad (15)$$

$$220 \quad R\omega_e = v_e = \sqrt{\frac{2GM}{R}} \quad \text{and} \quad \omega_e = \frac{v_e}{R} = \sqrt{\frac{2GM}{R^3}} \quad (16)$$

221 i.e Linear velocity of planet's rotation is equal to free particle's escape velocity. Without any external power or energy, test
222 particle gains escape velocity by virtue of planet's rotation. Note that if Earth completes one rotation in one hour then free
223 particles lying on the equator will get escape velocity. Now writing $M = \frac{4\pi}{3}R^3\rho_e$,

Requesting your kind and valuable (unbiased) review comments...

224
$$\omega_e = \frac{v_e}{R} = \sqrt{\frac{8\pi G \rho_e}{3}} \quad \text{Or} \quad \omega_e^2 = \frac{8\pi G \rho_e}{3} \quad (17)$$

225
$$\text{Density, } \rho_e = \frac{3\omega_e^2}{8\pi G} \quad (18)$$

226 In real time, this obtained density may or may not be equal to the actual density. But the ratio $\frac{8\pi G \rho_{real}}{3\omega_{real}^2}$ may have some
227 physical significance. The most important point to be noted here, is that, as far as dimensions and units are considered,
228 from equation (18), it is very clear that, proportionality constant being $\frac{3}{8\pi G}$,
229

230
$$\text{density} \propto (\text{angular velocity})^2 \quad (19)$$

231 Equation (18) is similar to “flat model concept” of cosmic “critical density”
232

233
$$\rho_c = \frac{3H_t^2}{8\pi G} \quad (20)$$

234 Comparing equations (18) and (20) dimensionally and conceptually, i.e.
235

236
$$\rho_e = \frac{3\omega_e^2}{8\pi G} \quad \text{with} \quad \rho_c = \frac{3H_t^2}{8\pi G} \quad (21)$$

237
$$H_t^2 \rightarrow \omega_e^2 \quad \text{and} \quad H_t \rightarrow \omega_e \quad (22)$$

238 It is very clear that, dimensions of ‘Hubble’s constant’ must be ‘radian/second’. In any physical system under study, for
239 any one ‘simple physical parameter’ there will not be two different units and there will not be two different physical
240 meanings. This is a simple clue and brings ‘cosmic rotation’ into picture. This is possible in a closed universe only. Cosmic
241 models that depend on this “critical density” may consider ‘angular velocity of the universe’ in the place of ‘Hubble’s
242 constant’. In the sense, with a great confidence ‘cosmic rotation’ can be included in the existing models of cosmology. Then
243 the term ‘critical density’ appears to be the ‘volume density’ of the closed and expanding universe. Thinking in this way,
244 considering ‘black hole geometry’ as the ‘eternal cosmic geometry’ and by assuming ‘constant light speed rotation’
245 throughout the cosmic evolution, at any time the currently believed cosmic ‘critical density’ can be shown to be the cosmic
246 black hole’s eternal ‘volume density’. Thus based on the Mach’s principle, ‘distance cosmic back ground’ can be quantified
247 in terms of ‘Hubble volume’ and ‘Hubble mass’.

248 **2.3 To reinterpret the Cosmic redshift**

249 Hubble initially interpreted red shifts [30] as a Doppler effect, due to the motion of the galaxies as they receded for our
250 location in the Universe[52]. He called it a ‘Doppler effect’ as though the galaxies were moving ‘through space’; that is
251 how some astronomers initially perceived it. This is different to what has now become accepted but observations alone
252 could not distinguish between the two concepts. In 1947 he [31] stated that: “The red shifts are more easily interpreted as
253 evidence of motion in the line of sight away from the earth – as evidence that the nebulae in all directions are rushing away
254 from us and that the farther away they are, the faster they are receding. This interpretation lends itself directly to theories of
255 expanding universe. The interpretation is not universally accepted, but even the most cautious of us admit that red shifts are
256 evidence of either an expanding universe or of some hitherto unknown principle of nature”. “Attempts have been made to
257 attain the necessary precision with the 100 inch, and the results appear to be significant. If they are valid, it seems likely
258 that the red-shifts may not be due to an expanding universe, and much of the current speculation on the structure of the
259 universe may require re-examination. The significant data, however, were necessarily obtained at the very limit of a single
260 instrument, and there were no possible means of checking the results by independent evidence. Therefore the results must
261 be accepted for the present as suggestive rather than definitive”. “We may predict with confidence that the 200 inch will
262 tell us whether the red shifts must be accepted as evidence of a rapidly expanding universe, or attributed to some new
263 principle in nature. Whatever may be the answer, the result may be welcomed as another major contribution to the
264 exploration of the universe.”

Requesting your kind and valuable (unbiased) review comments...

265 In this regard if one is willing to consider the proposed assumptions, in hydrogen atom emitted photon energy can be under
266 stood as follows.

- 267 1. As the cosmic time increases cosmic angular velocity and hence cosmic temperature both decrease. As a result, during
268 cosmic evolution, in hydrogen atom, binding energy increases in between proton and electron.
- 269 2. As cosmic temperature decreases, it requires more excitation energy to break the bond between electron and the proton.
270 In this way, during cosmic evolution, whenever it is excited, hydrogen atom emits photons with increased quantum of
271 energy.
- 272 3. Thus past light quanta emitted from old galaxy's excited hydrogen atom will have less energy and show a red shift with
273 reference to the current laboratory magnitude.
- 274 4. During journey light quanta will not lose energy and there will be no change in light wavelength.
- 275 5. Galactic photon energy in hydrogen atom when it was emitted can be estimated as follows.
276

277
$$E_t \cong \frac{hc}{\lambda_t} \cong \left(\frac{T_0}{T_t}\right) \left(\frac{hc}{\lambda_0}\right) \cong \left(\frac{T_0}{T_t}\right) E_0 \quad (23)$$

278 Here, λ_0 is the wavelength of photon in the laboratory.

279 E_t is the energy of received photon when it was emitted in the distant galaxy.

280 E_0 is the corresponding energy of photon in the current laboratory methods.

281 λ_t is the wavelength of emitted and received photon when it was emitted in the distant galaxy.

282 T_t is the cosmic temperature at the time when the photon was emitted and T_0 the current cosmic temperature.

283 In subsection 2.5 an attempt is made to understand the cosmological thermodynamic light emission mechanism in hydrogen
284 atom in a unified approach.

285 **2.4 To reinterpret the Hubble's Law**

286 Based on the assumptions it is possible to say that, during cosmic evolution, at any time, any galaxy will have revolution
287 speed as well as receding speed simultaneously and both can be expressed in the following way.
288

289
$$(V_G)_{revolution} \cong \left(\frac{r}{R_t}\right) c \cong rH_t \quad \text{where } r \leq \left(R_t \cong \frac{c}{H_t}\right) \quad (24)$$

290 r is the distance between galaxy and the cosmic center and R_t is the cosmic radius at time t .
291
292

293
$$\begin{aligned} (V_G)_{receding} &\cong \left(\frac{r}{R_t}\right) g_t \cong \left(\frac{r}{R_t}\right) \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} c \\ &\cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} rH_t \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^{-2} (V_G)_{revolution} \end{aligned} \quad (25)$$

294
295
$$\frac{(V_G)_{revolution}}{(V_G)_{receding}} \cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^2 \quad (26)$$

296 Please note that both the relations are independent of the observed redshift. This is for further study.
297
298

299 **2.5. To understand the cosmological thermodynamic light emission mechanism**

300 It is very tempting to make an analogy between the status of the cosmological 'Standard Model' and that of particle physics
301 [53]. In cosmology there are about 10 free parameters, each of which is becoming well determined, and with a great deal of
302 consistency between different measurements. However, none of these parameters can be calculated from a fundamental
303 theory, and so hints of the bigger picture, 'physics beyond the Standard Model,' are being searched for with ever more
304

Requesting your kind and valuable (unbiased) review comments...

305 ambitious experiments. Despite this analogy, there are some basic differences. For one thing, many of the cosmological
 306 parameters change with cosmic epoch, and so the measured values are simply the ones determined today, and hence they
 307 are not ‘constants,’ like particle masses for example (although they are deterministic, so that if one knows their values at
 308 one epoch, they can be calculated at another). Moreover, the parameter set is not as well defined as it is in the particle
 309 physics Standard Model; different researchers will not necessarily agree on which parameters should be considered as free,
 310 and the set can be extended as the quality of the data improves. In a more general sense, the cosmological ‘Standard Model’
 311 is much further from the underlying ‘fundamental theory,’ which will ultimately provide the values of the parameters from
 312 first principles. Nevertheless, any genuinely complete ‘theory of everything’ must include an explanation for the values of
 313 these cosmological parameters as well as the parameters of the Standard Model of particle physics. Current magnitude of
 314 Hubble constant [53-57] is (67.80 ± 0.77) km/sec/Mpc, (68.1 ± 1.2) km/sec/Mpc,
 315 (67.3 ± 1.2) km/sec/Mpc, (69.7 ± 2.0) km/sec/Mpc, (70.0 ± 2.2) km/sec/Mpc, (70.6 ± 3.3) km/sec/Mpc,
 316 (73.8 ± 2.4) km/sec/Mpc, and (72.5 ± 2.5) km/sec/Mpc. In a cosmological approach with various trial-error methods, at
 317 present in hydrogen atom, if $H_0 \cong 71$ km/sec/Mpc, Bohr radius [58] can be fitted as follows.
 318

$$\begin{aligned}
 (a_B)_0 &\cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{GM_0}{c^2} \right) \cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{2H_0} \right) \\
 &\cong \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{2H_0} \right) \cong \frac{1}{2} \left(\frac{4\pi\epsilon_0 G m_p^2}{e^2} \right) \left(\frac{c}{H_0} \right) \\
 &\cong 5.27225 \times 10^{-11} \text{ m.}
 \end{aligned}
 \tag{27}$$

320 $\left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right)$ is the electromagnetic and gravitational force ratio of proton. This relation seems to be very simple and needs
 321 no further derivation. Factor 2 may be connected with half of the current Hubble length $\left(\frac{1}{2} \frac{c}{H_0} \right)$. For any physicist or
 322 cosmologist it will be a very big surprise. Note that, this relation is free from the famous reduced Planck’s constant,
 323 electron rest mass and other arbitrary numbers or coefficients. After simplification and considering the ground state, it is
 324 possible to express the ground state potential energy of electron in the following way.
 325

$$\begin{aligned}
 (E_{\text{pot}})_0 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{4\pi\epsilon_0 G M_0} \right) \cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{1}{2} \frac{c}{H_0} \right)^{-1} \\
 &\cong -2 \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned}
 \tag{28}$$

327 Here $\left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)$ can be called as the current Hubble potential. Characteristic ground state kinetic energy of electron can be
 328 expressed in the following way.

$$\begin{aligned}
 (E_{\text{kin}})_0 &\cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{8\pi\epsilon_0 G M_0} \right) \cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{c^2}{2GM_0} \right) \\
 &\cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned}
 \tag{29}$$

330 Characteristic ground state total energy of electron can be expressed in the following way.
 331

$$\begin{aligned}
 (E_{\text{tot}})_0 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{8\pi\epsilon_0 G M_0} \right) \cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2}{4\pi\epsilon_0} \right) \left(\frac{c^2}{2GM_0} \right) \\
 &\cong - \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right)
 \end{aligned}
 \tag{30}$$

333 If $H_0 \cong 71$ km/sec/Mpc, $(E_{\text{tot}})_0 \cong -13.66$ eV. Based on this coincidence, this proposed new concept can be given some
 334

Requesting your kind and valuable (unbiased) review comments...

335 consideration and it can be suggested that the best value of H_0 lies in between 70 and 71 km/sec/Mpc. Unfortunately
 336 these relations seem to be independent of the reduced Planck's constant [59,60]. If one is willing to linkup these relations
 337 with the observed 'discrete' energy spectrum of the hydrogen atom, then the desired cosmological light emission
 338 mechanism can be developed in a unified picture. Considering the concept of stationary orbits and jumping nature of
 339 electron, emitted photon energy can be expressed in the following way.

$$340 \quad (E_{\text{photon}})_0 \cong \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \quad (31)$$

341 where $n_1 = n_2 \cong 1, 2, 3, \dots$ and $n_2 > n_1$. The best fit of H_0 can be obtained in the following way.

$$342 \quad \left. \begin{aligned} & \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \cong \frac{e^4 m_e}{32\pi^2 \epsilon_0^2 \hbar^2} \\ & \text{and } H_0 \cong \frac{G m_p^2 m_e c}{2\hbar^2} \cong 70.738 \text{ km/sec/Mpc} \end{aligned} \right\} \quad (32)$$

344 At any time in the past - in support of the proposed cosmological red shift interpretation, above relations can be re-
 345 expressed as follows.

$$347 \quad (E_{\text{pot}})_t \cong - \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 c^2}{4\pi\epsilon_0 G M_0} \right) \cong -2 \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (33)$$

$$348 \quad (E_{\text{kin}})_t \cong \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (34)$$

$$351 \quad (E_{\text{tot}})_t \cong - \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \quad (35)$$

352 This can be considered as the base for the 'cosmological thermodynamic light emission mechanism'. At any time in the
 353 past, at any galaxy, emitted photon energy can be expressed as follows.

$$354 \quad (E_{\text{photon}})_t \cong \left(\frac{T_0}{T_t} \right) \left(\frac{e^2}{4\pi\epsilon_0 G m_p^2} \right) \left(\frac{e^2 H_0}{4\pi\epsilon_0 c} \right) \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]_t \quad (36)$$

355 This issue is for further study. In a unified picture, with reference to the current cosmic temperature, electron's current
 356 quantum of angular momentum can be expressed as follows.

$$357 \quad \hbar \cong m_p \sqrt{\frac{G m_e c}{2 H_0}} \cong \frac{G m_p \sqrt{m_e M_0}}{c} \cong \hbar_0 \quad (37)$$

358 If atomic nuclear mass increases in integral multiples of the proton mass, then the observed discreteness of the reduced
 359 Planck's constant can be expressed as follows.

$$361 \quad n\hbar \cong \frac{G(n m_p) \sqrt{m_e M_0}}{c} \cong n\hbar_0 \quad (38)$$

362 where $n = 1, 2, 3, \dots$. This issue is also for further study. At any time in the past, hypothetically, in terms of the current and
 363 past 'primordial' cosmic temperatures, it is possible to express the cosmological 'variable quantum of angular momentum'
 364 of electron in the following way. Whether it is virtual or real or speculative - to be confirmed from further study.

Requesting your kind and valuable (unbiased) review comments...

365

$$\hbar_t \cong \sqrt{\frac{T_t}{T_0}} \cdot \hbar_0 \cong \sqrt{\frac{\lambda_t}{\lambda_0}} \cdot \hbar_0 \quad (39)$$

366

It may be noted that, throughout the cosmic evolution, Planck's constant and the Uncertainty constant both can be considered as 'constants'. Now the fundamental questions to be answered are –

367

368

369

370

371

372

373

374

- 1) Is reduced Planck's constant – an output of the atomic system?
- 2) Is the reduced Planck's constant – a cosmological variable?
- 3) Is the Planck's constant – a cosmological constant?
- 4) How to understand and how to consider the constancy of the Planck's constant along with the variable reduced Planck's constant?
- 5) Is the condition, $\hbar \rightarrow (h/2\pi)$ an indication of saturation or halt of cosmological expansion?

375

376

3. Connecting Cosmic Thermal and Physical Parameters

377

3.1 Cosmic Thermal energy density and Matter energy density

378

It may be noted that connecting CMBR energy density with Hubble's constant is really a very big task and mostly preferred in cosmology. At any given cosmic time, thermal energy density can be expressed with the following semi empirical relation.

379

380

381

$$aT_t^4 \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^{-2} \left(\frac{3H_t^2 c^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-2} \left(\frac{3H_t^2 c^2}{8\pi G} \right) \quad (40)$$

382

$$T_t \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-\frac{1}{2}} \left(\frac{3H_t^2 c^2}{8\pi G a} \right)^{\frac{1}{4}} \quad (41)$$

383

With a suitable derivation if above expression is obtained, then certainly the subject of black hole cosmology is put into main stream physics. Thus at present, if H_0 is close to 71 km/sec/Mpc, obtained CMBR temperature is 2.723 K [53-57]. For the time being this can be considered as a remarkable discovery and an accurate fit.

384

385

386

387

$$aT_0^4 \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-2} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{M_0}{M_s} \right) \right]^{-2} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \quad (42)$$

388

$$T_0 \cong \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-\frac{1}{2}} \left(\frac{3H_0^2 c^2}{8\pi G a} \right)^{\frac{1}{4}} \quad (43)$$

389

With reference to the current cosmic temperature, at any time in the past,

390

$$\frac{T_t}{T_0} \cong \frac{\left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-\frac{1}{2}} H_t}{\left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-\frac{1}{2}} H_0} \quad (44)$$

391

392

Using this relation, cosmic redshift data can be fitted. When the assumed CMBR temperature is 2999 K, estimated redshift is 1099 and is in very good agreement with the standard model of cosmology.

393

394

395

396

Mostly at the ending stage of expansion, rate of change in H_t will be practically zero and can be considered as practically constant. Thus at its ending stage of expansion, for the whole cosmic black hole as H_t practically remains constant, its

Requesting your kind and valuable (unbiased) review comments...

397 corresponding thermal energy density will be ‘the same’ throughout its volume. This ‘sameness’ may be the reason for the
 398 observed ‘isotropic’ nature of the current CMB radiation. With this coincidence it can be suggested that, at the beginning of
 399 cosmic evolution,

$$400 \quad aT_s^4 \cong \left(\frac{3H_s^2 c^2}{8\pi G} \right) \quad (45)$$

401 Matter-energy density can be considered as the geometric mean density of volume energy density and the thermal energy
 402 density and it can be expressed with the following semi empirical relation.

$$403 \quad (\rho_m)_t c^2 \cong \sqrt{\left(\frac{3H_t^2 c^2}{8\pi G} \right) (aT_t^4)} \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-1} \left(\frac{3H_t^2 c^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^{-1} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \quad (46)$$

404 Here one important observation to be noted is that, at any time

$$405 \quad \frac{3H_t^2}{8\pi G (\rho_m)_t} \cong \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right] \cong \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right] \quad (47)$$

406 Thus at present,

$$407 \quad (\rho_m)_0 \cong \frac{1}{c^2} \sqrt{\left(\frac{3H_0^2 c^2}{8\pi G} \right) (aT_0^4)} \cong \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-1} \left(\frac{3H_0^2}{8\pi G} \right) \cong \left[1 + \ln \left(\frac{M_0}{M_s} \right) \right]^{-1} \left(\frac{3H_0^2}{8\pi G} \right) \quad (48)$$

$$\cong 6.6 \times 10^{-32} \text{ gram / cm}^3$$

408 Based on the average mass-to-light ratio for any galaxy present matter density can be expressed with the following relation
 409 [61].

$$411 \quad (\rho_m)_0 \cong 1.5 \times 10^{-32} \eta h_0 \text{ gram/cm}^3 \quad (49)$$

412 Here $\eta \cong \left\langle \frac{M}{L} \right\rangle_{\text{galaxy}} / \left\langle \frac{M}{L} \right\rangle_{\text{sun}}$, $h_0 \cong H_0 / 100 \text{ Km/sec/Mpc} \cong 0.71$ Note that elliptical galaxies probably comprise about

413 60% of the galaxies in the universe and spiral galaxies thought to make up about 20% percent of the galaxies in the
 414 universe. Almost 80% of the galaxies are in the form of elliptical and spiral galaxies. For spiral galaxies, $\eta h_0^{-1} \cong 9 \pm 1$ and
 415 for elliptical galaxies, $\eta h_0^{-1} \cong 10 \pm 2$ For our galaxy inner part, $\eta h_0^{-1} \cong 6 \pm 2$. Thus the average ηh_0^{-1} is very close to 8 to 9
 416 and its corresponding matter density is close to $(6.0 \text{ to } 6.7) \times 10^{-32} \text{ gram/cm}^3$ and can be compared with the above proposed
 417 magnitude of $6.6 \times 10^{-32} \text{ gram/cm}^3$.

3.2 Age of the Growing Cosmic black hole

421 Age of the growing cosmic black hole can be assumed as the time taken to grow from the assumed Stoney scale to the
 422 current scale. At present,

$$423 \quad g_0 \cong \left(\frac{8\pi G a T_0^4}{3H_0^2 c^2} \right) c \cong \left[1 + \ln \left(\frac{M_0}{M_s} \right) \right]^{-2} c \cong \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-2} c \cong 14.66 \text{ km/sec} \quad (50)$$

424 Clearly speaking, at present, Hubble volume is growing at 14.66 km/sec in a decelerating trend. Starting from the Stoney
 425 scale, if the assumed growth rate is gradually decreasing, at any time average growth rate can be expressed as follows.
 427

$$428 \quad \frac{g_s + g_t}{2} \cong \frac{1}{2} \left\{ 1 + \left[1 + \ln \left(\frac{M_t}{M_s} \right) \right]^{-2} \right\} c \cong \frac{1}{2} \left\{ 1 + \left[1 + \ln \left(\frac{H_s}{H_t} \right) \right]^{-2} \right\} c \quad (51)$$

429 For the current scale, average growth rate can be expressed as follows.

$$431 \quad \frac{g_s + g_0}{2} \cong \frac{1}{2} \left\{ 1 + \left[1 + \ln \left(\frac{M_0}{M_s} \right) \right]^{-2} \right\} c \cong \frac{1}{2} \left\{ 1 + \left[1 + \ln \left(\frac{H_s}{H_0} \right) \right]^{-2} \right\} c \quad (52)$$

432 Time taken to reach from the Stoney scale to any assumed scale can be expressed as follows.
 433

Requesting your kind and valuable (unbiased) review comments...

434

435

$$\left(\frac{g_s + g_t}{2}\right)t \cong (R_t - R_s) \cong R_t \quad (53)$$

436

where, $R_t \gg \gg R_s$ and $R_s \approx 0$. Hence for the current scale,

437

438

$$\left(\frac{g_s + g_0}{2}\right)t_0 \cong (R_0 - R_s) \cong R_0 \cong \frac{c}{H_0} \quad (54)$$

439

$$t_0 \cong \left(\frac{g_s + g_0}{2}\right)^{-1} \frac{c}{H_0} \cong \left\{1 + \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2}\right\}^{-1} \frac{2}{H_0} \cong 27.496 \text{ Gyr.} \quad (55)$$

440

where $\left\{1 + \left[1 + \ln\left(\frac{H_s}{H_0}\right)\right]^{-2}\right\}^{-1} \cong 0.99995$. This proposal is for further study. Based on this proposal, after one second from

441

the Stoney scale, cosmic angular velocity is 2 rad/sec, growth rate is 29 km/sec and cosmic temperature is 3×10^9 K.

442

443

With reference to the current and past cosmic temperatures, at any time in the past, at any galaxy, for any hydrogen atom,

444

445

$$\frac{E_0}{E_t} \cong \frac{\lambda_t}{\lambda_0} \cong \frac{T_t}{T_0} \cong \left\{ \frac{\left[1 + \ln\left(\frac{H_s}{H_0}\right)\right] H_t}{\left[1 + \ln\left(\frac{H_s}{H_t}\right)\right] H_0} \right\}^{\frac{1}{2}} \cong \left\{ \frac{\left[1 + \ln\left(\frac{R_0}{R_s}\right)\right] R_0}{\left[1 + \ln\left(\frac{R_t}{R_s}\right)\right] R_t} \right\}^{\frac{1}{2}} \quad (56)$$

446

By guessing H_t , $(z_0 + 1)$ can be estimated. It seems to be a full and absolute definition for the cosmic redshift. Thus at any

447

time in the past,

448

$$\left(\frac{E_0}{E_t} - 1\right) \cong \left(\frac{\lambda_t}{\lambda_0} - 1\right) \cong \left(\frac{T_t}{T_0} - 1\right) \cong \left\{ \frac{\left[1 + \ln\left(\frac{H_s}{H_0}\right)\right] H_t}{\left[1 + \ln\left(\frac{H_s}{H_t}\right)\right] H_0} \right\}^{\frac{1}{2}} - 1 \cong \left\{ \frac{\left[1 + \ln\left(\frac{R_0}{R_s}\right)\right] R_0}{\left[1 + \ln\left(\frac{R_t}{R_s}\right)\right] R_t} \right\}^{\frac{1}{2}} - 1 \cong z_0 \quad (57)$$

449

Please see the following table-1 for the cosmic physical and thermal parameters. This table prepared with C++ program with reference to the observed 2.725 K. In this table:

450

451

452

Column-1 = Assumed cosmic angular velocity.

453

Column-2 = Estimated cosmic radius, from relation (7).

454

Column-3 = Estimated cosmic mass, from relation (7).

455

Column-4 = Estimated cosmic growth index, from relation (10).

456

Column-5 = Estimated cosmic growth rate, from relation (12).

457

Column-6 = Estimated cosmic time, from relation (53).

458

Column-7 = Estimated cosmic temperature, from relation (41)

459

Column-8 = Estimated cosmic redshift, from relation (57)

460

461

Table-1: Assumed Cosmic angular velocity and estimated other cosmic physical and thermal parameters

462

Assumed Cosmic Angular velocity	Estimated Cosmic radius	Estimated Cosmic mass	Cosmic Growth index $\cong \left[1 + \ln\left(\frac{H_s}{H_t}\right)\right]^2$	Estimated Cosmic Growth rate	Estimated Cosmic time	Estimated Cosmic temperature	Estimated Cosmic Redshift z_0
(rad/sec)	(meter)	(kg)	(number)	(km/sec)	(sec)	(K)	(number)
1.086E+44	2.761E-36	1.859E-09	1	299792	0.000E+00	2.237E+32	8.207E+31

Requesting your kind and valuable (unbiased) review comments...

2.305E+43	1.301E-35	8.759E-09	6.50173	46109.6	5.924E-44	6.455E+31	2.368E+31
2.305E+42	1.301E-34	8.759E-08	23.5461	12732.1	8.148E-43	1.480E+31	5.428E+30
2.305E+41	1.301E-33	8.759E-07	51.1943	5855.97	8.493E-42	3.853E+30	1.414E+30
2.305E+40	1.301E-32	8.759E-06	89.4463	3351.65	8.580E-41	1.060E+30	3.888E+29
2.305E+39	1.301E-31	8.759E-05	138.302	2167.66	8.615E-40	3.006E+29	1.103E+29
2.305E+38	1.301E-30	8.759E-04	197.762	1515.93	8.634E-39	8.692E+28	3.189E+28
2.305E+37	1.301E-29	8.759E-03	267.825	1119.36	8.645E-38	2.548E+28	9.347E+27
2.305E+36	1.301E-28	8.759E-02	348.492	860.256	8.653E-37	7.544E+27	2.768E+27
2.305E+35	1.301E-27	8.759E-01	439.763	681.714	8.658E-36	2.251E+27	8.258E+26
2.305E+34	1.301E-26	8.759E+00	541.638	553.492	8.662E-35	6.756E+26	2.479E+26
2.305E+33	1.301E-25	8.759E+01	654.116	458.317	8.665E-34	2.038E+26	7.477E+25
2.305E+32	1.301E-24	8.759E+02	777.199	385.735	8.667E-33	6.173E+25	2.265E+25
2.305E+31	1.301E-23	8.759E+03	910.885	329.122	8.668E-32	1.876E+25	6.883E+24
2.305E+30	1.301E-22	8.759E+04	1055.17	284.116	8.670E-31	5.719E+24	2.098E+24
2.305E+29	1.301E-21	8.759E+05	1210.07	247.748	8.671E-30	1.748E+24	6.411E+23
2.305E+28	1.301E-20	8.759E+06	1375.57	217.941	8.671E-29	5.352E+23	1.964E+23
2.305E+27	1.301E-19	8.759E+07	1551.67	193.207	8.672E-28	1.642E+23	6.025E+22
2.305E+26	1.301E-18	8.759E+08	1738.37	172.456	8.673E-27	5.048E+22	1.852E+22
2.305E+25	1.301E-17	8.759E+09	1935.68	154.877	8.673E-26	1.554E+22	5.701E+21
2.305E+24	1.301E-16	8.759E+10	2143.59	139.855	8.674E-25	4.790E+21	1.757E+21
2.305E+23	1.301E-15	8.759E+11	2362.11	126.917	8.674E-24	1.478E+21	5.424E+20
2.305E+22	1.301E-14	8.759E+12	2591.23	115.695	8.674E-23	4.568E+20	1.676E+20
2.305E+21	1.301E-13	8.759E+13	2830.96	105.898	8.675E-22	1.413E+20	5.184E+19
2.305E+20	1.301E-12	8.759E+14	3081.28	97.2947	8.675E-21	4.375E+19	1.605E+19
2.305E+19	1.301E-11	8.759E+15	3342.21	89.6987	8.675E-20	1.356E+19	4.973E+18
2.305E+18	1.301E-10	8.759E+16	3613.75	82.9588	8.675E-19	4.204E+18	1.542E+18
2.305E+17	1.301E-09	8.759E+17	3895.89	76.951	8.676E-18	1.305E+18	4.786E+17
2.305E+16	1.301E-08	8.759E+18	4188.63	71.5729	8.676E-17	4.052E+17	1.486E+17
2.305E+15	1.301E-07	8.759E+19	4491.98	66.7395	8.676E-16	1.259E+17	4.619E+16
2.305E+14	1.301E-06	8.759E+20	4805.93	62.3797	8.676E-15	3.915E+16	1.436E+16
2.305E+13	1.301E-05	8.759E+21	5130.48	58.4336	8.676E-14	1.218E+16	4.468E+15
2.305E+12	1.301E-04	8.759E+22	5465.64	54.8504	8.676E-13	3.791E+15	1.391E+15
2.305E+11	1.301E-03	8.759E+23	5811.41	51.5869	8.676E-12	1.180E+15	4.331E+14
2.305E+10	1.301E-02	8.759E+24	6167.77	48.6063	8.676E-11	3.678E+14	1.349E+14
2.305E+09	1.301E-01	8.759E+25	6534.74	45.8767	8.676E-10	1.146E+14	4.206E+13
2.305E+08	1.301E+00	8.759E+26	6912.31	43.3708	8.677E-09	3.575E+13	1.311E+13
2.305E+07	1.301E+01	8.759E+27	7300.49	41.0647	8.677E-08	1.115E+13	4.091E+12
2.305E+06	1.301E+02	8.759E+28	7699.27	38.9378	8.677E-07	3.480E+12	1.277E+12
2.305E+05	1.301E+03	8.759E+29	8108.66	36.9719	8.677E-06	1.086E+12	3.985E+11
2.305E+04	1.301E+04	8.759E+30	8528.65	35.1512	8.677E-05	3.392E+11	1.244E+11
2.305E+03	1.301E+05	8.759E+31	8959.24	33.4618	8.677E-04	1.059E+11	3.887E+10
2.305E+02	1.301E+06	8.759E+32	9400.43	31.8913	8.677E-03	3.310E+10	1.214E+10
2.305E+01	1.301E+07	8.759E+33	9852.23	30.4289	8.677E-02	1.035E+10	3.796E+09

Requesting your kind and valuable (unbiased) review comments...

2.305E+00	1.301E+08	8.759E+34	10314.6	29.0648	8.677E-01	3.234E+09	1.187E+09
2.305E-01	1.301E+09	8.759E+35	10787.6	27.7904	8.677E+00	1.011E+09	3.710E+08
2.305E-02	1.301E+10	8.759E+36	11271.3	26.598	8.677E+01	3.163E+08	1.161E+08
2.305E-03	1.301E+11	8.759E+37	11765.5	25.4807	8.677E+02	9.897E+07	3.631E+07
2.305E-04	1.301E+12	8.759E+38	12270.3	24.4324	8.677E+03	3.097E+07	1.136E+07
2.305E-05	1.301E+13	8.759E+39	12785.7	23.4475	8.677E+04	9.693E+06	3.556E+06
2.305E-06	1.301E+14	8.759E+40	13311.7	22.5209	8.677E+05	3.034E+06	1.113E+06
2.305E-07	1.301E+15	8.759E+41	13848.4	21.6482	8.677E+06	9.501E+05	3.486E+05
2.305E-08	1.301E+16	8.759E+42	14395.6	20.8253	8.677E+07	2.976E+05	1.092E+05
2.305E-09	1.301E+17	8.759E+43	14953.4	20.0484	8.677E+08	9.321E+04	3.419E+04
2.305E-10	1.301E+18	8.759E+44	15521.9	19.3142	8.677E+09	2.920E+04	1.071E+04
2.305E-11	1.301E+19	8.759E+45	16100.9	18.6196	8.677E+10	9.150E+03	3.356E+03
2.52E-12	1.19E+20	8.01E+46	16667.6	17.9865	7.94E+11	2998.85	1099.21
2.305E-12	1.301E+20	8.759E+46	16690.6	17.9618	8.677E+11	2.868E+03	1.051E+03
2.305E-13	1.301E+21	8.759E+47	17290.8	17.3382	8.677E+12	8.988E+02	3.288E+02
2.305E-14	1.301E+22	8.759E+48	17901.7	16.7466	8.677E+13	2.818E+02	1.024E+02
2.305E-15	1.301E+23	8.759E+49	18523.2	16.1847	8.677E+14	8.835E+01	3.141E+01
2.305E-16	1.301E+24	8.759E+50	19155.2	15.6507	8.677E+15	2.771E+01	9.164E+00
2.305E-17	1.301E+25	8.759E+51	19797.9	15.1427	8.677E+16	8.689E+00	2.188E+00
2.305E-18	1.301E+26	8.759E+52	20451.2	14.6589	8.677E+17	2.726E+00	0.000E+00

463


464

Please see the below graph for the cosmic growth index for ~ 61 values starting from 1 to 20451.2 of Column-4 in table-1.

465

466

Cosmic Growth Index



467

468

469

470

3.3. Direct fitting of the two current CMBR wavelengths

Requesting your kind and valuable (unbiased) review comments...

471 Note that the spectrum from Planck's law of black body radiation takes a different shape in the frequency domain from that
 472 of the wavelength domain, the frequency location of the peak emission does not correspond to the peak wavelength using
 473 the simple relationship between frequency, wavelength, and the speed of light. In other words, the peak wavelength and the
 474 peak frequency do not correspond. The frequency form of Wien's displacement law is derived using similar methods, but
 475 starting with Planck's law in terms of frequency instead of wavelength. The effective result is to substitute 3 for 5 in the
 476 equation for the peak wavelength. Thus it is possible to say that [62],
 477

$$478 \quad \sqrt{\frac{c}{\lambda_m f_m}} \cong \sqrt{1.75978} \cong 1.326567 \cong \frac{4}{3} \quad (58)$$

479 where λ_m and f_m are the peak wavelength in wavelength domain and peak frequency in frequency domain respectively.

480 Let λ_f is the wavelength corresponding to $\frac{dE_\nu}{d\nu}$ and E_ν is the total energy at all frequencies up to and including ν , at any
 481 given cosmic time. λ_m is the wavelength corresponding to $\frac{dE_\lambda}{d\lambda}$ and E_λ is the total energy at all wavelengths up to and
 482 including λ . Considering the observed CMBR wavelengths, it is possible to express both the wavelengths in the following
 483 way.

$$484 \quad [(\lambda_m)_t \text{ and } (\lambda_f)_t] \propto \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \quad (59)$$

485

$$486 \quad [(\lambda_m)_t \text{ and } (\lambda_f)_t] \propto \sqrt{\left(\frac{4\pi GM_t}{c^2}\right) \cdot \left(\frac{4\pi GM_S}{c^2}\right)} \quad (60)$$

487

488

489

Guessing in this way it is noticed that,

$$490 \quad (\lambda_f)_t \cong \left(\frac{4}{3}\right) \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G\sqrt{M_t M_S}}{c^2} \quad (61)$$

$$\cong \left(\frac{4}{3}\right) \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{4\pi G\sqrt{M_t M_S}}{c^2}$$

$$491 \quad (\lambda_m)_t \cong \left(\frac{3}{4}\right) \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G\sqrt{M_t M_S}}{c^2} \quad (62)$$

$$\cong \left(\frac{3}{4}\right) \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{4\pi G\sqrt{M_t M_S}}{c^2}$$

492

493

Thus it is possible to express both the wavelength relations in the following way.

$$494 \quad (\lambda_f, \lambda_m)_t \cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G\sqrt{M_t M_S}}{c^2} \quad (63)$$

$$\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{H_S}{H_t}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_t}} \cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{\frac{3H_t^2}{8\pi G(\rho_m)_t}} \cdot \frac{2\pi c}{\sqrt{H_S H_t}}$$

495

496

Alternatively geometric mean of $(\lambda_f, \lambda_m)_t$ can be expressed as follows.

Requesting your kind and valuable (unbiased) review comments...

497

$$\begin{aligned} \sqrt{(\lambda_m)_t (\lambda_f)_t} &\cong \sqrt{1 + \ln\left(\frac{M_t}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_t M_S}}{c^2} \\ &\cong \sqrt{1 + \ln\left(\frac{H_S}{H_t}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_t}} \cong \sqrt{\frac{3H_t^2}{8\pi G (\rho_m)_t}} \cdot \frac{2\pi c}{\sqrt{H_S H_t}} \end{aligned} \quad (64)$$

498
499

At present, if H_0 is close to 71 km/sec/Mpc,

500

$$\begin{aligned} (\lambda_f, \lambda_m)_0 &\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{M_0}{M_S}\right)} \cdot \frac{4\pi G \sqrt{M_0 M_S}}{c^2} \\ &\cong \left(\frac{4}{3}\right)^{\pm 1} \cdot \sqrt{1 + \ln\left(\frac{H_S}{H_0}\right)} \cdot \frac{2\pi c}{\sqrt{H_S H_0}} \cong (1.90 \text{ mm}, 1.069 \text{ mm}) \end{aligned} \quad (65)$$

501
502

With reference to $(\lambda_m)_t$ and Wien's displacement constant, from above relations $k_B T_t$ can be expressed as follows.

503

$$\begin{aligned} T_t &\cong \frac{2.898 \times 10^{-3}}{(\lambda_m)_t} \cong \left(\frac{hc}{4.965114 k_B}\right) \left(\frac{1}{(\lambda_m)_t}\right) \text{ and} \\ k_B T_t &\cong \left(\frac{4}{3x}\right) \sqrt{\left(1 + \ln\left(\frac{M_t}{M_S}\right)\right)^{-1} \left(\frac{M_t}{M_S}\right)} \cdot \left(\frac{hc^3}{4\pi G M_t}\right) \end{aligned} \quad (66)$$

504

where $x \cong 4.965114$.

505

$$k_B T_t \propto \left(\frac{hc^3}{4\pi G M_t}\right) \cong \frac{h H_t}{2\pi} \cong h \left(\frac{H_t}{2\pi}\right) \quad (67)$$

506
507

This relation may not be identical but similar to the famous Hawking's black hole temperature formula [63].

508

$$k_B T_t \propto \sqrt{\left(1 + \ln\left(\frac{M_t}{M_S}\right)\right)^{-1} \left(\frac{M_t}{M_S}\right)} \quad (68)$$

509
510
511

In this way in a very simple approach observed CMBR and the proposed Black hole universe concepts can be put into single frame of reference. Here the very interesting and strange observation is that, at present

512

$$\left(1 + \ln\left(\frac{M_0}{M_S}\right)\right)^{-1} \left(\frac{M_0}{M_S}\right) \cong \exp\left(\frac{1}{\alpha}\right) \quad (69)$$

513

where $\left(\frac{1}{\alpha}\right)$ is the inverse of the fine structure ratio. For any mathematician this seems to be a fun. For a cosmologist it

514
515

may be an accidental coincidence. For any physicist it is an astounding and exciting coincidence. Even though it depends upon one's own choice of scientific interest, from unification point of view, assuming it to be a cosmological variable it is

516

possible to express $\left(\frac{1}{\alpha}\right)$ in the following way.

517

$$\left(\frac{1}{\alpha}\right)_0 \cong \ln \left[\left(1 + \ln\left(\frac{M_0}{M_S}\right)\right)^{-1} \left(\frac{M_0}{M_S}\right) \right] \cong 137.047 \quad (70)$$

518

Here $\left(\frac{1}{\alpha}\right)_0$ may be considered as the current magnitude of 'inverse of the fine structure ratio. In atomic and nuclear physics,

519
520

the fine-structure ratio (α) is a fundamental physical constant namely the coupling constant characterizing the strength [64-66] of the electromagnetic interaction. Being a dimensionless quantity, it has a constant numerical value in all systems

Requesting your kind and valuable (unbiased) review comments...

521 of units. Note that, from unification point of view, till today role of dark energy or dark matter is unclear and undecided.
522 Their laboratory or physical existence is also not yet confirmed. In this critical situation this application or coincidence
523 can be considered as a key tool in particle cosmology. Based on the above heuristic observation and for the assumed initial
524 conditions of the universe, if $M_t \rightarrow M_S$, $\left(\frac{1}{\alpha}\right)_S \rightarrow 0$. Based on the relation (70), if one is willing to consider the cos-
525 mological variable nature of $\left(\frac{1}{\alpha}\right)$, relation (66) can be expressed as follows.

$$T_t \cong \sqrt{\left(\frac{1}{e^\alpha}\right)_t} \cdot \left(\frac{bc^2}{3\pi GM_t}\right) \quad (71)$$

528 At the beginning of cosmic evolution for the Stoney scale,

$$T_S \cong \left(\frac{bc^2}{3\pi GM_S}\right) \quad (72)$$

530 From now onwards, CMBR temperature can be called as '**Comic Black Hole's Thermal Radiation**' temperature and can
531 be expressed as '**CBHTR**' temperature. From ground based laboratory experiments, it is possible to measure the rate of
532 change in $\frac{d}{dt}\left(\frac{1}{\alpha_t}\right)$. Hence the absolute cosmic rate of expansion can be measured. Thus at any time based on

533 $\left[\frac{d}{dt}[(\lambda_m)_t \text{ and } (\lambda_f)_t], \frac{d}{dt}(T_t) \text{ and } \frac{d}{dt}(H_t)\right]$, the absolute cosmic rate of expansion can be confirmed. At present with
534 reference to $\left[\frac{d}{dt}[(\lambda_m)_0 \text{ and } (\lambda_f)_0], \frac{d}{dt}(T_0) \text{ and } \frac{d}{dt}(H_0)\right]$ current 'true' cosmic rate of expansion can be understood.

535 Drop in current 'cosmic temperature' can be considered as a measure of the current cosmic expansion and 'rate of decrease
536 in current cosmic temperature' can be considered as a measure of the current cosmic 'rate of expansion'. But if rate of
537 decrease in temperature is very small and is beyond the scope of current experimental verification, then the two possible
538 states are: a) cosmic temperature is decreasing at a very slow rate and universe is expanding at a very slow rate and b) there
539 is no 'observable' thermal expansion and there is no 'observable' cosmic expansion. If observed CMBR temperature is
540 2.725 K and is very low in magnitude and is very close to absolute zero, then thinking about and confirming the 'cosmic
541 acceleration' may not be reasonable. Similarly 'rate of decrease in current 'Hubble's constant' can be considered as a
542 measure of current cosmic 'rate of expansion'. If rate of decrease in current 'Hubble's constant is very small and is beyond
543 the scope of current experimental verification, then the two possible states are: a) current 'Hubble's constant is decreasing
544 at a very slow rate and current universe is expanding at a very slow rate and b) at present there is no 'observable' cosmic
545 expansion. Fortunately as per the Cobe/Planck satellite data current CMBR temperature is very smooth and isotropic. and
546 there is no data that refers to the rate of change in the current Hubble's constant. Hence it can be suggested that at present
547 there is no significant cosmic expansion. Even though this suggestion is completely against to the current notion of cosmic
548 acceleration [32,33], based on the proposed arguments, relations and observed data authors request the science
549 community to review the standard cosmology. If observed CMB radiation temperature is 2.725 K and is very low in
550 magnitude and is very close to absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be
551 reasonable.
552

553 **4. To understand the physical significance of large numbers in cosmology**

554
555 Great cosmologists proposed many interesting large numbers in cosmology [67-74]. Ultimately the essence of any
556 cosmological number or ratio is to connect the microscopic and macroscopic physical constants with a possible physical
557 meaning with in the 'evolving universe'. Clearly speaking large dimensionless constants and compound physical constants
558 must reflect an 'observable' intrinsic property of any natural physical phenomenon. Then only the real meaning of any
559 cosmological number can be explored. In this regard authors proposed many interesting relations in the previous sections of
560 this paper. Authors noticed that uncertainty relation or Planck's constant or reduced Planck's constant or inverse of the Fine
561 structure ratio or characteristic nuclear potential radius or rms radius of proton or classical radius of electron - play a

Requesting your kind and valuable (unbiased) review comments...

562 crucial role in the understanding the halt of cosmic expansion. The basic questions to be answered are: 1) The general idea
563 of large number coincidence is interesting, yet is there any observational proves? and 2) How Einstein's general theory of
564 relativity is fitted in the theory of the large cosmological numbers? In this regard the characteristic and key relation can be
565 expressed in the following way.

$$566 \quad \frac{c^3}{2GM_0} \cong H_0 \quad \text{Or} \quad \frac{c^3}{2GH_0} \cong M_0 \quad (73)$$

567 Here (M_0, H_0) can be considered as the current mass and current angular velocity of the black hole universe respectively.
568 By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.
569

$$570 \quad \frac{c^3}{2GM_{sat}} \cong H_{sat} \quad \text{Or} \quad \frac{c^3}{2GH_{sat}} \cong M_{sat} \quad (74)$$

571 Here (M_{sat}, H_{sat}) can be considered as the saturated mass and saturated angular velocity of the black hole universe at its
572 ending stage of expansion. Fortunately it is noticed that, $M_{sat} \cong M_0$ and $H_{sat} \cong H_0$. Authors strongly believe that the
573 following relations certainly help in understanding the mystery of the halting of the present cosmic expansion.
574

4.1 Role of the Uncertainty relation

575
576 It is noticed that,
577
578

$$579 \quad \frac{Gm_p m_e}{R_p H_0} \cong \frac{h}{4\pi} \quad (75)$$

580 Here $R_p \cong (0.84184 \text{ to } 0.87680) \text{ fm}$ is the rms radius of proton [75,76]. After re-arranging, it can be expressed in the
581 following way.
582

$$583 \quad \left(\frac{2Gm_p}{c^2 R_p} \right) \frac{m_e c^2}{H_0} \cong \left(\frac{2Gm_p}{c^2 R_p} \right) \left[m_e c \left(\frac{2\pi c}{H_0} \right) \right] \cong h \quad (76)$$

584 By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.
585
586

$$587 \quad H_{sat} \Rightarrow \frac{4\pi Gm_p m_e}{hR_p} \cong \frac{Gm_p m_e}{(h/4\pi)R_p} \quad (77)$$
$$\Rightarrow H_{sat} \cong (67.87 \text{ to } 70.69) \text{ km/sec/Mpc}$$

588 This is a remarkable fit and needs further study.
589

4.2 Role of the classical radius of electron

590
591 It is noticed that,
592
593

$$594 \quad \sqrt{\left(\frac{2G\sqrt{m_p m_e}}{c^2} \right) \left(\frac{c}{H_0} \right)} \cong \sqrt{\left(\frac{2G\sqrt{m_p m_e}}{c^2} \right) \left(\frac{2GM_0}{c^2} \right)} \cong \left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right) \quad (78)$$

595 $\left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right)$ is nothing but the presently believed classical radius of electron. In a broad picture or considering the
596 interaction in between proton and electron it is a very general idea to consider the geometric mean mass of proton and
597 electron. By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as
598 follows.

Requesting your kind and valuable (unbiased) review comments...

599

$$\left(\frac{c}{H_{sat}}\right) \Rightarrow \left(\frac{e^2}{4\pi\epsilon_0 m_e c^2}\right)^2 \left(\frac{c^2}{2G\sqrt{m_p m_e}}\right) \quad (79)$$

600

$$H_{sat} \Rightarrow \frac{2G\sqrt{m_p m_e}}{c} \left(\frac{4\pi\epsilon_0 m_e c^2}{e^2}\right)^2 \cong 67.533 \text{ km/sec/Mpc} \quad (80)$$

601

This is also a remarkable fit and needs further study.

602

4.3 Role of the characteristic nuclear potential radius

603

604

605

It is noticed that,

606

$$\frac{G\sqrt{M_0\sqrt{m_p m_e}}}{c^2} \cong \sqrt{\left(\frac{GM_0}{c^2}\right)\left(\frac{G\sqrt{m_p m_e}}{c^2}\right)} \cong 1.4 \times 10^{-15} \text{ m} \cong R_n \quad (81)$$

607

R_n is nothing but the presently believed characteristic nuclear potential radius [77] or the nuclear strong interaction range as proposed by Yukawa [78]. By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows [79-81].

608

609

610

611

$$\frac{G\sqrt{M_{sat}\sqrt{m_p m_e}}}{c^2} \Rightarrow R_n \quad (82)$$

612

$$H_{sat} \Rightarrow \frac{G\sqrt{m_p m_e}}{2cR_n^2} \quad (83)$$

613

This is also a remarkable coincidence and accuracy mainly depends upon the magnitude of the characteristic nuclear potential radius. Further study may reveal the mystery.

614

615

616

4.4 Role of the ‘inverse’ of the Fine structure ratio

617

618

619

620

Total thermal energy in the present Hubble volume can be expressed as follows.

621

$$(E_T)_0 \cong aT_0^4 \cdot \frac{4\pi}{3} \left(\frac{c}{H_0}\right)^3 \quad (84)$$

622

623

Thermal energy present in half of the current Hubble volume can be expressed as follows.

624

$$\frac{(E_T)_0}{2} \cong \frac{1}{2} \left[aT_0^4 \cdot \frac{4\pi}{3} \left(\frac{c}{H_0}\right)^3 \right] \quad (85)$$

625

626

If (c/H_0) is the present electromagnetic interaction range, then present characteristic Hubble potential can be expressed as

627

$$(E_e)_0 \cong \frac{e^2}{4\pi\epsilon_0 (c/H_0)} \cong \frac{e^2 H_0}{4\pi\epsilon_0 c} \quad (86)$$

628

629

630

If H_0 is close to 71 km/sec/Mpc and $T_0 \cong 2.725 \text{ K}$, it is noticed that,

631

$$\ln \sqrt{\frac{[(E_T)_0/2]}{(E_e)_0}} \cong 137.05 \quad (87)$$

Requesting your kind and valuable (unbiased) review comments...

632
633
634

By this time if the expanding black hole universe is coming to a halt, then above relation can be re-expressed as follows.

635

$$\ln \sqrt{\frac{[(E_T)_0/2]}{(E_e)_0}} \cong \ln \sqrt{\frac{[(E_T)_{sat}/2]}{(E_e)_{sat}}} \Rightarrow \left(\frac{1}{\alpha}\right) \quad (88)$$

636

$(E_T)_{sat}$ can be considered as the total thermal energy in the Hubble volume at the end of cosmic expansion.

637

$(E_e)_{sat}$ can be considered as the Hubble potential at the end of cosmic expansion.

638

639

640

5. To fit the nuclear charge radius and the Planck's constant

641

The subject of final unification is having a long history. After the nucleus was discovered [77] in 1908, it was clear that a new force was needed to overcome the electrostatic repulsion of the positively charged protons. Otherwise the nucleus could not exist. Moreover, the force had to be strong enough to squeeze the protons into a volume of size 10^{-15} meter. In general the word 'strong' is used since the strong interaction is the "strongest" of the four fundamental forces. Its observed strength is around 10^2 times that of the electromagnetic force, some 10^5 times as great as that of the weak force, and about 10^{39} times that of gravitation.

642

643

644

645

646

647

The aim of unification is to understand the relation that connects 'gravity', 'mass', 'charge' and the 'microscopic space-time curvature'. Many scientists addressed this problem in different ways [79-81]. The authors also made many attempts in their previously published papers [82-85]. Experimentally observed nuclear charge radius R_{ch} can be fitted with the following strange and simple unified relation.

648

649

650

651

652

$$R_{ch} \cong \sqrt{\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{2GM_s}{c^2}\right)} \cong 1.252 \text{ fermi} \quad (89)$$

653

Considering the rest energy of proton and 1.25 fermi, semi empirical mass formula energy coefficients can be fitted very easily.

654

655

656

$$\frac{R_{ch} c^2}{2GM_s} \cong \sqrt{\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right)} \quad (90)$$

657

Whether the expression $\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cong 90.62$ playing a 'key unified role' or 'only a fitting role' to be confirmed.

658

659

With a great accuracy the famous Planck's constant can be fitted with the following relation.

660

$$\begin{aligned} h &\cong \frac{1}{2} \ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \cdot \left(\sqrt{m_p m_e} \cdot c \cdot R_{ch}\right) \\ &\cong \ln\sqrt{\frac{e^2}{4\pi\epsilon_0 G m_p m_e}} \cdot \left(\sqrt{m_p m_e} \cdot c \cdot R_{ch}\right) \cong 6.63862 \times 10^{-34} \text{ J.sec} \end{aligned} \quad (91)$$

661

Recommended value of h is $6.6260695729 \times 10^{-34}$ J.sec and the error is 0.189%. Now above relation can be simplified into the following form [75].

662

663

664

$$h \cong \left[\ln\left(\frac{e^2}{4\pi\epsilon_0 G m_p m_e}\right) \right]^{3/2} \left(\frac{e^2}{4\pi\epsilon_0 c}\right) \quad (92)$$

665

Connecting quantum constants and gravity is really a very big task. At this juncture this relation can be given a chance. It casts a doubt on the independent existence of quantum mechanics. With this relation, obtained magnitude of the

666

667

Requesting your kind and valuable (unbiased) review comments...

668 gravitational constant is, $G \cong 7.48183566 \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \text{sec}^{-2}$. Independent of ‘length’, ‘force’ and other physical
669 considerations, with this relation order of magnitude of G can be confirmed from atomic physical constants. To proceed
670 further - at first the hierarchy of physical constants must be established and it needs further study and analysis.
671

672 **6. Conclusions**

673 **6.1 Need of the mass unit $M_S \cong \sqrt{e^2/4\pi\epsilon_0 G}$ in unification**

674 The basic idea of unification is – 1) To minimize the number of physical constants and to merge a group of different
675 fundamental constants into one compound physical constant with appropriate unified interpretation and 2) To merge and
676 minimize various branches of physics. In this regard instead of Planck mass, $M_S \cong \sqrt{e^2/4\pi\epsilon_0 G}$ can be considered as the
677 nature’s given true unified mass unit. Using this mass unit, proton-electron mass ratio and proton rest mass can be fitted in
678 the following way.
679
680

$$681 \quad \ln \sqrt{\frac{m_p}{m_e}} \cdot \left(\frac{m_p^2}{m_e}\right) \cong (M_S m_e^2)^{\frac{1}{3}} \quad (93)$$

$$682 \quad \ln \sqrt{\frac{m_p}{m_e}} \cdot \left(\frac{m_p}{m_e}\right) \cong \frac{(M_S m_e^2)^{\frac{1}{3}}}{m_p} \quad (94)$$

683 Here, lhs=6908.3745 and rhs=6899.7363. Accuracy can be improved with the following relation.
684
685

$$686 \quad \frac{(M_S m_e^2)^{\frac{1}{3}}}{m_p} \cong \left[\left(\frac{m_p}{m_e}\right) \ln \sqrt{\frac{m_p}{m_e}} \right] + \ln \left[\left(\frac{m_p}{m_e}\right) \ln \sqrt{\frac{m_p}{m_e}} \right] \quad (95)$$

687 Interesting observation is that $\ln \left[\frac{(M_S m_e^2)^{\frac{1}{3}}}{m_p} \right] \cong \ln(6900) \cong 8.84$ and is close to the presently believed inverse of the strong
688 coupling constant α_s [53]. From the above relation, magnitude of the gravitational constant [57,86,87] can be fitted in the
689 following way.
690

$$691 \quad \left. \begin{aligned} &\text{If } X \cong \left(\frac{m_p}{m_e}\right) \ln \sqrt{\frac{m_p}{m_e}} \text{ and } M_S \cong [X + \ln(X)]^3 \left(\frac{m_p^3}{m_e^2}\right) \\ &G \cong \frac{e^2}{4\pi\epsilon_0 M_S^2} \cong 6.672682478 \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \text{sec}^{-2} \end{aligned} \right\} \quad (96)$$

692 Now the strong coupling constant can be fitted with the following relation.
693
694

$$695 \quad \exp\left(\frac{1}{\ln(X)}\right) - 1 \cong \alpha_s \cong 0.11978 \quad (97)$$

696 **6.2 To consider the universe as a growing and light speed rotating primordial black hole**

698 If ‘black hole geometry’ is more intrinsic compared to the black hole ‘mass’ and ‘density’ parameters, if universe
699 constitutes so many galaxies and if each galaxy constitutes a central growing and fast spinning black hole then considering
700 universe as a ‘growing and light speed rotating primordial black hole’ may not be far away from reality. If universe is
701 having no black hole geometry - any massive body (which is bound to the universe) may not show a black hole structure.
702 That is black hole structure or geometry may be a subset of the cosmic geometry. At this juncture considering or rejecting

Requesting your kind and valuable (unbiased) review comments...

703 this proposal completely depends on the observed cosmic redshift. Based on the relations proposed in sections 2 and 4
704 observed cosmic redshift can be considered as a result of cosmological light emission mechanism. Authors are working on
705 the assumed Hubble volume and Hubble mass in different directions with different applications [1-13] that connect micro
706 physics and macro physics. Based on the proposed applications and shortcomings of the standard model of cosmology -
707 concepts of black hole cosmology may be given at least 99% priority.
708

709 **6.3 About the current cosmic black hole's deceleration**

710 In view of the applications proposed in sections (2) to (4) and with reference to the zero rate of change in inverse of the
711 fine structure ratio (from ground based experiments), zero rate of change in the 'current CMBR temperature' (from
712 Cobe/Planck satellite data) and zero rate of change in the 'current Hubble's constant' (from Cobe/Planck satellite data) it
713 can be suggested that, current cosmic expansion is almost all saturated and at present there is no significant cosmic
714 acceleration [47,48]. Clearly speaking, Stoney scale cosmic black hole's growth rate is equal to the speed of light and
715 current cosmic black hole is growing at 14.66 km/sec in a decelerating trend. It can be also be possible to suggest that
716 currently believed 'dark energy' is a pure, 'mathematical concept' and there exists no physical base behind its confirmation.
717 Now the key leftover things are nucleosynthesis and structure formation. Authors are working in this direction. As nuclear
718 binding energy was zero at the beginning of cosmic evolution, by considering the time dependent variable nature of
719 magnitudes of the semi empirical mass formula energy coefficients it is possible to show that, at the beginning of formation
720 of nucleons, nuclear stability is maximum for light atoms only. If so it can be suggested that, from the beginning of
721 formation of nucleons, in any galaxy, maximum scope is being possible only for the survival of light atoms and this may be
722 the reason for the accumulation and abundance of light atoms in large proportion.
723

724 **Acknowledgements**

725 The first author is indebted to professor K. V. Krishna Murthy, Chairman, Institute of Scientific Research on Vedas
726 (I-SERVE), Hyderabad, India and Shri K. V. R. S. Murthy, former scientist IICT (CSIR) Govt. of India, Director, Research
727 and Development, I-SERVE, for their valuable guidance and great support in developing this subject. Both the authors are
728 very much thankful to the anonymous referees for their valuable comments and kind suggestions in improving and bringing
729 this subject into current main stream physics research.
730

731 **References**

- 732
- 733 [1] U. V. S. Seshavatharam, S. Lakshminarayana. Basics of black hole cosmology – first critical scientific review.
734 Physical Science International journal, Vol-4, Issue-6, p.842-879. (2014)
 - 735 [2] Seshavatharam, U. V. S. & Lakshminarayana, S., On the Role of Hubble Volume in Black Hole Cosmology & Final
736 Unification. Prespacetime Journal, February 2014, Volume 5, Issue 2, pp. 148-173.
 - 737 [3] U. V. S. Seshavatharam. Physics of rotating and expanding black hole universe. Progress in Physics. April, p 7-14,
738 (2010).
 - 739 [4] U.V.S. Seshavatharam. The Primordial Cosmic Black Hole and the Cosmic Axis of Evil. International Journal of
740 Astronomy, 1(2): 20-37, (2012).
 - 741 [5] Seshavatharam, U. V. S, Lakshminarayana, S. Applications of Hubble Volume in Atomic Physics, Nuclear Physics,
742 Particle Physics, Quantum Physics and Cosmic Physics. Journal of Nuclear Physics, Material Sciences, Radiation and
743 Applications Vol. 1, No. 1, August 2013 pp. 45–60.
 - 744 [6] Seshavatharam U.V. S, Lakshminarayana. S, To confirm the existence of Black hole cosmology. International Journal
745 of Advanced Astronomy, 2 (1), 21-36, 2013
 - 746 [7] U. V. S. Seshavatharam, S. Lakshminarayana, Hubble Volume and the Fundamental Interactions, International Journal
747 of Astronomy, Vol. 1 No. 5, 2012, pp. 87-100.
 - 748 [8] U. V. S. Seshavatharam, S. Lakshminarayana, B.V.S.T. Sai. Is red shift an index of galactic 'atomic light emission'
749 mechanism? International Journal of Physics, Vol. 1, No.3, 49-64, (2013).
 - 750 [9] U. V. S. Seshavatharam, S. Lakshminarayana Microscopic Physical Phenomena in Black Hole Cosmos Rotating at
751 Light Speed. Prespacetime Journal. October 2013, Volume 4, Issue 9, pp. 884-922.
 - 752 [10] U. V. S. Seshavatharam, S. Lakshminarayana. Black Hole Cosmology: A Biological Boom. Journal of Astrobiology
753 and Outreach, 2014, 2:1. <http://dx.doi.org/10.4172/2332-2519.1000108>.
 - 754 [11] U. V. S. Seshavatharam, S. Lakshminarayana. Basic Interactions in Black Hole Cosmology. American Journal of
755 Astronomy and Astrophysics. Vol. 2, No. 1, 2013, pp. 6-17. doi: 10.11648/j.ajaa.20140201.12
 - 756 [12] U. V. S. Seshavatharam, S. Lakshminarayana. Friedman Cosmology: Reconsideration and New Results, International
757 Journal of Astronomy, Astrophysics and Space Science. Vol. 1, No. 2, 2014, pp. 16-26.

Requesting your kind and valuable (unbiased) review comments...

- 758 [13] Seshavatharam, U. V. S. and Lakshminarayana, S. The Reduced Planck's Constant, Mach's Principle, Cosmic
759 Acceleration and the Black Hole Universe. *Journal of Physical Science and Application*. Vol.2 (10) 441-447. (2012)
- 760 [14] Friedmann, A. On the Curvature of Space. *General Relativity and Gravitation* 31 (12): 1991-2000. 1999
- 761 [15] Pathria, R. K. The Universe as a Black Hole. *Nature* 240 (5379):298-299.doi:10.1038/240298a0 (1972).
- 762 [16] Good, I. J. Chinese universes. *Physics Today* 25 (7): 15. July. doi:10.1063/1.3070923 (1972).
- 763 [17] Joel Smoller and Blake Temple. Shock-wave cosmology inside a black hole. *Proc Natl Acad Sci U S A*. September 30;
764 100(20): 1121611218. (2003).
- 765 [18] Chul-Moon Yoo et al. Black Hole Universe. Time evolution. *Phys. Rev. Lett.* 111, 161102 (2013).
- 766 [19] Michael E. McCulloch. A Toy Cosmology Using a Hubble-Scale Casimir Effect. *Galaxies* 2014, 2, 81-88.
- 767 [20] T.X. Zhang and C. Frederic. Acceleration of black hole universe. *Astrophysics and Space Science*, Volume 349, Issue
768 1, pp 567-573. (2013).
- 769 [21] Zhang, Tianxi. Cosmic microwave background radiation of black hole universe. *Astrophysics and Space Science*,
770 Volume 330, Issue 1, pp 157-165. (2010).
- 771 [22] Zhang, Tianxi. Quasar Formation and Energy Emission in Black hole universe
772 <http://downloads.hindawi.com/journals/aa/2012/625126.pdf>. *Progress in Physics*, 3: 48-53, (2012).
- 773 [23] Poplawski, N. J. Radial motion into an Einstein-Rosen bridge. *Physics Letters B* 687 (23): 110-113. (2010).
- 774 [24] Poplawski, N. J. Big bounce from spin and torsion. *General Relativity and Gravitation* Vol. 44, No. 4 (2012) pp.
775 1007–1014.
- 776 [25] Poplawski, N. J. Energy and momentum of the Universe. *Class. Quantum Grav.* 31, 065005 (2014).
- 777 [26] Pourhasan R, Afshordi N and Mann R.B. Did a hyper black hole spawn the universe? *Nature - International weekly*
778 *journal of science*. 13 September 2013, doi:10.1038/nature.2013.13743, arXiv: 1309. 1487v2.
- 779 [27] Andy Gardner, Joseph P. Conlon. Cosmological natural selection and the purpose of the universe. *Complexity*.
780 Vol.18, Issue 5, pp48-56. 2013
- 781 [28] Smolin, L. Cosmological natural selection as the explanation for the complexity of the universe. *Physica A* 340, 705-
782 713. 2004.
- 783 [29] Hawking S.W. A Brief History of Time. Bantam Dell Publishing Group. 1988
- 784 [30] Hubble E. P, A relation between distance and radial velocity among extra-galactic nebulae, *PNAS*, 1929, vol. 15, 1929,
785 pp.168-173.
- 786 [31] Hubble, E.P, The 200-inch telescope and some problems it may solve. *PASP*, 59, pp153-167, 1947.
- 787 [32] J. A. Frieman et al. Dark energy and the accelerating universe. *Ann.Rev.Astron.Astrophys.*46, 2008, p 385.
- 788 [33] The Accelerating Universe. The Royal Swedish Academy of sciences. 2011 Nobel prize in physics.
789 www.nobelprize.org/nobel_prizes/physics/laureates/2011/advanced-physicsprize2011.pdf
- 790 [34] Hawking, S.W.; Ellis, G.F.R. (1973). *The Large-Scale Structure of Space-Time*. Cambridge University Press.
- 791 [35] G.J. Stoney, On the Physical Units of Nature. *Phil.Mag.* 11 (1881) 381-91.
- 792 [36] Michael J. Longo, Detection of a Dipole in the Handedness of Spiral Galaxies with Redshifts $z < 0.04$, *Phys. Lett. B* 699,
793 224-229 2011.
- 794 [37] S.-C. Su and M.-C. Chu. Is the universe rotating? *Astrophysical Journal*, 703 354. 2009.
- 795 [38] J. D. McEwen et al. Bayesian analysis of anisotropic cosmologies: Bianchi VIIIh and WMAP. *Mon. Not. R. Astron.*
796 *Soc.* 000, 1–15 (2013). arXiv:1303.3409v1.
- 797 [39] L. M. Chechin. On the Modern Status of the Universe Rotation Problem. *Journal of Modern Physics*, 2013, 4, 126-132.
- 798 [40] C Sivaram and Kenath Arun, Primordial Rotation of the Universe, Hydrodynamics, Vortices and Angular Momenta of
799 Celestial Objects. *The Open Astronomy Journal*, 2012, 5, 7-11
- 800 [41] Sidharth, B.G. Is the Universe Rotating? *Prespacetime Journal*. October 2010, Vol. 1, Issue 7, pp. 1168-1173.
- 801 [42] Marcelo Samuel Berman, Fernando de Mello Gomide. Local and Global Stability of the Universe. *Journal of Modern*
802 *Physics*, 2013, 4, 7-9
- 803 [43] Robert V Gentry. New Cosmic Center Universe Model Matches Eight of Big Bang's Major Predictions Without The
804 F-L Paradigm. CERN preprint, EXT-2003-022, 14 Apr 2003.
- 805 [44] G. Chapline et al. Tommy Gold Revisited: Why Does Not The Universe Rotate? *AIP Conf.Proc.*822:160-165, 2006.
806 <http://arxiv.org/abs/astro-ph/0509230>.
- 807 [45] Dmitri Rabounski. On the Speed of Rotation of Isotropic Space: Insight into the Redshift Problem. *The Abraham*
808 *Zelmanov Journal*, Vol. 2, 2009, 208-223.
- 809 [46] Kurt Godel. Rotating Universes in General Relativity Theory. *Proceedings of the international Congress of*
810 *Mathematicians in Cambridge*, 1: 175-81, 1950.
- 811 [47] S.W. Hawking. On the rotation of the universe. *Mon. Not. Royal. Astr. Soc.* 142, 129-141.1969.
- 812 [48] M. Novello and M. J. Reboucas. Rotating universe with successive causal and noncausal regions. *Phys. Rev. D* 19,
813 2850-2852 (1979)
- 814 [49] Barrow J D, Juszkiewicz R, Sonoda DH. Universal rotation - How large can it be? *Mon. Not. R. Astron. Soc.* 1985;
815 213: 917.

Requesting your kind and valuable (unbiased) review comments...

- 816 [50] Christopher S. Reynolds. Astrophysics: Black holes in a spin. *Nature*. 494, 432–433 (28 February 2013)
- 817 [51] U. V. S. Seshavatharam. Light speed rotating black holes: The special holes *International Journal of Advanced*
- 818 *Astronomy*, 1 (1) (2013) 13-20.
- 819 [52] Louis Marmet. On the Interpretation of Red-Shifts: A Quantitative Comparison of Red-Shift Mechanisms.
- 820 www.marmet.org/louis/index.html
- 821 [53] J. Beringer et al. Particle Data Group. *Phys. Rev. D*86, 010001 (2012)
- 822 [54] C. L. Bennett et al, Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and
- 823 Results. Submitted to *Astrophysical Journal Supplement Series*. <http://arxiv.org/abs/1212.5225v1>
- 824 [55] David N. Spergel et al. Planck Data Reconsidered. <http://arxiv.org/pdf/1312.3313.pdf>
- 825 [56] W. L. Freedman et al. Final Results from the Hubble Space Telescope Key Project to Measure the Hubble Constant.
- 826 *The Astrophysical Journal* 553 (1): 47-72. 2001.
- 827 [57] P.J. Mohr, B.N. Taylor, and D.B. Newell in [arXiv:1203.5425](http://arxiv.org/abs/1203.5425) and *Rev. Mod. Phys.* (to be published).
- 828 <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-phys-constants.pdf>
- 829 [58] N. Bohr. On the Constitution of Atoms and Molecules. (Part-1) *Philos. Mag.* 26, 1913, p 1.
- 830 [59] Abdus Salam. Einstein's Last Dream: The Space -Time Unification of Fundamental Forces, *Physics News*, 12(2):36,
- 831 June 1981.
- 832 [60] David Gross, Einstein and the search for Unification. *Current science*, Vol. 89, No. 2005, p 12.
- 833 [61] J. V. Narlikar. *Introduction to cosmology*. Cambridge Univ Press, 2002.
- 834 [62] Lianxi Ma et al. Two forms of Wien's displacement law. *Lat. Am. J. Phys. Educ.* Vol. 3, No. 3, Sept. 2009
- 835 [63] Hawking S.W. Particle creation by black holes. *Commun. Math. Phys.*, 1975, v.43, 199–220.
- 836 [64] Gaurab Ganguly et al. SeD Radical: A probe for measurement of time variation of Fine Structure Constant (α) and
- 837 Proton to Electron Mass Ratio (μ). <http://arxiv.org/pdf/1403.4061v2.pdf>.
- 838 [65] J.K. Webb et al. Indications of a spatial variation of the fine structure constant. *Physical Review letters*, 107 (19) 2011
- 839 [66] Srikanand R. et al., Time Variation of the Fine Structure Constant. *The Messenger*. No.116, 25-28 (2004)
- 840 [67] P. A. M. Dirac. The cosmological constants. *Nature*, 139, 1937, p 323.
- 841 [68] P. A. M. Dirac. A new basis for cosmology. *Proc. Roy. Soc. A* 165, 1938, p 199.
- 842 [69] Brandon Carter. Large number coincidences and the anthropic principle in cosmology. *General Relativity and*
- 843 *Gravitation*, Volume 43, Issue 11, pp 3225-3233, (2011)
- 844 [70] Ross A. McPherson. The Numbers Universe: An Outline of the Dirac/Eddington Numbers as Scaling Factors for
- 845 Fractal, Black Hole Universes. *EJTP* 5, No. 18 (2008) 81–94;
- 846 [71] Scott Funkhouser. A new large-number coincidence and a scaling law for the cosmological constant. *Proc. R. Soc. A* 8
- 847 *Mav* 2008 vol. 464 no. 20931345-1353:
- 848 [72] Barrow. J.D. *The Constants of Nature From Alpha to Omega-The Numbers that Encode the Deepest Secrets of the*
- 849 *Universe*. Pantheon Books. 2002:
- 850 [73] Gamov G. Numerology for the constants of nature. *Proceedings of the National Academy of Science U.S.A.*, 1968, v.
- 851 59(2). 313–318;
- 852 [74] Saibal Rav. Utpal Mukhopadhyay and Partha Pratim Ghosh. Large Number Hypothesis : A Review.
- 853 <http://arxiv.org/pdf/0705.1836.pdf>
- 854 [75] P.J. Mohr, B.N. Taylor, and D.B. Newell in [arXiv:1203.5425](http://arxiv.org/abs/1203.5425) and *Rev. Mod. Phys.*
- 855 <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-phys-constants.pdf>
- 856 [76] Michael O. Distler et al. The RMS Charge Radius of the Proton and Zemach Moments. *Phys. Lett.B.* 696: 343-
- 857 347,2011
- 858 [77] Geiger H and Marsden E. On a diffuse reaction of the particles. *Proc. Roy. Soc., Ser. A* 82: 495-500, 1909.
- 859 [78] H. Yukawa. On the Interaction of Elementary Particles. *Proc. Phys. Math. Soc. Jap.* 17 (48). 1935
- 860 [79] Recami E. Elementary Particles as Micro-Universes, and "Strong Black-holes": A Bi-Scale Approach to Gravitational
- 861 and Strong Interactions. Preprint NSF-ITP-02-94. posted in the arXives as the e-print physics/0505149, and references
- 862 therein.
- 863 [80] Salam A. and Sivaram C. Strong Gravity Approach to QCD and Confinement. *Mod. Phys. Lett.*, 1993, v. A8(4), 321-
- 864 326.
- 865 [81] Abdus Salam. Strong Interactions, Gravitation and Cosmology. Publ. in: NATO Advanced Study Institute, Erice,
- 866 June16-July 6, 1972 ; in: *High Energy Astrophysics and its Relation to Elementary Particle Physics*, 441-452 MIT
- 867 Press, Cambridge (1974).
- 868 [82] U. V. S. Seshavatharam and S. Lakshminarayana, Strong nuclear gravitational constant and the origin of nuclear
- 869 planck scale. *Progress in Physics*, vol. 3, July, 2010, p. 31-38.
- 870 [83] U. V. S. Seshavatharam and S. Lakshminarayana, Role of Avogadro number in grand unification. *Hadronic Journal*.
- 871 Vol-33, No 5, 2010 October. p 513.
- 872 [84] U. V. S. Seshavatharam and S. Lakshminarayana. Accelerating universe and the expanding atom. *Hadronic journal*,
- 873 35(3): 271, 2012.
- 874 [85] U. V. S. Seshavatharam and S. Lakshminarayana. Nucleus in Strong nuclear gravity. *Proceedings of the DAE Symp.*
- 875 *On Nucl. Phys.* 56: 302, 2011
- 876 [86] Terry Quinn, Harold Parks, Clive Speake and Richard Davis. An uncertain big G. *Phys.Rev. Lett.* 112.068103. (2013)

Requesting your kind and valuable (unbiased) review comments...

877

<http://dx.doi.org/10.1103/PhysRevLett.111.101102>.

878

[87] J. B. Fixler; G. T. Foster; J. M. McGuirk; M. A. Kasevich. Atom Interferometer Measurement of the Newtonian Constant of Gravity, Science 315 (5808): 74–77, (2007).

879

880

881

APPENDIX: Major shortcomings of modern big bang cosmology

882

1) It may be noted that, increased redshifts and increased distances forced Edwin Hubble to propose the Hubble's law. In fact there is no chance or scope or place for 'galaxy receding'. It is only our belief in its 'given' (Doppler shift based) interpretation. Even then, merely by estimating galaxy distance and without measuring galaxy receding speed, one cannot verify its acceleration. Clearly speaking: two mistakes are possible here. i) Assumed galaxy receding speed is not being measured and not being confirmed. ii) Without measuring and confirming the galaxy receding speed, how can one say and confirm that it (galaxy) is accelerating. It is really speculative.

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

2) If light is coming from the atoms of the gigantic galaxy, then redshift can also be interpreted as an index of the galactic cosmological atomic 'light emission mechanism'. In no way it seems to be connected with 'galaxy receding'.

3) According to the modern cosmological approach, bound systems like 'atoms' which are found to be the major constituents of galactic matter - will not change with cosmic expansion/acceleration. As per the present observational data this may be true. But it might be the result of ending stage of cosmic expansion. As the issue is directly related with unification it requires lot of research in basic physics to confirm. In this regard, without considering and without analysing the past data, one can not come to a conclusion. If one is willing to think in this direction observed galactic redshift data can be considered for this type of new analysis.

4) 'Rate of decrease in current 'Hubble's constant' can be considered as a measure of current cosmic 'rate of expansion'. If rate of decrease in current 'Hubble's constant is very small and is beyond the scope of current experimental verification, then the two possible states are: a) current 'Hubble's constant is decreasing at a very slow rate and current universe is expanding at a very slow rate and b) at present there is no 'observable' cosmic expansion. Without a proper confirmation procedure for the absolute cosmic expansion and guessing that current universe is expanding - cosmologists proposed and confirmed the existence of dark energy indirectly. It may not be reasonable. Quantitatively or at least qualitatively standard model of cosmology does not throw light on the generation and (normal) physical properties of 'dark energy'.

5) The standard Big Bang model tells us that the Universe exploded out of an infinitely dense point. But nobody knows what would have triggered this outburst: the known laws of physics cannot tell us what happened at that moment.

6) Really if there was a 'big bang' in the past, with reference to formation of the big bang as predicted by general theory of relativity and with reference to the cosmic expansion that takes place simultaneously in all directions at a uniform rate at that time about the point of big bang - 'point' of big bang can be considered as the centre or characteristic reference point of cosmic expansion in all directions. In this case, saying that there is no preferred direction in the expanding universe - may not be correct.

7) Either in the big bang or in the inflation, quantification of the initial assumed conditions seem to be poor, unclear and not linked with fundamental constants. The earliest phases of the Big Bang are subject to much speculation and inflation requires 'fine tuning'.

8) Standard cosmology does not give information on the origin of 'inflation'. Inflation is often called a period of accelerated expansion. With respect to 'no hair theorem' some similarities are there for cosmic inflation and black holes. Conceptually 'inflation' can be accommodated in any model of cosmology like open model or closed model.

9) A key requirement is that inflation must continue 'long enough' to produce the present observable universe from a single, small inflationary Hubble volume. Assuming a rapid rate of cosmic expansion and steady rate of time may not be reasonable. If space-time is interrelated then 'space' and 'time' both should simultaneously follow the momentary rapid exponential expansion. For example if space expands by a factor 10^{26} in size within a very 'short span', cosmic time should also increase in the same proportion. 'Time' seems to be a silent observer in the presently believed 'cosmic inflation'. It may not be reasonable.

10) There is no scientific evidence for the Friedmann's second assumption. We believe it only on the grounds of modesty.

11) Dimensionally it is perfectly possible to show that, the dimensions of Hubble's constant and angular velocity are same. If so considering Hubble's constant merely as an expansion parameter may not be correct.

12) Even though it was having strong footing, Mach's principle was not implemented successfully in standard cosmology. Clearly speaking the term "distance cosmic back ground" is not being defined and not being quantified in a physical approach.

13) At any given cosmic time, the product of 'critical density' and 'Hubble volume' gives a characteristic cosmic mass and it can be called as the 'Hubble mass'. Interesting thing is that, Schwarzschild radius of the 'Hubble mass' again

Requesting your kind and valuable (unbiased) review comments...

- 931 matches with the 'Hubble length'. Most of the cosmologists believe that this is merely a coincidence. Here the
932 researchers emphasize the fact that this coincidence is having deep connection with cosmic geometry and the
933 cosmological physical phenomena.
- 934 14) Somehow and by any reason, magnitude of the current Hubble mass being the same, hypothetically if volume density
935 approaches the current matter density, then Hubble length increases by a factor ~ 5 . Similarly if volume density
936 approaches the current thermal energy density, then Hubble length increases by a factor ~ 27 . These two numbers
937 can be compared with the presently believed first two of the three cosmological numbers 4.9%, 26.8% and 68.3%.
938 Based on this coincidence and as the currently believed third number $\sim 68\%$ is obtained from the relation $(100-$
939 $(4.9+26.8))\%$, its proposed existence seems to be ad-hoc.
- 940 15) If 'Planck mass' is the characteristic beginning 'mass scale' of the universe, then by substituting the geometric mean
941 mass of the present Hubble mass and the Planck mass in the famous Hawking's black hole temperature formula
942 automatically the observed 2.725 K can be fitted very accurately. Standard cosmology is not throwing any light on this
943 surprising coincidence.
- 944 16) If cosmic expansion is continuous and accelerating and redshift is a measure of cosmic expansion, then 'rate of
945 increase in redshift' can be considered as a measure of cosmic 'rate of expansion'. Then there is no possibility to
946 observe a 'constant' red shift. More over the current definition of red shift seems to be ad-hoc and not absolute. Hence
947 one may not be able to understand or confirm the actual cosmic rate of expansion.
- 948 17) Even though the whole physics strictly follows the 'constancy of speed of light', cosmic acceleration seems to violate
949 it. This is really doubtful.
- 950 18) Drop in current 'cosmic temperature' can be considered as a measure of the current cosmic expansion and 'rate of
951 decrease in current cosmic temperature' can be considered as a measure of the current cosmic 'rate of expansion'. But
952 if rate of decrease in temperature is very small and is beyond the scope of current experimental verification, then the
953 two possible states are: a) current cosmic temperature is decreasing at a very slow rate and current universe is
954 expanding at a very slow rate and b) at present there is no 'observable' thermal expansion and there is no 'observable'
955 cosmic expansion. If observed CMBR temperature is 2.725 K and is very low in magnitude and is very close to
956 absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be reasonable.
- 957 19) If observed cosmic microwave back ground radiation temperature is 2.725 K and is very low in magnitude and is very
958 close to absolute zero, then thinking about and confirming the 'cosmic acceleration' may not be reasonable.
- 959 20) In the standard model of cosmology, there is no clear cut information about the 'uniqueness' of the assumed 'dark
960 energy'. If its identification is not unique in nature, then different cosmology models can be developed with different
961 forms of 'dark energy'. If so understanding the absolute cosmic expansion rate with dark energy seems to be doubtful.
- 962 21) So far no ground based experiment confirmed the existence of dark energy. There is no single clue or evidence to any
963 of the natural physical properties of (the assumed) dark energy.
- 964 22) If 'Dark energy' is the major outcome of the 'accelerating universe', it is very important to note that - in understanding
965 the basic concepts of unification or other fundamental areas of physics, role of dark energy is very insignificant.
- 966 23) If existence of dark energy is true and dark energy is supposed to have a key role in the past and current cosmic
967 expansion, then it must have also played a key role in the beginning of cosmic evolution. In this regard no
968 information is available in standard cosmology.
- 969 24) Standard model of cosmology does not throw light on the generation and existence of atomic physical constants like
970 Planck's constant, reduced Planck's constant, inverse of fine structure ratio and nuclear charge radius etc. Clearly
971 speaking synthesis of elementary physical constants seem to be more important than the cosmological nucleosynthesis.
- 972 25) General theory of relativity does not throw any light on the 'mass generation' of charged particles. It only suggests
973 that space-time is curved near the massive celestial objects. More over it couples the cosmic (dust) matter with
974 geometry. But how matter/dust is created? Why and how elementary particle possesses both charge and mass? Such
975 types of questions are not being discussed in the frame work of general relativity.
- 976 26) Standard model of cosmology does not throw light on the charge-mass unification scheme of atomic particles. The
977 main object of unification is to understand the origin of elementary particles rest mass, magnetic moments and their
978 forces. Right now and till today 'string theory' with 4 + 6 extra dimensions is not in a position to explain the
979 unification of gravitational and non-gravitational forces. More clearly speaking it is not in a position to merge the
980 Planck scale and cosmic scale with the characteristic nuclear scale.
- 981 27) Either general theory of relativity or standard cosmology does not give any information on the applications of the
982 classical force limit (c^4/G) and the classical power limit (c^5/G). Compared to the hypothetical 'dark energy', with a
983 coefficient of unity, (c^4/G) can be considered as the cosmic vacuum force and (c^5/G) can be considered as the cosmic
984 vacuum power.

Requesting your kind and valuable (unbiased) review comments...

985 28) In Big bang model, confirmation of all the observations directly depend on the large scale galactic distances that are
986 beyond human reach and raise ambiguity in all respects. The subject of modern black hole physics is absolutely
987 theoretical. Advantage of Black hole cosmology lies in confirming its validity through the ground based atomic and
988 nuclear experimental results.
989

990 If one is willing to think in this new direction, certainly other hidden short comings can also be surfaced out. Most of the
991 modern cosmologists are enforced with 85 years old Hubble's interpretation. This is the time to re-interpret the Hubble's
992 law and to revise the basics of modern cosmology. Based on the proposed short comings the concepts of 'big bang
993 cosmology' can be relinquished and Black hole cosmology can be invoked for in-depth discussion.