

“If you had only one slide
to get your point across...”

The ‘One Slide’ Introduction to ***Generalized*** Quantum Impedances

Peter Cameron

“To understand the electron would be enough”
Einstein

The Essential Point

Impedances govern the flow of energy

Classical or quantum impedances,
mechanical or electromagnetic,
fermionic or bosonic,
topological or geometric,...

Impedances govern the flow of energy

This is not a theoretical musing

This is a fundamental concept, of universal applicability

Generalization of quantum impedances extends the concept from the Lorentz impedance of the quantum Hall effect and the near and far-field photon impedances to all quantum potentials and their associated forces and impedances

Define 'Impedance'


given The Essential Point,

'One Slide'

Impedance - a measure of the amplitude and **phase** of opposition to the flow of a current

- single force - electromagnetic [ohms] = [kg-m²/coul²-s]
or mechanical [kg/s] mass flow, deBroglie wave,...
- coupled forces - electromechanical, magnetomechanical,...
- for now stay with **electromagnetic**
- resistance, inductance, capacitance
- **distinctions** between classical and quantum
 - classical resistance is incoherent (heat, noise,...)
 - quantum resistance is phase-coherent (no dissipation)
 - **phase** is **not a single measurement observable** in quantum mechanics
 - scale invariant quantum impedances cannot transmit energy, communicate only quantum phase (not a single measurement observable)
- conjectures – all quantum impedances are topological?
all classical impedances are geometric?

conversion factor
from mechanical
to electrical
impedance



*EPR,
non-locality,...*



'One Slide'

Strong Decay

EM Decay

Weak Decay

$\pi_0 \rightarrow 2\gamma$	98.8%
$\pi_0 \rightarrow e^+e^-\gamma$	1.2%
$\eta \rightarrow 2\gamma$	39.3%
$\eta \rightarrow 3\pi^0$	32.6%
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.7%
$\eta \rightarrow \pi^+\pi^-\gamma$	4.6%
$\eta' \rightarrow \pi^+\pi^-\eta$	39.3%
$\eta' \rightarrow \rho^0\gamma$	
incl. $\pi^+\pi^-\gamma$	29.3%
$\eta' \rightarrow \pi^0\pi^0\eta$	21.6%
$\eta' \rightarrow \omega\gamma$	2.8%
$\eta' \rightarrow \gamma\gamma$	2.2%



- Horizontal axis in both plots is logarithmic
- Length scales are the same for both plots, and they are properly aligned
- Upper plot shows particle lifetimes multiplied by the speed of light, the coherence lengths (adapted from The Power of Alpha by Malcolm MacGregor)
- Plot at left is electron impedances, details at <http://redshift.vif.com/JournalFiles/V18N02PDF/V18N2CAM.pdf>
- Alpha-spaced coherence lengths of η' , η , and π_0 are at conjunctions of mode impedances, can couple to the photon for fast EM decay
- Location of π_0 at boundary of EM and weak decays is suggestive of the chiral anomaly
- Weak decays are mismatched to the photon
- Alpha-spaced coherence length alternation of fermions/bosons in weak decays appears to be related to parity violation and electroweak enhancement of charm/tau coherence lengths
- Clustering of superheavies (top, Higgs, Z, W) at the 10GeV coherence length is tantalizing
- Impedance junctions at the classical radius are related to mass quantization and MacGregor's 70MeV 'platform state'
- See also http://vixra.org/author/peter_cameron

Presentation Goals - Understand, then Explore

- **understand** the 'One Slide'
 - ~~essential point – impedances govern the flow of energy~~
 - **origin** of impedance network – two body problem & Mach's Principle
 - **origin** of the coherence length plot – Malcolm MacGregor
 - **implications** of their strong correlation – personal paradigm shift
- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - elementary particle spectrum, chiral anomaly, *axions, EDM...*
 - gravity – extend the model to the Planck particle
 - state reduction and non-locality *Italics connote material in preparation*
 - black hole information paradox – Rochester, Fields,...
 - weak measurement and time symmetry – Berlin
 - quantum interpretations – Berlin Optical Society of America sponsored and refereed
 - *dark matter*
 - *condensed matter*

Presentation Goals - Explore, then Understand

- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - elementary particle spectrum, chiral anomaly, axions, EDM...
 - gravity – extend the model to the Planck particle
 - state reduction and non-locality
 - black hole information paradox – Rochester, Fields,...
 - weak measurement and time symmetry – Berlin
 - quantum interpretations – Berlin
 - *dark matter* *Italics connote material in preparation*
 - *condensed matter*
- **understand** the One Slide
 - ~~essential point – impedances govern the flow of energy~~
 - **origin** of impedance network – two body problem & Mach's Principle
 - **origin** of the coherence length plot – Malcolm MacGregor
 - **implications** of their strong correlation – personal paradigm shift

[9] [viXra:1402.0064](#) replaced on 2014-06-10 07:16:24, (83 unique-IP downloads)

An Impedance Approach to the Chiral Anomaly

Authors: [Peter Cameron](#)

Category: [High Energy Particle Physics](#)

February 2014

[8] [viXra:1311.0143](#) replaced on 2013-12-11 22:25:43, (136 unique-IP downloads)

Quantum Interpretation of the Impedance Model

Authors: [Michaele Suisse](#), [Peter Cameron](#)

Category: [Quantum Physics](#)

November 2013

[7] [viXra:1310.0043](#) replaced on 2013-12-04 15:17:55, (237 unique-IP downloads)

Delayed Choice and Weak Measurement in the Nested Mach-Zehnder Interferometer

Authors: [Peter Cameron](#)

Category: [Quantum Physics](#)

October 2013

[6] [viXra:1306.0102](#) submitted on 2013-06-15 05:02:17, (122 unique-IP downloads)

Poster for the Rochester Conference on Quantum Optics and Information

Authors: [Peter Cameron](#)

Category: [Quantum Physics](#)

June 2013

[5] [viXra:1304.0078](#) replaced on 2013-04-20 08:41:41, (67 unique-IP downloads)

Abstract of Poster Accepted for the Fields Conference on Quantum Information

Authors: [Peter Cameron](#)

Category: [Quantum Gravity and String Theory](#)

April 2013

[4] [viXra:1304.0036](#) replaced on 2013-07-23 22:56:22, (215 unique-IP downloads)

A Possible Resolution of the Black Hole Information Paradox

Authors: [Peter Cameron](#)

Category: [Quantum Physics](#)

April 2013

[3] [viXra:1303.0039](#) replaced on 2013-05-11 14:40:00, (386 unique-IP downloads)

Quantum Impedances, Entanglement, and State Reduction

Authors: [Peter Cameron](#)

Category: [Quantum Physics](#)

March 2013

[2] [viXra:1211.0052](#) replaced on 2013-04-10 11:46:31, (263 unique-IP downloads)

Background Independent Relations Between Gravity and Electromagnetism

Authors: [Peter Cameron](#)

Category: [Quantum Gravity and String Theory](#)

November 2012

[1] [viXra:1207.0022](#) replaced on 2013-04-26 11:09:42, (310 unique-IP downloads)

Generalized Quantum Impedances: A Background Independent Model for the Unstable Particles

Authors: [Peter Cameron](#)

Category: [High Energy Particle Physics](#)

July 2012

Generalized Quantum Impedances: A Background Independent Model for the Unstable Particles

Peter Cameron*
Strongarm Studios
Mattituck, NY USA 11952

first posted July 2012

(Dated: April 26, 2013)

The discovery of exact impedance quantization in the quantum Hall effect was greatly facilitated by scale invariance. This letter explores the possibility that quantum impedances may be generalized, defined not just for the Lorentz force and the quantum Hall effect, but rather for all forces, resulting in a precisely structured network of scale dependent and scale invariant impedances. If the concept of generalized quantum impedances correctly describes the physical world, then such impedances govern how energy is transmitted and reflected, how the hydrogen atom is ionized by a 13.6eV photon, or why the π_0 branching ratio is what it is. An impedance model of the electron is presented, and explored as a model for the unstable particles.

INTRODUCTION

The model presented here [1–6] comprises

- quantization of electric and magnetic flux, charge, and dipole moment
- interactions between these three topologies - flux quantum, monopole, and dipole
- the photon
- confinement to a fundamental length, taken to be the Compton wavelength of the electron

Calculated transfer impedances of the interactions are presented as a function of spatial scale/energy. Possible roles for these impedances in the creation and structure of the unstable particles are discussed.




FIG. 1. Photon and electron impedances as a function of spatial scale as defined by photon energy

$$Z_M = Z_0 \left| \frac{1 + \frac{\lambda}{ir}}{1 + \frac{\lambda}{ir} + \left(\frac{\lambda}{ir}\right)^2} \right|$$



'One Slide'

Strong Decay

EM Decay

Weak Decay

$\pi_0 \rightarrow 2\gamma$	98.8%
$\pi_0 \rightarrow e^+e^-\gamma$	1.2%
$\eta \rightarrow 2\gamma$	39.3%
$\eta \rightarrow 3\pi^0$	32.6%
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.7%
$\eta \rightarrow \pi^+\pi^-\gamma$	4.6%
$\eta' \rightarrow \pi^+\pi^-\eta$	39.3%
$\eta' \rightarrow \rho^0\gamma$	
incl. $\pi^+\pi^-\gamma$	29.3%
$\eta' \rightarrow \pi^0\pi^0\eta$	21.6%
$\eta' \rightarrow \omega\gamma$	2.8%
$\eta' \rightarrow \gamma\gamma$	2.2%



- Horizontal axis in both plots is logarithmic
- Length scales are the same for both plots, and they are properly aligned
- Upper plot shows particle lifetimes multiplied by the speed of light, the coherence lengths (adapted from The Power of Alpha by Malcolm MacGregor)
- Plot at left is electron impedances, details at <http://redshift.vif.com/JournalFiles/V18N02PDF/V18N2CAM.pdf>
- Alpha-spaced coherence lengths of η' , η , and π_0 are at conjunctions of mode impedances, can couple to the photon for fast EM decay
- Location of π_0 at boundary of EM and weak decays is suggestive of the chiral anomaly
- Weak decays are mismatched to the photon
- Alpha-spaced coherence length alternation of fermions/bosons in weak decays appears to be related to parity violation and electroweak enhancement of charm/tau coherence lengths
- Clustering of superheavies (top, Higgs, Z, W) at the 10 GeV coherence length is tantalizing
- Impedance junctions at the classical radius are related to mass quantization and MacGregor's 70 MeV 'platform state'
- See also http://vixra.org/author/peter_cameron

[9] **viXra:1402.0064** replaced on 2014-06-10 07:16:24, (83 unique-IP downloads)

An Impedance Approach to the Chiral Anomaly

Authors: Peter Cameron

Category: High Energy Particle Physics

motivated by
Michael Creutz

February 2014

[8] **viXra:1311.0143** replaced on 2013-12-11 22:25:43, (136 unique-IP downloads)

Quantum Interpretation of the Impedance Model

Authors: Michael Suisse, Peter Cameron

Category: Quantum Physics

November 2013

[7] **viXra:1310.0043** replaced on 2013-12-04 15:17:55, (237 unique-IP downloads)

Delayed Choice and Weak Measurement in the Nested Mach-Zehnder Interferometer

Authors: Peter Cameron

Category: Quantum Physics

October 2013

[6] **viXra:1306.0102** submitted on 2013-06-15 05:02:17, (122 unique-IP downloads)

Poster for the Rochester Conference on Quantum Optics and Information

Authors: Peter Cameron

Category: Quantum Physics

June 2013

[5] **viXra:1304.0078** replaced on 2013-04-20 08:41:41, (67 unique-IP downloads)

Abstract of Poster Accepted for the Fields Conference on Quantum Information

Authors: Peter Cameron

Category: Quantum Gravity and String Theory

April 2013

[4] **viXra:1304.0036** replaced on 2013-07-23 22:56:22, (215 unique-IP downloads)

A Possible Resolution of the Black Hole Information Paradox

Authors: Peter Cameron

Category: Quantum Physics

April 2013

[3] **viXra:1303.0039** replaced on 2013-05-11 14:40:00, (386 unique-IP downloads)

Quantum Impedances, Entanglement, and State Reduction

Authors: Peter Cameron

Category: Quantum Physics

March 2013

[2] **viXra:1211.0052** replaced on 2013-04-10 11:46:31, (263 unique-IP downloads)

Background Independent Relations Between Gravity and Electromagnetism

Authors: Peter Cameron

Category: Quantum Gravity and String Theory

November 2012

[1] **viXra:1207.0022** replaced on 2013-04-26 11:09:42, (310 unique-IP downloads)

Generalized Quantum Impedances: A Background Independent Model for the Unstable Particles

Authors: Peter Cameron

Category: High Energy Particle Physics

July 2012

An Impedance Approach to the Chiral Anomaly

Peter Cameron^{*}
Strongarm Studios
Mattituck, NY USA 11952

(Dated: May 25, 2014)

The chiral potential is inverse square. The family of inverse square potentials includes the vector Lorentz potential of the quantum Hall and Aharonov-Bohm effects, and the centrifugal, Coriolis, and three body potentials. The associated impedances are scale invariant, quantum Hall being the most familiar. Modes associated with scale invariant impedances communicate only quantum phase, not an observable in a single quantum measurement. Modes associated with scale dependent impedances, including among others those of the $1/r$ monopole and $1/r^3$ dipole potentials, communicate both phase and energy. Making this clarifying distinction between phase (relative time) and energy explicit presents a new perspective on the anomaly. This approach is introduced via the Rosetta Stone of modern physics, the hydrogen atom. Precise impedance-based π^0 , η , and η' branching ratio calculations are presented as ratios of polynomials in powers of the fine structure constant, followed by discussion. Mass generation via chiral symmetry breaking is not addressed in the present paper. axion implications,...

INTRODUCTION

Anomalies may be defined as “...breakings of classical symmetries by quantum corrections, which arise when the regularizations needed to evaluate small fermion loop Feynman diagrams conflict with a classical symmetry of the theory.” [1] Stephen Adler

In a finite quantum theory chiral symmetry appears to be broken only by weak interactions. The presence of the anomaly in strong and electromagnetic quantum field theory (QFT) calculations [1-8] seems to be an inevitable result of the regularization needed to remove infinities before mass and charge renormalizations can be accomplished. However, one has a choice - in the presence of the anomaly either chiral symmetry or gauge invariance must be broken.

gauge invariant (no covariant derivative) and finite (no renormalization)

The impedance approach is gauge invariant.

Gauge invariance is built in. Complex impedances shift phases. Complex quantum impedances shift quantum phases. The scale invariant impedance associated with the chiral potential [9, 10] communicates quantum phase and only quantum phase [11-13]. No need for the covariant derivative. One need only take the appropriate impedances into account.

The phase-only character of inverse square potentials, their incapacity to do work, is emphasized in the related case of the centrifugal potential of the free Schroedinger particle by Holstein [14]. The symmetry is understood to be scale invariance (unbroken sans regularization).

The impedance approach is finite

Impedance is a geometric concept, depends on size and shape. In the limit of the small, the point/singularity is infinitely mismatched to you and I. We cannot share energy with it. While presumably equally decoupled, the quantum limit of the large is more subtle, in the emergent realm of the classical, and ultimately the cosmological. In both limits, small and large, divergences are removed by the impedance mismatches. Regularization and renormalization are not necessary.

The anomaly does not arise in the impedance approach, a result of the finiteness and gauge invariance.

π^0 branching ratio calculation – experimental values in parentheses

$$Z_{\gamma\gamma} = \frac{1}{\frac{1}{Z_0} + \frac{1}{Z_0}} = \frac{Z_0}{2} = 188.37 \Omega \quad (1)$$

and that of the $e^+e^-\gamma$ mode as

$$Z_{ee\gamma} = \frac{1}{\frac{1}{R_H} + \frac{1}{R_H} + \frac{4\alpha^2}{Z_0}} = \frac{Z_0}{4\alpha^2 + 4\alpha} = 12813 \Omega \quad (2)$$


where $R_H = \frac{Z_0}{2\alpha}$ is the quantum Hall resistance, so that

$$Z_{\pi^0} = \frac{1}{\frac{1}{Z_{\gamma\gamma}} + \frac{1}{Z_{ee\gamma}}} = \frac{Z_0}{4\alpha^2 + 4\alpha + 2} = 185.64 \Omega \quad (3)$$

and the branching ratios are for photons
no α in the numerator

$$\Gamma_{\gamma\gamma} = \frac{Z_{\pi^0}}{Z_{\gamma\gamma}} = \frac{1}{2\alpha^2 + 2\alpha + 1} = 0.9855 \text{ (0.988)} \quad (4)$$

$$\Gamma_{ee\gamma} = \frac{Z_{\pi^0}}{Z_{ee\gamma}} = \frac{2\alpha^2 + 2\alpha}{2\alpha^2 + 2\alpha + 1} = 0.0145 \text{ (0.012)} \quad (5)$$





'One Slide'

Strong Decay

EM Decay

Weak Decay

$\pi_0 \rightarrow 2\gamma$	98.8%
$\pi_0 \rightarrow e^+e^-\gamma$	1.2%
$\eta \rightarrow 2\gamma$	39.3%
$\eta \rightarrow 3\pi^0$	32.6%
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.7%
$\eta \rightarrow \pi^+\pi^-\gamma$	4.6%
$\eta' \rightarrow \pi^+\pi^-\eta$	39.3%
$\eta' \rightarrow \rho^0\gamma$	
incl. $\pi^+\pi^-\gamma$	29.3%
$\eta' \rightarrow \pi^0\pi^0\eta$	21.6%
$\eta' \rightarrow \omega\gamma$	2.8%
$\eta' \rightarrow \gamma\gamma$	2.2%



- Horizontal axis in both plots is logarithmic
- Length scales are the same for both plots, and they are properly aligned
- Upper plot shows particle lifetimes multiplied by the speed of light, the coherence lengths (adapted from The Power of Alpha by Malcolm MacGregor)
- Plot at left is electron impedances, details at <http://redshift.vif.com/JournalFiles/V18N02PDF/V18N2CAM.pdf>
- Alpha-spaced coherence lengths of η' , η , and π_0 are at conjunctions of mode impedances, can couple to the photon for fast EM decay
- Location of π_0 at boundary of EM and weak decays is suggestive of the chiral anomaly
- Weak decays are mismatched to the photon
- Alpha-spaced coherence length alternation of fermions/bosons in weak decays appears to be related to parity violation and electroweak enhancement of charm/tau coherence lengths
- Clustering of superheavies (top, Higgs, Z, W) at the 10 GeV coherence length is tantalizing
- Impedance junctions at the classical radius are related to mass quantization and MacGregor's 70 MeV 'platform state'
- See also http://vixra.org/author/peter_cameron

eta and eta' branching ratios
calculated from impedance matches

$$\Gamma_{\gamma\gamma} = \frac{Z_\eta}{Z_{\gamma\gamma}} = \frac{8}{20\alpha^2 + 68\alpha + 19} \quad (14)$$

$$= 0.410 \text{ (0.393)}$$

no factor of α in the numerator
true for all photon branches

$$\Gamma_{3\pi^0} = \frac{Z_\eta}{Z_{3\pi^0}} = \frac{3(4\alpha^2 + 4\alpha + 2)}{20\alpha^2 + 68\alpha + 19} \quad (15)$$

$$= 0.312 \text{ (0.326)}$$


$$\Gamma_{\pi\pi\pi^0} = \frac{Z_\eta}{Z_{\pi\pi\pi^0}} = \frac{2(4\alpha^2 + 20\alpha + 2)}{20\alpha^2 + 68\alpha + 19} \quad (16)$$

$$= 0.220 \text{ (0.227)}$$

$$\Gamma_{\pi\pi\gamma} = \frac{Z_\eta}{Z_{\pi\pi\gamma}} = \frac{16\alpha + 1}{20\alpha^2 + 68\alpha + 19} \quad (17)$$

$$= 0.057 \text{ (0.046)}$$

excellent fit to the data (however, factors of 2)



eta .393 .326 .227 .046

model .410 .312 .220 .057

eta' .393 .293 .216 .050

DISCUSSION

Historical Perspective on Quantum Impedances

Impedances govern the flow of energy. This is a fundamental concept of universal applicability. Historically, it has been overlooked in quantum theory.

The 1980 discovery [22] of a new fundamental constant of nature, the Nobel Prize discovery of exact impedance quantization in the quantum Hall effect, was greatly facilitated by scale invariance. This classically peculiar impedance is topological, the measured resistance being independent of the size or shape of the Hall bar. Prior to that discovery, impedance quantization was more implied than explicit in the literature [23–28].

In the 1959 thesis of Bjorken [25] is an approach summarized [29] as “...an analogy between Feynman diagrams and electrical circuits, with Feynman parameters playing the role of resistance, external momenta as current sources, and coordinate differences as voltage drops. Some of that found its way into section 18.4 of...” the canonical text [26]. As presented there, the units of the Feynman parameter are [sec/kg], the units of *mechanical conductance* [15]. Form factors are proportional to conductances, inversely proportional to resistances.

With the confusion that resulted from interpreting conductance as resistance, and more importantly lacking the concept of quantized impedance, the anticipated intuitive advantage of the circuit analogy [26] was lost and the possibility of the jump from well-considered analogy to a photon-electron impedance model was not realized.

Like the first Rochester Conference on Coherence and Quantum Optics in 1960, the 1963 paper/thesis by Vernon and Feynman [27] on the “Interaction of Systems” was motivated by the invention of the maser. It is a particularly suggestive combination of the languages of the electrical engineer and the physicist. The authors devoted a thesis to the concepts needed for impedance matching to the maser. However, lacking again was the explicit concept of quantized impedance in the maser.

Had exact impedance quantization been discovered in 1950 rather than 1980, one wonders whether the impedance concept might have found its way into the foundation of QED at that time, before it was set in the bedrock, to underpin rather than illuminate electroweak theory, QCD, and gravity [12, 30–39].

Presentation Goals - Understand, then Explore

- **understand** the One Slide
 - ~~essential point~~ – impedances govern the flow of energy
 - **origin** of impedance network – two body problem & Mach's Principle
 - **origin** of the coherence length plot – Malcolm MacGregor
 - **implications** of their strong correlation – personal paradigm shift
- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - gravity – extend the model to the Planck particle
 - state reduction and non-locality
 - black hole information paradox – Rochester, Fields,...
 - weak measurement and time symmetry – Berlin
 - quantum interpretations – Berlin
 - *dark matter*
 - *condensed matter*

Italics connote material in preparation

July 24, 1975

THE TWO BODY PROBLEM AND MACH'S PRINCIPLE

Peter Cameron

2210 Water Street

Port Huron, Michigan 48060

The classical analysis of the two-body problem is frequently complicated by the introduction of a system of co-ordinates which is independent of either of the bodies. The validity of such an analysis rests upon the premise that the co-ordinate frame does not interact with the physical system via any known physical laws, and that one is therefore free to choose whatever reference frame seems most useful.


A strong epistemological argument might be advanced against this reasoning. If sufficiently rigorous constraints are placed upon the spatial properties of the interacting bodies, the introduction of an independent observer will have a radical effect upon the form of the equations which



vibratory
piledriver/extractor

the foundation of all
that followed

It took 35 years to
understand where it fits



3

what waves in the deBroglie wave?

$$F = d(mv)/dt = mv^2/r + v dm/dt$$

The first term has no meaning and must be discarded. The second term would also seem to be meaningless. We have no reason to suspect that m_1 varies in time, and nothing in our initial conditions seems to require that m_1 be a point mass, a circumstance which would deprive us of the ability to observe radial velocity. Either we accept the second force term as counter-balancing the gravitational attraction or we regard the whole situation as senseless. Nothing in the initial conditions requires that the problem is senseless, so we write

$$v_{\text{rad}} dm_1/dt = Gm_1m_2/r^2 \quad \text{or} \\ dm_1/dt = (Gm_1m_2/v_{\text{rad}})(1/r^2) \quad (1)$$

In writing this we note that it was necessary to take $v=v_{\text{rad}}$ to maintain the co-linearity of forces. The quantity Gm_1m_2/v_{rad} has units of angular momentum, which suggests

$$dm_1/dt = L/r^2 \quad (2)$$

system composed of elementary particles. The Bohr model of the hydrogen atom is a familiar example. As before, we consider uniform circular motion ($n=1$), we consider the proton to be the center of mass, and we require that relativistic corrections be negligible and that the intrinsic angular momentum and magnetic fields of the particles be ignored. The law of force is

$$dp/dt = q^2/4\pi\epsilon_0 r^2$$

Following the line of reasoning previously developed, we write this as

$$dm_e/dt = (q^2/4\pi\epsilon_0 v_{rad})(1/r^2) \quad \text{conversion factor [kg/sec] to [ohms]} \quad (3)$$

which for the Bohr atom $n=1$

$$dm_e/dt = \hbar/r^2$$

$$Z_{centri} := \frac{h}{\lambda_{bar_e}^2} \cdot \frac{\lambda_{bar_e}^2}{e^2} = \frac{h}{e^2} = 2.5812807554 \times 10^4 \text{ ohm}$$

a result which is similar in form to the previously analyzed gravitationally bound system.

this is the quantum 'centripetal impedance', equal to the quantum Hall impedance

a challenge – find the near-field photon impedance
in any of the standard grad school E&M texts

reminder – impedances govern
the flow of energy

Photon Impedance Match to a Single Free Electron

Peter Cameron
Brookhaven National Laboratory
Upton, NY 11973
cameron@bnl.gov

It is not surprising that consideration of impedance matching the photon to the electron, or more specifically to the quantum of resistance at the length scale defined by the mass and angular momentum of the electron, has been long ignored in quantum electrodynamics. Conceptually the development of QED preceded the discovery of ‘exact quantization’ and the associated von Klitzing constant by many decades. Additionally, the relevance of the resistance quantum to photon interactions with a single free electron has only recently begun to be appreciated. In this note we offer a simple presentation of such an impedance match, briefly discuss the unexpected emergence of the fine structure constant from these simple first principles, and suggest how the procedure can be inverted to deliver a first principles calculation of the mass of the electron.

Possible Origin of the 70MeV Mass Quantum

Peter Cameron
Brookhaven National Laboratory
Upton, NY 11973
cameron@bnl.gov

The absence of three fundamental entities from the experimental evidence is notable. The search for two, the magnetic monopole and the electric dipole moment, is extensively documented in the literature. The third, the electric flux quantum, is remarkably absent. One is tempted to speculate that these circumstances are related, and that exploration of the electric flux quantum might shed light upon, and beyond, the absence of the magnetic monopole and the electric dipole. This note presents a tentative early effort to find a guidepost or two along the road to such an exploration, or at least a compass that permits the choice of direction. What emerges is a possible origin of the 70MeV platform state. While documentation of this mass quantum (it is simply the mass of the electron divided by the fine structure constant) in the literature is not so starkly absent as for the electric flux quantum, it is surprisingly sparse.

Magnetic and Electric Flux Quanta: the Pion Mass

Peter Cameron cameron@bnl.gov
Brookhaven National Laboratory
Upton, NY 11973

'paradox' is in reference to
origin of Bohr magneton

The angular momentum of the magnetic flux quantum is balanced by that of the associated supercurrent, such that in condensed matter the resultant angular momentum is zero. The notion of a flux quantum in free space is not so simple, needing both magnetic and electric flux quanta to propagate the stable dynamic structure of the photon. Considering these flux quanta at the scale where quantum field theory becomes essential, at the scale defined by the reduced Compton wavelength of the electron, exposes variants of a paradox that apparently has not been addressed in the literature. Leaving the paradox unresolved in this note, reasonable electromagnetic rationales are presented that permit to calculate the masses of the electron, muon, pion, and nucleon with remarkable accuracy. The calculated mass of the electron is correct at the nine significant digit limit of experimental accuracy, the muon at a part in one thousand, the pion at two parts in ten thousand, and the nucleon at seven parts in one hundred thousand. The accuracy of the pion and nucleon mass calculations reinforces the unconventional common notion that the strong force is electromagnetic in origin.

Electron Impedances

Peter Cameron
Brookhaven National Laboratory
Upton, NY 11973
cameron@bnl.gov

It is only recently, and particularly with the quantum Hall effect and the development of nanoelectronics, that impedances on the scale of molecules, atoms and single electrons have gained attention. In what follows the possibility that characteristic impedances might be defined for the photon and the single free electron is explored in some detail, the premise being that the concepts of electrical and mechanical impedances are relevant to the elementary particle. The scale invariant quantum Hall impedance is pivotal in this exploration, as is the two body problem and Mach's principle.

To understand the electron would be enough - Einstein

MECHANICAL IMPEDANCE

While the concept of electrical impedance is comfortably familiar to any electrical engineer and many physicists, mechanical impedance [1] is more obscure. It is defined as [19]

$$Z_{mech} = \frac{F}{v}$$

where F is the applied force and v the resulting velocity. The form is similar to Newton's second law written as

$$m = \frac{F}{a}$$

where m is the mass to which the force is applied and a is the resulting acceleration.

Taking the force F to be, for example, the centrifugal force

$$F_{centri} = \frac{mv^2}{r}$$

gives the centrifugal impedance

$$Z_{centri} = \frac{mv}{r}$$

where r is the radius of curvature of the path of the mass acted upon by this force.

From the 'Unstable Particles' note

The centrifugal force is in some sense a mechanical equivalent of the vector Lorentz force present in the quantum Hall effect. Like the Lorentz force, it is velocity dependent. Unlike velocity dependent forces other than the Lorentz and centrifugal forces, it is not dissipative. Like the Lorentz force, it is perpendicular to the direction of motion, and hence can do no work.

Defining v by the deBroglie relation $v = \frac{h}{mr}$ yields the simple form

$$Z_{centri} = \frac{h}{r^2}$$

The units of mechanical impedance are [kg/s], those of electrical impedance [ohm] = [(kg/s)(m/Coul)²]. Taking the second term on the right hand side, the line charge density term, to be a conversion factor between mechanical and electrical impedances and the charge to be the charge quantum e gives

$$Z_{centri} = \frac{h}{r^2} \frac{r^2}{e^2} = \frac{h}{e^2} \simeq 25\,812.8\Omega$$

This impedance is numerically and symbolically identical to the scale invariant quantum Hall impedance, and is plotted in figure 2 (green dots).

The method presented in the above example can be used to calculate quantum impedances for forces other than the centrifugal and vector Lorentz forces. The impedance plot of figure 2 shows results from such calculations [2].

Presentation Goals - Explore, then Understand

- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - **gravity – extend the model to the Planck particle**
 - **state reduction and non-locality**
 - **black hole information paradox – Rochester, Fields,...**
 - weak measurement and time Symmetry – Berlin
 - quantum interpretations – Berlin
 - *dark matter* *Italics connote material in preparation*
 - *condensed matter*
- **understand** the One Slide
 - ~~essential point – impedances govern the flow of energy~~
 - ~~origin of impedance network – two body problem & Mach's Principle~~
 - **origin** of the coherence length plot – Malcolm MacGregor
 - **implications** of their strong correlation – personal paradigm shift

[9] [viXra:1402.0064](#) replaced on 2014-06-10 07:16:24, (83 unique-IP downloads)

An Impedance Approach to the Chiral Anomaly

Authors: Peter Cameron

Category: High Energy Particle Physics

February 2014

[8] [viXra:1311.0143](#) replaced on 2013-12-11 22:25:43, (136 unique-IP downloads)

Quantum Interpretation of the Impedance Model

Authors: Michaele Suisse, Peter Cameron

Category: Quantum Physics

November 2013

[7] [viXra:1310.0043](#) replaced on 2013-12-04 15:17:55, (237 unique-IP downloads)

Delayed Choice and Weak Measurement in the Nested Mach-Zehnder Interferometer

Authors: Peter Cameron

Category: Quantum Physics

October 2013

[6] [viXra:1306.0102](#) submitted on 2013-06-15 05:02:17, (122 unique-IP downloads)

Poster for the Rochester Conference on Quantum Optics and Information

Authors: Peter Cameron

Category: Quantum Physics

June 2013

[5] [viXra:1304.0078](#) replaced on 2013-04-20 08:41:41, (67 unique-IP downloads)

Abstract of Poster Accepted for the Fields Conference on Quantum Information

Authors: Peter Cameron

Category: Quantum Gravity and String Theory

April 2013

[4] [viXra:1304.0036](#) replaced on 2013-07-23 22:56:22, (215 unique-IP downloads)

A Possible Resolution of the Black Hole Information Paradox

Authors: Peter Cameron

Category: Quantum Physics

this
is synthesis of
these

April 2013

[3] [viXra:1303.0039](#) replaced on 2013-05-11 14:40:00, (386 unique-IP downloads)

Quantum Impedances, Entanglement, and State Reduction

Authors: Peter Cameron

Category: Quantum Physics

March 2013

[2] [viXra:1211.0052](#) replaced on 2013-04-10 11:46:31, (263 unique-IP downloads)

Background Independent Relations Between Gravity and Electromagnetism

Authors: Peter Cameron

Category: Quantum Gravity and String Theory

November 2012

[1] [viXra:1207.0022](#) replaced on 2013-04-26 11:09:42, (310 unique-IP downloads)

Generalized Quantum Impedances: A Background Independent Model for the Unstable Particles

Authors: Peter Cameron

Category: High Energy Particle Physics

July 2012

Background Independent Relations between Gravity and Electromagnetism

Peter Cameron

Planck particle model
Is electron model at
Planck length



Received: date / Accepted: date

first posted on vixra November 2012

The model presented here [1–6] comprises


- quantization of electric and magnetic flux, charge, and dipole moment
- interactions between these three topologies - flux quantum, monopole, and dipole
- the photon
- confinement to a fundamental length, taken to be the Compton wavelength of the Planck particle

Calculated transfer impedances of the interactions are presented as a function of spatial scale/energy. Possible

Abstract As every circuit designer knows, the flow of energy is governed by impedance matching. Classical or quantum impedances, mechanical or electromagnetic, fermionic or bosonic, topological,... To understand the flow of energy it is essential to understand the relations between the associated impedances. The connection between electromagnetism and gravitation can be made explicit by examining the impedance mismatch between the electrically charged Planck particle and the electron. This mismatch is shown to be the ratio of the gravitational and electromagnetic forces between these particles.

Keywords background independence · scale invariance · quantum impedance · network theory · scattering matrix · near field · Planck particle · state reduction · information theory

Figure 4 of the Rochester Conference Poster



impedance continuity at
event horizon suggests
impedance model sees
gravity in 'flat space'

G is gone – one less fundamental constant
it cancels out in the impedance calculation
What is the origin of 10.23 offset?

Quantum Impedances, Entanglement, and State Reduction

Peter Cameron*
Strongarm Studios
Mattituck, NY USA 11952

first draft posted
March 2013

(Dated: May 11, 2013)

The measurement problem, the mechanism of quantum state reduction, has remained an open question for nearly a century. The 'quantum weirdness' of the problem was highlighted by the introduction of the Einstein-Podolsky-Rosen paradox in 1935. Motivated by Bell's Theorem nonlocality was first experimentally observed in 1972 by Clauser and Freedman in the entangled states of an EPR experiment, and is now an accepted fact. Special relativity requires that no energy is transferred in the nonlocal collapse of these entangled two-body wavefunctions, that no work is done, no information communicated. In the family of quantum impedances those which are scale invariant, the Lorentz and centrifugal impedances, satisfy this requirement. This letter explores their role in the collapse of the wave function.

A Possible Resolution of the Black Hole Information Paradox

Peter Cameron

Strongarm Studios, PO Box 1030, Mattituck, NY 11952

petethepop@aol.com

Abstract: Nonlocal reduction of entangled states is clarified by considering the role of background independent scale-invariant quantum impedances in decay/decoherence of unstable elementary particles, providing simple resolution of the black hole information paradox.

© 2013 Optical Society of America

OCIS codes: 030.1640 Coherence, 270.5585 Quantum information and processing

*OSA refereed, as was
Berlin Conference*

1. Introduction

Decay of the unstable particles offers the possibility of informing nonlocal reduction of entangled states. Both follow from phase decoherence (with the resultant complication that phase is not an observable in state reduction). Unlike entangled states, where unitary evolution of the two (or more) body wave function requires nonlocal phase coherence, in the case of the unstable particles the essential coherence is self-coherence.

4. The Black Hole Information Paradox

An earlier note [8] calculated the impedance mismatch between the electron and the Planck particle. This mismatch is precisely equal to the ratio of the gravitational and electromagnetic forces between these two particles, indicating that the quantum impedance approach is valid at the event horizon, and perhaps beyond, to the singularity (which is completely decoupled by the infinitely large impedance mismatch at the dimensionless 'point').

As regards the paradox, if the scale invariant impedances are valid at the event horizon and responsible for nonlocal state reduction, and the holographic principle applies, then the paradox is removed.

Presentation Goals - Understand, then Explore

- **understand** the One Slide
 - ~~essential point~~ — impedances govern the flow of energy
 - ~~origin~~ of impedance network — two body problem & Mach's Principle
 - **origin of the coherence length plot** — Malcolm MacGregor
 - **implications** of their strong correlation — personal paradigm shift
- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - ~~gravity — extend the model to the Planck particle~~
 - ~~state reduction and non-locality~~
 - ~~black hole information paradox — Rochester, Fields,...~~
 - weak measurement and time symmetry — Berlin
 - quantum interpretations — Berlin
 - *dark matter*
 - *condensed matter*

Italics connote material in preparation

The 36 metastable particles with lifetime $\tau > 10^{-21}$ sec


UQ = unpaired-quark cluster
PQ = paired-quark or radiative cluster

● unpaired meson
○ paired meson
▲ unpaired baryon
☆ excited state
◆ lepton

$$\tau_i / \tau_{\pi^\pm} = \alpha^{x_i}$$

Higher mass s, c, b flavor clusters

Low mass (LM) particles
($m < 1 \text{ GeV}/c^2$)



Copyrighted Material

α

THE POWER OF

Electron Elementary Particle Generation
with α -Quantized Lifetimes and Masses

MALCOLM H. MAC GREGOR


Copyrighted Material

[19] M. H. MacGregor, "The Fine-Structure Constant as a Universal Scaling Factor", Lett. Nuovo Cimento **1**, 759-764 (1971)

[20] M. H. MacGregor, "The Electromagnetic Scaling of Particle Lifetimes and Masses", Lett. Nuovo Cimento **31**, 341-346 (1981)

[21] M. H. MacGregor, The Power of Alpha, World Scientific (2007) see also <http://70mev.org/alpha/>


MacGregor's lifetime plot with his factor of 2, 3, and 4 corrections



from 2011 – making the connection between impedances and lifetimes

Full Spectrum (neutrino?) of Unstable Particle Lifetimes

From previous slide with horizontal axis reversed



Electromagnetic decays are possible only when the impedance match is reasonably good, Unstable particles that exist on longer time/length scales than the Rydberg cannot impedance match to the photon, require parity violation and the neutrino



'One Slide'

Strong Decay

EM Decay

Weak Decay

$\pi_0 \rightarrow 2\gamma$	98.8%
$\pi_0 \rightarrow e^+e^-\gamma$	1.2%
$\eta \rightarrow 2\gamma$	39.3%
$\eta \rightarrow 3\pi^0$	32.6%
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.7%
$\eta \rightarrow \pi^+\pi^-\gamma$	4.6%
$\eta' \rightarrow \pi^+\pi^-\eta$	39.3%
$\eta' \rightarrow \rho^0\gamma$	
incl. $\pi^+\pi^-\gamma$	29.3%
$\eta' \rightarrow \pi^0\pi^0\eta$	21.6%
$\eta' \rightarrow \omega\gamma$	2.8%
$\eta' \rightarrow \gamma\gamma$	2.2%



- Horizontal axis in both plots is logarithmic
- Length scales are the same for both plots, and they are properly aligned
- Upper plot shows particle lifetimes multiplied by the speed of light, the coherence lengths (adapted from The Power of Alpha by Malcolm MacGregor)
- Plot at left is electron impedances, details at <http://redshift.vif.com/JournalFiles/V18N02PDF/V18N2CAM.pdf>
- Alpha-spaced coherence lengths of η' , η , and π_0 are at conjunctions of mode impedances, can couple to the photon for fast EM decay
- Location of π_0 at boundary of EM and weak decays is suggestive of the chiral anomaly
- Weak decays are mismatched to the photon
- Alpha-spaced coherence length alternation of fermions/bosons in weak decays appears to be related to parity violation and electroweak enhancement of charm/tau coherence lengths
- Clustering of superheavies (top, Higgs, Z, W) at the 10 GeV coherence length is tantalizing
- Impedance junctions at the classical radius are related to mass quantization and MacGregor's 70 MeV 'platform state'
- See also http://vixra.org/author/peter_cameron

Presentation Goals - **Explore**, then Understand

- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - ~~gravity—extend the model to the Planck particle~~
 - ~~state reduction and non-locality~~
 - ~~black hole information paradox—Rochester, Fields,...~~
 - **weak measurement and time Symmetry – Berlin**
 - **quantum interpretations – Berlin**
 - *dark matter*
 - *condensed matter*
- **understand** the One Slide
 - ~~**essential point** – impedances govern the flow of energy~~
 - ~~**origin** of impedance network—two body problem & Mach's Principle~~
 - ~~**origin** of the coherence length plot—Malcolm MacGregor~~
 - **implications** of their strong correlation – personal paradigm shift

Delayed Choice and Weak Measurement in the Nested Mach-Zehnder Interferometer

*accepted for presentation at the
2014 Berlin Conference on
Quantum Information and Measurement*

Peter Cameron
Strongarm Studios
Mattituck, NY USA 11952
petethepop@aol.com


*OSA refereed, as was the
Rochester Conference. This paper
was motivated by Lev Vaidman's
excellent talk at Rochester.*

Abstract: This note discusses interpretation of recent weak nested interferometer measurements in terms of state vectors traveling both forward and backward in time. A compatible quantum impedance interpretation is presented. Delayed choice variants are proposed.

OCIS codes: 000.2658, 270.5585

5. Conclusion

There appears to be a connection between invariant impedances, weak measurement, and time symmetry. Invariant impedances transfer no energy, only phase. Weak measurement measures phase, which is acausal. Thus the impedance model is compatible with a TSVF of state vectors coupled by invariant impedances only. A test of the TSVF interpretation would then look for f1 and f2 with insertion of BS2 after the photon has passed (!) and before $t=3\sqrt{2}$.



Viewpoint: What Can we Say about a Photon's Past?

Jeff Lundeen, Department of Physics, University of Ottawa, MacDonald Hall, 150 Louis Pasteur Road, Ottawa, Ontario K1N 6N5, Canada

Published December 9, 2013 | Physics 6, 133 (2013) | DOI: 10.1103/Physics.6.133

An experiment demonstrates that even when physicists think a quantum particle has followed a single path it might not have.

Quantum Interpretation of the Impedance Model

*accepted for presentation at the
2014 Berlin Conference on
Quantum Information and
Measurement*

Michaele Suisse and Peter Cameron
Strongarm Studios
Mattituck, NY USA 11952
michaele.suisse@gmail.com, petethepop@aol.com

*OSA refereed, as was the
Rochester Conference*

Abstract: Quantum Interpretations try to explain emergence of the world we observe from formal quantum theory. Impedances govern the flow of energy, are helpful in such attempts. We include quantum impedances in comparisons of selected interpretations.
OCIS codes: 000.6800, 270.5585

Index	Interpretation	Authors	non-local?	probabilistic?	hidden variables?	wavefcn real?	wavefcn collapse?	universal wavefcn?	observer role?	unique history?
30	Objective Collapse	GRW 1986, Penrose 1989	Yes	Yes	No	Yes	Yes	No	No	Yes
30	Transactional	Cramer 1986	Yes	Yes	No	Yes	Yes	No	No	Yes
30	Quantum Impedances	Cameron & Suisse 2013	Yes	Yes	No	Yes	Yes	No	No	Yes
25	Relational	Rovelli 1994	No	Yes	No	No	Yes	No	No	agnostic
23	Quantum Logic	Birkhoff 1936	agnostic	agnostic	No	agnostic	No	No	No	Yes
17	Ithaca	Mermin 1996	No	Yes	No	No	No	No	No	No
15	Consistent Histories	Griffiths 1984	No	agnostic	No	agnostic	No	No	No	No
15	Copenhagen	Bohr & Heisenberg 1927	No	Yes	No	No	Yes	No	Yes	Yes
9	Qbism	Caves, Fuchs, Schack 2002	No	Yes	No	No	Yes	No	Yes	No
6	Orthodox	von Neumann 1932	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
-3	Many Worlds	Everett 1957	No	No	No	Yes	No	Yes	No	No
-18	de Broglie – Bohm	de Broglie 1927, Bohm 1952	Yes	No	Yes	Yes	No	Yes	No	Yes

Fig. 1. Comparison of the Interpretations. The Index parameter quantifies the strength of agreement between a given interpretation and the rest of the table. Values in the Index column are calculated by adding a point for entries that agree with a given interpretation, subtracting for entries that disagree, and giving half values for the agnostics.

Presentation Goals - **Explore**, then Understand

- **explore** the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - ~~gravity—extend the model to the Planck particle~~
 - ~~state reduction and non-locality~~
 - ~~black hole information paradox—Rochester, Fields,...~~
 - ~~weak measurement and time Symmetry—Berlin~~
 - ~~quantum interpretations—Berlin~~
 - **dark matter** *Italics connote material in preparation*
 - **condensed matter**
- **understand** the One Slide
 - ~~essential point - impedances govern the flow of energy~~
 - ~~origin of impedance network—two body problem & Mach's Principle~~
 - ~~origin of the coherence length plot—Malcolm MacGregor~~
 - **implications** of their strong correlation – personal paradigm shift

from the Electron Impedances note

	'spinor' flux quantum	monopole charge quantum	dipole dipole quantum
electric	dark	observable	dark
magnetic	observable	dark	observable

the constituents of the impedance model

from the chiral anomaly note

'Dark' Modes and Anomalies

The impedance plot of figure 2 is not complete.

Absent are the longitudinal dipole-dipole impedances, the longitudinal and transverse charge-dipole impedances (the charge-dipole impedances are a subset of the scale invariant three body impedances), and the Coriolis impedance. There may be others, and likely are. Given the spin dependence of the weak interaction, one would expect that adding the longitudinal impedances to the figure would give additional insight into the weak decays, probably essential for instance in impedance-based calculations of those branching ratios.

Present in the plot are several impedances that (excepting the unstable particle spectrum) are absent in our observations of the world, do not couple to the photon,

namely those associated with the electric flux quantum, magnetic monopole, and electric dipole. Figure 6 shows the alternation with topological complexity.


We see the magnetic flux quantum, electric monopole, and magnetic dipole in the stable particles which comprise our physical world, but not their electromagnetic complements. It seems that the only place we see these 'dark' components is in the unstable particle spectrum. This broken symmetry is partially understood in terms of the relative strengths of the magnetic and electric charge quanta[12, 88], and might have a not-yet-obvious role in the chiral anomaly.

[88] T. Datta, "The fine structure constant, magnetic monopoles, and the Dirac charge quantization condition", Lett. Nuovo Cim. **37** 2 (1983)
<http://link.springer.com/article/10.1007>

bonus - EDM $\mu_B = \frac{e\hbar}{2m_e c}$

a product of electric charge **e** and distance **h/mc** define EDM, not MDM.


in near field of the observables the electric flux quantum is coupling to magnetic impedance, and magnetic flux quantum to electric impedance.



Observable + Dark

[88] T. Datta, "The fine structure constant, magnetic monopoles, and the Dirac charge quantization condition", Lett. Nuovo Cim. 37 2 (1983)
<http://link.springer.com/article/10.1007>

Observable Only




condensed matter

- if the concept of generalized quantum impedances is eventually perceived and accepted to be of some value, the **practical realizations** will of course be in **condensed matter**.
- a good place to start is perhaps impedance matching all (or as many as possible) of the electron impedances to the graphene lattice
- helpful references might be found in the bibliography of the chiral anomaly note, as shown here and on the following pages

[86] L. Brillouin, Wave Propagation in Periodic Structures - Electric Filters and Crystal Lattices, 2nd Ed., McGraw Hill (1946).

[87] R. Collins, "The Doppler Origin of deBroglie Waves", APS preprint server (1998)

- at right is a plot of 'relevant papers per year'



of the references on the following three slides, where relevant is taken to mean potentially useful in understanding possible roles of generalized quantum impedances in condensed matter. Coverage ends mid-2012.

Trending upwards nicely.

- [40] A. Khondker et.al, "Transmission Line Analogy of Resonance Tunneling: The Generalized Impedance Concept," J. Appl. Phys., **63** (10) 5191-5193 (1988)
- [41] S. Luryi, "Quantum Capacitance Devices", Appl. Phys. Lett. **52** 6 (1988)
- [42] S. Girvin, "Quantum Fluctuations and the Single-Junction Coulomb blockade", PRL **64** 26 (1990)
- [43] S. Kabir et.al, "Application of Quantum Mechanical Wave Impedance in the Solution of the Schrodinger Equation in Quantum Wells", Solid-State Electronics **34** (12), 1466-1468 (1991)
- [44] J. Rosner, "The Smith Chart and Quantum Mechanics" Am. J. Phys. **61** 310 (1993) <http://dx.doi.org/10.1119/1.17262>
- [45] H. Tung and C. Lee, "An Energy Bandpass Filter Using Superlattice Structures", IEEE Jour. Quant. Elect., **32** (3) 507-512 (1996)
- [46] V. Privmana et.al, "Quantum computation in quantum-Hall systems", Phys. Lett. A **239** (3) 141146 (1998) <http://arxiv.org/abs/quant-ph/9707017>
- [47] C. Chang and C. Kou, "Electron-wave quantum well energy band-pass filters", J. Phys. D **32** 139146 (1999)
- [48] Y. Imry and R. Landauer, "Conductance Viewed as Transmission", Rev. Mod. Phys. **72** (2) 306-312 (1999)
- [49] T. Datta and R. Tsu, "Quantum Wave Resistance of Schrodinger Functions" (2003) <http://arxiv.org/abs/cond-mat/0311479>
- [50] F. Delahaye and B. Jeckelmann, "Revised technical guidelines for reliable dc measurements of the quantized Hall resistance", Metrologia **40** 217 (2003)
- [51] K. von Klitzing, "25 Years of Quantum Hall Effect", Seminaire Poincare **2** 1-16 (2004) <http://www.bourbaphy.fr/klitzing.pdf>
- [52] F. Fischer and M. Grayson, "Influence of voltmeter impedance on quantum Hall measurements" Jour. Appl. Phys. **98** 013710 (2005)
- [53] K. Pham, "Probing the intrinsic shot noise of a Luttinger Liquid through impedance matching" (2005) <http://arxiv.org/abs/cond-mat/0504389>
- [54] Y. Katayama, "Physical and Circuit Modeling of Coupled Open Quantum Systems" Proc. 5th IEEE Conf. on Nanotechnology, Nagoya (2005) <http://arxiv.org/abs/cond-mat/0504389>
- [55] J. Gabelli et.al, "Violation of Kirchoffs Laws for a Coherent RC Circuit", Science **313** 499-502 (2006)
- [56] P. Burke et.al, "Quantitative theory of nanowire and nanotube antenna performance", IEEE Trans. Nanotech. **5** (4) 314-334 (2006) <http://arxiv.org/pdf/cond-mat/0408418.pdf>
- [57] S. Corlevi, "Quantum effects in nanoscale Josephson junction circuits", KTH PhD thesis, Stockholm (2006)
- [58] C. Day, "Tiny oscillating circuit exhibits new quantization of electrical conductance", Physics Today (Sep 2006) http://www.phys.ens.fr/~placais/publication/2006_PhysicsToday_C.DayN&V.pdf

- [59] E. Nelin, "Impedance model for quantum-mechanical barrier problems", *Physics-Uspekhi* **50** 3 293 (2007) http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=ufn&paperid=440&option_lang=eng
- [60] J. Seminario, "Quantum Current-Voltage Relation for a Single Electron", *J. Phys. B* **40** F275F276 (2007)
- [61] K. Novoselov et.al, "Room-Temperature Quantum Hall Effect in Graphene", *Science* **315** (5817) 1379 (2007) <http://arxiv.org/abs/cond-mat/0702408>
- [62] D. Ferry, "Semiconductor Device Scaling: Physics, Transport, and the Role of Nanowires", *Int. Jour. High Speed Electronics and Systems* **17** 3 (2007)
- [63] K. Pham, "Exact Green's Functions and Bosonization of a Luttinger Liquid Coupled to Impedances" (2007), *Prog. Th. Phys.* **118** 3 (2007) <http://arxiv.org/abs/0709.0475>
- [64] S. Shevchenko, "Impedance measurement technique for quantum systems", *EPJ B* **61** 187-191 (2008) <http://arxiv.org/abs/0708.0464>
- [65] R. Schoelkopf and S. Girvin, "Wiring up quantum systems", *Nature* **451** 664-669 (2008)
- [66] B. Zeldovich, "Impedance and parametric excitation of oscillators", *Physics-Uspekhi* **51** (5) 465-484 (2008)
- [67] A. Compston et.al, "A Fundamental Limit on Antenna Gain for Electrically Small Antennas", *IEEE Sarnoff Symposium*, Princeton (2008) <http://ieeexplore.ieee.org/xpl/articleDetails.jsp?arnumber=4520113>
- [68] H. Torres-Silva, "New Interpretation of the Atomic Spectra of the Hydrogen Atom: A Mixed Mechanism of Classical LC Circuits and Quantum Wave-Particle Duality", *Revista Chilena de Ingeniera* **16** (numero especial) 24-30 (2008)
- [69] R. Tsu and T. Datta, "Conductance and Wave Impedance of Electrons", *Prog. in Electromagnetics Res. Symp.*, Hangzhou, China, (2008)
- [70] A. Villegas et.al, "Longwave Phonon Tunneling Using an Impedance Concept", *PIERS Online*, **4** (2) 227-230 (2008)
- [71] E. Nelin, "Impedance Characteristics of Crystal-like Structures", *Tech. Phys.* **54** (7) 953-957 (2009)
- [72] E. Nelin, "Resonance Parameters of Double-Barrier Structures", *Tech. Phys. Lett.* **35** (5) 443-445 (2009)
- [73] D. Timofeev et.al, "Electronic Refrigeration at the Quantum Limit", *PRL* **102** 200801 (2009) <http://arxiv.org/abs/0902.2584>

- [74] M. Afzelius and C. Simon, "Impedance-matched cavity quantum memory", Phys. Rev. A **82** 022310 (2010) <http://arxiv.org/abs/1004.2469>
- [75] C. Mora and H. LaHur, "Universal resistances of the quantum resistance-capacitance circuit", Nature Phys. **6** 697701 (2010) <http://arxiv.org/abs/0911.4908>
- [76] J. Griffet et.al, "Impedance of a Nanoantenna and a Single Quantum Emitter", PRL **105** 117701 (2010)
- [77] D. Yoh et.al, "Single-walled carbon nanotubes as excitonic optical wires", Nature Nanotechnology **6** 51-56 (2011) <http://www.nature.com/nnano/journal/v6/n1/full/nnano.2010.248.html>
- [78] V. Sorger et.al, "Experimental demonstration of low-loss optical waveguiding at deep sub-wavelength scales", Nature Communications **2** 331 (2011)
- [79] R. Ottens et.al, "Near-Field Radiative Heat Transfer between Macroscopic Planar Surfaces" PRL **107** 014301 (2011) <http://arxiv.org/abs/1103.2389>
- [80] R. Tsu, Superlattice to Nanoelectronics, 2nd Ed., Elsevier, Chapter 11 (2011)
- [81] L. Novotny, "From near-field optics to optical antennas", Physics Today p.47 (July 2011) <http://www.optics.rochester.edu/workgroups/novotny/papers/novotny11b.pdf>
- [82] K. Ando, "Electrically tunable spin injector free from the impedance mismatch problem", Nature Materials **10** 655659 (2011) <http://www.optics.rochester.edu/workgroups/novotny/papers/novotny11b.pdf>
- [83] D. Ferry, "Ohm's Law in a Quantum World", Science **335** (6064) 45-46 (Jan 2012) <http://www.asu.edu/news/Science-2012-Ferry-45-6.pdf>
- [84] C. Bruot et.al, "Mechanically controlled molecular orbital alignment in single molecule junctions", Nature Nanotechnology **7** 3540 (2012) <http://www.nature.com/nnano/journal/v7/n1/full/nnano.2011.212.html>
- [85] H. Vasquez et.al, "Probing the conductance superposition law in single-molecule circuits with parallel paths", Nature Nanotechnology (2012)
- [86] L. Brillouin, Wave Propagation in Periodic Structures - Electric Filters and Crystal Lattices, 2nd Ed., McGraw Hill (1946).
- [87] R. Collins, "The Doppler Origin of deBroglie Waves", APS preprint server (1998)

Presentation Goals - Understand, then Explore

- **understand** the One Slide
 - ~~essential point~~ — impedances govern the flow of energy
 - ~~origin~~ of impedance network — two body problem & Mach's Principle
 - ~~origin~~ of the coherence length plot — Malcolm MacGregor
 - *implications of their strong correlation — personal paradigm shift*
- ~~explore~~ the Implications ~ ten reasons one might want to know about quantum impedances:
 - ~~elementary particle spectrum, chiral anomaly, axions, EDM...~~
 - ~~gravity — extend the model to the Planck particle~~
 - ~~state reduction and non-locality~~
 - ~~black hole information paradox — Rochester, Fields,...~~
 - ~~weak measurement and time symmetry — Berlin~~
 - ~~quantum interpretations — Berlin~~
 - *dark matter*
 - *condensed matter*

Italics connote material in preparation

Thank You for Your Time and Attention