

# **Electro-Magnetic Field Equation and Wave Function in Rindler spacetime**

**Sangwha-Yi**

**Department of Math , Taejon University 300-716**

## **ABSTRACT**

In the general relativity theory, we find the electro-magnetic field transformation and the electro-magnetic field equation (Maxwell equation) in Rindler spacetime. We find the electro-magnetic wave equation and the electro-magnetic wave function in Rindler space-time. Specially, this article say the uniqueness of the accelerated frame because the accelerated frame can treat electro-magnetic field equation.

**PACS Number:**04,04.90.+e,03.30, 41.20

**Key words:**General relativity theory,

Rindler spacetime,

Electro-magnetic field transformation,

Electro-magnetic field equation

Wave function

Wave equation

**e-mail address:**sangwhal@nate.com

**Tel:**051-624-3953

## 1. Introduction

In the general relativity theory, our article's aim is that we find the electro-magnetic field equation in Rindler space-time.

The Rindler coordinate is

$$ct = \left( \frac{c^2}{a_0} + \xi^1 \right) \sinh\left( \frac{a_0 \xi^0}{c} \right)$$

$$x = \left( \frac{c^2}{a_0} + \xi^1 \right) \cosh\left( \frac{a_0 \xi^0}{c} \right) - \frac{c^2}{a_0}, y = \xi^2, z = \xi^3 \quad (1)$$

In this time, the tetrad  $\theta^a_\mu$  is

$$d\tau^2 = dt^2 - \frac{1}{c^2} [dx^2 + dy^2 + dz^2]$$

$$= -\frac{1}{c^2} \eta_{ab} \frac{\partial x^a}{\partial \xi^\mu} \frac{\partial x^b}{\partial \xi^\nu} d\xi^\mu d\xi^\nu$$

$$= -\frac{1}{c^2} \eta_{ab} \theta^a_\mu \theta^b_\nu d\xi^\mu d\xi^\nu = -\frac{1}{c^2} g_{\mu\nu} d\xi^\mu d\xi^\nu, \quad \theta^a_\mu = \frac{\partial x^a}{\partial \xi^\mu} \quad (2)$$

$$\theta^a_0(\xi^0) = \frac{\partial x^a}{\partial \xi^0} = ((1 + \frac{a_0}{c^2} \xi^1) \cosh(\frac{a_0 \xi^0}{c}), (1 + \frac{a_0}{c^2} \xi^1) \sinh(\frac{a_0 \xi^0}{c}), 0, 0) \quad (3)$$

About  $y$ -axis's and  $z$ -axis's orientation

$$\theta^a_2(\xi^0) = \frac{\partial x^a}{\partial \xi^2} = (0, 0, 1, 0), \quad \theta^a_3(\xi^0) = \frac{\partial x^a}{\partial \xi^3} = (0, 0, 0, 1) \quad (4)$$

The other unit vector  $\theta^a_1(\xi^0)$  is

$$\theta^a_1(\xi^0) = \frac{\partial x^a}{\partial \xi^1} = (\sinh(\frac{a_0 \xi^0}{c}), \cosh(\frac{a_0 \xi^0}{c}), 0, 0) \quad (5)$$

Therefore,

$$\begin{aligned} cdt &= c \cosh(\frac{a_0 \xi^0}{c}) d\xi^0 (1 + \frac{a_0}{c^2} \xi^1) + \sinh(\frac{a_0 \xi^0}{c}) d\xi^1 \\ &= c \cosh(\frac{a_0 \xi^0}{c}) d\xi^0 + \sinh(\frac{a_0 \xi^0}{c}) d\xi^1 \\ dx &= c \sinh(\frac{a_0 \xi^0}{c}) d\xi^0 (1 + \frac{a_0}{c^2} \xi^1) + \cosh(\frac{a_0 \xi^0}{c}) d\xi^1 \end{aligned}$$

$$= c \sinh\left(\frac{a_0 \xi^0}{c}\right) d\xi^0 + \cosh\left(\frac{a_0 \xi^0}{c}\right) d\xi^1, dy = d\xi^2 = d\xi^2, dz = d\xi^3 = d\xi^3 \quad (6)$$

The vector transformation is

$$V^\mu = \frac{\partial x^\mu}{\partial x^\alpha} V^\alpha, \quad U_\mu = \frac{\partial x^\alpha}{\partial x^\mu} U_\alpha \quad (7)$$

Therefore, the transformation of the electro-magnetic 4-vector potential  $(\phi, \vec{A}) = A^\alpha$  is

$$A^\alpha = \frac{\partial x^\alpha}{\partial x^\mu} A^\mu = \frac{\partial x^\alpha}{\partial \xi^\mu} A_\xi^\mu = e^\alpha_\mu A_\xi^\mu, \quad e^\alpha_\mu = \frac{\partial x^\alpha}{\partial \xi^\mu}$$

$$dx^\alpha = \frac{\partial x^\alpha}{\partial x^\mu} dx^\mu = \frac{\partial x^\alpha}{\partial \xi^\mu} d\xi^\mu = e^\alpha_\mu d\xi^\mu, \quad e^\alpha_\mu = \frac{\partial x^\alpha}{\partial \xi^\mu} \quad (8)$$

Hence, the transformation of the electro-magnetic 4-vector potential  $(\phi, \vec{A})$  in inertial frame and the electro-magnetic 4-vector potential  $(\phi_\xi, \vec{A}_\xi)$  in uniformly accelerated frame is

$$\begin{aligned} & \left( \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \phi = 4\pi\rho \\ & \left( \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \vec{A} = \frac{4\pi}{c} \vec{j} \\ & \text{4-vector } (c\rho, \vec{j}) = \rho_0 \frac{dx^\alpha}{d\tau} \\ & \phi = \cosh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \sinh\left(\frac{a_0 \xi^0}{c}\right) A_\xi^1 \\ & = \cosh\left(\frac{a_0 \xi^0}{c}\right) \hat{\phi}_\xi + \sinh\left(\frac{a_0 \xi^0}{c}\right) \hat{A}_\xi^1 \\ & A_x = \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \cosh\left(\frac{a_0 \xi^0}{c}\right) A_\xi^1 \\ & = \sinh\left(\frac{a_0 \xi^0}{c}\right) \hat{\phi}_\xi + \cosh\left(\frac{a_0 \xi^0}{c}\right) \hat{A}_\xi^1 \\ & A_y = A_{\xi^2} = \hat{A}_{\xi^2}, A_z = A_{\xi^3} = \hat{A}_{\xi^3} \end{aligned} \quad (9)$$

$$g = \begin{pmatrix} -(1 + \frac{a_0 \xi^1}{c^2})^2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \eta = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$e^a{}_\mu e_b{}^\mu = \delta^a{}_b, \quad e^a{}_\mu e_a{}^\nu = \delta_\mu{}^\nu$$

$$e^a{}_\mu e^b{}_\nu \eta_{ab} = g_{\mu\nu} \rightarrow A^T \eta A = g$$

$$e_a{}^\mu e_b{}^\nu g_{\mu\nu} = \eta_{ab} \rightarrow (A^T)^{-1} g A^{-1} = (A^T)^{-1} A^T \eta A A^{-1} = \eta$$

$$e^a{}_\mu = \eta^{ab} g_{\mu\nu} e_b{}^\nu \rightarrow \eta^{-1} (A^T)^{-1} A^T \eta A = A = \eta^{-1} (A^T)^{-1} g \quad (10)$$

$$\begin{pmatrix} cd t \\ dx \\ dy \\ dz \end{pmatrix} = \begin{pmatrix} \cosh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0 \xi^1}{c^2}) & \sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ \sinh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0 \xi^1}{c^2}) & \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} cd\xi^0 \\ d\xi^1 \\ d\xi^2 \\ d\xi^3 \end{pmatrix}$$

$$= A \begin{pmatrix} cd\xi^0 \\ d\xi^1 \\ d\xi^2 \\ d\xi^3 \end{pmatrix}$$

$$= \begin{pmatrix} \cosh(\frac{a_0 \xi^0}{c}) & \sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ \sinh(\frac{a_0 \xi^0}{c}) & \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} cd\hat{\xi}^0 \\ d\hat{\xi}^1 \\ d\hat{\xi}^2 \\ d\hat{\xi}^3 \end{pmatrix}$$

(11)

$$\begin{aligned}
e_\mu^\alpha = \frac{\partial \xi^\alpha}{\partial x^\mu} &= A^{-1} = \begin{pmatrix} \frac{c\partial\xi^0}{\partial t} & \frac{c\partial\xi^0}{\partial x} & \frac{c\partial\xi^0}{\partial y} & \frac{c\partial\xi^0}{\partial z} \\ \frac{\partial\xi^1}{\partial t} & \frac{\partial\xi^1}{\partial x} & \frac{\partial\xi^1}{\partial y} & \frac{\partial\xi^1}{\partial z} \\ \frac{\partial\xi^2}{\partial t} & \frac{\partial\xi^2}{\partial x} & \frac{\partial\xi^2}{\partial y} & \frac{\partial\xi^2}{\partial z} \\ \frac{\partial\xi^3}{\partial t} & \frac{\partial\xi^3}{\partial x} & \frac{\partial\xi^3}{\partial y} & \frac{\partial\xi^3}{\partial z} \end{pmatrix} \\
&= \begin{pmatrix} \cosh(\frac{a_0\xi^0}{c}) & -\sinh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ \frac{(1+\frac{a_0\xi^1}{c^2})}{c^2} & \frac{(1+\frac{a_0\xi^1}{c^2})}{c^2} & 0 & 0 \\ -\sinh(\frac{a_0\xi^0}{c}) & \cosh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{12}
\end{aligned}$$

$$\begin{aligned}
\begin{pmatrix} \frac{1}{c}\frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix} &= (A^{-1})^T \begin{pmatrix} \frac{1}{c}\frac{\partial}{\partial\xi^0} \\ \frac{\partial}{\partial\xi^1} \\ \frac{\partial}{\partial\xi^2} \\ \frac{\partial}{\partial\xi^3} \end{pmatrix} = (A^T)^{-1} \begin{pmatrix} \frac{1}{c}\frac{\partial}{\partial\xi^0} \\ \frac{\partial}{\partial\xi^1} \\ \frac{\partial}{\partial\xi^2} \\ \frac{\partial}{\partial\xi^3} \end{pmatrix} \\
&= \begin{pmatrix} \cosh(\frac{a_0\xi^0}{c}) & -\sinh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ \frac{(1+\frac{a_0\xi^1}{c^2})}{c^2} & \frac{(1+\frac{a_0\xi^1}{c^2})}{c^2} & 0 & 0 \\ -\sinh(\frac{a_0\xi^0}{c}) & \cosh(\frac{a_0\xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{c}\frac{\partial}{\partial\xi^0} \\ \frac{\partial}{\partial\xi^1} \\ \frac{\partial}{\partial\xi^2} \\ \frac{\partial}{\partial\xi^3} \end{pmatrix}
\end{aligned}$$

$$= \begin{pmatrix} \cosh(\frac{a_0 \xi^0}{c}) & -\sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ -\sinh(\frac{a_0 \xi^0}{c}) & \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{c} \frac{\partial}{\partial \xi^0} \\ \frac{\partial}{\partial \xi^1} \\ \frac{\partial}{\partial \xi^2} \\ \frac{\partial}{\partial \xi^3} \end{pmatrix} \quad (13)$$

$$\begin{aligned} \frac{1}{c} \frac{\partial}{\partial t} &= \frac{c \partial \xi^0}{c \partial t} \frac{1}{c} \frac{\partial}{\partial \xi^0} + \frac{\partial \xi^1}{c \partial t} \frac{\partial}{\partial \xi^1} \\ &= \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \\ &= \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \hat{\xi}^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \hat{\xi}^1} \\ \frac{\partial}{\partial x} &= \frac{c \partial \xi^0}{\partial x} \frac{1}{c} \frac{\partial}{\partial \xi^0} + \frac{\partial \xi^1}{\partial x} \frac{\partial}{\partial \xi^1} \\ &= -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \\ &= -\sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \hat{\xi}^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \hat{\xi}^1} \\ \frac{\partial}{\partial y} &= \frac{\partial}{\partial \xi^2} = \frac{\partial}{\partial \hat{\xi}^2}, \quad \frac{\partial}{\partial z} = \frac{\partial}{\partial \xi^3} = \frac{\partial}{\partial \hat{\xi}^3} \\ \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 &= \frac{1}{c^2 (1 + \frac{a_0 \xi^1}{c^2})^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_{\xi}^2 = \frac{1}{c^2} (\frac{\partial}{\partial \hat{\xi}^0})^2 - \nabla_{\hat{\xi}}^2 \\ \vec{\nabla} &= (\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}), \quad \vec{\nabla}_{\xi} = (\frac{\partial}{\partial \xi^1}, \frac{\partial}{\partial \xi^2}, \frac{\partial}{\partial \xi^3}), \quad \vec{\nabla}_{\hat{\xi}} = (\frac{\partial}{\partial \hat{\xi}^1}, \frac{\partial}{\partial \hat{\xi}^2}, \frac{\partial}{\partial \hat{\xi}^3}) \end{aligned} \quad (14)$$

## 2. Electro-magnetic Field in the Rindler space-time

The electro-magnetic field  $(\vec{E}, \vec{B})$  is in the inertial frame,

$$\vec{E} = -\vec{\nabla}\phi - \frac{\partial \vec{A}}{\partial t}, \vec{B} = \vec{\nabla} \times \vec{A} \quad (15)$$

$$E_x = -\frac{\partial \phi}{\partial x} - \frac{\partial A_x}{\partial t}$$

$$= -\left[ -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot \left[ \cosh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0 \xi^1}{c^2})\phi_\xi + \sinh(\frac{a_0 \xi^0}{c})A_{\xi^1} \right]$$

$$= -\left[ -\frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot \left[ \sinh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0 \xi^1}{c^2})\phi_\xi + \cosh(\frac{a_0 \xi^0}{c})A_{\xi^1} \right]$$

$$= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^1}}{\partial \xi^0} - (1 + \frac{a_0 \xi^1}{c^2}) \frac{\partial \phi_\xi}{\partial \xi^1} - 2\phi_\xi \frac{a_0}{c^2}$$

$$= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^1} \left[ (1 + \frac{a_0}{c^2} \xi^1)^2 \phi_\xi \right] - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^1}}{\partial \xi^0}$$

$$= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^1} \left[ (1 + \frac{a_0}{c^2} \xi^1) \hat{\phi}_\xi \right] - \frac{\partial \hat{A}_{\xi^1}}{\partial \xi^0} \quad (16)$$

$$E_y = -\frac{\partial \phi}{\partial y} - \frac{\partial A_y}{\partial t} = -\frac{\partial}{\partial \xi^2} \left[ \cosh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0}{c^2} \xi^1)\phi_\xi + \sinh(\frac{a_0 \xi^0}{c})A_{\xi^1} \right]$$

$$= -\left[ -\frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] A_{\xi^2}$$

$$= -(1 + \frac{a_0 \xi^1}{c^2}) \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial \phi_\xi}{\partial \xi^2} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial A_{\xi^2}}{\partial \xi^0}$$

$$+ \sinh(\frac{a_0}{c} \xi^0) \left[ \frac{\partial A_{\xi^2}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^2} \right]$$

$$\begin{aligned}
&= \cosh\left(\frac{a_0}{c}\xi^0\right) \left[ -\frac{1}{(1+\frac{a_0}{c^2}\xi^1)} \frac{\partial}{\partial\xi^2} [\phi_\xi (1+\frac{a_0\xi^1}{c^2})^2] - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial A_{\xi^2}}{c\partial\xi^0} \right. \\
&\quad \left. + \sinh\left(\frac{a_0}{c}\xi^0\right) \left[ \frac{\partial A_{\xi^2}}{\partial\xi^1} - \frac{\partial A_{\xi^1}}{\partial\xi^2} \right] \right] \\
&= \cosh\left(\frac{a_0}{c}\xi^0\right) \left[ -\frac{1}{(1+\frac{a_0}{c^2}\xi^1)} \frac{\partial}{\partial\xi^2} [\hat{\phi}_\xi (1+\frac{a_0\xi^1}{c^2})] - \frac{\partial \hat{A}_{\xi^2}}{c\partial\xi^0} \right. \\
&\quad \left. + \sinh\left(\frac{a_0}{c}\xi^0\right) \left[ \frac{\partial \hat{A}_{\xi^2}}{\partial\xi^1} - \frac{\partial \hat{A}_{\xi^1}}{\partial\xi^2} \right] \right] \tag{17} \\
E_z &= -\frac{\partial\phi}{\partial z} - \frac{\partial A_z}{c\partial t} = -\frac{\partial}{\partial\xi^3} \left[ \cosh\left(\frac{a_0\xi^0}{c}\right) (1+\frac{a_0}{c^2}\xi^1)\phi_\xi + \sinh\left(\frac{a_0\xi^0}{c}\right) A_{\xi^1} \right] \\
&\quad - \left[ \frac{\cosh\left(\frac{a_0\xi^0}{c}\right)}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial}{c\partial\xi^0} - \sinh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial}{\partial\xi^1} \right] A_{\xi^3} \\
&= -(1+\frac{a_0\xi^1}{c^2}) \cosh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial\phi_\xi}{\partial\xi^3} - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \cosh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial A_{\xi^3}}{c\partial\xi^0} \\
&\quad + \sinh\left(\frac{a_0}{c}\xi^0\right) \left[ \frac{\partial A_{\xi^3}}{\partial\xi^1} - \frac{\partial A_{\xi^1}}{\partial\xi^3} \right] \\
&= \cosh\left(\frac{a_0}{c}\xi^0\right) \left[ -\frac{1}{(1+\frac{a_0}{c^2}\xi^1)} \frac{\partial}{\partial\xi^3} [\phi_\xi (1+\frac{a_0\xi^1}{c^2})^2] - \frac{1}{(1+\frac{a_0\xi^1}{c^2})} \frac{\partial A_{\xi^3}}{c\partial\xi^0} \right. \\
&\quad \left. + \sinh\left(\frac{a_0}{c}\xi^0\right) \left[ \frac{\partial A_{\xi^3}}{\partial\xi^1} - \frac{\partial A_{\xi^1}}{\partial\xi^3} \right] \right] \\
&= \cosh\left(\frac{a_0}{c}\xi^0\right) \left[ -\frac{1}{(1+\frac{a_0}{c^2}\xi^1)} \frac{\partial}{\partial\xi^3} [\hat{\phi}_\xi (1+\frac{a_0\xi^1}{c^2})] - \frac{\partial \hat{A}_{\xi^3}}{c\partial\xi^0} \right]
\end{aligned}$$

$$+ \sinh\left(\frac{a_0}{c} \xi^0\right) \left[ \frac{\partial \hat{A}_{\xi^3}}{\partial \xi^1} - \frac{\partial \hat{A}_{\xi^1}}{\partial \xi^3} \right] \quad (18)$$

$$B_x = \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = \frac{\partial A_{\xi^3}}{\partial \xi^2} - \frac{\partial A_{\xi^2}}{\partial \xi^3} = \frac{\partial \hat{A}_{\xi^3}}{\partial \xi^2} - \frac{\partial \hat{A}_{\xi^2}}{\partial \xi^3} \quad (19)$$

$$B_y = \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} = \frac{\partial A_x}{\partial \xi^3} - \frac{\partial A_{\xi^3}}{\partial x}$$

$$= \frac{\partial}{\partial \xi^3} \left[ \sinh\left(\frac{a_0 \xi^0}{c}\right) \left(1 + \frac{a_0}{c^2} \xi^1\right) \phi_\xi + \cosh\left(\frac{a_0 \xi^0}{c}\right) A_{\xi^1} \right]$$

$$\begin{aligned} & - \left[ - \frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^3} \\ & = \cosh\left(\frac{a_0}{c} \xi^0\right) \left[ \frac{\partial A_{\xi^1}}{\partial \xi^3} - \frac{\partial A_{\xi^3}}{\partial \xi^1} \right] \\ & - \sinh\left(\frac{a_0}{c} \xi^0\right) \left[ - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^3} \left[ \phi_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right)^2 \right] - \frac{1}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial A_{\xi^3}}{\partial \xi^0} \right] \\ & = \cosh\left(\frac{a_0}{c} \xi^0\right) \left[ \frac{\partial \hat{A}_{\xi^1}}{\partial \xi^3} - \frac{\partial \hat{A}_{\xi^3}}{\partial \xi^1} \right] \\ & - \sinh\left(\frac{a_0}{c} \xi^0\right) \left[ - \frac{1}{\left(1 + \frac{a_0}{c^2} \xi^1\right)} \frac{\partial}{\partial \xi^3} \left[ \hat{\phi}_\xi \left(1 + \frac{a_0 \xi^1}{c^2}\right) \right] - \frac{\partial \hat{A}_{\xi^3}}{\partial \xi^0} \right] \end{aligned} \quad (20)$$

$$B_z = \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = \frac{\partial A_{\xi^2}}{\partial x} - \frac{\partial A_x}{\partial \xi^2}$$

$$\begin{aligned} & = \left[ - \frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{\left(1 + \frac{a_0 \xi^1}{c^2}\right)} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] A_{\xi^3} \end{aligned}$$

$$\begin{aligned}
& -\frac{\partial}{\partial \xi^2} [\sinh(\frac{a_0 \xi^0}{c})(1 + \frac{a_0}{c^2} \xi^1) \phi_\xi + \cosh(\frac{a_0 \xi^0}{c}) A_{\xi^1}] \\
& = \cosh(\frac{a_0}{c} \xi^0) [\frac{\partial A_{\xi^2}}{\partial \xi^1} - \frac{\partial A_{\xi^1}}{\partial \xi^2}] \\
& + \sinh(\frac{a_0}{c} \xi^0) [-\frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^2} [\phi_\xi (1 + \frac{a_0 \xi^1}{c^2})^2] - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^2}}{c \partial \xi^0}] \\
& = \cosh(\frac{a_0}{c} \xi^0) [\frac{\partial \hat{A}_{\xi^2}}{\partial \xi^1} - \frac{\partial \hat{A}_{\xi^1}}{\partial \hat{\xi}^2}] \\
& + \sinh(\frac{a_0}{c} \xi^0) [-\frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^2} [\hat{\phi}_\xi (1 + \frac{a_0 \xi^1}{c^2})] - \frac{\partial \hat{A}_{\xi^2}}{c \partial \hat{\xi}^0}] \quad (21)
\end{aligned}$$

Hence, we can define the electro-magnetic field  $(\vec{E}_\xi, \vec{B}_\xi)$  in Rindler spacetime.

$$\begin{aligned}
\vec{E}_\xi &= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \vec{\nabla}_\xi \{\phi_\xi (1 + \frac{a_0 \xi^1}{c^2})^2\} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial \vec{A}_\xi}{c \partial \xi^0} \\
&= -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \vec{\nabla}_{\hat{\xi}} \{\hat{\phi}_\xi (1 + \frac{a_0 \xi^1}{c^2})\} - \frac{\partial \vec{\hat{A}}_\xi}{c \partial \hat{\xi}^0} \\
\vec{B}_\xi &= \vec{\nabla}_\xi \times \vec{A}_\xi = \vec{\nabla}_{\hat{\xi}} \times \vec{\hat{A}}_\xi \\
\text{In this time, } \vec{\nabla}_\xi &= (\frac{\partial}{\partial \xi^1}, \frac{\partial}{\partial \xi^2}, \frac{\partial}{\partial \xi^3}), \vec{A}_\xi = (A_{\xi^1}, A_{\xi^2}, A_{\xi^3}) \\
\vec{\nabla}_{\hat{\xi}} &= (\frac{\partial}{\partial \hat{\xi}^1}, \frac{\partial}{\partial \hat{\xi}^2}, \frac{\partial}{\partial \hat{\xi}^3}), \vec{\hat{A}}_\xi = (\hat{A}_{\xi^1}, \hat{A}_{\xi^2}, \hat{A}_{\xi^3}) \quad (22)
\end{aligned}$$

We obtain the transformation of the electro-magnetic field.

$$E_x = -\frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^1} \{\phi_\xi (1 + \frac{a_0 \xi^1}{c^2})^2\} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial A_{\xi^1}}{c \partial \xi^0}$$

$$\begin{aligned}
&= -\frac{1}{(1+\frac{a_0\xi^1}{c^2})}\frac{\partial}{\partial\xi^1}\{\hat{\phi}_\xi(1+\frac{a_0\xi^1}{c^2})\} - \frac{\partial\hat{A}_{\xi^1}}{c\partial\xi^0} = E_{\xi^1}, \\
E_y &= E_{\xi^2} \cosh(\frac{a_0\xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0\xi^0}{c}), \\
E_z &= E_{\xi^3} \cosh(\frac{a_0\xi^0}{c}) - B_{\xi^2} \sinh(\frac{a_0\xi^0}{c}) \\
B_x &= B_{\xi^1}, \\
B_y &= B_{\xi^2} \cosh(\frac{a_0\xi^0}{c}) - E_{\xi^3} \sinh(\frac{a_0\xi^0}{c}) \\
B_z &= B_{\xi^3} \cosh(\frac{a_0\xi^0}{c}) + E_{\xi^2} \sinh(\frac{a_0\xi^0}{c})
\end{aligned} \tag{23}$$

Hence,

$$E_x = E_{\xi^1}, B_x = B_{\xi^1},$$

$$\begin{pmatrix} E_y \\ B_y \\ E_z \\ B_z \end{pmatrix} = H \begin{pmatrix} E_{\xi^2} \\ B_{\xi^2} \\ E_{\xi^3} \\ B_{\xi^3} \end{pmatrix} \tag{24}$$

$$H = \begin{pmatrix} \cosh(\frac{a_0\xi^0}{c}) & 0 & 0 & \sinh(\frac{a_0\xi^0}{c}) \\ 0 & \cosh(\frac{a_0\xi^0}{c}) & -\sinh(\frac{a_0\xi^0}{c}) & 0 \\ 0 & -\sinh(\frac{a_0\xi^0}{c}) & \cosh(\frac{a_0\xi^0}{c}) & 0 \\ \sinh(\frac{a_0\xi^0}{c}) & 0 & 0 & \cosh(\frac{a_0\xi^0}{c}) \end{pmatrix}$$

The inverse-transformation of the electro-magnetic field is

$$E_{\xi^1} = E_x, B_{\xi^1} = B_x$$

$$\begin{pmatrix} E_{\xi^2} \\ B_{\xi^2} \\ E_{\xi^3} \\ B_{\xi^3} \end{pmatrix} = H^{-1} \begin{pmatrix} E_y \\ B_y \\ E_z \\ B_z \end{pmatrix}$$

$$H^{-1} = \begin{pmatrix} \cosh(\frac{a_0 \xi^0}{c}) & 0 & 0 & -\sinh(\frac{a_0 \xi^0}{c}) \\ 0 & \cosh(\frac{a_0 \xi^0}{c}) & \sinh(\frac{a_0 \xi^0}{c}) & 0 \\ 0 & \sinh(\frac{a_0 \xi^0}{c}) & \cosh(\frac{a_0 \xi^0}{c}) & 0 \\ -\sinh(\frac{a_0 \xi^0}{c}) & 0 & 0 & \cosh(\frac{a_0 \xi^0}{c}) \end{pmatrix} \quad (25)$$

$$E_{\xi^1} = E_x, B_{\xi^1} = B_x$$

$$\begin{aligned} E_{\xi^2} &= E_y \cosh(\frac{a_0 \xi^0}{c}) - B_z \sinh(\frac{a_0 \xi^0}{c}), \\ B_{\xi^2} &= B_y \cosh(\frac{a_0 \xi^0}{c}) + E_z \sinh(\frac{a_0 \xi^0}{c}) \\ E_{\xi^3} &= E_z \cosh(\frac{a_0 \xi^0}{c}) + B_y \sinh(\frac{a_0 \xi^0}{c}) \\ B_{\xi^3} &= B_z \cosh(\frac{a_0 \xi^0}{c}) - E_y \sinh(\frac{a_0 \xi^0}{c}) \end{aligned} \quad (26)$$

### 3. Electro-magnetic Field Equation(Maxwell Equation) in the Rindler space-time

Maxwell equation is

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho \quad (27-i)$$

$$\vec{\nabla} \times \vec{B} = \frac{\partial \vec{E}}{c \partial t} + \frac{4\pi}{c} \vec{j} \quad (27-ii)$$

$$\vec{\nabla} \cdot \vec{B} = 0 \quad (27-iii)$$

$$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{c \partial t} \quad (27-iv)$$

$$1. \vec{\nabla} \cdot \vec{E} = 4\pi\rho$$

$$E_x = E_{\xi^1},$$

$$\begin{aligned}
E_y &= E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right), \\
E_z &= E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
4\pi\rho &= \frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z} \\
&= \left[ -\frac{\sinh\left(\frac{a_0 \xi^0}{c}\right)}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh\left(\frac{a_0 \xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] E_{\xi^1} \\
&\quad + \frac{\partial}{\partial \xi^2} [E_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&\quad + \frac{\partial}{\partial \xi^3} [E_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)] \\
&= \cosh\left(\frac{a_0}{c} \xi^0\right) (\vec{\nabla}_\xi \cdot \vec{E}_\xi) + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[ \frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial E_{\xi^1}}{\partial \xi^0} \right] \\
&= \cosh\left(\frac{a_0}{c} \xi^0\right) (\vec{\nabla}_\xi \cdot \vec{E}_\xi) + \sinh\left(\frac{a_0}{c} \xi^0\right) \left[ \frac{\partial B_{\hat{\xi}^3}}{\partial \hat{\xi}^2} - \frac{\partial B_{\hat{\xi}^2}}{\partial \hat{\xi}^3} - \frac{\partial E_{\xi^1}}{\partial \hat{\xi}^0} \right] \quad (28) \\
2 \cdot \vec{\nabla} \times \vec{B} &= \frac{\partial \vec{E}}{c \partial t} + \frac{4\pi}{c} \vec{j} \\
B_x &= B_{\xi^1} \\
B_y &= B_{\xi^2} \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_z &= B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
\text{X-component} &= \frac{\partial B_z}{\partial y} - \frac{\partial B_y}{\partial z} \\
&= \frac{\partial}{\partial \xi^2} [B_{\xi^3} \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_{\xi^2} \sinh\left(\frac{a_0 \xi^0}{c}\right)]
\end{aligned}$$

$$\begin{aligned}
& -\frac{\partial}{\partial \xi^3} [B_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) - E_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})] \\
& = \cosh(\frac{a_0 \xi^0}{c}) [\frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3}] + \sinh(\frac{a_0 \xi^0}{c}) [\frac{\partial E_{\xi^2}}{\partial \xi^2} + \frac{\partial E_{\xi^3}}{\partial \xi^3}] \\
& = \frac{\partial E_x}{c \partial t} + \frac{4\pi}{c} j_x \\
& = \left[ \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{c \partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] E_{\xi^1} + \frac{4\pi}{c} j_x
\end{aligned}$$

Hence,

$$\begin{aligned}
& \frac{4\pi}{c} j_x \\
& = \sinh(\frac{a_0 \xi^0}{c}) (\vec{\nabla}_{\xi} \cdot \vec{E}_{\xi}) + \cosh(\frac{a_0 \xi^0}{c}) [\frac{\partial B_{\xi^3}}{\partial \xi^2} - \frac{\partial B_{\xi^2}}{\partial \xi^3} - \frac{1}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial E_{\xi^1}}{c \partial \xi^0}] \\
& = \sinh(\frac{a_0 \xi^0}{c}) (\vec{\nabla}_{\hat{\xi}} \cdot \vec{E}_{\xi}) + \cosh(\frac{a_0 \xi^0}{c}) [\frac{\partial B_{\xi^3}}{\partial \hat{\xi}^2} - \frac{\partial B_{\hat{\xi}^2}}{\partial \hat{\xi}^3} - \frac{\partial E_{\hat{\xi}^1}}{c \partial \hat{\xi}^0}] \quad (29)
\end{aligned}$$

$$\text{Y-component) } \frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x}$$

$$= \frac{\partial B_{\xi^1}}{\partial \xi^3}$$

$$\begin{aligned}
& - \left[ -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{c \partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) + E_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})] \\
& = \frac{\partial E_y}{c \partial t} + \frac{4\pi}{c} j_y \\
& = \left[ \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{c \partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})]
\end{aligned}$$

$$\begin{aligned}
& + \frac{4\pi}{c} j_y \\
\frac{4\pi}{c} j_y &= \frac{\partial B_{\xi^1}}{\partial \xi^3} - \frac{\partial B_{\xi^3}}{\partial \xi^1} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{a_0}{c^2} B_{\xi^3} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^2}}{\partial \partial \xi^0} \\
&= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^3} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \{B_{\xi^3} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^2}}{\partial \partial \xi^0} \\
&= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^3} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^1} \{B_{\xi^3} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{\partial E_{\xi^2}}{\partial \partial \hat{\xi}^0} \\
(30) \quad & \text{Z-component) } \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \\
&= \left[ -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) - E_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})] \\
&\quad - \frac{\partial B_{\xi^1}}{\partial \xi^2} \\
&= \frac{\partial E_z}{\partial \partial t} + \frac{4\pi}{c} j_z \\
&= \left[ \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) - B_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})] \\
&\quad + \frac{4\pi}{c} j_z \\
\frac{4\pi}{c} j_z &= \frac{\partial B_{\xi^2}}{\partial \xi^1} - \frac{\partial B_{\xi^1}}{\partial \xi^2} + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{a_0}{c^2} B_{\xi^2} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^3}}{\partial \partial \xi^0} \\
&= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \{B_{\xi^2} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^2} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial E_{\xi^3}}{\partial \partial \xi^0}
\end{aligned}$$

$$= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^1} \{B_{\xi^2} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^2} \{B_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{\partial E_{\xi^3}}{c \partial \hat{\xi}^0}$$

(31)

$$3. \vec{\nabla} \cdot \vec{B} = 0$$

$$\vec{\nabla} \cdot \vec{B} = \frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z}$$

$$= \left[ - \frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] B_{\xi^1}$$

$$+ \frac{\partial}{\partial \xi^2} [B_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) - E_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})]$$

$$+ \frac{\partial}{\partial \xi^3} [B_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) + E_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})]$$

$$= \cosh(\frac{a_0 \xi^0}{c}) (\vec{\nabla}_{\xi} \cdot \vec{B}_{\xi}) + \sinh(\frac{a_0 \xi^0}{c}) \left[ - \left( - \frac{\partial E_{\xi^2}}{\partial \xi^3} + \frac{\partial E_{\xi^3}}{\partial \xi^2} \right) - \frac{\partial B_{\xi^1}}{c \partial \hat{\xi}^0} \right] = 0$$

(32)

$$4. \vec{\nabla} \times \vec{E} = - \frac{\partial \vec{B}}{c \partial t}$$

$$E_x = E_{\xi^1},$$

$$E_y = E_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0 \xi^0}{c}),$$

$$E_z = E_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) - B_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})$$

$$\text{X-component) } \frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z}$$

$$= \frac{\partial}{\partial \xi^2} [E_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) - B_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})]$$

$$- \frac{\partial}{\partial \xi^3} [E_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})]$$

$$= \cosh\left(\frac{a_0}{c}\xi^0\right)\left[\frac{\partial E_{\xi^3}}{\partial \xi^2} - \frac{\partial E_{\xi^2}}{\partial \xi^3}\right] - \sinh\left(\frac{a_0\xi^0}{c}\right)\left[\frac{\partial B_{\xi^2}}{\partial \xi^2} + \frac{\partial B_{\xi^3}}{\partial \xi^3}\right]$$

$$= -\frac{\partial B_x}{c\partial t}$$

$$= -\left[\frac{\cosh\left(\frac{a_0\xi^0}{c}\right)}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial}{c\partial \xi^0} - \sinh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial}{\partial \xi^1}\right] B_{\xi^1}$$

Hence,

$$-\sinh\left(\frac{a_0\xi^0}{c}\right)(\vec{\nabla}_{\xi} \cdot \vec{B}_{\xi}) + c \circ s \frac{a_0\xi^0}{c} \left[ \left(\frac{\partial E_{\xi^3}}{\partial \xi^2} - \frac{\partial E_{\xi^2}}{\partial \xi^3}\right) + \frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial B_{\xi^1}}{c\partial \xi^0} \right]$$

$$= -\sinh\left(\frac{a_0\xi^0}{c}\right)(\vec{\nabla}_{\xi} \cdot \vec{B}_{\xi}) + \cosh\left(\frac{a_0\xi^0}{c}\right)\left[\left(\frac{\partial E_{\xi^3}}{\partial \xi^2} - \frac{\partial E_{\xi^2}}{\partial \xi^3}\right) + \frac{\partial B_{\xi^1}}{c\partial \xi^0}\right] = 0 \quad (33)$$

Y-component)  $\frac{\partial E_x}{\partial z} - \frac{\partial E_z}{\partial x}$

$$= \frac{\partial E_{\xi^1}}{\partial \xi^3}$$

$$- \left[ -\frac{\sinh\left(\frac{a_0\xi^0}{c}\right)}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial}{c\partial \xi^0} + \cosh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^3} \cosh\left(\frac{a_0\xi^0}{c}\right) - B_{\xi^2} \sinh\left(\frac{a_0\xi^0}{c}\right)]$$

$$= -\frac{\partial B_y}{c\partial t}$$

$$= -\left[\frac{\cosh\left(\frac{a_0\xi^0}{c}\right)}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial}{c\partial \xi^0} - \sinh\left(\frac{a_0\xi^0}{c}\right) \frac{\partial}{\partial \xi^1}\right] \cdot [B_{\xi^2} \cosh\left(\frac{a_0\xi^0}{c}\right) - E_{\xi^3} \sinh\left(\frac{a_0\xi^0}{c}\right)]$$

$$\frac{\partial E_{\xi^1}}{\partial \xi^3} - \frac{\partial E_{\xi^3}}{\partial \xi^1} - \frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{a_0}{c^2} E_{\xi^3} + \frac{1}{\left(1 + \frac{a_0\xi^1}{c^2}\right)} \frac{\partial B_{\xi^2}}{c\partial \xi^0}$$

$$= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^3} \{E_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \{E_{\xi^3} (1 + \frac{a_0 \xi^1}{c^2})\} + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial B_{\xi^2}}{\partial \xi^0}$$

$$= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^3} \{E_{\xi^1} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^1} \{E_{\xi^3} (1 + \frac{a_0 \xi^1}{c^2})\} + \frac{\partial B_{\xi^2}}{\partial \hat{\xi}^0} = 0$$

(34)

$$\text{Z-component) } \frac{\partial E_y}{\partial x} - \frac{\partial E_x}{\partial y}$$

$$= \left[ -\frac{\sinh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} + \cosh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [E_{\xi^2} \cosh(\frac{a_0 \xi^0}{c}) + B_{\xi^3} \sinh(\frac{a_0 \xi^0}{c})]$$

$$- \frac{\partial E_{\xi^1}}{\partial \xi^2}$$

$$= - \frac{\partial B_z}{\partial \hat{\xi}^0}$$

$$= - \left[ \frac{\cosh(\frac{a_0 \xi^0}{c})}{(1 + \frac{a_0 \xi^1}{c^2})} \frac{\partial}{\partial \xi^0} - \sinh(\frac{a_0 \xi^0}{c}) \frac{\partial}{\partial \xi^1} \right] \cdot [B_{\xi^3} \cosh(\frac{a_0 \xi^0}{c}) + E_{\xi^2} \sinh(\frac{a_0 \xi^0}{c})]$$

$$\frac{\partial E_{\xi^2}}{\partial \xi^1} - \frac{\partial E_{\xi^1}}{\partial \xi^2} + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{a_0}{c^2} E_{\xi^2} + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial B_{\xi^3}}{\partial \xi^0}$$

$$= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^1} \{E_{\xi^2} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \xi^2} \{E_{\xi^1} (1 + \frac{a_0 \xi^1}{c^2})\} + \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial B_{\xi^3}}{\partial \xi^0}$$

$$= \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^1} \{E_{\xi^2} (1 + \frac{a_0}{c^2} \xi^1)\} - \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)} \frac{\partial}{\partial \hat{\xi}^2} \{E_{\xi^1} (1 + \frac{a_0 \xi^1}{c^2})\} + \frac{\partial B_{\xi^3}}{\partial \hat{\xi}^0} = 0$$

(35)

Therefore, we obtain the electro-magnetic field equation by Eq (28)-Eq(35) in Rindler spacetime .

$$\vec{\nabla}_{\xi} \cdot \vec{E}_{\xi} = \vec{\nabla}_{\hat{\xi}} \cdot \vec{E}_{\xi} = 4\pi\rho_{\xi} (1 + \frac{a_0 \xi^1}{c^2}) \quad (36-i)$$

$$\begin{aligned} \frac{1}{(1+\frac{a_0\xi^1}{c^2})}\vec{\nabla}_\xi \times \{\vec{B}_\xi(1+\frac{a_0\xi^1}{c^2})\} &= \frac{1}{(1+\frac{a_0\xi^1}{c^2})}\frac{\partial \vec{E}_\xi}{c\partial \xi^0} + \frac{4\pi}{c}\vec{j}_\xi \\ \frac{1}{(1+\frac{a_0\xi^1}{c^2})}\vec{\nabla}_{\hat{\xi}} \times \{\vec{B}_\xi(1+\frac{a_0\xi^1}{c^2})\} &= \frac{\partial \vec{E}_\xi}{c\partial \hat{\xi}^0} + \frac{4\pi}{c}\vec{j}_\xi \end{aligned} \quad (36\text{-ii})$$

$$\vec{\nabla}_\xi \cdot \vec{B}_\xi = \vec{\nabla}_{\hat{\xi}} \cdot \vec{B}_\xi = 0 \quad (36\text{-iii})$$

$$\begin{aligned} \frac{1}{(1+\frac{a_0\xi^1}{c^2})}\vec{\nabla}_\xi \times \{\vec{E}_\xi(1+\frac{a_0\xi^1}{c^2})\} &= -\frac{1}{(1+\frac{a_0\xi^1}{c^2})}\frac{\partial \vec{B}_\xi}{c\partial \xi^0} \\ \frac{1}{(1+\frac{a_0\xi^1}{c^2})}\vec{\nabla}_{\hat{\xi}} \times \{\vec{E}_\xi(1+\frac{a_0\xi^1}{c^2})\} &= -\frac{\partial \vec{B}_\xi}{c\partial \hat{\xi}^0} \end{aligned} \quad (36\text{-iv})$$

$$\vec{E}_\xi = (E_{\xi^1}, E_{\xi^2}, E_{\xi^3}), \vec{B}_\xi = (B_{\xi^1}, B_{\xi^2}, B_{\xi^3}),$$

$$\vec{\nabla}_\xi = (\frac{\partial}{\partial \xi^1}, \frac{\partial}{\partial \xi^2}, \frac{\partial}{\partial \xi^3}), \vec{\nabla}_{\hat{\xi}} = (\frac{\partial}{\partial \hat{\xi}^1}, \frac{\partial}{\partial \hat{\xi}^2}, \frac{\partial}{\partial \hat{\xi}^3})$$

Hence, the transformation of 4-vector  $(c\rho, \vec{j}) = \rho_0 \frac{dx^\alpha}{d\tau}$  is

$$\begin{aligned} \rho &= \rho_\xi(1+\frac{a_0\xi^1}{c^2})\cosh(\frac{a_0\xi^0}{c}) + \frac{j_{\xi^1}}{c}\sinh(\frac{a_0\xi^0}{c}) \\ j_x &= j_{\xi^1}\cosh(\frac{a_0\xi^0}{c}) + c\rho_\xi(1+\frac{a_0}{c^2}\xi^1)\sinh(\frac{a_0\xi^0}{c}), \quad j_y = j_{\xi^2}, j_z = j_{\xi^3} \end{aligned}$$

$$\text{In this time, 4-vector } (c\rho_\xi, \vec{j}_\xi) = \rho_0 \frac{d\xi^\alpha}{d\tau} \quad (37)$$

#### 4. Electro-magnetic wave equation in Rindler space-time

The electro-magnetic wave function is

$$E_x = E_{x0} \sin \Phi, E_y = E_{y0} \sin \Phi, E_z = E_{z0} \sin \Phi$$

$$B_x = B_{x0} \sin \Phi, B_y = B_{y0} \sin \Phi, B_z = B_{z0} \sin \Phi$$

$$E_{\xi^1} = E_x, B_{\xi^1} = B_x$$

$$E_{\xi^1} = E_{x0} \sin \Phi', B_{\xi^1} = B_{x0} \sin \Phi'$$

$$\begin{aligned}
E_{\xi^2} &= E_y \cosh\left(\frac{a_0 \xi^0}{c}\right) - B_z \sinh\left(\frac{a_0 \xi^0}{c}\right), \\
&= (E_{y0} \sin \Phi') \cosh\left(\frac{a_0 \xi^0}{c}\right) - (B_{z0} \sin \Phi') \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_{\xi^2} &= B_y \cosh\left(\frac{a_0 \xi^0}{c}\right) + E_z \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
&= (B_{y0} \sin \Phi') \cosh\left(\frac{a_0 \xi^0}{c}\right) + (E_{z0} \sin \Phi') \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
E_{\xi^3} &= E_z \cosh\left(\frac{a_0 \xi^0}{c}\right) + B_y \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
&= (E_{z0} \sin \Phi') \cosh\left(\frac{a_0 \xi^0}{c}\right) + (B_{y0} \sin \Phi') \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
B_{\xi^3} &= B_z \cosh\left(\frac{a_0 \xi^0}{c}\right) - E_y \sinh\left(\frac{a_0 \xi^0}{c}\right) \\
&= (B_{z0} \sin \Phi') \cosh\left(\frac{a_0 \xi^0}{c}\right) - (E_{y0} \sin \Phi') \sinh\left(\frac{a_0 \xi^0}{c}\right)
\end{aligned} \tag{38}$$

$$\begin{aligned}
\Phi &= \omega(t - l \frac{x}{c} - m \frac{y}{c} - n \frac{z}{c}), \\
\Phi' &= \omega' (\hat{\xi}^0 - l' \frac{\hat{\xi}^1}{c} - m' \frac{\hat{\xi}^2}{c} - n' \frac{\hat{\xi}^3}{c}) \\
l'^2 + m'^2 + n'^2 &= 1 \quad , l'^2 + m'^2 + n'^2 = 1
\end{aligned} \tag{39}$$

Hence,

$$\begin{aligned}
&[\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_{\xi}^2] E_{\xi^1} \\
&= [\frac{1}{c^2} (\frac{\partial}{\partial \hat{\xi}^0})^2 - \nabla_{\hat{\xi}}^2] E_{\xi^1} = 0 \\
&[\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_{\xi}^2] B_{\xi^1} \\
&= [\frac{1}{c^2} (\frac{\partial}{\partial \hat{\xi}^0})^2 - \nabla_{\hat{\xi}}^2] B_{\xi^1} = 0
\end{aligned}$$

$$\begin{aligned}
& \left[ \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left( \frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] E_y \\
&= \left[ \frac{1}{c^2} \left( \frac{\partial}{\partial \hat{\xi}^0} \right)^2 - \nabla_{\hat{\xi}}^2 \right] E_y = 0 \\
& \left[ \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left( \frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] B_y \\
&= \left[ \frac{1}{c^2} \left( \frac{\partial}{\partial \hat{\xi}^0} \right)^2 - \nabla_{\hat{\xi}}^2 \right] B_y = 0 \\
& \left[ \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left( \frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] E_z \\
&= \left[ \frac{1}{c^2} \left( \frac{\partial}{\partial \hat{\xi}^0} \right)^2 - \nabla_{\hat{\xi}}^2 \right] E_z = 0 \\
& \left[ \frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} \left( \frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] B_z \\
&= \left[ \frac{1}{c^2} \left( \frac{\partial}{\partial \hat{\xi}^0} \right)^2 - \nabla_{\hat{\xi}}^2 \right] B_z = 0
\end{aligned} \tag{40}$$

The electro-magnetic wave equation is in vacuum

$$\begin{aligned}
& \vec{\nabla}_{\xi} \times \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \vec{\nabla}_{\xi} \times \{ \vec{E}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \} \\
&= \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{\nabla}_{\xi} \times \{ \vec{E}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \} + \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \vec{\nabla}_{\xi} \times \vec{\nabla}_{\xi} \times \{ \vec{E}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \} \\
&= \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{E}_{\xi} \\
&+ \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{\nabla}_{\xi} \times \vec{E}_{\xi} \\
&+ \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \vec{\nabla}_{\xi} \times \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{E}_{\xi} \\
&+ \left( 1 + \frac{a_0}{c^2} \xi^1 \right)^2 \vec{\nabla}_{\xi} \times \vec{\nabla}_{\xi} \times \vec{E}_{\xi} \\
&= \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{\nabla}_{\xi} \left( 1 + \frac{a_0}{c^2} \xi^1 \right) \times \vec{E}_{\xi} + \left( 1 + \frac{a_0}{c^2} \xi^1 \right)^2 \vec{\nabla}_{\xi} \times \vec{\nabla}_{\xi} \times \vec{E}_{\xi}
\end{aligned}$$

$$\begin{aligned}
&= [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{E}_\xi] \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) - [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1)] \vec{E}_\xi \\
&\quad + (1 + \frac{a_0}{c^2} \xi^1)^2 [\vec{\nabla}_\xi (\vec{\nabla}_\xi \cdot \vec{E}_\xi) - \nabla_\xi^2 \vec{E}_\xi] \\
&= -\frac{1}{c} \frac{\partial}{\partial \xi^0} [\vec{\nabla}_\xi \times \{\vec{B}_\xi (1 + \frac{a_0 \xi^1}{c^2})\}] = -\frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \vec{E}_\xi,
\end{aligned}$$

In this time,  $\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) = (\frac{a_0}{c^2}, 0, 0)$  (41)

Hence,

$$\begin{aligned}
&\vec{\nabla}_\xi \times (1 + \frac{a_0}{c^2} \xi^1) \vec{\nabla}_\xi \times \{\vec{E}_\xi (1 + \frac{a_0}{c^2} \xi^1)\} + \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \vec{E}_\xi \\
&= [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{E}_\xi] \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) - [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1)] \vec{E}_\xi \\
&\quad + (1 + \frac{a_0}{c^2} \xi^1)^2 [\vec{\nabla}_\xi (\vec{\nabla}_\xi \cdot \vec{E}_\xi) - \nabla_\xi^2 \vec{E}_\xi] + \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \vec{E}_\xi \\
&= \frac{a_0^2}{c^4} (E_{\xi^1}, 0, 0) - \frac{a_0^2}{c^4} (E_{\xi^1}, E_{\xi^2}, E_{\xi^3}) + (1 + \frac{a_0}{c^2} \xi^1)^2 [\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_\xi^2] \vec{E}_\xi \\
&= \frac{a_0^2}{c^4} (0, -E_{\xi^2}, -E_{\xi^3}) + (1 + \frac{a_0}{c^2} \xi^1)^2 [\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_\xi^2] \vec{E}_\xi \\
&= \frac{a_0^2}{c^4} (0, -E_{\xi^2}, -E_{\xi^3}) + (1 + \frac{a_0}{c^2} \xi^1)^2 [\frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_\xi^2] \vec{E}_\xi \\
&= \vec{0} \tag{42}
\end{aligned}$$

Hence, the magnetic wave equation is in vacuum

$$\begin{aligned}
&\vec{\nabla}_\xi \times (1 + \frac{a_0}{c^2} \xi^1) \vec{\nabla}_\xi \times \{\vec{B}_\xi (1 + \frac{a_0}{c^2} \xi^1)\} + \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \vec{B}_\xi \\
&= [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{B}_\xi] \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) - [\vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1) \cdot \vec{\nabla}_\xi (1 + \frac{a_0}{c^2} \xi^1)] \vec{B}_\xi \\
&\quad + (1 + \frac{a_0}{c^2} \xi^1)^2 [\vec{\nabla}_\xi (\vec{\nabla}_\xi \cdot \vec{B}_\xi) - \nabla_\xi^2 \vec{B}_\xi] + \frac{1}{c^2} (\frac{\partial}{\partial \xi^0})^2 \vec{B}_\xi \\
&= \frac{a_0^2}{c^4} (0, -B_{\xi^2}, -B_{\xi^3}) + (1 + \frac{a_0}{c^2} \xi^1)^2 [\frac{1}{c^2} \frac{1}{(1 + \frac{a_0}{c^2} \xi^1)^2} (\frac{\partial}{\partial \xi^0})^2 - \nabla_\xi^2] \vec{B}_\xi
\end{aligned}$$

$$\begin{aligned}
&= \frac{a_0^2}{c^4} (0, -B_{\xi^2}, -B_{\xi^3}) + (1 + \frac{a_0}{c^2} \xi^1)^2 \left[ \frac{1}{c^2} \left( \frac{\partial}{\partial \xi^0} \right)^2 - \nabla_{\xi}^2 \right] \vec{B}_{\xi} \\
&= \vec{0}
\end{aligned} \tag{43}$$

The electromagnetic wave function, Eq(38),Eq(39) satisfy the electromagnetic wave equation, Eq(42),Eq(43).

## 5. Conclusion

We find the electro-magnetic field transformation and the electro-magnetic equation in uniformly accelerated frame.

Generally, the coordinate transformation of accelerated frame is

$$\begin{aligned}
(I) \quad ct &= \left( \frac{c^2}{a_0} + \xi^1 \right) \sinh \left( \frac{a_0 \xi^0}{c} \right) \\
x &= \left( \frac{c^2}{a_0} + \xi^1 \right) \cosh \left( \frac{a_0 \xi^0}{c} \right) - \frac{c^2}{a_0} , y = \xi^2 , z = \xi^3
\end{aligned} \tag{44}$$

$$\begin{aligned}
(II) \quad ct &= \frac{c^2}{a_0} \exp \left( \frac{a_0}{c^2} \xi^1 \right) \sinh \left( \frac{a_0 \xi^0}{c} \right) \\
x &= \frac{c^2}{a_0} \exp \left( \frac{a_0}{c^2} \xi^1 \right) \cosh \left( \frac{a_0 \xi^0}{c} \right) - \frac{c^2}{a_0} , y = \xi^2 , z = \xi^3
\end{aligned} \tag{45}$$

Hence, this article say the accelerated frame is Rindler coordinate (I) that can treat electro-magnetic field equation.

## Reference

- [1] S.Weinberg,Gravitation and Cosmology(John wiley & Sons,Inc,1972)
- [2] W.Rindler, Am.J.Phys.**34**.1174(1966)
- [3] P.Bergman,Introduction to the Theory of Relativity(Dover Pub. Co.,Inc., New York,1976),Chapter V
- [4] C.Misner, K.Thorne and J. Wheeler, Gravitation(W.H.Freedman & Co.,1973)
- [5] S.Hawking and G. Ellis,The Large Scale Structure of Space-Time(Cambridge University Press,1973)
- [6] R.Adler,M.Bazin and M.Schiffer,Introduction to General Relativity(McGraw-Hill,Inc.,1965)
- [7] A.Miller, Albert Einstein's Special Theory of Relativity(Addison-Wesley Publishing Co., Inc., 1981)
- [8] W.Rindler, Special Relativity(2nd ed., Oliver and Boyd, Edinburg,1966)
- [9] [Massimo Pauri](#), [Michele Vallisneri](#), "Marzke-Wheeler coordinates for accelerated observers in special relativity":Arxiv:gr-qc/0006095(2000)

[10]A. Einstein, " Zur Elektrodynamik bewegter Körper", Annalen der Physik. 17:891(1905)