

ORTHOGONAL PARALLEL MCMC METHODS FOR SAMPLING AND OPTIMIZATION

L. Martino^{*}, V. Elvira[†], D. Luengo[‡], J. Corander[◇], F. Louzada^{*}

^{*} Institute of Mathematical Sciences and Computing, Universidade de São Paulo, São Carlos (Brazil).

[◇] Dep. of Mathematics and Statistics, University of Helsinki, Helsinki (Finland).

[†] Dep. of Signal Theory and Communic., Universidad Carlos III de Madrid, Leganés (Spain).

[‡] Dep. of Circuits and Systems Engineering, Universidad Politécnica de Madrid, Madrid (Spain).

ABSTRACT

Monte Carlo (MC) methods are widely used in statistics, signal processing and machine learning. A well-known class of MC methods are Markov Chain Monte Carlo (MCMC) algorithms. In order to foster better exploration of the state space, specially in high-dimensional applications, several schemes employing multiple parallel MCMC chains have been recently introduced. In this work, we describe a novel parallel interacting MCMC scheme, called *orthogonal MCMC* (O-MCMC), where a set of “vertical” parallel MCMC chains share information using some “horizontal” MCMC techniques working on the entire population of current states. More specifically, the vertical chains are led by random-walk proposals, whereas the horizontal MCMC techniques employ independent proposals, thus allowing an efficient combination of global exploration and local approximation. The interaction is contained in these horizontal iterations. Within the analysis of different implementations of O-MCMC, novel schemes for reducing the overall computational cost of parallel multiple try Metropolis (MTM) chains are also presented. Furthermore, a modified version of O-MCMC for optimization is provided by considering parallel simulated annealing (SA) algorithms. We also discuss the application of O-MCMC in a big data framework. Numerical results show the advantages of the proposed sampling scheme in terms of efficiency in the estimation, as well as robustness in terms of independence with respect to initial values and parameter choice.

Index Terms— Parallel Markov Chain Monte Carlo (MCMC), Parallel Multiple Try Metropolis, Adaptive Multiple Try Metropolis, Block Independent Metropolis, Parallel Simulated Annealing.

1. INTRODUCTION

Monte Carlo (MC) methods are widely employed in different fields, for computational inference and stochastic optimization [1, 2, 4, 5]. Markov Chain Monte Carlo (MCMC) methods [5] are well-known MC methodologies to draw random samples and compute efficiently integrals involving a complicated multidimensional target probability density function (pdf), $\pi(\mathbf{x})$ with $\mathbf{x} \in \mathcal{D} \subseteq \mathbb{R}^{d_x}$. MCMC techniques only need to be able to evaluate the target pdf, but the difficulty of diagnosing and speeding up the convergence has driven intensive research effort in this field. For instance, several adaptive MCMC methods have been developed in order to determine adequately the shape and spread of the proposal density which is used to generate candidate samples, within the MCMC scheme [3, 6, 5, 7]. Nevertheless, guaranteeing the theoretical convergence is still an issue in most of the cases. Moreover, in a single specific (long) run, the generated chain can remain trapped in a local mode and, in this scenario, the adaptation could even slow down the convergence. Thus, in order to speed up the exploration of the state space (and specially to deal with high-dimensional applications [8]), several schemes employing parallel chains have been recently proposed [2, 7], as well as multiple try and interacting schemes [9], but the problem is still far from being solved. The interest in the parallel computation can be also originated by other motivations. For instance, several authors have studied the parallelization of MCMC algorithms which have traditionally been implemented in an iterative non-parallel fashion, in order to reduce their computation time [10, 11]. Furthermore, parallel MCMC schemes are required in a big data problems, where, for instance, the complete posterior distribution is split in different partial sub-posteriors [12, 13, 14, 15, 16].

In this work, we focus mainly on the implementation of parallel MCMC chains in order to foster the exploration of the state space and improve the overall performance.¹ More specifically, we present a novel family of parallel MCMC schemes,

¹A preliminary version of this work has been published in [17]. With respect to that paper, here we propose different novel interacting schemes for exchanging information among the chains, analyze the theoretical basis of proposed approach and discuss certain relationships with other techniques, in detail. Several variants are presented for reducing the overall computational cost and for applying O-MCMC in optimization problems. Furthermore, we provide more exhaustive numerical simulations.

called orthogonal MCMC (O-MCMC) algorithms, where N different chains are independently run and, at some iterations, they exchange information using another MCMC technique applied on the entire cloud of current states. Assuming that all the MCMC techniques used yield chains converging to the target pdf, the ergodicity property is guaranteed: the whole kernel is still valid, since it is obtained as the multiplication of ergodic kernels with the same invariant pdf. Fixing a given computational cost, this computing effort can be divided in N parallel processes but, at some iteration, information among the chains is exchanged in order to enhance the overall mixing. The novel O-MCMC scheme is able to combine efficiently both the random-walk and the independent proposal approaches, as both strategies have advantages and drawbacks. On the one hand, random-walk proposal pdfs are often used when there is no specific information about the target, since this approach turns out to be more explorative than using a fixed proposal. On the other hand, a well-chosen independent proposal density usually provides less correlation among the samples in the generated chain. The novel method can mix both approaches efficiently: the parallel “vertical” chains (based on random-walk proposals) move around as “free explorers” roaming the state space, whereas the “horizontal” MCMC technique (applied over the population of current states and based on independent proposals) works as a “park ranger”, redirecting “lost explorers” towards the “beaten track” according to the target pdf. Unlike in [18], the exchange of information occurs taking always into account the whole population of current states, instead of applying *crossover* or *exchange* schemes between specific pairs of chains. Furthermore, in this work, tempering of the target pdf is not considered for sampling purposes (but, it is employed for optimization). Hence, in this sense our approach resembles the nonreversible parallel MH algorithms described in [19, 20] where the whole population of states is also updated jointly at the times of interaction, pursuing non-reversibility instead of tempering as a means to accelerate convergence towards posterior mode regions. However, both tempering and crossovers could also be easily implemented within an O-MCMC framework.

Another contribution of the work is the computational improvement provided by novel parallel implementations of MCMC techniques using multiple candidates at each iteration. We present two novel schemes for parallel Multiple try Metropolis (MTM) [21, 22, 23] chains in order to reduce the overall computational cost in the same fashion of [10], saving generated samples, target evaluations and multinomial sampling steps. One of them is an extended version, using several candidates, of the Block Independent Metropolis presented in [10]. The ergodicity of both schemes is guaranteed. These novel parallel MTM techniques are employed as horizontal methods in O-MCMC. The corresponding O-MCMC scheme (using a novel parallel MTM method) can be also interpreted as an MTM algorithm employing an *adaptive* proposal density. This pdf is a mixture of N components: the adaptation of the location parameters of the N components is driven by the vertical parallel chains (note that the outputs of these chains are also used in the estimation). Furthermore, we describe a modified version of O-MCMC for solving optimization problems, considering parallel Simulated Annealing algorithms [24, 25, 26] for the vertical movements. The application for big data problems is also discussed. Numerical simulations show that O-MCMC exhibits both flexibility and robustness with respect to the initialization and parameterization of the proposals.

The paper is structured as follows. Section 2 summarizes the general framework and the aim of the work. Section 3 describes a generic complete O-MCMC scheme, whereas Sections 4 and 5 provide different specific examples of vertical and horizontal movements, respectively. Section 7 provides different numerical results. Finally, we finish with some remarks in Section 8.

2. BAYESIAN INFERENCE PROBLEM

In many applications, we aim at inferring a variable of interest given a set of observations or measurements. Let us denote the variable of interest by $\mathbf{x} \in \mathcal{D} \subseteq \mathbb{R}^{d_x}$, and let $\mathbf{y} \in \mathbb{R}^{d_y}$ be the observed data. The posterior pdf is then

$$\bar{\pi}(\mathbf{x}) \propto p(\mathbf{x}|\mathbf{y}) = \frac{\ell(\mathbf{y}|\mathbf{x})g(\mathbf{x})}{Z(\mathbf{y})}, \quad (1)$$

where $\ell(\mathbf{y}|\mathbf{x})$ is the likelihood function, $g(\mathbf{x})$ is the prior pdf and $Z(\mathbf{y})$ is the model evidence (a.k.a. marginal likelihood). In general, $Z(\mathbf{y})$ is unknown, so we consider the corresponding unnormalized target function,

$$\pi(\mathbf{x}) = \ell(\mathbf{y}|\mathbf{x})g(\mathbf{x}) \propto \bar{\pi}(\mathbf{x}). \quad (2)$$

In general, the study of the posterior density $\bar{\pi}(\mathbf{x})$ is impossible analytically (for instance, integrals involving $\bar{\pi}(\mathbf{x})$ are typically intractable), so that numerical approximations are required. Our goal is to approximate efficiently $\bar{\pi}(\mathbf{x})$ employing a cloud of random samples. In general, a direct method for drawing independent samples from $\bar{\pi}(\mathbf{x})$ is not available and alternative approaches (e.g., MCMC algorithms) are needed. The only required assumption is to be able of evaluating the unnormalized target function $\pi(\mathbf{x})$.

3. O-MCMC ALGORITHMS: GENERAL OUTLINE

Let us consider N parallel “vertical” chains, $\{\mathbf{x}_{n,t}\}_{n=1}^N$ with $t \in \mathbb{N}$, generated by different MCMC techniques with *random-walk* proposal pdfs $q_n(\mathbf{x}|\mathbf{x}_{n,t-1}) = q_n(\mathbf{x} - \mathbf{x}_{n,t-1})$, i.e., $\mathbf{x}_{n,t-1}$ plays the role of a location parameter for the density $q_n(\mathbf{x})$ in the next iteration. Let us denote the population of current states at the t -th iteration as

$$\mathcal{P}_t = \{\mathbf{x}_{1,t}, \mathbf{x}_{2,t}, \dots, \mathbf{x}_{N,t}\}.$$

At certain selected iteration t , such that $t = mT_V$ where T_V is a constant and $m = 1, \dots, M$, we apply another MCMC technique taking into account the entire population of state \mathcal{P}_{t-1} , yielding a new cloud of samples \mathcal{P}_t . In this “horizontal” transitions, the different chains share information. The horizontal MCMC technique uses an proposal pdf $\varphi(\mathbf{x})$, which is independent from the previous states differently from the random walk proposals employed in the vertical MCMC chains. The general O-MCMC approach is represented graphically in Figure 1 and summarized below:

1. **Initialization:** Set $t = 1$. Choose the N initial states,

$$\mathcal{P}_0 = \{\mathbf{x}_{1,0}, \mathbf{x}_{2,0}, \dots, \mathbf{x}_{N,0}\},$$

the total number of iterations, T , and three positive integer values $M, T_V, T_H \in \mathbb{N} \setminus \{0\}$ such that $M(T_V + T_H) = T$.

2. **For $m=1, \dots, M$:**

- (a) **Vertical period:** For

$$t = (m-1)(T_V + T_H) + 1, \dots, mT_V + (m-1)T_H,$$

run N independent MCMC techniques starting from $\mathbf{x}_{n,t-1} \in \mathcal{P}_{t-1}$, thus obtaining $\mathbf{x}_{n,t}$, for $n = 1, \dots, N$, and then a new population of states $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \mathbf{x}_{2,t}, \dots, \mathbf{x}_{N,t}\}$.

- (b) **Horizontal period:** For

$$t = mT_V + (m-1)T_H + 1, \dots, m(T_V + T_H),$$

apply an MCMC approach taking in account the entire population \mathcal{P}_{t-1} at each step t , for generating the next cloud \mathcal{P}_t .

3. **Output:** Return the $T = M(T_V + T_H)$ samples contained in all the sets $\mathcal{P}_t, t = 1, \dots, T$.

One vertical period contains T_V iterations of the chains whereas, in one horizontal period we have T_H iterations. Hence, given $t = (m-1)(T_V + T_H)$, after one cycle of vertical and horizontal steps we have $t = m(T_V + T_H)$. The total number of cycles (or epochs)² is $M = \frac{T}{T_V + T_H}$. The ergodicity is guaranteed if the vertical and horizontal steps produce ergodic chains with invariant density $\pi(\mathbf{x})$. See, Appendix A for further details. Relationships with other techniques, for instance [27, 28], are discussed in Section 5.4. In the following two sections, we discuss different specific possible implementations of O-MCMC.

4. VERTICAL MOVEMENTS

In this section, we describe different suitable implementations of the vertical parallel chains. Although it is not strictly necessary, we suggest using random walk proposal densities in the vertical chains. The idea is to exploit predominantly the explorative behaviors of the independent parallel MCMC methods. Therefore, we consider proposal of type $q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$ where $\mathbf{x}_{n,t-1}$ plays the role of a location parameter. For instance, a sample $\mathbf{x}' \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$ can be expressed as

$$\mathbf{x}' = \mathbf{x}_{n,t-1} + \boldsymbol{\xi}_{n,t}, \quad (3)$$

where $\boldsymbol{\xi}_{n,t}$ has pdf $q(\boldsymbol{\xi})$ with zero mean and covariance matrix \mathbf{C}_n . Another more sophisticated possibility is to include the gradient information of the target within the proposal pdf as suggest in the *Metropolis-Adjusted Langevin Algorithm* (MALA) [29]. Namely, in this case a sample $\mathbf{x}' \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$,

$$\mathbf{x}' = \mathbf{x}_{n,t-1} + \frac{\epsilon}{2} \nabla \log [\pi(\mathbf{x}_{n,t-1})] + \boldsymbol{\xi}_{n,t}, \quad (4)$$

²One cycle, or epoch, includes one the vertical period and one horizontal period.




Fig. 1. A graphical representation of the O-MCMC approach. After T_V vertical transitions, then T_H horizontal steps are performed.

where $\xi_{n,t}$ has pdf $q(\xi)$ with zero mean, covariance matrix $\mathbf{C}_n = \epsilon \mathbb{I}_{d_x \times d_x}$ and $\mathbb{I}_{d_x \times d_x}$ is the unit matrix and $\nabla f(\mathbf{x})$ denotes the gradient of a generic function $f(\mathbf{x})$. This second alternative can be particularly useful in high-dimensional spaces, although it inevitably increases the probability of the chain becoming trapped in one mode of the target, in a multi-modal scenario. Thus, the joint application of N parallel chains appears very appropriate in this scenario. Moreover, the application of the O-MCMC scheme facilitates the jumps among different modes. Clearly, the random walk proposal density $q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$ can be applied within different MCMC kernels. The simplest possibility consists in using a *Metropolis-Hastings* (MH) algorithm [5]. For each $n = 1, \dots, N$ and for a given time step t , one MH update of the n -th chain is obtained as

1. Draw $\mathbf{x}' \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$.
2. Set $\mathbf{x}_{n,t} = \mathbf{x}'$ with probability

$$\alpha_n = \min \left[1, \frac{\pi(\mathbf{x}') q_n(\mathbf{x}_{n,t-1}|\mathbf{x}')}{\pi(\mathbf{x}_{n,t-1}) q_n(\mathbf{x}'|\mathbf{x}_{n,t-1})} \right],$$

otherwise, with probability $1 - \alpha_n$, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$.

Other possible schemes can be used as alternative to MH: for instance, two famous alternatives are the *Multiple Try Metropolis* (MTM) [21, 23] and the *Delayed Rejection Metropolis* [30] techniques.

5. HORIZONTAL MOVEMENTS

As described above, at each iteration t , the vertical chains return a population $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$. When $t = mT_V + (m-1)T_H$, with $m \in \{1, \dots, M\}$, i.e., after T_V vertical transitions, then T_H horizontal steps are performed. The purpose of these horizontal MCMC transitions is to share information among the N different chains, improving the global mixing. In the following, we consider two different general approaches for sharing the information among the chains:

- In the first one, a population-based MCMC algorithm is applied. The states of vertical chains contained in \mathcal{P}_t are used as initial population. Furthermore, the population-based MCMC scheme takes in account all the current population for making decisions about the next population. The simple case describes previously, i.e., the direct application of an MH scheme in the extended space, is an specific example of this approach.
- In the second one, the initial population \mathcal{P}_t is also used for building a suitable proposal density $\psi(\mathbf{x})$. This pdf ψ is employed by the N parallel MCMC chains for yielding the next populations $\mathcal{P}_{t+1}, \dots, \mathcal{P}_{t+T_H}$. More specifically, in this work, we suggest to construct $\psi(\mathbf{x})$ as a mixture of N pdfs, each one centered in $\mathbf{x}_{n,t} \in \mathcal{P}_t$.

Next we show several specific examples. In all the different cases, for the horizontal movements, we consider the use of independent proposal pdfs, unlike for the vertical ones, where we have suggested the use of random walk proposals.

Table 1. Sample Metropolis-Hastings (SMH) algorithm for horizontal transitions in O-MCMC.

1. For $t = mT_V + (m - 1)T_H + 1, \dots, m(T_V + T_H)$:	
(a) Draw $\mathbf{x}_{0,t-1} \sim \varphi(\mathbf{x})$.	
(b) Choose a “bad” sample $\mathbf{x}_{k,t-1}$ in \mathcal{P}_{t-1} , i.e., $k \in \{1, \dots, N\}$, according to the <i>inverse</i> of the importance sampling weights, i.e., with probability	
	$\gamma_k = \frac{\frac{\varphi(\mathbf{x}_{k,t-1})}{\pi(\mathbf{x}_{k,t-1})}}{\sum_n \frac{\varphi(\mathbf{x}_{n,t-1})}{\pi(\mathbf{x}_{n,t-1})}}, \quad k = 1, \dots, N.$
(c) Accept the new population,	
with probability	$\mathcal{P}_t = \{\mathbf{x}_{1,t} = \mathbf{x}_{1,t-1}, \dots, \mathbf{x}_{k,t} = \mathbf{x}_{0,t-1}, \dots, \mathbf{x}_{N,t} = \mathbf{x}_{N,t-1}\},$
	$\alpha = \frac{\sum_{n=1}^N \frac{\varphi(\mathbf{x}_{n,t-1})}{\pi(\mathbf{x}_{n,t-1})}}{\sum_{i=0}^N \frac{\varphi(\mathbf{x}_{i,t-1})}{\pi(\mathbf{x}_{i,t-1})} - \min_{0 \leq i \leq N} \frac{\varphi(\mathbf{x}_{i,t-1})}{\pi(\mathbf{x}_{i,t-1})}}. \quad (6)$
Otherwise, set $\mathcal{P}_t = \mathcal{P}_{t-1}$.	

5.1. Population-based approach

Here, we consider a generalized target density,

$$\bar{\pi}_g(\mathbf{x}_1, \dots, \mathbf{x}_N) \propto \prod_{n=1}^N \pi(\mathbf{x}_n), \quad (5)$$

where each marginal, $\pi(\mathbf{x}_n)$ with $n = 1, \dots, N$ and $\mathbf{x}_n \in \mathcal{D} \subseteq \mathbb{R}^{d_x}$, coincides with the target pdf in Eq. (2). The idea is that the horizontal MCMC transitions leave invariant the extended target $\bar{\pi}_g$. Namely, after a “burn-in” period, the samples in the population $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$ are distributed according to $\bar{\pi}_g$. The simplest possible population-based scheme consists in employing an standard Metropolis-Hastings (MH) algorithm directly in the extended domain, $\mathcal{D}^N \subseteq \mathbb{R}^{d_x \times N}$, with target $\bar{\pi}_g(\mathbf{x}_1, \dots, \mathbf{x}_N)$, generating (block) transitions from $\mathcal{P}_{t-1} = \{\mathbf{x}_{1,t-1}, \dots, \mathbf{x}_{N,t-1}\}$ to $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$. However, the probability of accepting a new population becomes negligible as N grows. An alternative example of possible population-based scheme, we describe the *Sample Metropolis-Hastings* (SMH) method [31, Chapter 4]. At each iteration, the underlying idea of SMH is replacing one “bad” sample in the population with a “better” one, according to a certain suitable probability. The new sample, candidate of be incorporated in the population, is generated from an independent proposal pdf $\varphi(\mathbf{x})$. The algorithm is designed so that, after a “burn-in” period t_b , the elements in $\mathcal{P}_{t'}$ ($t' > t_b$) are distributed according to $\bar{\pi}_g$ in Eq. (5). Table 1 shows the details of the algorithm.

Let us remark that the difference between \mathcal{P}_{t-1} and \mathcal{P}_t is at most one sample, and the acceptance probability, $0 \leq \alpha \leq 1$, depends on the entire population, $\mathbf{x}_{n,t-1}$ for $n = 1, \dots, N$ and the proposed one, $\mathbf{x}_{0,t-1}$. At each step, the sample chosen to be replaced is selected according to a probability the inverse of the corresponding importance weight. The ergodicity can be proved considering the extended density $\bar{\pi}_g$ as target pdf. For further details see considerations in [31, Chapter 4]. Note also that the SMH algorithm becomes the standard MH method for $N = 1$. Hence, for $N = 1$ the specific O-MCMC implementation using SMH consists of applying alternatively two MH kernels with different types of proposals: a random walk proposal, $q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$, and an independent one, $\varphi(\mathbf{x})$. This is a well-known scheme (cf. [5, 31]), which can be seen as a particular case of the O-MCMC family of algorithms.

5.2. Mixture-based approach

An alternative approach consists in defining the following mixture of pdfs, updated each T_V vertical transitions,

$$\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t) = \frac{1}{N} \sum_{n=1}^N \varphi_n(\mathbf{x}|\mathbf{x}_{n,t}), \quad \text{where } t = mT_V + (m - 1)T_H, \quad (7)$$

where $m = 1, \dots, M$ and each $\mathbf{x}_{n,t} \in \mathcal{P}_t$ plays the role of location parameter of the n -th component of the mixture, φ_n . Observe that $\psi(\mathbf{x})$ changes at the beginning of the horizontal iterations, considering the last vertical states, but remains fixed




Fig. 2. A graphical representation of the mixture-based strategy. The mixture $\psi(\mathbf{x})$ is formed by N components, $\varphi_n(\mathbf{x}|\mathbf{x}_{n,t})$, where $\mathbf{x}_{n,t} \in \mathcal{P}_t$ plays the role of location parameter.




Fig. 3. A schematic representation of basic horizontal scheme described in Table 2. One specific transition of one specific chain is represented with the probability $\alpha_n = \omega(\mathbf{x}') \wedge \omega(\mathbf{x}_{n,t-1})$, showing the two possible future states at the t -th iteration, of the n -th chain.

during horizontal iterations. Thus, during the complete O-MCMC run, we employ M different mixtures ψ 's, one for each horizontal period, so that a more adequate notation would be $\psi_m(\mathbf{x})$ instead of $\psi(\mathbf{x})$. However, for simplifying the notation, we keep $\psi(\mathbf{x})$.

Figure 2 provides for a graphical representation. We employ $\psi(\mathbf{x})$ within N independent MCMC schemes as an independent proposal density, namely independent from the previous state of the chain. The underlying idea is using the information in \mathcal{P}_t , with $t = mT_V + (m-1)T_H$, to build a good proposal function for performing N independent MCMC processes. The theoretical motivation is that, after the burn-in periods, the vertical chains have converged to the target so that we can write $\mathbf{x}_{n,t} \sim \bar{\pi}(\mathbf{x})$ for $n = 1, \dots, N$. Then, $\psi(\mathbf{x})$ in Eq. (7) can be interpreted as a kernel density estimation of $\bar{\pi}$ where φ_n play the role of kernel functions.

As first example of this strategy, we consider the application of MH transitions. The procedure is shown in Table 2. At each iteration t , one sample \mathbf{x}' is generated from $\psi(\mathbf{x})$ and then N different MH tests are performed. Then, T_H transitions of N parallel chains are performed.

Alternatively, a different sample \mathbf{x}'_n , drawn from $\psi(\mathbf{x})$, can be tested for each chain, as shown in Table 3. Hence, N different samples are drawn at each iterations (instead of only one) but, after building $\psi(\mathbf{x}|\mathcal{P}_t)$, the process could be completely parallelized. The variant in Table 3 provides in general better performance, although at the expense of a increasing computational cost, in terms of evaluations of the target and number of generated samples. However, the methodology called *block independent MH* [10], proposed exactly for reducing the computational effort recycling generated samples and target evaluations, can be also employed. For clarifying that, consider for simplicity $T_H = N$. Step 2(a) in Table 2 could be modified drawing N independent samples $\mathbf{x}'_1, \dots, \mathbf{x}'_N$ from $\psi(\mathbf{x})$ and, at each iteration t , a different circular permutation of the set $\{\mathbf{x}'_1, \dots, \mathbf{x}'_N\}$ could be tested in the different N acceptance MH tests³. Finally, observe that the procedure in Table 2 presents certain similarities with the Normal Kernel Coupler (NKC) method introduced in [32]. Clearly, NKC-type algorithms can be also employed as alternative populated-based approaches.

More advanced techniques can be also modified and used as horizontal methods. Specifically, the adaptation to this scenario

³For further clarifications, see the extension of this scheme for a Multiple Try Metropolis method described in Section 5.2.1.

Table 2. Basic mixture scheme for horizontal transitions in O-MCMC.

1. Build $\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t)$ as in Eq. (7) where $t = mT_V + (m - 1)T_H$.
2. For $t = mT_V + (m - 1)T_H + 1, \dots, m(T_V + T_H)$:
 - (a) Draw $\mathbf{x}' \sim \psi(\mathbf{x})$.
 - (b) For $n = 1, \dots, N$:
 - i. Set $\mathbf{x}_{n,t} = \mathbf{x}'$, with probability

$$\alpha_n = \min \left[1, \frac{\pi(\mathbf{x}')\psi(\mathbf{x}_{n,t-1})}{\pi(\mathbf{x}_{n,t-1})\psi(\mathbf{x}')} \right] = \omega(\mathbf{x}') \wedge \omega(\mathbf{x}_{n,t-1}).$$

Otherwise, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$. We have denoted $\omega(\mathbf{x}) = \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}$ and $a \wedge b = \min[a, b]$, with $a, b \in \mathbb{R}$.

- (c) Set $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$.

Table 3. Variant of the basic mixture scheme for horizontal transitions in O-MCMC.

1. Build $\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t)$ as in Eq. (7) where $t = mT_V + (m - 1)T_H$.
2. For $t = mT_V + (m - 1)T_H + 1, \dots, m(T_V + T_H)$:
 - (a) For $n = 1, \dots, N$:
 - i. Draw $\mathbf{x}'_n \sim \psi(\mathbf{x})$.
 - ii. Set $\mathbf{x}_{n,t} = \mathbf{x}'_n$, with probability

$$\alpha_n = \min \left[1, \frac{\pi(\mathbf{x}'_n)\psi(\mathbf{x}_{n,t-1})}{\pi(\mathbf{x}_{n,t-1})\psi(\mathbf{x}'_n)} \right] = \omega(\mathbf{x}'_n) \wedge \omega(\mathbf{x}_{n,t-1}).$$

Otherwise, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$. We have denoted $\omega(\mathbf{x}) = \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}$ and $a \wedge b = \min[a, b]$, with $a, b \in \mathbb{R}$.

- (b) Set $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$.

of multiple try schemes is particularly interesting. For instance, we adjust two special cases⁴ of the *Ensemble MCMC* [33] and *Multiple Try Metropolis* methods [21, 34, 23] for fitting them within O-MCMC. Tables 4 and 5 summarize them. In both cases, $L \geq 1$ different i.i.d. samples $\mathbf{z}_1, \dots, \mathbf{z}_L$ are drawn from $\psi(\mathbf{x})$. In the parallel Ensemble MCMC (P-EnM) scheme, at each iteration t , one resampling step per chain is performed, considering the set of $L + 1$ samples $\{\mathbf{z}_1, \dots, \mathbf{z}_L, \mathbf{x}_{n,t-1}\}$, $n = 1, \dots, N$ (using clearly importance weights). In the parallel MTM (P-MTM) scheme, at each iteration t , N resampling steps are performed considering the set of L candidates $\{\mathbf{z}_1, \dots, \mathbf{z}_L\}$ and then the new possible states are tested (i.e., accepted or not) according to suitable acceptance probabilities α_n , $n = 1, \dots, N$, involving also the previous states $\mathbf{x}_{n,t-1}$.

The ergodicity of both schemes is discussed in Appendix C. The algorithms in Tables 4-5 are obtained by a rearrangement of the basic schemes in [33, 21, 34] in order to generate, at each iteration t , N new states for N independent parallel chains. The new states of the N chains are selected filtering the same set of candidates $\{\mathbf{z}_1, \dots, \mathbf{z}_L\}$, drawn from the same independent proposal pdf ψ . Note that, with respect to a *standard* parallel approach, they require less evaluations of the target pdf: at each iteration, the algorithms in Tables 4-5 require L new evaluations of the target instead of NL (as occurs in a standard parallel approach). For further explanations, see Appendix C.1.1 and Figure 8. With $L = 1$, the algorithm in Table 4 coincides with the application of N parallel MH methods with Barker's acceptance rule [35]. The algorithm in Table 5 with $L = 1$ coincides with the scheme presented in Table 2. However, a number of tries $L \geq N$ is suggested.

5.2.1. Block Independent Multiple Try Metropolis algorithm

Previously, we have pointed out that with the scheme in Table 5 only L evaluations of the target are required at each iteration, instead of NL as standard parallel approach. The proposed scheme in Table 5 can be also modified in the same fashion of the block independent MH method [10] in order to and reducing the number of multinomial sampling steps, without jeopardizing the ergodicity of the parallel chains. We remark that the corresponding technique, called *Block Independent Multiple Try*

⁴They are special cases of the corresponding algorithms, since an independent proposal pdf φ is used.

Table 4. Parallel Ensemble MCMC (P-EnM) scheme for horizontal transitions in O-MCMC.

1. Build $\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t)$ as in Eq. (7) where $t = mT_V + (m - 1)T_H$.
2. For $t = mT_V + (m - 1)T_H + 1, \dots, m(T_V + T_H)$:
 - (a) Draw L possible i.i.d. candidates $\mathbf{z}_1, \dots, \mathbf{z}_L \sim \psi(\mathbf{x})$.
 - (b) For $n = 1, \dots, N$:
 - i. Set $\mathbf{x}_{n,t} = \mathbf{z}_k \in \{\mathbf{z}_1, \dots, \mathbf{z}_L\}$, i.e., with $k \in \{1, \dots, L\}$, with probability

$$\alpha_k = \frac{\frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{x}_{n,t-1})}{\psi(\mathbf{x}_{n,t-1})}} \quad k = 1, \dots, L, \quad (8)$$

or, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$ with probability

$$\alpha_{L+1} = 1 - \sum_{k=1}^L \alpha_k = \frac{\frac{\pi(\mathbf{x}_{n,t-1})}{\psi(\mathbf{x}_{n,t-1})}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{x}_{n,t-1})}{\psi(\mathbf{x}_{n,t-1})}}. \quad (9)$$

- ii. Set $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$.

Table 5. Parallel Multiple Try Metropolis (P-MTM) scheme for horizontal transitions in O-MCMC.

1. Build $\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t)$ as in Eq. (7) where $t = mT_V + (m - 1)T_H$.
2. For $t = mT_V + (m - 1)T_H + 1, \dots, m(T_V + T_H)$:
 - (a) Draw L possible i.i.d. candidates $\mathbf{z}_1, \dots, \mathbf{z}_L \sim \psi(\mathbf{x})$.
 - (b) Draw N independent samples $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$ such that $\mathbf{z}_{k_n} \in \{\mathbf{z}_1, \dots, \mathbf{z}_L\}$, i.e., with $k_n \in \{1, \dots, L\}$ and $n = 1, \dots, N$, with probability

$$\beta_{k_n} = \frac{\frac{\pi(\mathbf{z}_{k_n})}{\psi(\mathbf{z}_{k_n})}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)}}. \quad (10)$$

Namely, resample N times the samples in the set $\{\mathbf{z}_1, \dots, \mathbf{z}_L\}$ with probability β_k , $k = 1, \dots, L$.

- (c) For $n = 1, \dots, N$:
 - i. Set $\mathbf{x}_{n,t} = \mathbf{z}_{k_n}$ with probability

$$\alpha_n = \min \left[1, \frac{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} - \frac{\pi(\mathbf{z}_{k_n})}{\psi(\mathbf{z}_{k_n})} + \frac{\pi(\mathbf{x}_{n,t-1})}{\psi(\mathbf{x}_{n,t-1})}} \right]. \quad (11)$$

Otherwise, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$ (with probability $1 - \alpha_n$).

- (d) Set $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$.

Metropolis (BI-MTM), can be always employed when N parallel independent MTM are applied (even outside the O-MCMC scheme, clearly) in order to reduce the overall computational cost. Here, let us assume that the value N is such that the number of total transitions of one chain, T_H , can be divided in $B = \frac{T_H}{N} \in \mathbb{N}$ blocks. The idea is based on using N circular permutations of the resampled set $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$, i.e.,

$$\begin{aligned} \mathcal{V}_1 &= \{\mathbf{v}_{1,1} = \mathbf{z}_{k_1}, \mathbf{v}_{2,1} = \mathbf{z}_{k_2}, \dots, \mathbf{v}_{N-1,1} = \mathbf{z}_{k_{N-1}}, \mathbf{v}_{N,1} = \mathbf{z}_{k_N}\}, \\ \mathcal{V}_2 &= \{\mathbf{v}_{1,2} = \mathbf{z}_{k_N}, \mathbf{v}_{2,2} = \mathbf{z}_{k_1}, \dots, \mathbf{v}_{N-1,2} = \mathbf{z}_{k_{N-2}}, \mathbf{v}_{N,2} = \mathbf{z}_{k_{N-1}}\}, \\ &\vdots \\ \mathcal{V}_N &= \{\mathbf{v}_{1,N} = \mathbf{z}_{k_2}, \mathbf{v}_{2,N} = \mathbf{z}_{k_3}, \dots, \mathbf{v}_{N-1,N} = \mathbf{z}_{k_N}, \mathbf{v}_{N,N} = \mathbf{z}_{k_1}\}, \end{aligned} \quad (12)$$

where each set \mathcal{V}_n denotes one the N possible circular permutations of $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$. In order to preserve the ergodicity, each \mathbf{z}_{k_j} 's is drawn from a different set of tries $\mathcal{S}_j = \{\mathbf{z}_1^{(j)}, \dots, \mathbf{z}_L^{(j)}\}$. More specifically, before a *block* of N iterations, NL




Fig. 4. A graphical representation of one block within the BI-MTM technique, described in Table 6. One specific transition of one MTM chain is represented with the probability $\alpha_n(\mathbf{x}_{n,t-1}, \mathbf{v}_{n,j})$, showing the two possible future states at the t -th iteration, of the n -th chain. One block is formed by N transitions.

tries are drawn from $\psi(\mathbf{x})$, yielding N different sets $\mathcal{S}_j = \{z_1^{(j)}, \dots, z_L^{(j)}\}$, $j = 1, \dots, N$, each one containing L elements. Then, one sample z_{k_j} is resampled from each \mathcal{S}_j with probability proportional to the corresponding importance weight, and the circular permutations in Eq. (12) are created considering $\{z_{k_1}, \dots, z_{k_N}\}$. The complete BI-MTM algorithm is detailed in Table 6 and further considerations are provided in Appendix C. In Table 6, we have denoted the acceptance probability as $\alpha_n(\mathbf{x}_{n,t-1}, \mathbf{v}_{n,j})$ for remarking the two possible future states of the n -th chain at the t -th iteration. Figure 4 depicts a schematic sketch of the different steps of one block within the BI-MTM algorithm. Moreover, Figure 8 provides a graphical comparison among different parallel MTM approaches. BI-MTM requires only N multinomial sampling steps for each block, i.e., N iterations, instead of N^2 as in P-MTM in Table 5. Moreover, BI-MTM is completely parallelizable. Indeed, alternatively one could draw NLT_H samples from $\psi(\mathbf{x})$, perform NT_H multinomial sampling steps within NT_H different sets, and then run the T_H parallel iterations of the N chains, i.e., one unique block, using circular permutations of the NT_H resampled samples (previously obtained). The reduction of computation cost is obtained at the expense of a moderate decrease of the performance.

5.3. Computational Cost

In general, the most costly steps are the evaluation of the target pdf, depending on the model or the number of data. The number of evaluations E_H of the target, in one horizontal period, are $E_H = T_H$ for SMH in Table 1, whereas $E_H = LT_H$ in P-EnM and P-MTM (considering, in all cases, only the new evaluations at each iteration, the others can be automatically reused). Using SMH, T_H multinomial sampling steps are performed, each one over a population of N samples. In P-EnM and P-MTM, NT_H multinomial sampling steps are required (with $N > 1$), each one over a set of L samples. The total number of evaluations of the target $E_T = M(E_V + E_H)$, including the vertical transitions, is $E_T = M(NT_V + T_H)$ when the SMH is employed in the horizontal steps, or $E_T = M(NT_V + LT_H)$ when P-EnM and P-MTM are employed. Furthermore, in BI-MTM, we have again $E_T = M(NT_V + LT_H)$ but only T_H multinomial sampling steps. Note also that in a standard parallel multiple try approach we would have $E_H = NLT_H$ evaluations of the target and NT_H multinomial sampling steps, each one over a set of L samples. Finally, we remark that, using SMH, we perform one acceptance test in step, i.e., T_H in one horizontal period. Using a multiple candidates scheme, we employ NT_H acceptance test in one horizontal period. All these considerations are summarized in Table 7. For further details and observations, see Appendix C.1.1.

5.4. Relationship with other techniques

The techniques in Table 4 and 5 are particularly interesting since they involve the use of resampling steps without jeopardizing the ergodicity of the resulting global O-MCMC process. Moreover, the SMH algorithm in Table 1 employs an *inverted* resampling scheme since a sample in the population is chosen to be replaced with probability proportional to the inverse of the importance weights. Other methodologies in literature employ a combination of MCMC iterations and resampling steps, for instance, *iterated batch importance sampler* (IBIS) and *sequential Monte Carlo* (SMC) samplers for a static scenario [27, 28]. Their underlying idea could be interpreted as belonging to the O-MCMC philosophy: in these methodologies, the resampling steps are applied as an ‘‘horizontal’’ approach for interchanging information within the population. The resampling procedure generates

Table 6. Block Independent Multiple Try Metropolis (BI-MTM) algorithm for N parallel chains.

1. Let N be the total number of parallel MTM chains and T_H be the total number of iterations of each chain, such that $\frac{T_H}{N} \in \mathbb{N}$. Choose a number of tries L . Set $t_0 = mT_V + (m-1)T_H$ if BI-MTM is used within O-MCMC, otherwise set $t_0 = 0$.
2. For each block $b = 1, \dots, B = \frac{T_H}{N}$ do:

- (a) Draw NL possible i.i.d. candidates $\mathbf{z}_1^{(h)}, \dots, \mathbf{z}_L^{(h)} \sim \psi(\mathbf{x})$ with $h = 1, \dots, N$.
- (b) Draw one sample \mathbf{z}_{k_h} from each set $\mathcal{S}_h = \{\mathbf{z}_1^{(h)}, \dots, \mathbf{z}_L^{(h)}\}$, $h = 1, \dots, N$, with probability

$$\beta_\ell^{(h)} = \frac{\frac{\pi(\mathbf{z}_\ell^{(h)})}{\psi(\mathbf{z}_\ell^{(h)})}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell^{(h)})}{\psi(\mathbf{z}_\ell^{(h)})}}.$$

Thus, finally we have N different samples, $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$, such that each $\mathbf{z}_{k_h} \in \mathcal{S}_h$.

- (c) Create the circular permutations $\mathbf{v}_{n,j} \in \{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$ defined as in Eq. (12).
- (d) For $t = (b-1)N + 1 + t_0, \dots, bN + t_0$ (i.e., exactly N transitions):
 - i. Set $j = t - (b-1)N - t_0$ (so that $j = 1, \dots, N$, in one block).
 - ii. For $n = 1, \dots, N$:
 - A. Set $\mathbf{x}_{n,t} = \mathbf{v}_{n,j}$, with probability

$$\alpha_n(\mathbf{x}_{n,t-1}, \mathbf{v}_{n,j}) = \min \left[1, \frac{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell^{(j)})}{\psi(\mathbf{z}_\ell^{(j)})}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell^{(j)})}{\psi(\mathbf{z}_\ell^{(j)})} - \frac{\pi(\mathbf{v}_{n,j})}{\psi(\mathbf{v}_{n,j})} + \frac{\pi(\mathbf{x}_{n,t-1})}{\psi(\mathbf{x}_{n,t-1})}} \right].$$

Otherwise, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$.

- iii. Set $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$.

Table 7. Computational cost of O-MCMC given different horizontal schemes. Recall that $M = \frac{T}{T_V + T_H}$.

Computational features	SMH	P-EnM and P-MTM	BI-MTM	Standard Parallel MTM chains
E_H	T_H	LT_H	LT_H	NLT_H
$E_T = M(E_V + E_H)$	$M(NT_V + T_H)$	$M(NT_V + LT_H)$	$M(NT_V + LT_H)$	$M(NT_V + NLT_H)$
Total number of multinomial sampling steps	MT_H	MNT_H	MT_H	MNT_H
Cardinality of set for the multinomial sampling	N	L	L	L
Total number of acceptance tests	$M(NT_V + T_H)$	$M(NT_V + NT_H)$	$M(NT_V + NT_H)$	$M(NT_V + NT_H)$

samples from a particle approximation

$$\hat{\pi}^{(L)}(\mathbf{x}) = \sum_{\ell=1}^L \beta_\ell \delta(\mathbf{x} - \mathbf{z}_\ell), \quad (13)$$

of the measure of $\bar{\pi}(\mathbf{x})$, where $\mathbf{z}_\ell \sim \psi(\mathbf{x})$ (or, similarly, $\mathbf{z}_\ell \sim q_\ell(\mathbf{x})$ [36]) and β_ℓ are defined in Eq. (10) in Table 5, with $\ell = 1, \dots, L$. The quality of this approximation improves as the number L of samples grows. However, for a finite value of L there exists a discrepancy which can produce problems in the corresponding sampling algorithm. For further details see Appendix B. One main issue is the loss in diversity in the population.

This problem is reduced drastically in O-MCMC since the ergodicity is ensured in both, vertical and horizontal movements. Clearly, this improvement in the performance is obtained at the expense of an increase of the computational cost. For instance, let us consider the use of SMH in horizontal transitions. The cloud of samples is not impoverished by the application of SMH, even if a poor choice of the proposal $\varphi(\mathbf{x})$ is made. In the worst case, the newly proposed samples are always discarded and computational time is wasted. In the best case, a proposal located in a low probability region can jump close to a mode of the target. Clearly, In the mixture multiple try approach, it is better to choice $L \geq N$ for fostering the safeguard of the diversity.

Moreover, in the mixture approach, the mixture $\psi(\mathbf{x}) = \psi(\mathbf{x}|\mathcal{P}_t)$ is built using the states in \mathcal{P}_t as location parameters, and then it does not change for the next T_H horizontal steps. Thus, the information contained of the state $\{\mathbf{x}_{n,t}\}_{n=1}^N \in \mathcal{P}_t$ is employed in the next T_H iterations even if some state is not well-located. For clarifying this point, consider for instance the basic scheme in Table 2. The mixture $\psi(\mathbf{x}) = \psi(\mathbf{x}|\mathcal{P}_t)$ does not change so that the information provided by the population $\mathcal{P}_t = \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$ at the iteration t is still used in the iterations $t+1, \dots, t+T_H$. Namely, using a “particle filtering slang”, in these T_H steps, “no particles are killed” where the “particles” in this case are the N states $\{\mathbf{x}_{n,t}\}_{n=1}^N \in \mathcal{P}_t$ at the t -th iteration. This feature is also the main difference between the scheme in Table 2 and the NKC-type methods [32], where one component of the mixture is relocated after each iteration.

5.5. Joint adaptation of the proposal densities

Let us denote as \mathbf{C}_n and $\mathbf{\Lambda}_n$ the covariance matrices of the vertical and horizontal proposal pdfs, respectively. In order to design an algorithm as robust as possible, we suggest keeping the scale parameters \mathbf{C}_n fixed for the vertical proposal pdfs $q_n(\mathbf{x}|\mathbf{x}_{n,t-1}, \mathbf{C}_n)$, to avoid a loss of diversity within the set of chosen variances. However, if desired, they could be adapted easily as suggested in [7]. On the other hand, we suggest of adapting the scale parameters of the horizontal proposal pdfs φ_n , $n = 1, \dots, N$, since it is less delicate since a poor choice of the φ_n 's entails an increase in the computational cost but the diversity in the cloud of samples is always preserved. Several strategies have been proposed in [3, 37] and [7], for adapting proposal functions online within MCMC schemes. For the sake of simplicity, we discuss separately the cases of population-based or the mixture-based approaches.

- *Adaptation within SMH*: in this case, the strategies in [37, 7] are appropriate. Thus, After a training period $T_{train} < T$, all the generated samples (i.e., for each $t > T_{train}$ and from all the chains) can be used to adapt the location and scale parameters of proposal pdf $\varphi(\mathbf{x})$. Namely, denoting $\varphi_t(\mathbf{x}) = \varphi(\mathbf{x}; \boldsymbol{\mu}_t, \mathbf{\Lambda}_t)$, we can use the following approach:
 - If $t \leq T_{train}$: set $\boldsymbol{\mu}_t = \boldsymbol{\mu}_0$, $\mathbf{\Lambda}_t = \mathbf{\Lambda}_0$ (where $\boldsymbol{\mu}_0$ and $\mathbf{\Lambda}_0$ are the initial choices).
 - If $t > T_{train}$: set $\boldsymbol{\mu}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N \mathbf{x}_{n,j}$, and $\mathbf{\Lambda}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N (\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)(\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)^\top + \mathbf{C}$, where \mathbf{C} is a chosen covariance matrix. The empirical mean and covariance matrix estimators can be also computed recursively [3].
- *Adaptation of the mixture $\psi(\mathbf{x})$* : the methods in Section 5.2 employ a mixture $\psi(\mathbf{x}) = \frac{1}{N} \sum_{n=1}^N \varphi_n(\mathbf{x})$. In this case, each components $\varphi_{n,t}(\mathbf{x}) = \varphi_{n,t}(\mathbf{x}; \boldsymbol{\mu}_t, \mathbf{\Lambda}_t)$ should be adapted, jointly with the weights of the mixture. A possible (and simple) adaptation scheme is provided in [3] where all the parameters of the mixture are updated online. The method in [3] can easily reformulated for a framework with parallel chains. In this case, the states of the parallel chains are divided in N different clusters according to the euclidean distance between them and location parameters of the N components in the mixture $\psi(\mathbf{x})$. Then, new centroids (i.e., location parameters), covariances matrices and weights are updated according to the mean, covariances and cardinality of each cluster, respectively.

6. O-MCMC FOR OPTIMIZATION AND BIG DATA CONTEXT

6.1. Interacting Parallel Simulated Annealing

We can be easily modified the O-MCMC schemes converting them in stochastic optimization algorithms. Indeed, it is possible replacing the N vertical MH chains with N parallel *simulated annealing* (SA) methods [24, 25]. Let us denote as $\gamma_{n,t} \in (0, +\infty)$ a finite scale parameter that is decreasing function of t approaching zero for $t \rightarrow +\infty$, i.e.,

$$\begin{cases} \gamma_{n,t} \geq \gamma_{n,t+1} \geq \dots \geq \gamma_{n,t+\tau} > 0, \\ \lim_{t \rightarrow +\infty} \gamma_{n,t} = 0, \end{cases} \quad (14)$$

for $n = 1, \dots, N$. Moreover, for the sake of simplicity, we consider symmetric proposal functions $q_n(\mathbf{y}|\mathbf{x}) = q_n(\mathbf{x}|\mathbf{y})$. Then, one transition of n -th SA is described below:

1. Draw $\mathbf{x}' \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$.
2. Set $\mathbf{x}_{n,t} = \mathbf{x}'$ with probability

$$\alpha_n = \min \left[1, \frac{[\pi(\mathbf{x}')]^{\frac{1}{\gamma_{n,t}}}}{[\pi(\mathbf{x}_{n,t-1})]^{\frac{1}{\gamma_{n,t}}}} \right] = \min \left[1, \left(\frac{\pi(\mathbf{x}')}{\pi(\mathbf{x}_{n,t-1})} \right)^{\frac{1}{\gamma_{n,t}}} \right],$$

otherwise, with probability $1 - \alpha_n$, set $\mathbf{x}_{n,t} = \mathbf{x}_{n,t-1}$.

Above, with respect to the MH algorithm, we have replaced the target $\pi(\mathbf{x}) > 0$ with $[\pi(\mathbf{x})]^{\frac{1}{\gamma_{n,t}}} > 0$ with modes that become sharper and narrower when we reduce the scale parameter $\gamma_{n,t}$. Note that the movements such $\pi(\mathbf{x}') > \pi(\mathbf{x}_{n,t-1})$ are always accepted whereas, when $\pi(\mathbf{x}') < \pi(\mathbf{x}_{n,t-1})$, they are accepted with probability $P_d = \left(\frac{\pi(\mathbf{x}')}{\pi(\mathbf{x}_{n,t-1})}\right)^{\frac{1}{\gamma_{n,t}}} \in (0, 1]$. This probability $P_d \rightarrow 0$ vanishes to zero as $\gamma_{n,t} \rightarrow 0$ (guaranteeing the convergence to the global maximum when $t \rightarrow +\infty$). In the same fashion, for the horizontal steps we also consider the modified extended target,

$$\bar{\pi}_g(\mathbf{x}_1, \dots, \mathbf{x}_N) \propto \prod_{n=1}^N [\pi(\mathbf{x}_n)]^{\frac{1}{\gamma_{n,t}}}, \quad (15)$$

so that all the presented schemes, previously described, can be automatically applied. Several possible decreasing functions $\gamma_{n,t}$ has been suggested in [24, 26, 25].

6.2. O-MCMC for big data and data-tempered distributions

In a Big Data context, the posterior density $\bar{\pi}(\mathbf{x}) \propto \pi(\mathbf{x})$ is typically split in different partial posteriors, $\bar{\pi}_n(\mathbf{x}) \propto \pi_n(\mathbf{x})$, with $n = 1, \dots, N$, each one considering a disjoint sub-sets of the observed data and such that

$$\bar{\pi}(\mathbf{x}) \propto \prod_{n=1}^N \pi_n(\mathbf{x}),$$

The general idea is to generate samples from the partial posteriors $\bar{\pi}_n(\mathbf{x})$, and then to combine them in some way for approximating the complete target $\bar{\pi}(\mathbf{x})$ [13]. Here, we remark that ‘‘O-MCMC philosophy’’ can be applied in this framework, for improving the mixing of each chain. Thus, let us consider the application of N parallel (vertical) MCMC chains, addressing one different partial target $\bar{\pi}_n(\mathbf{x})$. Since each observed data provides information about the same phenomenon, an interaction among the N parallel chains can be improve the mixing of different MCMC techniques. In this context, the use of a mixture pdf as described in the O-MCMC approach of Section 5.2, appears appropriate for providing a cooperative interaction among the parallel chains. Using the similar observations, O-MCMC can be apply within the so called ‘‘data point tempered’’ techniques [27] where a sequence of P posteriors, $\pi_1(\mathbf{x}), \pi_2(\mathbf{x}), \dots, \pi_P(\mathbf{x})$, with an increasing number of data, are considered (typically, the last one contains all the data, i.e., $\pi_P(\mathbf{x}) = \pi(\mathbf{x})$).

7. NUMERICAL SIMULATIONS

7.1. Multimodal target distribution

In this section, we consider a bivariate multimodal target pdf, which is itself a mixture of 5 Gaussian pdfs, i.e.,

$$\bar{\pi}(\mathbf{x}) = \pi(\mathbf{x}) = \frac{1}{5} \sum_{i=1}^5 \mathcal{N}(\mathbf{x}; \boldsymbol{\nu}_i, \mathbf{G}_i), \quad \mathbf{x} \in \mathbb{R}^2, \quad (16)$$

with means $\boldsymbol{\nu}_1 = [-10, -10]^\top$, $\boldsymbol{\nu}_2 = [0, 16]^\top$, $\boldsymbol{\nu}_3 = [13, 8]^\top$, $\boldsymbol{\nu}_4 = [-9, 7]^\top$, and $\boldsymbol{\nu}_5 = [14, -14]^\top$, and with covariance matrices $\mathbf{G}_1 = [2, 0.6; 0.6, 1]$, $\mathbf{G}_2 = [2, -0.4; -0.4, 2]$, $\mathbf{G}_3 = [2, 0.8; 0.8, 2]$, $\mathbf{G}_4 = [3, 0; 0, 0.5]$, and $\mathbf{G}_5 = [2, -0.1; -0.1, 2]$. The target pdf $\pi(\mathbf{x})$ has then 5 different modes. We apply O-MCMC to estimate the expected value $E[\mathbf{X}]$ of $\mathbf{X} \sim \bar{\pi}(\mathbf{x})$ (the true mean is $E[\mathbf{X}] = [1.6, 1.4]^\top$) using different values for the number of parallel chains $N \in \{5, 100, 1000\}$. Furthermore, we choose deliberately a ‘‘bad’’ initialization to test the robustness of the algorithm. Specifically, we set $\mathbf{x}_{n,0} \sim \mathcal{U}([-4, 4] \times [-4, 4])$ for $n = 1, \dots, N$. This initialization is ‘‘bad’’ in the sense that it does not cover the modes of $\pi(\mathbf{x})$. In all cases, we consider $q_n(\mathbf{x}|\mathbf{x}_{n,t-1}) = \mathcal{N}(\mathbf{x}; \mathbf{x}_{n,t-1}, \mathbf{C}_n)$ using the same isotropic covariance matrix, $\mathbf{C}_n = \sigma^2 \mathbf{I}_2$, for every proposal of the vertical chains. We test different values of $\sigma \in \{2, 5, 10, 70\}$ to gauge the performance of O-MCMC. In O-MCMC, we consider the application of SMH and P-MTM as horizontal techniques, as described below. In both cases, we adapt the covariance matrices of the proposal pdfs as suggested in Section 5.5.

- *O-MCMC with SMH*: As horizontal proposal, we use again a Gaussian pdf, $\varphi_t(\mathbf{x}) = \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_t, \boldsymbol{\Lambda}_t)$ where $\boldsymbol{\mu}_t$ and $\boldsymbol{\Lambda}_t$ are adapted online: namely, $\boldsymbol{\mu}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N \mathbf{x}_{n,j}$, and $\boldsymbol{\Lambda}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N (\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)(\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)^\top + \boldsymbol{\Lambda}_0$, where

$\boldsymbol{\mu}_0 = [0, 0]^\top$, $\boldsymbol{\Lambda}_0 = \lambda^2 \mathbf{I}_2$ with $\lambda = 2$.⁵ As remarked in Section 5.5, this adaptive procedure is quite robust since employs samples generated by different parallel chains [7]. Furthermore, we fix $T = 4000$ and $T_H = T_V$. We test different values of $T_V \in \{1, 100\}$ and, as a consequence, $M = \frac{T}{T_V + T_H} = \frac{T}{2T_V} \in \{20, 2000\}$.⁶ Recall that the total number of evaluations of the targets in O-MCMC with SMH is $E_T = M(NT_V + T_H) = \frac{T}{2}(N + 1)$ (see Section 5.3).

- *O-MCMC with P-MTM*: We also test the O-MCMC scheme with P-MTM as horizontal technique. In this case, the (independent) proposal pdf is the mixture $\psi(\mathbf{x}) = \psi_m(\mathbf{x}|\mathcal{P}_t) = \frac{1}{N} \sum_{n=1}^N \varphi_n(\mathbf{x}|\mathbf{x}_{n,t}, \boldsymbol{\Lambda}_t)$ with $t = mT_V + (m - 1)T_H$, $\boldsymbol{\Lambda}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N (\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)(\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)^\top + \boldsymbol{\Lambda}_0$, where $\boldsymbol{\mu}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N \mathbf{x}_{n,j}$, $\boldsymbol{\Lambda}_0 = 4\mathbf{I}_2$ (for all $n = 1, \dots, N$). We consider different number of tries $L = \{5, 50\}$ and set again $T_V = T_H$. In this case, the number of evolution of the target is $E_T = M(NT_V + LT_H) = \frac{T}{2}(N + L)$ (see Section 5.3).

Comparison with independent parallel chains (IPCs). We compare the performance of O-MCMC with the application of IPCs, namely, only vertical independent transitions. Therefore, we can infer the benefit of applying the horizontal interaction. For a fair comparison, in IPCs we use the same MH kernels, i.e., with the same proposals q_n 's, and we keep fixed the total number of evaluations of the target E_T in both cases, O-MCMC and IPCs. Note that $E_T = NT'$ in IPCs where N is the number of chains and T' the total number of iterations for each one. We test different values of N . Tables 8 and 9 show the Mean Square Error (MSE), averaged among the two dimensions, in the estimation of the expected value $E[\mathbf{X}] = [1.6, 1.4]^\top$, averaged over 200 independent runs. O-MCMC with SMH always outperforms IPCs, specially for small σ and N . O-MCMC shows a much more stable behavior w.r.t. the parameter choice σ . For large scale parameters ($\sigma \in \{10, 70\}$) and a large number of chains ($N \in \{100, 1000\}$), the MSE of IPCs approaches the MSE of O-MCMC. A possible explanation is that the interaction is particularly useful with small N and a wrong choice of σ , whereas the use of large number of chains such as $N = 100$ or $N = 1000$ is enough, in this bidimensional example, for obtaining good performance. Moreover, O-MCMC with SMH presents an anomalous behavior when the variance of the vertical proposal pdfs is $\sigma = 2$. In this specific case, i.e., only for $\sigma = 2$, the MSE seems increases with N . However, note that O-MCMC provides the lower MSE, in any cases, comparing with the same computational effort E_T (with the exception of O-MCMC with P-MTM and $\sigma = 10$).

Comparison with Population Monte Carlo (PMC). We also compare with the standard PMC technique [38], described in Appendix B. We use $N \in \{100, 500, 2000\}$ and $T = 2000$ for PMC, so that the total number of evaluations of the target is $E_T = NT \in \{2 \cdot 10^5, 10 \cdot 10^5, 40 \cdot 10^5\}$. The proposal pdfs used in PMC are the same that we apply for the vertical chains in O-MCMC, i.e., $q_n(\mathbf{x}|\mathbf{x}_{n,t-1}) = \mathcal{N}(\mathbf{x}; \mathbf{x}_{n,t-1}, \mathbf{C}_n)$ using again the same covariance matrix, $\mathbf{C}_n = \sigma^2 \mathbf{I}_2$, for $n = 1, \dots, N$ and $\sigma \in \{2, 5, 10, 70\}$. We have considered a higher number of E_T for PMC with respect to O-MCMC, since O-MCMC involves several acceptance tests which are not contained in PMC. Thus, in order to provide a comparison as fair as possible, we allow a greater number of evaluations of the target, E_T , for PMC. Table 10 shows the MSE (mean of the MSEs of each component) of the O-MCMC schemes and the PMC method, for estimating $E[\mathbf{X}]$. We can see that the O-MCMC schemes, even with less E_T , provide lower MSEs with the exception of the cases corresponding to $\sigma = 10$.

N	O-MCMC with SMH						Independent parallel chains (IPCs)		
	5		100		1000		5	100	1000
T_V	1	100	1	100	1	100	—	—	—
$\sigma = 2$	1.4881	2.3649	1.7515	2.9146	5.6803	6.1354	28.7856	8.2925	7.3543
$\sigma = 5$	1.4989	2.1724	1.4512	1.7089	1.3606	1.4825	13.0602	2.2842	1.8373
$\sigma = 10$	1.1769	1.4034	0.1062	0.1129	0.0142	0.0139	2.4443	0.1247	0.0128
$\sigma = 70$	1.8175	2.0730	0.3554	0.3483	0.2866	0.2815	5.4897	0.5469	0.3264
T	4000						2400	2020	2002
E_T	$12 \cdot 10^3$		$2.02 \cdot 10^5$		$20.02 \cdot 10^5$		$12 \cdot 10^3$	$2.02 \cdot 10^5$	$20.02 \cdot 10^5$

Table 8. Mean Square Error (MSE) in the estimation of the mean of the target, using O-MCMC with SMH and IPCs, considering different values of σ and T_V (recall, we set $T_V = T_H$). The total number of evaluations of the target E_T is the same for O-MCMC (where $E_T = \frac{T}{2}(N + 1)$ since $T_V = T_H$) and IPCs (where $E_T = NT$).

⁵We set $T_{train} = T_V$, i.e., the adaptation starts after that the samples of the first vertical period are collected. Thus, before of the first horizontal step, $\varphi_t(\mathbf{x})$ has been already updated.

⁶We use all the generated samples in the estimation without removing any “burn-in” period.

	O-MCMC with P-MTM				Independent parallel chains (IPCs)	
	5		50		5	50
N	5		50		5	50
T_V	1		1		–	–
L	5	50	5	50	–	–
$\sigma = 2$	1.3907	1.1421	1.3156	0.7678	17.1352	3.5950
$\sigma = 5$	1.6159	0.9074	1.4011	1.0072	12.5791	2.3277
$\sigma = 10$	1.5738	0.8634	1.0982	0.8379	0.9403	0.1134
T	4000				4000	
E_T	$2 \cdot 10^4$	$11 \cdot 10^4$	$11 \cdot 10^4$	$20 \cdot 10^4$	$2 \cdot 10^4$	$20 \cdot 10^4$

Table 9. Mean Square Error (MSE) in the estimation of the mean of the target, using O-MCMC with P-MTM and IPCs, considering different values of σ and T_V (recall, we set $T_V = T_H$). The total number of evaluations of the target E_T is the same for O-MCMC (where $E_T = \frac{T}{2}(N + L)$ since $T_V = T_H$) and IPCs (where $E_T = NT$).

N	O-MCMC with SMH			O-MCMC with P-MTM		PMC		
	5	100	1000	5	50	100	500	2000
$\sigma = 2$	1.4881	1.7515	5.6803	1.3907	1.3156	48.11	35.1772	28.4326
$\sigma = 5$	1.4989	1.4512	1.3606	1.6159	1.4011	2.5998	2.3230	1.9153
$\sigma = 10$	1.1769	0.1062	0.0142	1.5738	1.0982	0.0512	0.0141	0.0054
$\sigma = 70$	1.8175	0.3554	0.2866	2.0185		2.3963	0.8252	0.1161
T	4000					2000		
E_T	$12 \cdot 10^3$	$2.02 \cdot 10^5$	$20.02 \cdot 10^5$	$2 \cdot 10^4$	$11 \cdot 10^4$	$2 \cdot 10^5$	$10 \cdot 10^5$	$40 \cdot 10^5$

Table 10. Mean Square Error (MSE) in the estimation of the mean of the target, using O-MCMC ($T_V = T_H = 1$ and $L = 5$ for P-MTM) and the standard PMC method [38].

7.2. Spectral analysis: estimating the frequencies of a noisy multi-sinusoidal signal

Many problems in science and engineering require dealing with a noisy multi-sinusoidal signal, whose general form is given by

$$y_c(\tau) = A_0 + \sum_{i=1}^S A_i \cos(2\pi f_i \tau + \phi_i) + r(\tau), \quad \tau \in \mathbb{R}, \quad (17)$$

where A_0 is a constant term, S is the number of sinusoids, $\{A_i\}_{i=1}^S$ is the set of amplitudes, $\{2\pi f_i\}_{i=1}^S$ are the frequencies, $\{\phi_i\}_{i=1}^S$ their phases, and $r(\tau)$ is an additive white Gaussian noise (AWGN) term. The estimation of the parameters of this signal is required by many applications in signal processing [39, 40], in control (where a multi-harmonic disturbance is often encountered in industrial plants) [41, 42] or in digital communications (where multiple narrowband interferers can be roughly modeled as sinusoidal signals) [43, 44]. Let us assume that we have K equispaced points from $y_c(\tau)$, obtained discretizing $y_c(\tau)$ with a period $T_s < \frac{\pi}{\max_{1 \leq i \leq S} 2\pi f_i}$ (in order to fulfill the sampling theorem [45]):

$$y[k] = A_0 + \sum_{i=1}^S A_i \cos(\Omega_i k + \phi_i) + r[k], \quad k = 1, \dots, K, \quad (18)$$

where $y[k] = y_c(kT_s)$ for $k = 0, 1, \dots, K - 1$, $\Omega_i = 2\pi f_i T_s$ for $i = 1, \dots, S$, and $r[k] \sim \mathcal{N}(0, \sigma_w^2)$. Our goal is applying the O-MCMC-type algorithms to provide an accurate estimate of the set of unknown frequencies, $\{\Omega_i\}_{i=1}^S$ or merely $\{f_i\}_{i=1}^S$. For keeping the notation of the rest of the work, we define the vector of possible frequencies as $\mathbf{x} \in \mathbb{R}^S$. Thus, given a fixed considering the hyper-rectangular domain $\mathcal{D} = [0, \frac{1}{2}]^S$ (it is straightforward to note the periodicity outside \mathcal{D}), and a uniform prior on \mathcal{D} , the posterior distribution given K data is $\bar{\pi}(\mathbf{x}) \propto \exp(-V(\mathbf{x}))$, where

$$V(\mathbf{x}) = V(x_1, \dots, x_S) = \frac{1}{2\sigma_w^2} \sum_{k=1}^K \left(y[k] - A_0 - \sum_{i=1}^S A_i \cos(x_i k + \phi_i) \right)^2 \mathbb{I}_{\mathcal{D}}(\mathbf{x}). \quad (19)$$

We have denoted $\mathbb{I}_{\mathcal{D}}(\mathbf{x})$ the indicator function such that $\mathbb{I}_{\mathcal{D}}(\mathbf{x}) = 1$ if $\mathbf{x} \in \mathcal{D}$ and $\mathbb{I}_{\mathcal{D}}(\mathbf{x}) = 0$ if $\mathbf{x} \notin \mathcal{D}$. Moreover, for the sake of simplicity, we have assumed also that S and σ_w^2 are known. Furthermore, we set $A_0 = 0$, $A_i = A = 1$ and $\phi_i = 0$.⁷ Note that the problem is symmetric with respect to the hyperplane $x_1 = x_2 = \dots = x_S$ (and, in general, multimodal). Bidimensional examples of $V(\mathbf{x}) = \log \pi(\mathbf{x})$ are depicted in Figure 5. We apply O-MCMC, comparing with IPCs, in two different types of experiments described briefly below. In all cases, we set $\mathbf{x}_{n,0} \sim \mathcal{U}(\mathcal{D})$ for $n = 1, \dots, N$, $T_H = T_V = 1$, and consider the proposals $q_n(\mathbf{x}|\mathbf{x}_{n,t-1}) = \mathcal{N}(\mathbf{x}; \mathbf{x}_{n,t-1}, \mathbf{C}_n)$ with $\mathbf{C}_n = \sigma^2 \mathbf{I}_S$, $n = 1, \dots, N$, for the vertical chains (\mathbf{I}_S is the unit matrix of dimension $S \times S$).

First experiment. We set $\mathbf{f} = [f_1 = 0.1, f_2 = 0.3]^\top$ and generate $K = 10$ synthetic data from the model. Since in this case, $S = 2$ and $\mathcal{D} = [0, \frac{1}{2}]^2$, it is possible to approximate the posterior with a very thin grid and compute the first 4 non-central moments and, as a consequence, we can compare the performance of different Monte Carlo sampling methods. Then, we test O-MCMC-SMH with the horizontal proposal $\varphi(\mathbf{x}) = \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}, \boldsymbol{\Lambda})$ where $\boldsymbol{\mu} = [0.25, 0.25]^\top$ and $\boldsymbol{\Lambda} = \sigma^2 \mathbf{I}_2$, i.e., uses the same σ considered for the vertical chains (recall that $\mathbf{C}_n = \sigma^2 \mathbf{I}_2$). We set the total number of target evaluations $E_T = M(N + 1) \in \{2730, 5450, 10.9 \cdot 10^3\}$. For a fair comparison, we consider N independent parallel chains (IPCs) choosing T such that $E'_T = NT$ is equal to E_T , i.e., $E'_T = E_T$ (see Section 5.3). We test different values of $\sigma \in [0.05, 0.5]$ and $N \in \{2, 5, 10\}$. We test the combinations of number of chains N and epochs M (T for IPCs) in order to keep fixed E_T . The Relative Error (RE) in the estimation, averaged over 500 independent runs, is shown in Figure 6. We can observe that O-MCMC (solid line) outperforms IPCs (dashed line) providing lower REs. The performance becomes similar as the computational effort E_T grows since the state space in the first experiment, $\mathcal{D} = [0, \frac{1}{2}]^2$, is small enough for allowing an exhaustive exploration of \mathcal{D} by independent chains.

Second experiment. We test O-MCMC in higher dimension considering $S = 4$, i.e., $\mathcal{D} = [0, \frac{1}{2}]^4$. We fix $\mathbf{f} = [f_1 = 0.1, f_2 = 0.2, f_3 = 0.3, f_4 = 0.4]^\top$. In this experiment, we consider an optimization problem for finding the global mode of π with $K = 30$ observations. With $K = 30$ observations, the main mode is enough tight around \mathbf{f} , so that we consider \mathbf{f} as true localization of the global mode. For simplicity and for breaking the symmetry, we restrict the search to the simplex contained in \mathcal{D} with vertices at $[0, 0, 0, 0]^\top$, $[\frac{1}{2}, 0, 0, 0]^\top$, $[\frac{1}{2}, \frac{1}{2}, 0, 0]^\top$, $[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 0]^\top$ and $[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}]^\top$. We test O-MCMC-PMTM considering again Gaussian horizontal proposals in the mixture ψ , with $\boldsymbol{\Lambda} = \lambda^2 \mathbf{I}_4$ for all n . We test $\lambda = 0.1$ and $\lambda = \sigma$ (where σ is employed in the covariance matrices $\mathbf{C}_n = \sigma^2 \mathbf{I}_4$ of the vertical chains). Moreover, we test the adaptation of $\boldsymbol{\Lambda}$, i.e., $\boldsymbol{\Lambda}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N (\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)(\mathbf{x}_{n,j} - \boldsymbol{\mu}_t)^\top + \boldsymbol{\Lambda}_0$, where $\boldsymbol{\mu}_t = \frac{1}{Nt} \sum_{j=1}^t \sum_{n=1}^N \mathbf{x}_{n,j}$, $\boldsymbol{\Lambda}_0 = 0.02 \mathbf{I}_4$, for all $n = 1, \dots, N$. We set $N = 20$ as number of chains, $L = 20$ as number of tries, and $E_T \approx 8700$ as total number of evaluations of the target. For a fair comparison, we again consider N and T for IPCs such that $E'_T = NT$ is equal to E_T , i.e., $E'_T = E_T$. The vertical proposal pdfs are the same than those for the IPCs scheme. Furthermore, we apply a data-tempering approach [27] described in Section 6, employing a sequence of 29 target pdfs π_i each one considering an increasing number of observations $K_i = 2 + (i - 1)$ with $i = 1, \dots, 29$. The computational effort E_T is distributed uniformly in each π_i . We compute the Relative Error (RE) of the last states of the N chains with respect to the true vector \mathbf{f} . Figure 7 depicts the curves of the RE versus different values of $\sigma \in [0.05, 0, 5]$. We can observe that O-MCMC-PMTM always outperforms IPCs in this optimization problem.

8. CONCLUSIONS

In this work, we have introduced a novel family of MCMC algorithms, named Orthogonal MCMC schemes, that incorporates “horizontal” MCMC transitions to share information among a cloud of parallel “vertical” MCMC chains. We have described different alternatives for exchanging information among independent parallel chains. Compared to the fully independent parallel chains approach, the novel interacting techniques show a more robust behavior with respect to the parameterization and better performance for different number of chains. One reason is that the novel algorithms provide a good trade-off between the use of an independent or random walk proposal density, i.e., between local and global explorations. We have considered two different approaches for the interaction among the chains: in the first one, a MCMC technique over the entire population is directly applied whereas, in the second one, the initial population \mathcal{P}_t is used for building a suitable mixture density $\psi(\mathbf{x})$ employed as proposal function in the horizontal transitions. This second approach can be interpreted as an adaptive MCMC scheme where the location parameters of the N components of mixture $\psi(\mathbf{x})$ are updated driven by N parallel MCMC chains. The outputs of these parallel chains are also employed in the approximation of the target. Furthermore, we have designed different parallel Multiple Try Metropolis (P-MTM) schemes using an independent proposal pdf, where the drawn candidates are “recycled” in order to reduce the overall computational cost. Finally, we have described two modified versions of O-MCMC for optimization and the application to big data problems. The ergodicity of all the proposed methodologies have been discussed and several

⁷Let us remark that the estimation of all these parameters would make the inference harder, but can be easily incorporated into our algorithm.




Fig. 5. Several examples of function $V(\mathbf{x}) = \log \pi(\mathbf{x})$ with $S = 2$, given different realizations of the measurements $y[1], \dots, y[K]$. In Figures (a)-(d)-(e)-(f), we set $\mathbf{f} = [\frac{1}{6}, \frac{1}{3}]^\top$ and consider $K = 10, 15, 20, 30$, respectively. In Figures (b)-(c), we set $\mathbf{f} = [0.1, 0.3]^\top$ and $K = 10$. Black dotted points shows all the states generated throughout an O-MCMC-PMTM run with $N = 10, L = 10$ and $T = 500$. The initial states are chosen uniformly within $\mathcal{D} = [0, \frac{1}{2}]^\top$.




Fig. 6. Relative Error (averaged over 500 runs) in the first experiment for O-MCMC-SMH (solid line) and IPCs (dashed line) with different computational effort E_T . Note that O-MCMC always outperforms IPCs. Note also that their performance becomes similar as the overall computational cost E_T grows (due to the small size of the state space, $\mathcal{D} = [0, \frac{1}{2}]^2$).

numerical simulations have been provided showing the advantages of the novel approach.




Fig. 7. Relative Error (averaged over 500 runs) in the second experiment (dimension $S = 4$) searching the global maximum with O-MCMC (solid line) and IPCs (dashed line). In the O-MCMC-PMTM scheme, we use different $\lambda = 0.1$ and $\lambda = \sigma$, for the covariance matrices $\Lambda_n = \lambda^2 \mathbf{I}_4$ of the horizontal proposal pdfs in the mixture ψ . Moreover, we test an adapted covariance matrices for the horizontal proposal pdfs (see Figure (c)).

9. ACKNOWLEDGEMENTS

This work has been supported by the ERC grant 239784 and AoF grant 251170, the Spanish government through the CONSOLIDER-INGENIO 2010 program (CSD2008-00010) and AGES (S2010/BMD-2422), OTOSiS (TEC2013-41718-R), and DISSECT (TEC2012-38058-C03-01) projects, the BBVA Foundation through the MG-FIAR project, obtained in “Primera Convocatoria de Ayudas Fundación BBVA a Investigadores, Innovadores y Creadores Culturales”, by the Grant 2014/23160-6 of São Paulo Research Foundation (FAPESP) and by the Grant 305361/2013-3 of National Council for Scientific and Technological Development (CNPq).

10. REFERENCES

- [1] A. Doucet and X. Wang, “Monte Carlo methods for signal processing,” *IEEE Signal Processing Magazine*, vol. 22, no. 6, pp. 152–170, Nov. 2005.
- [2] J. Ye, A. Wallace, and J. Thompson, “Parallel Markov chain Monte Carlo computation for varying-dimension signal analysis,” in *Proc. EUSIPCO 2009*, Glasgow (Scotland), 24–28 Aug. 2009, pp. 2673–2677.
- [3] D. Luengo and L. Martino, “Fully adaptive Gaussian mixture Metropolis-Hastings algorithm,” in *Proc. ICASSP 2013*, Vancouver (Canada), 2013, pp. 6148–6152.
- [4] L. Martino, V. Elvira, D. Luengo, and J. Corander, “An adaptive population importance sampler: Learning from the uncertainty,” *IEEE Transactions on Signal Processing*, vol. 63, no. 16, pp. 4422–4437, 2015.
- [5] C. P. Robert and G. Casella, *Monte Carlo Statistical Methods*, Springer, 2004.
- [6] L. Martino, J. Read, and D. Luengo, “Independent doubly adaptive rejection Metropolis sampling within Gibbs sampling,” *IEEE Transactions on Signal Processing*, vol. 63, no. 12, pp. 3123–3138, 2015.
- [7] R. Craiu, J. Rosenthal, and C. Yang, “Learn from the neighbor: Parallel-chain and regional adaptive MCMC,” *Journal of the American Statistical Association*, vol. 104, no. 448, pp. 1454–1466, 2009.
- [8] W. J. Fitzgerald, “Markov chain Monte Carlo methods with applications to signal processing,” *Signal Processing*, vol. 81, no. 1, pp. 3–18, January 2001.
- [9] R. Casarin, R. V. Craiu, and F. Leisen, “Interacting multiple try algorithms with different proposal distributions,” *Statistics and Computing*, vol. 23, no. 2, pp. 185–200, 2013.
- [10] P. Jacob, C. P. Robert, and M. H. Smith, “Using parallel computation to improve Independent Metropolis-Hastings based estimation,” *Journal of Computational and Graphical Statistics*, vol. 3, no. 20, pp. 616–635, 2011.

- [11] B. Calderhead, “A general construction for parallelizing Metropolis-Hastings algorithms,” *Proceedings of the National Academy of Sciences of the United States of America (PNAS)*, vol. 111, no. 49, pp. 17408–17413, 2014.
- [12] R. Bardenet, A. Doucet, and C. Holmes, “On Markov chain Monte Carlo methods for tall data,” *arXiv:1505.02827*, 2015.
- [13] W. Neiswanger, C. Wang, and E. Xing, “Asymptotically exact, embarrassingly parallel MCMC,” *arXiv:1311.4780*, pp. 1–16, 21 Mar. 2014.
- [14] X. Wang and D. B. Dunson, “Parallelizing MCMC via Weierstrass sampler,” *arXiv:1312.4605v2*, 25 May 2014.
- [15] X. Wang, F. Guo, K. A. Heller, and D. B. Dunson, “Parallelizing MCMC with random partition trees,” *arXiv:1311.4780*, 2015.
- [16] Steven L. Scott, Alexander W. Blocker, Fernando V. Bonassi, Hugh A. Chipman, Edward I. George, and Robert E. McCulloch, “Bayes and big data: The consensus Monte Carlo algorithm,” in *EFaBBayes 250th conference*, 2013, vol. 16.
- [17] L. Martino, V. Elvira, D. Luengo, A. Artes, and J. Corander, “Orthogonal MCMC algorithms,” *IEEE Workshop on Statistical Signal Processing (SSP)*, pp. 364–367, June 2014.
- [18] A. Jasra, D. A. Stephens, and C. C. Holmes, “On population-based simulation for static inference,” *Statistics and Computing*, vol. 17, no. 3, pp. 263–279, 2007.
- [19] J. Corander, M. Gyllenberg, and T. Koski, “Bayesian model learning based on a parallel MCMC strategy,” *Statistics Computing*, vol. 16, pp. 355–362, 2006.
- [20] J. Corander, M. Ekdahl, and T. Koski, “Parallel interacting mcmc for learning of topologies of graphical models,” *Data Mining and Knowledge Discovery*, vol. 17, no. 3, pp. 431–456, 2008.
- [21] J. S. Liu, F. Liang, and W. H. Wong, “The multiple-try method and local optimization in Metropolis sampling,” *Journal of the American Statistical Association*, vol. 95, no. 449, pp. 121–134, March 2000.
- [22] L. Martino, V. P. Del Olmo, and J. Read, “A multi-point Metropolis scheme with generic weight functions,” *Statistics and Probability Letters*, vol. 82, no. 7, pp. 1445–1453, 2012.
- [23] L. Martino and J. Read, “On the flexibility of the design of multiple try Metropolis schemes,” *Computational Statistics*, vol. 28, no. 6, pp. 2797–2823, 2013.
- [24] S. K. Kirkpatrick, C. D. Gelatt Jr., and M. P. Vecchi, “Optimization by simulated annealing,” *Science*, vol. 220, no. 4598, pp. 671–680, May 1983.
- [25] M. Locatelli, “Convergence of a simulated annealing algorithm for continuous global optimization,” *Journal of Global Optimization*, vol. 18, pp. 219–234, 2000.
- [26] F. Liang, Y. Cheng, and G. Lin, “Simulated stochastic approximation annealing for global optimization with a square-root cooling schedule,” *Journal of the American Statistical Association*, vol. 109, no. 506, pp. 847–863, 2014.
- [27] N. Chopin, “A sequential particle filter for static models,” *Biometrika*, vol. 89, pp. 539–552, 2002.
- [28] P. Del Moral, Arnaud Doucet, and A. Jasra, “Sequential Monte Carlo samplers,” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, vol. 68, no. 3, pp. 411–436, 2006.
- [29] G. Roberts and R. Tweedie, “Exponential convergence of Langevin distributions and their discrete approximations,” *Bernoulli*, vol. 2, no. 4, pp. 341–363, 1996.
- [30] A. Mira, “On Metropolis-Hastings algorithms with delayed rejection,” *Metron*, vol. LIX, no. 3-4, pp. 231–241, 2001.
- [31] F. Liang, C. Liu, and R. Carroll, *Advanced Markov Chain Monte Carlo Methods: Learning from Past Samples*, Wiley Series in Computational Statistics, England, 2010.
- [32] G. R. Warnes, “The Normal Kernel Coupler: An adaptive Markov Chain Monte Carlo method for efficiently sampling from multi-modal distributions,” *Technical Report*, 2001.

- [33] R. Neal, “MCMC using ensembles of states for problems with fast and slow variables such as Gaussian process regression,” *arXiv:1101.0387*, 2011.
- [34] J. S. Liu, *Monte Carlo Strategies in Scientific Computing*, Springer, 2004.
- [35] A. A. Barker, “Monte Carlo calculations of the radial distribution functions for a proton-electron plasma,” *Australian Journal of Physics*, vol. 18, pp. 119–133, 1965.
- [36] Víctor Elvira, Luca Martino, David Luengo, and Mónica F Bugallo, “Efficient multiple importance sampling estimators,” *Signal Processing Letters, IEEE*, vol. 22, no. 10, pp. 1757–1761, 2015.
- [37] Heikki Haario, Eero Saksman, and Johanna Tamminen, “An adaptive Metropolis algorithm,” *Bernoulli*, vol. 7, no. 2, pp. 223–242, April 2001.
- [38] O. Cappé, A. Guillin, J. M. Marin, and C. P. Robert, “Population Monte Carlo,” *Journal of Computational and Graphical Statistics*, vol. 13, no. 4, pp. 907–929, 2004.
- [39] Petre Stoica, “List of references on spectral line analysis,” *Signal Processing*, vol. 31, no. 3, pp. 329–340, 1993.
- [40] Hing-Cheung So, Kit Wing Chan, Yiu-Tong Chan, and KC Ho, “Linear prediction approach for efficient frequency estimation of multiple real sinusoids: algorithms and analyses,” *Signal Processing, IEEE Transactions on*, vol. 53, no. 7, pp. 2290–2305, 2005.
- [41] Marc Bodson and Scott C Douglas, “Adaptive algorithms for the rejection of sinusoidal disturbances with unknown frequency,” *Automatica*, vol. 33, no. 12, pp. 2213–2221, 1997.
- [42] Alexey A Bobtsov and AA Pyrkin, “Cancellation of unknown multiharmonic disturbance for nonlinear plant with input delay,” *International Journal of Adaptive Control and Signal Processing*, vol. 26, no. 4, pp. 302–315, 2012.
- [43] Catharina Carlemalm, H Vincent Poor, and Andrew Logothetis, “Suppression of multiple narrowband interferers in a spread-spectrum communication system,” *Selected Areas in Communications, IEEE Journal on*, vol. 18, no. 8, pp. 1365–1374, 2000.
- [44] K Deergha Rao, MNS Swamy, and Eugene I Plotkin, “A nonlinear adaptive filter for narrowband interference mitigation in spread spectrum systems,” *Signal processing*, vol. 85, no. 3, pp. 625–635, 2005.
- [45] J. G. Proakis, *Digital Communications*, McGraw-Hill, Singapore, 1995.
- [46] S. Brooks, A. Gelman, G. Jones, and X. L. Meng, *Handbook of Markov Chain Monte Carlo*, Chapman and Hall CRC Press, 2011.

A. STATIONARY DISTRIBUTION OF O-MCMC

A.1. Analysis for the mixture-based approach

Let us consider two MCMC kernels, $K_n^{(V)}(\mathbf{y}|\mathbf{x})$, $K_n^{(H)}(\mathbf{z}|\mathbf{y})$ with $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathcal{D} \in \mathbb{R}^{d_x}$, corresponding to the n -th chain for the vertical and horizontal steps, respectively. We assume $\bar{\pi}(\cdot)$ is the invariant density of both chains. Namely, we consider MCMC techniques which steps are summarized in the two conditional probabilities $K_n^{(V)}(\mathbf{y}|\mathbf{x})$ and $K_n^{(H)}(\mathbf{z}|\mathbf{y})$, such that

$$\int_{\mathcal{D}} K_n^{(V)}(\mathbf{y}|\mathbf{x})\bar{\pi}(\mathbf{x})d\mathbf{x} = \bar{\pi}(\mathbf{y}), \quad \int_{\mathcal{D}} K_n^{(H)}(\mathbf{z}|\mathbf{y})\bar{\pi}(\mathbf{y})d\mathbf{y} = \bar{\pi}(\mathbf{z}).$$

For the sake of simplicity, we tackle a simpler case where $K_n^{(V)}$, $K_n^{(H)}$ are used sequentially, once each. Namely, we consider the sequential application of $K_n^{(V)}$ and $K_n^{(H)}$, i.e, first $\mathbf{y}' \sim K_n^{(V)}(\mathbf{y}|\mathbf{x})$ and then draw $\mathbf{z}' \sim K_n^{(H)}(\mathbf{z}|\mathbf{y}')$, i.e., the probability of transition from \mathbf{z} to \mathbf{x}

$$T(\mathbf{z}|\mathbf{x}) = \int_{\mathcal{D}} K_n^{(H)}(\mathbf{z}|\mathbf{y})K_n^{(V)}(\mathbf{y}|\mathbf{x})d\mathbf{y}. \quad (20)$$

The target $\bar{\pi}$ is also invariant w.r.t. $T(\mathbf{z}|\mathbf{x})$ [46, Chapter 1]. Indeed, we can write

$$\begin{aligned}
\int_{\mathcal{D}} T(\mathbf{z}|\mathbf{x})\bar{\pi}(\mathbf{x})d\mathbf{x} &= \int_{\mathcal{D}} \left[\int K_n^{(H)}(\mathbf{z}|\mathbf{y})K_n^{(V)}(\mathbf{y}|\mathbf{x})d\mathbf{y} \right] \bar{\pi}(\mathbf{x})d\mathbf{x}, \\
&= \int_{\mathcal{D}} K_n^{(H)}(\mathbf{z}|\mathbf{y}) \left[\int K_n^{(V)}(\mathbf{y}|\mathbf{x})\bar{\pi}(\mathbf{x})d\mathbf{x} \right] d\mathbf{y}, \\
&= \int_{\mathcal{D}} K_n^{(H)}(\mathbf{z}|\mathbf{y})\bar{\pi}(\mathbf{y})d\mathbf{y}, \\
&= \bar{\pi}(\mathbf{z}),
\end{aligned} \tag{21}$$

that is the definition of invariant pdf w.r.t. $T(\mathbf{z}|\mathbf{x})$.

A.2. Analysis for the population-based approach

Considering now an extended state space $\mathbb{R}^{d_x \times N}$, we can interpret that O-MCMC yields a unique chain in $\mathbb{R}^{d_x \times N}$. Namely, one population of states at the t -th iteration represents one extended state of this unique chain. Here, we show that this chain, generated by O-MCMC, has the extended target density

$$\bar{\pi}_g(\mathbf{x}_1, \dots, \mathbf{x}_N) \propto \prod_{n=1}^N \pi(\mathbf{x}_n),$$

as invariant pdf. We can use similar arguments employed for the mixture-based schemes, considering now a population of current states, i.e.,

$$\mathcal{P}_{t-1} = \{\mathbf{x}_{1,t-1}, \dots, \mathbf{x}_{N,t-1}\}.$$

We denote the vertical MCMC kernels as $K_n(\mathbf{x}_{n,t}|\mathbf{x}_{n,t-1})$ with $\bar{\pi}$ invariant, whereas one horizontal kernel $K_H(\mathcal{P}_t|\mathcal{P}_{t-1})$ ⁸ with invariant pdf the following extended target pdf

$$\bar{\pi}_g(\mathcal{P}) \propto \pi_g(\mathbf{x}_1, \dots, \mathbf{x}_N) = \prod_{n=1}^N \pi(\mathbf{x}_n).$$

The complete kernel of the orthogonal procedure, formed by one vertical and one orthogonal step, is

$$T(\mathcal{P}_t|\mathcal{P}_{t-2}) = \int_{\mathcal{D}^N} K_H(\mathcal{P}_t|\mathcal{P}_{t-1}) \left[\prod_{n=1}^N K_n(\mathbf{x}_{n,t-1}|\mathbf{x}_{n,t-2}) \right] \prod_{n=1}^N d\mathbf{x}_{n,t-1}.$$

In this case, we can write

$$\begin{aligned}
&\int_{\mathcal{D}^N} T(\mathcal{P}_t|\mathcal{P}_{t-2})\bar{\pi}_g(\mathcal{P}_{t-2})d\mathcal{P}_{t-2} = \\
&= \int_{\mathcal{D}^N} \int_{\mathbb{R}^N} K_H(\mathcal{P}_t|\mathcal{P}_{t-1}) \prod_{n=1}^N K_n(\mathbf{x}_{n,t-1}|\mathbf{x}_{n,t-2}) \prod_{n=1}^N \bar{\pi}(\mathbf{x}_{n,t-2}) \prod_{n=1}^N d\mathbf{x}_{n,t-1} \prod_{n=1}^N d\mathbf{x}_{n,t-2}. \\
&= \int_{\mathcal{D}^N} K_H(\mathcal{P}_t|\mathcal{P}_{t-1}) \prod_{n=1}^N \bar{\pi}(\mathbf{x}_{n,t-1}) \prod_{n=1}^N d\mathbf{x}_{n,t-1} = \int_{\mathbb{R}^N} K_H(\mathcal{P}_t|\mathcal{P}_{t-1})\bar{\pi}_g(\mathcal{P}_{t-1})d\mathcal{P}_{t-1} = \bar{\pi}_g(\mathcal{P}_t).
\end{aligned}$$

Namely, the kernel $T(\mathcal{P}_t|\mathcal{P}_{t-2})$ has $\bar{\pi}_g$ as invariant density. This result can be easily extended when T_V vertical and T_H horizontal transitions are applied, using the same arguments. Note that the generated chain preserves the pdf π as shown previously, but in general is not reversible [46, Section 1.12.7].

⁸For the sake of simplicity, we abuse of the notation using here the set \mathcal{P}_t as a vector.

B. DISTRIBUTION AFTER RESAMPLING

Resampling procedures are employed in different Monte Carlo techniques such as Population Monte Carlo (PMC), Iterated Batch Importance Sampler (IBIS) and, more generally, in Sequential Monte Carlo (SMC) methods for a static scenario [38, 27, 28]. For simplicity, let us consider here a standard PMC-type scheme. In PMC, N different proposal pdfs q_1, \dots, q_N are employed at each iteration. Starting from $\{\mathbf{x}_{1,0}, \dots, \mathbf{x}_{N,0}\}$, the basic PMC scheme consists of the following steps:

1. For $t = 1, \dots, T$:

(a) For $n = 1, \dots, N$, draw one sample $\mathbf{x}_{n,t}$ from q_n , i.e.,

$$\mathbf{x}_{n,t} \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1}),$$

(b) Draw N independent samples $\{\mathbf{z}_1, \dots, \mathbf{z}_N\}$ such that each $\mathbf{z}_n \in \{\mathbf{x}_{1,t}, \dots, \mathbf{x}_{N,t}\}$, with $n = 1, \dots, N$, with probability

$$\beta_n = \frac{\frac{\pi(\mathbf{x}_{n,t})}{q_n(\mathbf{x}_{n,t})}}{\sum_{n=1}^N \frac{\pi(\mathbf{x}_{n,t})}{q_n(\mathbf{x}_{n,t})}}. \quad (22)$$

(c) Set $\mathbf{x}_{n,t} = \mathbf{z}_n$.

The step 2(b) corresponds to resample (with replacement) N times the population $\{\mathbf{x}_{n,t}\}_{n=1}^N$. Note that the weights in Eq. (22) are the same used in (10). For the sake of simplicity, since here we consider a generic iteration t , let us simplify the notation denoting as $\mathbf{x}_n = \mathbf{x}_{n,t} \sim q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$ ($1 \leq n \leq N$, $1 \leq t \leq T$), and as $q_n(\mathbf{x}) = q_n(\mathbf{x}|\mathbf{x}_{n,t-1})$. Moreover, we define as

$$\mathbf{m}_{-n} = [\mathbf{x}_1, \dots, \mathbf{x}_{n-1}, \mathbf{x}_{n+1}, \dots, \mathbf{x}_N],$$

the matrix containing all the samples except for the n -th. Let us also denote as $\mathbf{z} \in \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$, a generic sample after applying one multinomial resampling step. Hence, the distribution of \mathbf{z} is given by

$$\phi(\mathbf{z}) = \int_{\mathcal{D}^N} \hat{\pi}^{(N)}(\mathbf{z}|\mathbf{x}_1, \dots, \mathbf{x}_N) \left[\prod_{n=1}^N q_n(\mathbf{x}_n) \right] d\mathbf{x}_1 \dots d\mathbf{x}_N, \quad (23)$$

where

$$\hat{\pi}^{(N)}(\mathbf{z}|\mathbf{x}_1, \dots, \mathbf{x}_N) = \sum_{j=1}^N \beta_j \delta(\mathbf{z} - \mathbf{x}_j), \quad (24)$$

and β_j are given in Eq. (22). Here, with the notation $\hat{\pi}^{(N)}(\mathbf{z}|\mathbf{x}_1, \dots, \mathbf{x}_N)$ we have remarked the dependence on the generated samples \mathbf{x}_n 's in order to facilitate the understanding of Eq. (23). Then, after some straightforward rearrangements, Eq. (23) can be rewritten as

$$\phi(\mathbf{z}) = \sum_{j=1}^N \left(\int_{\mathcal{D}^{N-1}} \frac{\frac{\pi(\mathbf{x}_j)}{q_j(\mathbf{x}_j)}}{\sum_{n=1}^N \frac{\pi(\mathbf{x}_n)}{q_n(\mathbf{x}_n)}} \left[\prod_{\substack{n=1 \\ n \neq j}}^N q_n(\mathbf{x}_n) \right] d\mathbf{m}_{-j} \right) \delta(\mathbf{z} - \mathbf{x}_j). \quad (25)$$

Finally, we can write

$$\phi(\mathbf{z}) = \pi(\mathbf{z}) \sum_{j=1}^N \int_{\mathcal{D}^{N-1}} \frac{1}{N \hat{Z}} \left[\prod_{\substack{n=1 \\ n \neq j}}^N q_n(\mathbf{x}_n) \right] d\mathbf{m}_{-j}, \quad (26)$$

where $\hat{Z} = \frac{1}{N} \sum_{n=1}^N \frac{\pi(\mathbf{x}_n)}{q_n(\mathbf{x}_n)}$ is the estimate of the normalizing constant of the target obtained by using the importance sampling technique. When $N \rightarrow \infty$, then $\hat{Z} \rightarrow Z$ [5], and thus $\phi(\mathbf{z}) \rightarrow \frac{1}{Z} \pi(\mathbf{z}) = \bar{\pi}(\mathbf{z})$. Clearly, there exists a discrepancy between $\phi(\mathbf{z})$ and $\bar{\pi}(\mathbf{z})$.

C. ERGODICITY OF THE PARALLEL SCHEMES BASED ON MULTIPLE CANDIDATES

Similarly as in PMC, the parallel Ensemble MCMC (P-EnM) and Multiple Try Metropolis (P-MTM) schemes in Tables 4-5 are based on the particle approximations of the measure of the target. In both cases, L independent samples $\mathbf{z}_1, \dots, \mathbf{z}_L$ drawn from $\psi(\mathbf{x})$, i.e.,

$$\mathbf{z}_\ell \sim \psi(\mathbf{x}), \quad (27)$$

for $\ell = 1, \dots, L$. Below, we show that P-EnM and P-MTM yield reversible chains with stationary density the generalized pdf $\bar{\pi}_g$, proving the detailed balance condition is satisfied [5].

C.1. Parallel Multiple Try Metropolis

In P-MTM, we can define the particle approximation based on the set $\{\mathbf{z}_1, \dots, \mathbf{z}_L\}$, i.e.,

$$\hat{\pi}^{(L)}(\mathbf{z}) = \sum_{\ell=1}^L \beta_\ell \delta(\mathbf{z} - \mathbf{z}_\ell), \quad (28)$$

where the normalized weights β_ℓ 's are given in Eq (10). Note that, the expression above coincides with Eq. (24). Let us also denote as the matrix

$$\mathbf{m}_{-k} = [\mathbf{z}_1, \dots, \mathbf{z}_{k-1}, \mathbf{z}_{k+1}, \dots, \mathbf{z}_L],$$

containing all the samples \mathbf{z}_ℓ 's with the exception of \mathbf{z}_k . We denote as $K_n(\mathbf{x}_{n,t} | \mathbf{x}_{n,t-1})$ is the MTM kernel of n -th chain, namely, $K_n(\mathbf{z} | \mathbf{x})$ is the probability of the n -th chain of jumping from the state $\mathbf{x} = \mathbf{x}_{t-1}$ to $\mathbf{z} = \mathbf{z}_k \in \{\mathbf{z}_1, \dots, \mathbf{z}_L\}$ (for simplicity, we consider here only the case $\mathbf{z} \neq \mathbf{x}$). Note that all the \mathbf{z}_ℓ 's are both drawn and resampled independently (see steps 2(a) and 2(b) in Table 5). Thus, the conditional probability $K_n(\mathbf{z} | \mathbf{x})$ can be expressed as

$$\begin{aligned} K_n(\mathbf{z} = \mathbf{z}_k | \mathbf{x}) &= \sum_{\ell=1}^L K_n(\mathbf{z}_k | \mathbf{x}, k = \ell), \\ &= L \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1}^L \psi(\mathbf{z}_\ell) \right] \hat{\pi}_{MTM}^{(L)}(\mathbf{z}_k) \alpha_n(\mathbf{x}, \mathbf{z}_k | \mathbf{m}_{-k}) d\mathbf{m}_{-k}, \quad \text{for } \mathbf{z} \neq \mathbf{x}. \end{aligned} \quad (29)$$

where the function α_n is given in Eq. (11) and we have considered the case \mathbf{x} and \mathbf{z} (the case, $\mathbf{z} = \mathbf{x}$ is straightforward). The factor L is due of the exchangeability among the L random candidates. Thus, we can also write

$$\begin{aligned} \bar{\pi}(\mathbf{x}) K_n(\mathbf{z}_k | \mathbf{x}) &= L \bar{\pi}(\mathbf{x}) \psi(\mathbf{z}_k) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1; \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \beta_k \alpha_n(\mathbf{x}, \mathbf{z}_k | \mathbf{m}_{-k}) d\mathbf{m}_{-k}, \\ &= L \bar{\pi}(\mathbf{x}) \psi(\mathbf{z}_k) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1; \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \frac{\frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)}} \alpha_n(\mathbf{x}, \mathbf{z}_k | \mathbf{m}_{-k}) d\mathbf{m}_{-k}, \\ &= \frac{L}{Z} \pi(\mathbf{x}) \pi(\mathbf{z}_k) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1; \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \frac{1}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)}} \alpha_n(\mathbf{x}, \mathbf{z}_k | \mathbf{m}_{-k}) d\mathbf{m}_{-k}, \end{aligned} \quad (30)$$

where we have also used the equality $\bar{\pi}(\mathbf{x}) = \frac{1}{Z} \pi(\mathbf{x})$. Replacing

$$\alpha_n(\mathbf{x}, \mathbf{z}_k | \mathbf{m}_{-k}) = \min \left[1, \frac{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} - \frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)} + \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}} \right],$$

in the expression (30) and with some simple rearrangements, we obtain

$$\begin{aligned} \bar{\pi}(\mathbf{x}) K_n(\mathbf{z}_k | \mathbf{x}) &= \frac{L}{Z} \pi(\mathbf{x}) \pi(\mathbf{z}_k) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1; \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \\ &\min \left[\frac{1}{\sum_{\ell \neq k}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)}}, \frac{1}{\sum_{\ell \neq k}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}} \right] d\mathbf{m}_{-k}. \end{aligned} \quad (31)$$




Fig. 8. A graphical representation of the several parallel MTM schemes with $N = 2$ chains and $L = 3$ tries. The BI-MTM scheme in (d) requires only 6 evaluations of the target pdf and 2 multinomial sampling steps considering two iterations, $t - 1$ and t .

We can observe that, in equation above, we can exchange the position of the variables \mathbf{x} so that \mathbf{z}_k and the expression does not change. So that we can write

$$\bar{\pi}(\mathbf{x})K_n(\mathbf{z}_k|\mathbf{x}) = \bar{\pi}(\mathbf{z}_k)K_n(\mathbf{x}|\mathbf{z}_k), \quad (32)$$

for all $n = 1, \dots, N$. The expression above is the so-called *detailed balance condition* [5]: since it holds for all n , the complete horizontal MTM process has $\bar{\pi}_g$ as invariant pdf.

C.1.1. Important observations and Block Independent MTM

First of all, note that with respect to a standard parallel multiple try approach, the novel P-MTM scheme generates only L candidates at each iteration, instead of NL samples. Indeed, P-MTM “recycles” the samples $\mathbf{z}_1, \dots, \mathbf{z}_L$ from the independent proposal pdf $\psi(\mathbf{x})$, using them in all the N chains. Namely, in P-MTM, at one iteration, the different MTM chains share the same set of tries. However, looking a single chain, each time L new samples are drawn from $\psi(\mathbf{x})$ so that the chain is driven exactly from a standard (valid) MTM kernel. Figures 8(a) and (b) compare graphically the standard parallel MTM approach and to P-MTM scheme (with $N = 2$ chains and $L = 3$ tries). Observe that, in Figure 8(a), 12 new evaluations of the target are needed whereas only 6, in Figure 8(b).

Using the same arguments, the method remains valid if only one resampling step is performed at each iteration, providing one \mathbf{z}^* : in this case the same \mathbf{z}^* is tested in the different acceptance tests of the N parallel MTM chains, at the same iteration (exactly as in Table 2 and Fig. 3 for MH kernels). Figure 8(c) shows this case. In order to reduce the possible loss of the diversity, since several chains could jump at the same new state \mathbf{z}^* , an alternative strategy can be employed: the Block Independent MTM (BI-MTM) algorithm described in Table 6. Since the proposal ψ is independent and then fixed, before a block of N transitions, we can draw NL tries from $\psi(\mathbf{x})$. Then, we can divide them in N sets \mathcal{S}_j , with $j = 1, \dots, N$ and select one sample from each set, obtaining $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$ with $\mathbf{z}_{k_j} \in \mathcal{S}_j$. Then, we use N different permutations of $\{\mathbf{z}_{k_1}, \dots, \mathbf{z}_{k_N}\}$ for performing N iterations of the N parallel chains, providing a better mixing with respect to the case in Figure 8(c). This strategy, i.e., the BI-MTM scheme, is perfectly equivalent to the previous one, shown in Figure 8(c), from a theoretical and computational point of view. BI-MTM is represented graphically in Figure 8(d).

C.2. Parallel Ensemble MCMC

Let us consider now the method in Table 4. In this case, the particle approximation is

$$\begin{aligned}\hat{\pi}_n^{(L+1)}(\mathbf{z}) &= \sum_{\ell=1}^L \alpha_\ell \delta(\mathbf{z} - \mathbf{z}_\ell) + \alpha_{L+1} \delta(\mathbf{z} - \mathbf{x}_{n,t-1}) \\ &= \sum_{\ell=1}^{L+1} \alpha_\ell \delta(\mathbf{z} - \mathbf{z}_\ell), \quad \text{where } \mathbf{z}_{L+1} = \mathbf{x}_{n,t-1},\end{aligned}\tag{33}$$

In this case, for a given $n = 1, \dots, N$, the conditional probability $K_n(\mathbf{z} = \mathbf{z}_k | \mathbf{x})$, where $\mathbf{x} = \mathbf{x}_{n,t-1}$ and $\mathbf{z}_k \in \{\mathbf{z}_1, \dots, \mathbf{z}_L, \mathbf{z}_{L+1} = \mathbf{x}_{n,t-1}\}$, is given by

$$\begin{aligned}K_n(\mathbf{z}_k | \mathbf{x}) &= \sum_{\ell=1}^L K_n(\mathbf{z}_k | \mathbf{x}, k = \ell), \\ &= L \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1}^L \psi(\mathbf{z}_\ell) \right] \hat{\pi}_n^{(L+1)}(\mathbf{z}_k) d\mathbf{m}_{-k}, \quad \text{for } \mathbf{z} \neq \mathbf{x}.\end{aligned}\tag{34}$$

After some simple rearrangements (similarly in P-MTM) and using the formula of the weights in Eq. (8), we obtain

$$\begin{aligned}\bar{\pi}(\mathbf{x}) K_n(\mathbf{z}_k | \mathbf{x}) &= L \bar{\pi}(\mathbf{x}) \psi(\mathbf{z}_k) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1, \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \frac{\frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)}}{\sum_{\ell=1}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}} d\mathbf{m}_{-k}, \\ &= \frac{L}{Z} \pi(\mathbf{x} \pi(\mathbf{z}_k)) \int_{\mathcal{D}^{L-1}} \left[\prod_{\ell=1, \ell \neq k}^L \psi(\mathbf{z}_\ell) \right] \frac{1}{\sum_{\ell=1; \ell \neq k}^L \frac{\pi(\mathbf{z}_\ell)}{\psi(\mathbf{z}_\ell)} + \frac{\pi(\mathbf{z}_k)}{\psi(\mathbf{z}_k)} + \frac{\pi(\mathbf{x})}{\psi(\mathbf{x})}} d\mathbf{m}_{-k}.\end{aligned}\tag{35}$$

Observing the last equation, we can clearly replace the variable \mathbf{x} with \mathbf{z}_k and vice versa, without changing the expression. Hence, finally we obtain

$$\bar{\pi}(\mathbf{x}) K_n(\mathbf{z}_k | \mathbf{x}) = \bar{\pi}(\mathbf{z}_k) K_n(\mathbf{x} | \mathbf{z}_k),$$

for all $n = 1, \dots, N$, that is the detailed balance condition. For further considerations, see App. C.1.1 above.