

# A simple model of quantization: an approach from chaos

<sup>1</sup>Moisés Domínguez-Espinosa, <sup>1</sup>Jaime Meléndez-Martínez

<sup>1</sup>Facultad de Ciencias, Universidad Nacional Autónoma de México, Avenida Universidad 3000, Ciudad de México, 04510. E-mail: moi\_de@ciencias.unam.mx March 28, 2016

**Abstract.** There is a paradigm in Quantum Mechanics that explains quantization through normal vibration modes called *Eigenstates* that arise from Schrödinger wave equation. In this contribution we propose an alternative methodology of quantization by using basic concepts of mechanics and chaos from which a *Toy Model* is built.

## 1. Motivation

Let us assume that a pair of particles interact with quantum noise [3][11] such that they are perturbed in the form of *kicks* [1][10] and besides, these particles attract each another due to a central force. The Lagrangian that describes this phenomenon consists of one term associated to the central force acting on the total mass of the system and the *Ansatz* that models the complex interaction between the quantum noise and the particles:

$$\mathcal{L} = \frac{1}{2}(m\dot{r}^2 + mr^2\dot{\theta}^2) - V(r) - K\theta \sin(\Omega + \xi\dot{\theta}) \sum_{j=-\infty}^{\infty} \delta(t - jT). \quad (1)$$

Where  $j \in \mathbb{Z}$ ,  $-\pi \leq \Omega \leq \pi$  and,  $K, \xi$  are parameters that will be defined later on, and  $T$  is the perturbation period.

From Euler-Lagrange equations [4] we have

$$\frac{d}{dt}(m\dot{r}) - mr\dot{\theta}^2 + \frac{\partial}{\partial r}V(r) + \epsilon(K, \xi(m, r), T) = 0 \quad (2)$$

$$\frac{d}{dt}(mr^2\dot{\theta}) + K \sin(\Omega + \xi\dot{\theta}) \sum_{j=-\infty}^{\infty} \delta(t - jT) = 0 \quad (3)$$

with  $\epsilon(K, \xi(m, r), T) \ll \frac{\partial}{\partial r}V(r)$ .

Let us assume that  $r$  is a constant, thus from equation (3) we have

$$\ddot{\theta} = -\frac{K}{mr^2} \sin(\Omega + \xi\dot{\theta}) \sum_{j=-\infty}^{\infty} \delta(t - jT) \quad (4)$$

$$\dot{\theta}_{j+1} = \dot{\theta}_j - \frac{K}{mr^2} \sin(\Omega + \xi\dot{\theta}_j). \quad (5)$$

Taking the fixed points [5] in (5),  $\dot{\theta}_{j+1} = \dot{\theta}_j = \dot{\theta}^*$  then

$$\sin(\Omega + \xi\dot{\theta}^*) = 0.$$

Thus, the fixed points are

$$\dot{\theta}_n^* = \frac{n\pi - \Omega}{\xi} \quad n \in \mathbb{Z}. \quad (6)$$

In order to obtain the stable fixed points we take [5]

$$|f'(\dot{\theta}^*)| < 1 \quad (7)$$

where

$$f'(\dot{\theta}^*) = 1 - \frac{K\xi}{mr^2} \cos(\Omega + \xi\dot{\theta}^*). \quad (8)$$

Taking the  $n$  even stable fixed points in equation (6) we have

$$\dot{\theta}_n^* = \frac{2n\pi - \Omega}{\xi} \quad n \in \mathbb{Z} \quad (9)$$

and from (7), (8) and (9):

$$0 < \frac{K\xi}{mr^2} < 2. \quad (10)$$

Taking  $\frac{K\xi}{mr^2} = 2\pi\mu_0$  and  $0 < \mu_0 < \frac{1}{\pi}$  we can write equation (9) as:

$$\dot{\theta}_n^* = \frac{(2n\pi - \Omega)K}{2\pi\mu_0 mr^2} \quad n \in \mathbb{Z}. \quad (11)$$

Now, if  $\Omega = 0$  and  $K = \mu_0 H$  in equation (11) then

$$L_n = nH$$

## 2. Experimental consequences

An interesting consequence from the stability condition is that, if  $K = \mu_0 \hbar$  in (10) where  $\hbar$  is the Planck's constant, and taking  $r$  as in Bohr's model [7][8][9]  $r^2 = \frac{n^4 \hbar^4}{m^2 k^2 e^4}$  where  $k$  is the Coulomb constant, we obtain

$$0 < \frac{2\pi\mu_0 Rc}{n^4 \nu_\xi} < 1 \quad (12)$$

where  $R = 1.0972 \times 10^7 m^{-1}$  is the Rydberg's constant,  $c$  is the speed of light, which leads to  $Rc \sim 10^{15} Hz$  and  $\nu_\xi = \frac{1}{\xi}$ .

Table 1  $n$  vs  $\frac{2Rc}{n^4}$

$n$	$\frac{2Rc}{n^4} [Hz]$
1	$6.580 \times 10^{15}$
2	$4.112 \times 10^{14}$
3	$8.123 \times 10^{13}$
4	$2.570 \times 10^{13}$
5	$1.053 \times 10^{13}$
6	$5.077 \times 10^{12}$
7	$2.740 \times 10^{12}$
8	$1.606 \times 10^{12}$
9	$1.003 \times 10^{12}$
10	$6.580 \times 10^{11}$
11	$4.494 \times 10^{11}$
12	$3.173 \times 10^{11}$
13	$2.304 \times 10^{11}$
14	$1.713 \times 10^{11}$

If  $\xi(m, r)$  is a constant then  $\epsilon(K, \xi(m, r), T)$  must be zero.

## 3. Fixed points of $M$ period: an interesting observation

We can scale the system to a convenient scale, if in equation (5) we take  $\Omega = 0$ ,  $\xi = 2\pi [s]$ ,  $mr^2 = 1 [kg \cdot m^2]$  and  $H = 1 [J \cdot s]$  we have

$$\dot{\theta}_{j+1} = \dot{\theta}_j - K \sin(2\pi\dot{\theta}_j)$$

in this case  $K = \mu_0$ , and  $0 < K < \frac{1}{\pi}$ .

If we increase  $K$  beyond period one we obtain the following stable period *cascaes* [6] (see Figure 1). This diagram is also known as a bifurcation diagram.<sup>1</sup>




Figure 1  $\dot{\theta}$  vs  $K$

These periodic points follow an order established by Sharkovskii's theorem [2]:

$$3 < 5 < 7 < \dots < 2 \cdot 3 < 2 \cdot 5 < 2 \cdot 7 < \dots < 2^N \cdot 3 < 2^N \cdot 5 < 2^N \cdot 7 < \dots < 8 < 4 < 2 < 1 \quad N \in \mathbb{N}$$

<sup>1</sup> Bifurcation diagrams were first discovered by Robert May.

Let us consider the following subset

$$\dots < 2 \cdot 3 < 2 \cdot 5 < 2 \cdot 7 < \dots < 2 < 1 = \dots < 6 < 10 < 14 < \dots < 2 < 1$$

which coincidentally emerges in the range of energies of the different chemical elements (see Figures 2 and 3). Thus, to analytically calculate these periods is necessary to estimate the compositions for each corresponding period [5],  $f^M = x$ , i.e.:

$$f^6 = f \circ f \circ f \circ f \circ f \circ f$$

$$f^{10} = f \circ f$$

$$f^{14} = f \circ f$$

⋮

$$f^2 = f \circ f$$




Figure 2 Energy levels




Figure 3 Sharkovskii's order

for which would exist  $K_{s\dots fdp}$  or  $\mu_{s\dots fdp}$  to be determined.

These fixed points of higher periods may, for example, propagate through the energy levels according to the following modified equation from Bohr's model

$$E = \frac{mk^2 Z^2 e^4}{2\hbar^2 (n \pm \Omega_{s\dots fdp} \pm \zeta_{s\dots fdp})^2}$$

where  $\Omega_{s\dots fdp}$  and  $\zeta_{s\dots fdp}$  would be related to the fixed points of periods 2, ..., 14, 10, 6 possibly to the sub-level  $s, \dots, f, d, p$ .

#### 4. Discussion and Conclusions

Even so, it would be necessary to take into account relativistic effects, and extra dimension factors as well as the effects of the spin of the electron as is currently carried out by modern quantum mechanics to describe the fine-structure; however this represents a simplified model.

In this work we present a methodology based on basic concepts of stability to quantize the angular momentum. It was also shown that, in order for this model to be stable, it is sufficient condition that the model is immersed in noise with the same

frequency of that of the background radiation frequency and, that the order of the energy levels agree with the order given by Sharkovskii's theorem. Finally, we can apply this methodology to other scales.

### 5. Acknowledgments

Moisés Domínguez Espinosa would like to express his deepest appreciation to Dr. Luis de la Peña for his comments to improve the quality of this work. Also thanks UNAM, Facultad de Ciencias for the support received.

### References

- [1] H.G. Schuster & W. Just. *Deterministic Chaos An Introduction*. Wiley-VCH, Weinheim, 2004.
- [2] Ricard V. Solé & Susanna C. Manrubia. *Orden y Caos en sistemas complejos*. Fundamentos. Universitat Politècnica de Catalunya, Barcelona, 2001.
- [3] Ilya Prigogine. *¿Tan sólo una ilusión? Una exploración del orden al caos*. Tusquets. Barcelona, 2009.
- [4] Goldstein Herbert. *Classical Mechanics*. Pearson Education. California, 2002.
- [5] Strogatz Steven H. *Nonlinear Dynamics and Chaos*. Perseus Books. USA, 1994.
- [6] David Ruelle. *Casualidad y Caos*. UNAM, México, 2003.
- [7] Olsen D. James and McDonald Kirk T. *Classical Lifetime of a Bohr Atom*. Joseph Henry Laboratories, Princeton University, NJ, 2005.
- [8] Rohrlich F. The dynamics of a charged sphere and the electron. Department of Physics, Syracuse, New York. *Am. J. Phys.*, Vol. 65, No. 11, November 1997.
- [9] Paul A. Tipler. *Modern Physics*. W.H. Freeman. USA, 2008.
- [10] Kadanoff Leo P. *From Periodic Motion to Unbounded Chaos: Investigations of the Simple Pendulum*. *Physica Scripta*. University of Chicago 1984.
- [11] Luis de la Peña, Ana María Cetto, Andrea Valdés Hernández, *The Emerging Quantum*, Springer, 2015.