

Global in Time Solvability of Incompressive NSIVP in the Whole Space abstract

Global in time solvability of incompressive Navier-Stokes initial value problem in the whole space is proved using time transformation analysis.

Introduction

Navier-Stokes equations was derived as fundamental equations for hydromechanics by Navier^[1] and Stokes^[2]. Hereafter, its solvability is studied for a long time, but, due to the difficulty based on nonlinearity of the equations, even limited to incompressive fluid, not enough results for solvability have gotten over 170 years.

With regard to NSIVP for incompressible fluid, local in time solvability and global in time solvability for small initial value, including well-posedness, are known by means of arguments based on analysis started by Leray^[3], Hopf^[4] or Kiselev-Ladyzhenskaya^[5], Ito^[6], Kato-Fujita^[7], and enhanced by Kato^[8], Giga-Miyakawa^[9]. But, global in time solvability for initial value not only for small value has not been known.

In the present paper, global in time solvability in the whole space for initial value not only for small value is proved using time transformation analysis. Meanwhile, global in time solvability in periodic space for initial value not only for small value is proved as well using time transformation analysis, in another paper^[11].

This gives positive answer to the whole space version in the CMI millenium problem^[10] related to Navier-Stokes equations, and similar results for more comprehensive initial values.

Overview

As main result, NSIVP in the whole space \mathbf{R}^n , $n \geq 3$ has global in time classical solution, for initial value not only for small value. The global in time solution is well-posed, which means this is unique, smooth and continuous to initial value.

Also, there is a decreasing upper limit function of initial value for norm of solution.

The proof consists of local in time analysis, and global in time analysis based on a priori estimation. Analysis based on a priori estimation is typical global in time analysis. Basic analysis has application limit and due to this limit, global in time solvability of incompressive Navier-Stokes initial value problem has not been proved. In the present argument, to overcome this limit, time transformation is used. By this time transformation analysis, effective area with regard to norm upper limit estimation is expanded to overcome the limit.

With regard to local in time analysis, as the first step, for initial value $\mathbf{a} \in \mathbf{L}^Q, Q \in (n, \infty)$, the existence of existing time T_M and solution $\mathbf{u} \in \mathbf{L}^Q, t \in [0, T_M]$ is proved, and for initial value $\mathbf{a} \in \mathbf{L}^2 \cap \mathbf{L}^q, q \in (n, \infty]$, the existence of existing time T_M and solution $\mathbf{u} \in \mathbf{L}^2 \cap \mathbf{L}^q, t \in [0, T_M]$ is proved.

And next, a priori energy nonincreasing is proved, and $\mathbf{L}^r, r \in [2, \infty]$ -norm of solution and derivative of solution has nonincreasing or decreasing upper limit function. After that, based on local in time solvability and a priori estimation, global in time solvability is proved. Time transformation analysis are repeatedly used through these process not limited in the step of proving global in time solvability.

1. preliminary

Definition 1 (function space)

For time point $t \in [0, \infty)$, time interval $TI \subseteq [0, \infty]$ and Lebesgue space $\mathbf{L}^q(\mathbf{R}^n), q \in [1, \infty], n \geq 3$, functional space $\mathbf{L}_t^q(\mathbf{R}^n), \mathbf{L}_{TI}^q(\mathbf{R}^n)$ are defined as follows.

$$\|\varphi\|_{\mathbf{L}_t^q(\mathbf{R}^n)} = \|\varphi_t\|_{\mathbf{L}^q(\mathbf{R}^n)}, \mathbf{L}_t^q(\mathbf{R}^n) = \{\varphi \mid \|\varphi\|_{\mathbf{L}_t^q(\mathbf{R}^n)} < \infty\}$$

$$\|\varphi\|_{\mathbf{L}_{TI}^q(\mathbf{R}^n)} = \sup_{t \in TI} \|\varphi_t\|_{\mathbf{L}^q(\mathbf{R}^n)}, \quad \mathbf{L}_{TI}^q(\mathbf{R}^n) = \{\varphi \mid \|\varphi\|_{\mathbf{L}_{TI}^q(\mathbf{R}^n)} < \infty\}$$

Here, function over spatial space φ_t is function over time and space φ with fixed time t .

These functional space $\mathbf{L}_t^q(\mathbf{R}^n), \mathbf{L}_{TI}^q(\mathbf{R}^n)$ are Banach space.

Hereafter, these are noted simply by $\mathbf{L}_t^q, \mathbf{L}_{TI}^q$.

Moreover, function spaces $\mathbf{L}^{[p,q]}, \mathbf{L}^{(p,q)}$ are defined as follows, providing $1 \leq p < q \leq \infty$.

$$\mathbf{L}^{[p,q]} = \bigcap_{r \in [p,q]} \mathbf{L}^r, \quad \mathbf{L}^{(p,q)} = \bigcap_{r \in (p,q)} \mathbf{L}^r \quad \square$$

Difinition 2 (multiple index)

In the present paper, size of multiple index $\alpha = (\alpha_1, \dots, \alpha_n)_{\in (\{0\} \cup \mathbf{N})^n}$ is defined by $|\alpha| = \alpha_1 + \dots + \alpha_n$, and product of powers of spatial variable correspond to α is defined by $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$, derivative in regard as spatial variable correspond to α is defined by $\partial^\alpha = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n}$.

Difinition 3 (Helmholtz discomposition)

For a function φ , its nondivergent component $\mathcal{P}\varphi$ and nonrotational component $\overline{\mathcal{P}}\varphi$ are defined as follows if these exist, and Helmholtz decomposition means decomposition of φ to $\mathcal{P}\varphi, \overline{\mathcal{P}}\varphi$.

$$\varphi = \mathcal{P}\varphi + \overline{\mathcal{P}}\varphi, \quad \partial \cdot \mathcal{P}\varphi = 0, \quad (\mathcal{P}\varphi, \overline{\mathcal{P}}\varphi)_{L^2} = 0$$

If there is series of priodic functions, each of which has Helmholtz composition $\{\varphi_k\}_{k \in \mathbf{N}}, \varphi_k = \mathcal{P}\varphi_k + \overline{\mathcal{P}}\varphi_k$ and serieses of Helmholtz components $\{(\mathcal{P}\varphi_k, \overline{\mathcal{P}}\varphi_k)\}_{k \in \mathbf{N}}$ have limits $(\mathcal{P}\varphi, \overline{\mathcal{P}}\varphi)$, nondivergent component of φ is defined by $\mathcal{P}\varphi$ and nonrotational component of φ is defined by $\overline{\mathcal{P}}\varphi$, and Helmholtz decomposition means decomposition of φ to $\mathcal{P}\varphi, \overline{\mathcal{P}}\varphi$.

Generally for function space \mathbf{X} (like $\mathbf{L}^q, \mathbf{L}_{TI}^q$ above), function space $\mathcal{P}\mathbf{X}$ is defined as follows.

$$\mathcal{P}\mathbf{X} = \{\varphi \in \mathbf{X} \mid \varphi = \mathcal{P}\varphi\} \quad \square$$

Difinition 4 (heat kernel)

Heat kernel K corresponds to heat equation $\partial_t f - \nu \Delta f = 0$ is defined by following expression.

$$K(t, \mathbf{x}) = \frac{1}{\sqrt{4\pi\nu t}^n} \exp\left(-\frac{\mathbf{x}^2}{4\nu t}\right)$$

Difinition 5 (initial value)

Basic condition for initial value function \mathbf{a} is defined as follows, providing $q \in (n, \infty]; m \geq 2$.

$$(1.1) \quad \mathbf{a} \in \mathcal{P}\mathbf{L}^{[2,q]}, \quad \partial^\alpha \mathbf{a} \in \mathbf{L}^{[2,q]}, \quad |\alpha| \leq m$$

Difinition 6 (NSIVP)

NSIVP (Navier-Stokes initial value problem) is defined as follows.

$$(1.2) \quad \mathbf{u} \in \mathcal{P}\mathbf{L}_{(0,\infty)}^{[2,q]}$$

$$\partial_t \mathbf{u} - \nu \Delta \mathbf{u} + (\mathbf{u} \cdot \partial) \mathbf{u} + \frac{1}{\rho} \partial p = \mathbf{0}, \quad , (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

$$\mathbf{u}(0, \mathbf{x}) = \mathbf{a}(\mathbf{x}), \quad , \mathbf{x} \in \mathbf{R}^n$$

Following integral equation is equivalent to NSIVP(1.2), providing existence of derivatives of solution.

$$(1.3) \quad \mathbf{u}_t = K_t * \mathbf{a} - \int_0^t d\tau \mathcal{P}(\partial K_{t-\tau} * \mathbf{u}_\tau \mathbf{u}_\tau), \quad , (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

For funtion $\Phi = \Phi(t)$ with property $\Phi(0) = 0, \Phi(t) > 0, \partial_t \Phi(t) = \varphi(t)$ and function $\mathbf{f} = \mathbf{f}(t, \mathbf{x})$,

defining $\mathbf{f}^\Phi(t, \mathbf{x}) = \mathbf{f}(\Phi(t), \mathbf{x})$, time-transformed solution \mathbf{u}^Φ for original solution \mathbf{u} is defined. Then, following time-transformed-equation for \mathbf{u}^Φ is equivalent to original equation.

$$(1.4) \quad \mathbf{u}_t^\Phi = K_t^\Phi * \mathbf{a} - \int_0^t d\tau \mathcal{P}(\partial K_{t-\tau}^\Phi * \varphi_\tau \cdot \mathbf{u}_\tau^\Phi \mathbf{u}_\tau^\Phi) \quad , (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

Difinition 7 (linear term and nonlinear term of solution)

In case of the solution \mathbf{u} for NSIVP (1.2) is decomposed to 2 terms according to integral equation (1.3) or time transformed integral equation (1.4), term $\mathbf{u}^{(L)}$ is defined as linear term of the solution and term $\mathbf{u}^{(NL)}$ is defined as nonlinear term of the solution.

$$\begin{aligned} \mathbf{u}_t^{(L)} &= K_t * \mathbf{a} \quad , \quad \mathbf{u}_t^{(NL)} = - \int_0^t d\tau \mathcal{P}(\partial K_{t-\tau} * \cdot \mathbf{u}_\tau \mathbf{u}_\tau) \\ \mathbf{u}_t^{\Phi(L)} &= K_t^\Phi * \mathbf{a} \quad , \quad \mathbf{u}_t^{\Phi(NL)} = - \int_0^t d\tau \mathcal{P}(\partial K_{t-\tau}^\Phi * \varphi_\tau \cdot \mathbf{u}_\tau^\Phi \mathbf{u}_\tau^\Phi) \end{aligned} \quad \square$$

2.local in time solvability analysis

In this chapter, local in time solvability of NSIVPs is proved. First local in time solvability of integral equation (1.4) is proved, and second smoothness of its solution is proved, then local in time solvability of NSIVPs is proved. Using strict contracting map based fixed point theorem, local in time solvability of integral equation is proved.

proposition 1 (local in time solvability of integral equation)

There exist decreasing function $TM = TM(\xi)$ of positive number ξ , for arbitrary initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^q$, $q \in (n, \infty]$, integral equation (1.3) has a solution $\mathbf{u} \in \mathbf{L}_{[0, TM]}^q$, $TM = TM(\|\mathbf{a}\|_{\mathbf{L}^q})$. \square

proof

Map Ψ and set S_λ are defined as follows, for $T \in (0, \infty]$, $t \in [0, T]$ and $\lambda \in (0, \infty)$.

$$\begin{aligned} \Psi \mathbf{f}_t &= K_t * \mathbf{a} - \int_0^t d\tau \mathcal{P}(\partial K_{t-\tau} * \cdot \mathbf{f}_\tau \mathbf{f}_\tau) \\ S_\lambda &= \{ \mathbf{f} \in \mathbf{L}_{[0, T]}^q \mid \|\mathbf{f}\|_{\mathbf{L}_{[0, T]}^q} \leq \lambda \} \end{aligned}$$

Then following estimation relations are derived for $\mathbf{f}, \mathbf{g} \in S_\lambda$, providing $\chi(t) = \nu^{-1}(\nu t)^\beta$, $\beta = \frac{1}{2}(1 - \frac{n}{q})$.

$$\begin{aligned} \|\Psi \mathbf{f}\|_{\mathbf{L}_t^q} &\leq C_1 \|\mathbf{a}\|_{\mathbf{L}^q} + C_2 \chi(t) \|\mathbf{f}\|_{\mathbf{L}_{[0, T]}^q}^2 \\ \|\Psi \mathbf{f} - \Psi \mathbf{g}\|_{\mathbf{L}_t^q} &\leq C_2 \chi(t) (\|\mathbf{f}\|_{\mathbf{L}_{[0, T]}^q} + \|\mathbf{g}\|_{\mathbf{L}_{[0, T]}^q}) \|\mathbf{f} - \mathbf{g}\|_{\mathbf{L}_{[0, T]}^q} \end{aligned}$$

Constant $\Theta_{\in(0,1)}$ and $TM(\xi), T_M, \lambda$ are defined as follows.

$$\begin{aligned} TM(\xi) &= \Theta \frac{1}{\nu} \left(\frac{\nu}{4C_1 C_2 \xi} \right)^{\frac{1}{\beta}}, \quad T_M = TM(\|\mathbf{a}\|_{\mathbf{L}^q}) \\ \lambda &= \frac{1}{2C_2 \chi(T_M)} \left(1 - \sqrt{1 - 4C_1 C_2 \|\mathbf{a}\|_{\mathbf{L}^q} \chi(T_M)} \right) \end{aligned}$$

Then, following relations are derived.

$$\begin{aligned} \chi(T_M) &= \Theta^\beta (4C_1 C_2 \|\mathbf{a}\|_{\mathbf{L}^q})^{-1} \\ \lambda &\leq 2C_1 \|\mathbf{a}\|_{\mathbf{L}^q} \end{aligned}$$

Also, there exist constant $C_{\in(0,1)}$, and following relations are derived for each time $t \in [0, T_M]$.

$$\begin{aligned} C_1 \|\mathbf{a}\|_{\mathbf{L}^q} + C_2 \chi(t) \lambda^2 &< \lambda \\ 2C_2 \chi(t) \lambda &\leq C \end{aligned}$$

Therefore, based on estimation relations above, following relations are derived for $\mathbf{f}, \mathbf{g} \in S_\lambda$.

$$\Psi \mathbf{f} \in S_\lambda$$

$$\|\Psi \mathbf{f} - \Psi \mathbf{g}\|_{\mathbf{L}_{[0, T_M]}^q} \leq C \|\mathbf{f} - \mathbf{g}\|_{\mathbf{L}_{[0, T_M]}^q}$$

These mean that map Ψ gives strict contradicting map over set S_λ .

Therefore, by fixed point theorem, map Ψ has a fixed point \mathbf{u} in set S_λ .

This gives $\Psi \mathbf{u}_t = \mathbf{u}_t, t \in [0, T_M]$ and therefore \mathbf{u} is a solution of integral equation (1.3) over time interval $[0, T_M]$.

Moreover, based on aforesated arguement, following relations are derived, so $\mathbf{u} \in \mathbf{L}_{[0, T_M]}^q$.

$$\begin{aligned} \|\mathbf{u}\|_{\mathbf{L}_{[0, T_M]}^q} &\leq 2C_1 \|\mathbf{a}\|_{\mathbf{L}^q} \\ \|\mathbf{u}^{(NL)}\|_{\mathbf{L}_{[0, T_M]}^q} &\leq C_1 \|\mathbf{a}\|_{\mathbf{L}^q} \quad \square \end{aligned}$$

Note (estimation of upper limit for nonlinear term)

Estimation of upper limit of $\|\mathcal{P}(\partial K_{t-\tau} * \mathbf{f}_\tau \mathbf{f}_\tau)\|_{\mathbf{L}^q, q \in (n, \infty]}$ in the proof above is based on following relations.

$$\begin{aligned} \|\mathcal{P}_{ij} \partial_k K_{t-\tau} * f_{k\tau} f_{j\tau}\|_{L^q} &\leq \|\mathcal{P}_{ij} \partial_k K_{t-\tau}\|_{L^q} \|f_{k\tau}\|_{L^q} \|f_{j\tau}\|_{L^q} \\ \|\mathcal{P}_{ij} \partial_k K\|_{L^q} &\leq c_Q \left\| \left(1 - \frac{\xi_i \xi_j}{\xi^2}\right) \xi_k \hat{K} \right\|_{L^q} \\ &\leq 2c_Q \|\xi_k \hat{K}\|_{L^q} = C(\nu t)^{-\frac{n}{2} \frac{1}{q} - \frac{1}{2}}; 1 = \frac{1}{q} + \frac{1}{Q}, Q \in [1, 2] \end{aligned}$$

Here \mathcal{P}_{ij} s are components correspond to spatial coordinates of \mathcal{P} s, $\xi = (\xi_1, \dots, \xi_n)$ is n -dimensional Fourier variable, \hat{K} is Fourier transformed function of heat kernel K ; the former expression is confirmed by means of Young's inequality and the latter expression is confirmed by means of Hausdolf-Young inequality. \square

proposition 2 (regularity of local in time solution of integral equation)

Local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}_{[0, T_M]}^q$ of integral equation (1.3) based on proposition 1 for initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2, q], q \in (n, \infty]}$ has following characteristics (2.1-4) and regular, providing $|\alpha| \geq 1$ and, as for (2.2)(2.4), $q < \infty$.

$$\begin{aligned} (2.1) \quad \mathbf{u} &\in \mathbf{L}_{[0, T_M]}^r, r \in [2, q] \\ (2.2) \quad \mathbf{u} &\in \mathbf{L}_{(0, T_M]}^r, r \in (q, \infty) \quad ; \quad \sup_{t \in (0, T_M]} t^{\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{u}\|_{\mathbf{L}_t^r} < \infty \\ (2.3) \quad \partial^\alpha \mathbf{u} &\in \mathbf{L}_{(0, T_M]}^r, r \in [2, q] \quad ; \quad \sup_{t \in (0, T_M]} t^{\frac{|\alpha|}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} < \infty \\ (2.4) \quad \partial^\alpha \mathbf{u} &\in \mathbf{L}_{(0, T_M]}^r, r \in (q, \infty) \quad ; \quad \sup_{t \in (0, T_M]} t^{\frac{n}{2}(\frac{1}{q} - \frac{1}{r}) + \frac{|\alpha|}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} < \infty \end{aligned}$$

Moreover, local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}_{[0, T_M]}^q$ of integral equation (1.3) based on proposition 1 for initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2, q], q \in (n, \infty]}$, $\partial^\alpha \mathbf{a} \in \mathcal{P}\mathbf{L}^{[2, q], q \in (n, \infty]}, |\alpha| \leq m$ has following characteristics (2.5) adding to (2.1-4) above.

$$(2.5) \quad \partial^\alpha \mathbf{u} \in \mathbf{L}_{[0, T_M]}^r, r \in [2, q], |\alpha| \leq m \quad \square$$

proof

(1) First, characteristics (2.1) are proved as follows.

Following expression is confirmed for time $t \in [0, T_M]$, providing $r \in [2, q]$.

$$\|\mathbf{u}\|_{\mathbf{L}_t^r} \leq C_1 \|\mathbf{a}\|_{\mathbf{L}^r} + C_2 \|\mathbf{u}\|_{\mathbf{L}_{[0, T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2} \frac{1}{q} - \frac{1}{2}} \|\mathbf{u}\|_{\mathbf{L}_\tau^r}$$

Here, set A, X and operator \mathcal{K} as follows, providing $t \in (0, T_M]$.

$$\begin{aligned} A &= C_1 \|\mathbf{a}\|_{\mathbf{L}^r} \\ X_t &= \|\mathbf{u}\|_{\mathbf{L}_t^r} \end{aligned}$$

$$\mathcal{K}f_t = C_2 \|\mathbf{u}\|_{\mathbf{L}^q_{[0, T_M]}} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} f_\tau$$

Then, the expression above can be expressed by following relation.

$$X \leq A + \mathcal{K}X$$

Iterating use of this relation, following expression is confirmed for $k \in \mathbf{N}$.

$$X \leq \sum_{j=0}^k \mathcal{K}^j A + \mathcal{K}^{k+1} X$$

Then, for $j \in \mathbf{N}$ and $t \in [0, T_M]$, following expression is derived, providing $B(x, y)$ s are beta functions, $\Gamma(x)$ s are gamma functions and $\beta = \frac{1}{2} \left(1 - \frac{n}{q}\right)$.

$$\begin{aligned} \mathcal{K}^j A &\leq A (c_2 \|\mathbf{u}\|_{\mathbf{L}^q_{[0, T_M]}} \nu^{-1} (\nu t)^\beta)^j \prod_{k=0}^{j-1} B(\beta, 1 + k\beta) \\ &\leq A \frac{1}{\Gamma(1 + j\beta)} (C_2 \Gamma(\beta) \|\mathbf{a}\|_{\mathbf{L}^q} \nu^{-1} (\nu t)^\beta)^j \end{aligned}$$

$$\mathcal{K}^j X \leq \|\mathbf{u}\|_{\mathbf{L}^q_{[0, T_M]}} \frac{1}{\Gamma(1 + j\beta)} (C_2 \Gamma(\beta) \|\mathbf{a}\|_{\mathbf{L}^q} \nu^{-1} (\nu t)^\beta)^j$$

Therefore, limit of $\mathcal{K}^k X$ as k approaches infinity equals 0, and following relations are confirmed, provided $t \in [0, T_M]$.

$$X \leq U^{(r)}(\|\mathbf{a}\|_{\mathbf{L}^r}, t) < \infty$$

$$U^{(r)}(\|\mathbf{a}\|_{\mathbf{L}^r}, t) = C_1 \|\mathbf{a}\|_{\mathbf{L}^r} \sum_{j=0}^{\infty} \frac{1}{\Gamma(1 + j\beta)} (C_2 \Gamma(\beta) \|\mathbf{a}\|_{\mathbf{L}^q} \nu^{-1} (\nu t)^\beta)^j$$

This concludes following result.

$$\sup_{t \in [0, T_M]} \|\mathbf{u}\|_{\mathbf{L}^r_t} < \infty$$

This is what is to be proved.

(2) Next, charactricites (2.2) are proved as follows.

In case of $r \in (q, \infty]$, by setting $\Phi(t) = t^\varepsilon, \varepsilon \in (0, (\frac{n}{2}(\frac{2}{q} - \frac{1}{r}) + \frac{1}{2}) - 1)$, following characteristics is confirmed under condition $\Phi_t \in (0, T_M]$.

$$\begin{aligned} \|\mathbf{u}^\Phi\|_{\mathbf{L}^r_t} &\leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^q} + c_2 \|\mathbf{u}\|_{\mathbf{L}^q_{[0, T_M]}}^2 \int_0^t d\tau (\nu \Phi_{t-\tau})^{-\frac{n}{2}(\frac{2}{q} - \frac{1}{r}) - \frac{1}{2}} \varphi_\tau \\ &\leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^q} + C_2 \nu^{-1} (\nu \Phi_t)^{\frac{1}{2} - \frac{n}{2}(\frac{2}{q} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^q}^2 \end{aligned}$$

Therefore following expressions are confirmed.

$$\begin{aligned} \|\mathbf{u}\|_{\mathbf{L}^r_t} &\leq C_1 (\nu t)^{-\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^q} + C_2 \nu^{-1} (\nu t)^{\frac{1}{2} - \frac{n}{2}(\frac{2}{q} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^q}^2 \\ (\nu t)^{\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{u}\|_{\mathbf{L}^r_t} &\leq C_1 \|\mathbf{a}\|_{\mathbf{L}^q} + C_2 \nu^{-1} (\nu t)^{\frac{1}{2}(1 - \frac{n}{q})} \|\mathbf{a}\|_{\mathbf{L}^q}^2 \\ (\nu t)^{\frac{n}{2}(\frac{1}{q} - \frac{1}{r})} \|\mathbf{u}\|_{\mathbf{L}^r_t} &\leq U^{(q)}(\|\mathbf{a}\|_{\mathbf{L}^q}) \end{aligned}$$

This is what is to be proved.

(3) Next, charactricites (2.3) are proved as follows.

In case of $r \in [2, q]$, by setting $Q = \min\{q, 2r\}, \Phi(t) = t^\varepsilon, \varepsilon \in (0, (\frac{n}{2}(\frac{2}{Q} - \frac{1}{r}) + \frac{1}{2} + \frac{|\alpha|}{2}) - 1)$, following expression is confirmed under condition $\Phi_t \in (0, T_M]$.

$$\|\partial^\alpha \mathbf{u}^\Phi\|_{\mathbf{L}^r_t} \leq C_1 (\nu \Phi_t)^{-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^r} + C_2 \|\mathbf{u}\|_{\mathbf{L}^Q_{[0, T_M]}}^2 \int_0^t d\tau (\nu \Phi_{t-\tau})^{-\frac{n}{2}(\frac{2}{Q} - \frac{1}{r}) - \frac{1}{2} - \frac{|\alpha|}{2}} \varphi_\tau$$

$$\begin{aligned}
&= C_1(\nu\Phi_t)^{-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^r} + C_2\nu^{-1}(\nu\Phi_t)^{\frac{1}{2}-\frac{n}{2}(\frac{2}{Q}-\frac{1}{r})-\frac{|\alpha|}{2}} b_\varepsilon \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^Q}^2 \\
b_\varepsilon &= \varepsilon B\left(1 - \varepsilon\left(\frac{n}{2}\left(\frac{2}{Q} - \frac{1}{r}\right) + \frac{1}{2} + \frac{|\alpha|}{2}\right), \varepsilon\right) \\
\|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} &\leq C_1(\nu t)^{-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^r} + C_2\nu^{-1}(\nu t)^{\frac{1}{2}-\frac{n}{2}(\frac{2}{Q}-\frac{1}{r})-\frac{|\alpha|}{2}} b_\varepsilon \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^Q}^2 \\
(\nu t)^{\frac{|\alpha|}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} &\leq U^{(r,\alpha)}(\|\mathbf{a}\|_{\mathbf{L}^r})
\end{aligned}$$

This is what is to be proved.

(4) Next, charactricitics (2.4) are proved as follows.

In case of $r \in (q, \infty]$, by setting $\Phi(t) = t^\varepsilon, \varepsilon \in (0, (\frac{n}{2}(\frac{2}{q}-\frac{1}{r}) + \frac{1}{2} + \frac{|\alpha|}{2})^{-1})$, following expressin is confirmed under condition $\Phi_t \in (0, T_M]$.

$$\begin{aligned}
\|\partial^\alpha \mathbf{u}^\Phi\|_{\mathbf{L}_t^r} &\leq C_1(\nu\Phi_t)^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{r})-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^q} + c_2 \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q}^2 \int_0^t d\tau (\nu\Phi_{t-\tau})^{-\frac{n}{2}(\frac{2}{q}-\frac{1}{r})-\frac{1}{2}-\frac{|\alpha|}{2}} \varphi_\tau \\
&\leq C_1(\nu\Phi_t)^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{r})-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^q} + C_2\nu^{-1}(\nu\Phi_t)^{\frac{1}{2}-\frac{n}{2}(\frac{2}{q}-\frac{1}{r})-\frac{|\alpha|}{2}} b_\varepsilon \|\mathbf{a}\|_{\mathbf{L}^q}^2 \\
b_\varepsilon &= \varepsilon B\left(1 - \varepsilon\left(\frac{n}{2}\left(\frac{2}{q} - \frac{1}{r}\right) + \frac{1}{2} + \frac{|\alpha|}{2}\right), \varepsilon\right) \\
\|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} &\leq C_1(\nu t)^{-\frac{n}{2}(\frac{1}{q}-\frac{1}{r})-\frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^q} + C_2\nu^{-1}(\nu t)^{\frac{1}{2}-\frac{n}{2}(\frac{2}{q}-\frac{1}{r})-\frac{|\alpha|}{2}} b_\varepsilon \|\mathbf{a}\|_{\mathbf{L}^q}^2 \\
(\nu t)^{\frac{n}{2}(\frac{1}{q}-\frac{1}{r})+\frac{|\alpha|}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} &\leq U^{(q,\alpha)}(\|\mathbf{a}\|_{\mathbf{L}^q})
\end{aligned}$$

This is what is to be proved.

(5) At last, charactricitics (2.5) are proved as follows.

Here, (2.5) $_{|\alpha|\leq k}$ for $k = 1, \dots, m$ are confirmed by mathematical induction. In case of $k = 1$, for α s that satisfy $|\alpha| = 1$, following expression is confirmed.

$$\|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} \leq C_1 \|\partial^\alpha \mathbf{a}\|_{\mathbf{L}^r} + C_2 \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_\tau^r}$$

Here, set A, X and operator \mathcal{K} as follows for time $t \in (0, T_M]$.

$$A = C_1 \|\partial^\alpha \mathbf{a}\|_{\mathbf{L}^r}$$

$$X_t = \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r}$$

$$\mathcal{K}f_t = C_2 \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} f_\tau$$

Then, like arguement in (1), following results are confirmed for α s that satisfy $|\alpha| = 1$.

$$\sup_{t \in [0, T_M]} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} < \infty$$

Therefore (2.5) $_{|\alpha|=1}$ is confirmed.

In case of (2.5) $_{|\alpha|\leq k \leq m-1}$ is valid, for α s that satisfy $|\alpha| = k + 1$, following relations are confirmed, providing $\beta + \gamma = \alpha$; $|\beta|, |\gamma| \leq k$ for β, γ as regards to summation.

$$\begin{aligned}
\|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} &\leq C_1 \|\partial^\alpha \mathbf{a}\|_{\mathbf{L}^r} + C_2 \sum_{\beta, \gamma} c_{\beta, \gamma} \|\partial^\beta \mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} \|\partial^\gamma \mathbf{u}\|_{\mathbf{L}_\tau^r} \\
&\quad + C_2 \|\mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_\tau^r}
\end{aligned}$$

Here, A, X and operator \mathcal{K} are defined as follows for time $t \in (0, T_M]$.

$$A = C_1 \|\partial^\alpha \mathbf{a}\|_{\mathbf{L}^r} + C_2 \sum_{\beta, \gamma} c_{\beta, \gamma} \|\partial^\beta \mathbf{u}\|_{\mathbf{L}_{[0,T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2}\frac{1}{q}-\frac{1}{2}} \|\partial^\gamma \mathbf{u}\|_{\mathbf{L}_\tau^r}$$

$$X_t = \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r}$$

$$\mathcal{K}f_t = C_2 \|\mathbf{u}\|_{\mathbf{L}_{[0, T_M]}^q} \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2} \frac{1}{q} - \frac{1}{2}} f_\tau$$

Then, like argument in (1), following results are confirmed for α s that satisfy $|\alpha| = k+1$.

$$\sup_{t \in [0, T_M]} \|\partial^\alpha \mathbf{u}\|_{\mathbf{L}_t^r} < \infty$$

Therefore, (2.5) $_{|\alpha|=k+1}$ is confirmed.

These above are what are to be proved. □

Theorem 1(Local in time solvability of NSIVP)

NSIVP(1.2) has a local in time solution. □

proof

According to **proposition 2**, integral equation (1.3) has a local in time solution that can be partially differentiated arbitraly times. Then integral equation (1.3) is equivalent to following integral equation.

$$\mathbf{u}_t = K_t * \mathbf{a} - \int_0^t d\tau K_{t-\tau} * \mathcal{P}((\mathbf{u}_\tau \cdot \partial) \mathbf{u}_\tau)$$

Therefore, this integral equation has local in time solution which can be partially derivated arbitraly times. Then, this integral equation is equivalent to NSIVP(1.2). Therefore, NSIVP(1.2) has a local in time solution. □

3. A priori estimation

Here, a priori estimation for NSIVP are confirmed. This corresponds to energy nonincreasing.

proposition 3(energy nonincreasing)

Following a priori estimation is confirmed for local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}_{[0, T_M]}^{[2, q]}$ of NSIVP for initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2, q], q \in (n, \infty]}$.

$$(3.1) \quad \|\mathbf{u}\|_{\mathbf{L}_{[0, T_M]}^2} \leq \|\mathbf{a}\|_{\mathbf{L}^2} \quad \square$$

proof

Solution \mathbf{u} of NSIVP with initial value \mathbf{a} satisfies following equation, providing $t \in [0, T_M]$.

$$(\partial_t - \nu \Delta) \mathbf{u}^2 = -2\nu \partial \mathbf{u} \partial \mathbf{u} - \partial \left(\mathbf{u} \left(\mathbf{u}^2 + \frac{2}{\rho} p \right) \right), \quad \partial \mathbf{u} \partial \mathbf{u} = \sum_{i, j} \partial_i u_j \partial_i u_j$$

$$\mathbf{u}_t^2 = K_t * \mathbf{a}^2 - 2 \int_0^t d\tau K_{t-\tau} * \nu \partial \mathbf{u}_\tau \partial \mathbf{u}_\tau - \int_0^t d\tau \partial \left(K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right)$$

$$\leq K_t * \mathbf{a}^2 - \int_0^t d\tau \partial \left(K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right)$$

Therefore, following expression is confirmed for $k \in \{0\} \cup \mathbf{N}$ and $B_r = \{ \mathbf{x} \in \mathbf{R}^n \mid x \leq r \}$.

$$\int_k^{k+1} dr \int_{B_r} dV \mathbf{u}_t^2 \leq \int_k^{k+1} dr \int_{B_r} dV K_t * \mathbf{a}^2 - \int_0^t d\tau \int_k^{k+1} dr \int_{B_r} dV \partial \left(K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right)$$

Then, according to mean value theorem, for each $k \in \{0\} \cup \mathbf{N}$ there exist $r_k \in (k, k+1)$, and following expression is confirmed.

$$\lim_{k \rightarrow \infty} \int_k^{k+1} dr \int_{B_r} dV \mathbf{u}_t^2 = \lim_{k \rightarrow \infty} \int_{B_{r_k}} dV \mathbf{u}_t^2 = \|\mathbf{u}_t\|_2^2$$

Similarly, following equation is confirmed.

$$\lim_{k \rightarrow \infty} \int_k^{k+1} dr \int_{B_r} dV K_t * \mathbf{a}^2 = \|K_t * \mathbf{a}^2\|_1 \leq \|\mathbf{a}\|_2^2$$

On the other hand, following equation are confirmed, providing $\partial B_r = \{\mathbf{x} \in \mathbf{R}^n \mid x = r\}$, $\Delta B_k = \{\mathbf{x} \in \mathbf{R}^n \mid k \leq x \leq k+1\}$.

$$\begin{aligned} & \left| \int_k^{k+1} dr \int_{B_r} dV \partial \left(K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right) \right| = \left| \int_k^{k+1} dr \int_{\partial B_r} dS K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right| \\ & \leq \int_k^{k+1} dr \int_{\partial B_r} dS K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) = \int_{\Delta B_k} dV K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \\ & \sum_{k=0}^{\infty} \int_{\Delta B_k} dV K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) = \int_{\mathbf{R}^n} dV K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \\ & = \left\| K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right\|_1 \leq \left\| \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{2}{\rho} p_\tau \right) \right\|_1 \leq C \|\mathbf{u}_\tau\|_3^3 < \infty \end{aligned}$$

Here, following relation is used.

$$\|u p\|_1 \leq \|u\|_3 \|p\|_{\frac{3}{2}} = \|u\|_3 \|\rho \Delta^{-1} \partial \partial (u u)\|_{\frac{3}{2}} \leq C \|u\|_3^3$$

Therefore, following relations are confirmed.

$$\begin{aligned} & \lim_{k \rightarrow \infty} \int_{\Delta B_k} dV K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{1}{\rho} p_\tau \right) = 0 \\ & \lim_{k \rightarrow \infty} \int_k^{k+1} dr \int_{B_r} dV \partial \left(K_{t-\tau} * \mathbf{u}_\tau \left(\mathbf{u}_\tau^2 + \frac{1}{\rho} p_\tau \right) \right) = 0 \end{aligned}$$

Therefore, following relation is confirmed.

$$\|\mathbf{u}_t\|_2^2 \leq \|\mathbf{a}\|_2^2$$

This is what is to be proved. \square

proposition 4(estimation of upper limit of solution)

Following a priori estimation for $\mathbf{L}_t^r, r \in [2, \infty]$ norm of local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}_{[0, T_M]}^{[2, q]}$ of NSIVP (1.2) with initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2, q], q \in (n, \infty)}$ is confirmed.

$$\begin{aligned} (3.2) \quad & \|\mathbf{u}^{(L)}\|_{\mathbf{L}_t^r} \leq C_1^{(r)} (\nu t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^2} \\ & \|\mathbf{u}^{(NL)}\|_{\mathbf{L}_t^r} \leq C_2^{(r)} \nu^{-1} (\nu t)^{-\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2}} \|\mathbf{a}\|_{\mathbf{L}^2}^2 \\ (3.3) \quad & \|\partial^\alpha \mathbf{u}^{(L)}\|_{\mathbf{L}_t^r} \leq C_1^{(\alpha, r)} (\nu t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r}) - \frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^2} \\ & \|\partial^\alpha \mathbf{u}^{(NL)}\|_{\mathbf{L}_t^r} \leq C_2^{(\alpha, r)} \nu^{-1} (\nu t)^{-\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2} - \frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^2}^2 \end{aligned}$$

proof

By setting $\Phi_t = t^\varepsilon, \varepsilon \in (0, (\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2}) - 1)$ or $\Phi_t = t^\varepsilon, \varepsilon \in (0, (\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2} + \frac{|\alpha|}{2}) - 1)$, based on time transformed integral equation (1.4) and (3.1), following equations are confirmed.

$$\begin{aligned} & \|\mathbf{u}_t^{(L)\Phi}\|_{\mathbf{L}^r} \leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})} \|\mathbf{a}\|_{\mathbf{L}^2} \\ & \|\mathbf{u}_t^{(NL)\Phi}\|_{\mathbf{L}^r} \leq c_2 \int_0^t d\tau (\nu \Phi_{t-\tau})^{-\frac{n}{2}(1 - \frac{1}{r}) - \frac{1}{2}} \varphi_\tau \|\mathbf{a}\|_{\mathbf{L}^2}^2 = C_2 \nu^{-1} (\nu \Phi_t)^{-\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2}} \|\mathbf{a}\|_{\mathbf{L}^2}^2 \\ & \|\partial^\alpha \mathbf{u}_t^{(L)\Phi}\|_{\mathbf{L}^r} \leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r}) - \frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^2} \\ & \|\partial^\alpha \mathbf{u}_t^{(NL)\Phi}\|_{\mathbf{L}^r} \leq c_2 \int_0^t d\tau (\nu \Phi_{t-\tau})^{-\frac{n}{2}(1 - \frac{1}{r}) - \frac{1}{2} - \frac{|\alpha|}{2}} \varphi_\tau \|\mathbf{a}\|_{\mathbf{L}^2}^2 = C_2 \nu^{-1} (\nu \Phi_t)^{-\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2} - \frac{|\alpha|}{2}} \|\mathbf{a}\|_{\mathbf{L}^2}^2 \end{aligned}$$

This is what is to be confirmed. \square

4. analysis of global in time solvability

proposition 5 (global in time solvability of NSIVP)

In regard to NSIVP with initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2,q],q \in (n,\infty)}$, following results are confirmed.

(1) For a local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,q]_{[0,T]}}$, there exist expanded time $\Delta T = \Delta T(\|\mathbf{a}\|_{\mathbf{L}^2}, T)$ and expanded local in time solution $\mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,q]_{[0,T+\Delta T]}}$. Moreover, $\Delta T(\|\mathbf{a}\|_{\mathbf{L}^2}, T)$ is increasing function of T .

(2) There exists global in time solution \mathbf{u} . This satisfies $\mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,q]_{[0,\infty)}} \cap \mathcal{P}\mathbf{L}^{(q,\infty)_{(0,\infty)}}$, $\partial^\alpha \mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,\infty]_{(0,\infty)}}$. \square

proof

(1) is confirmed as follows.

Resetting the endpoint time of finite time interval of a solution $\mathbf{u}_{t,t \in (0,T)}, T \in (0,\infty)$ as initial time resetting the solution at endpoint time as initial value, and using solvability analysis (**proposition 1**), expanded solution is constructed. Based on a priori estimation of expanded solution at time $t = T : \|\mathbf{u}_T\|_{\mathbf{L}^q} \leq U_2^{(q)}(\|\mathbf{a}\|_{\mathbf{L}^2}, T)_{,q \in (n,\infty)}$ (**proposition 4**), for existence time ΔT of the expanded solution, $\Delta T \geq TM(U_2^{(q)}(\|\mathbf{a}\|_{\mathbf{L}^2}, T))$ is confirmed. Because $U_2^{(q)}(\|\mathbf{a}\|_{\mathbf{L}^2}, t)$ is a decreasing function of t and $TM(\xi)$ is decreasing function of ξ , $TM(U_2^{(q)}(\|\mathbf{a}\|_{\mathbf{L}^2}, T))$ is increasing function of T .

(2) is confirmed as follows.

Based on local in time solvability of NSIVP (**proposition 1**), solution of NSIVP in finite time interval is expandable ((1)above) and its expanded existing time increase to the length of existing time interval ((1)above), by iterating expansion of solution, for arbitrary length of interval, solution can be constructed. Therefore NSIVP is global in time solvable.

Moreover, by means of **proposition 2**, $\mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,q]_{[0,\infty)}} \cap \mathcal{P}\mathbf{L}^{(q,\infty)_{(0,\infty)}}$, $\partial^\alpha \mathbf{u} \in \mathcal{P}\mathbf{L}^{[2,\infty]_{(0,\infty)}}$ are confirmed. \square

proposition 6 (asymptotic attenuation characteristic and global in time boundedness of solution of NSIVP)

For global in time solution \mathbf{u} of NSIVP(1.2) with initial value $\mathbf{a} \in \mathcal{P}\mathbf{L}^{[2,q],q \in (n,\infty)}$, following results are confirmed.

(1) asymptotic attenuation characteristic

Solution \mathbf{u} and its partial derivatives in regard to spatial coordinates have asymptotic attenuation characteristic for $t \rightarrow \infty$ as follows.

$$(4.1) \quad \begin{aligned} \|\mathbf{u}^{(L)}\|_{\mathbf{L}_t^r} &= O(t^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{r})}) \\ \|\mathbf{u}^{(NL)}\|_{\mathbf{L}_t^r} &= O(t^{-\frac{n}{2}(1-\frac{1}{r})+\frac{1}{2}}) \end{aligned}$$

$$(4.2) \quad \begin{aligned} \|\partial^\alpha \mathbf{u}^{(L)}\|_{\mathbf{L}_t^r} &= O(t^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{r})-\frac{|\alpha|}{2}}) \\ \|\partial^\alpha \mathbf{u}^{(NL)}\|_{\mathbf{L}_t^r} &= O(t^{-\frac{n}{2}(1-\frac{1}{r})+\frac{1}{2}-\frac{|\alpha|}{2}}) \end{aligned}$$

(2) global in time boundedness

Solution \mathbf{u} is global in time bounded as follows.

$$(4.3) \quad \|\mathbf{u}\|_{\mathbf{L}_{[0,\infty)}^r} < \infty, r \in [2,q] \quad \square$$

proof

(1) asymptotic attenuation characteristic

Based on **proposition 5** and, (3.2) in regard to (4.1), (3.3) in regard to (4.2), these are

confirmed.

(2) global in time boundedness

Based on **proposition 1**, $\|\mathbf{u}^{(NL)}\|_{\mathbf{L}^q} \leq C\|\mathbf{a}\|_{\mathbf{L}^q, t \in [0, T_M]}$ is confirmed, and based on **proposition 3** $\|\mathbf{u}^{(NL)}\|_{\mathbf{L}^2} = \|\mathbf{u} - \mathbf{u}^{(L)}\|_{\mathbf{L}^2}$ is confirmed; and interpolating these, following relation is confirmed.

$$\|\mathbf{u}^{(NL)}\|_{\mathbf{L}^r} \leq C_1^{(r)} \|\mathbf{a}\|_{\mathbf{L}^q}^{\theta_q} \|\mathbf{a}\|_{\mathbf{L}^2}^{\theta_2}, \theta_q = \frac{\frac{1}{2} - \frac{1}{r}}{\frac{1}{2} - \frac{1}{q}}, \theta_2 = \frac{\frac{1}{r} - \frac{1}{q}}{\frac{1}{2} - \frac{1}{q}}, t \in [0, T_M], r \in [2, q]$$

On the other hand, following relation is confirmed based on **proposition 5** and **proposition 4**.

$$\|\mathbf{u}^{(NL)}\|_{\mathbf{L}^r} \leq c_2^{(r)} \nu^{-1} (\nu t)^{-\gamma} \|\mathbf{a}\|_{\mathbf{L}^2}^2 \leq c_2^{(r)} \nu^{-1} (\nu T_M)^{-\gamma} \|\mathbf{a}\|_{\mathbf{L}^2}^2, t \in [T_M, \infty), r \in [2, q]$$

Therefore, with expression of T_M in the proof of **proposition 1**, following relation that gives (4.3) is confirmed.

$$\begin{aligned} \|\mathbf{u}^{(NL)}\|_{\mathbf{L}^r} &\leq \max\{C_1^{(r)} \|\mathbf{a}\|_{\mathbf{L}^q}^{\theta_q} \|\mathbf{a}\|_{\mathbf{L}^2}^{\theta_2}, c_2^{(r)} \nu^{-1} (\nu T_M)^{-\gamma} \|\mathbf{a}\|_{\mathbf{L}^2}^2\} \\ &\leq \max\{C_1^{(r)} \|\mathbf{a}\|_{\mathbf{L}^q}^{\theta_q} \|\mathbf{a}\|_{\mathbf{L}^2}^{\theta_2}, C_2^{(r)} \nu^{-1-\gamma\beta} \|\mathbf{a}\|_{\mathbf{L}^q}^{\gamma\beta} \|\mathbf{a}\|_{\mathbf{L}^2}^2\}, \gamma\beta = \frac{\gamma}{\beta} \end{aligned}$$

proposition 7 (equable continuous on initial value and uniqueness of solution of NSIVP)

Global in time solution of NSIVP(1.2) correspond to initial value $\mathbf{a}_{\in \mathcal{P}\mathbf{L}^{[2, q]}}$ depends equably and continuously on initial value \mathbf{a} , moreover and is unique. \square

proof

For 2 solutions $\mathbf{u}^{(1)}, \mathbf{u}^{(2)}$ of NSIVP which correspond to 2 initial values $\mathbf{a}^{(1)}, \mathbf{a}^{(2)} \in \mathcal{P}\mathbf{L}^{[2, q]}$ respectively, following expressions are confirmed, providing $r \in [2, q]$ and $t \in [0, T_M]$.

$$\begin{aligned} \mathbf{u}_t^{(1)} - \mathbf{u}_t^{(2)} &= K_t * (\mathbf{a}^{(1)} - \mathbf{a}^{(2)}) - \int_0^t d\tau \mathcal{P} \partial K_{t-\tau} * (\cdot \mathbf{u}_\tau^{(1)} \mathbf{u}_\tau^{(1)} - \cdot \mathbf{u}_\tau^{(2)} \mathbf{u}_\tau^{(2)}) \\ \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} &\leq C_1 \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^r} \\ &\quad + C_2 (\|\mathbf{u}^{(1)}\|_{\mathbf{L}_{[0, T_M]}^q} + \|\mathbf{u}^{(2)}\|_{\mathbf{L}_{[0, T_M]}^q}) \int_0^t d\tau (\nu(t-\tau))^{-\frac{n}{2} \frac{1}{q} - \frac{1}{2}} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_\tau^r} \end{aligned}$$

Therefore, similarly as in proof of **proposition 2** (1), following expressions are confirmed, providing $t \in [0, T_M]$.

$$\begin{aligned} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} &\leq C_1 \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^r} U^{(r)}(t) \\ \sup_{t \in [0, T_M]} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} &\leq C_1 \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^r} U^{(r)}(T_M) \\ U^{(r)}(t) &= \sum_{j=0}^{\infty} \frac{1}{\Gamma(1+j\beta)} (C_2 \Gamma(\beta) (\|\mathbf{u}^{(1)}\|_{\mathbf{L}_{[0, T_M]}^q} + \|\mathbf{u}^{(2)}\|_{\mathbf{L}_{[0, T_M]}^q}) \nu^{-1} (\nu t)^\beta)^j \end{aligned}$$

On the other hand, similarly as in proof of **proposition 4**, following expressions are confirmed, providing $r \in [2, q]$ and $t > 0$.

$$\begin{aligned} \|\mathbf{u}_t^{(1)\Phi} - \mathbf{u}_t^{(2)\Phi}\|_{\mathbf{L}^r} &\leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})} \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2} \\ &\quad + c_2 (\|\mathbf{a}^{(1)}\|_{\mathbf{L}^2} + \|\mathbf{a}^{(2)}\|_{\mathbf{L}^2}) \int_0^t d\tau (\nu \Phi_{t-\tau})^{-\frac{n}{2}(1 - \frac{1}{r}) - \frac{1}{2}} \varphi_\tau \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2} \\ &\leq C_1 (\nu \Phi_t)^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})} \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2} \\ &\quad + C_2 \nu^{-1} (\nu \Phi_t)^{-\frac{n}{2}(1 - \frac{1}{r}) + \frac{1}{2}} (\|\mathbf{a}^{(1)}\|_{\mathbf{L}^2} + \|\mathbf{a}^{(2)}\|_{\mathbf{L}^2}) \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2} \\ \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} &\leq \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2} U_2^{(r)}(t) \end{aligned}$$

Therefore, following expression is confirmed, providing $t \in [T_M, \infty)$.

$$\sup_{t \in [T_M, \infty)} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} \leq \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^2 U_2^{(r)}}(T_M)$$

Therefore, following expression is confirmed.

$$\sup_{t \in [0, \infty)} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} \leq \|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^r} \max\{U^{(r)}(T_M), U_2^{(r)}(T_M)\}$$

Therefore providing $\|\mathbf{a}^{(1)} - \mathbf{a}^{(2)}\|_{\mathbf{L}^r} \rightarrow 0$, $\sup_{t \in [0, \infty)} \|\mathbf{u}^{(1)} - \mathbf{u}^{(2)}\|_{\mathbf{L}_t^r} \rightarrow 0$ is confirmed. Namely,

solutions equably and continuously depends on initial values.

By setting $\mathbf{a}^{(1)} = \mathbf{a}^{(2)}$ in arguement above, $\mathbf{u}^{(1)} = \mathbf{u}^{(2)}$ is confirmed. Namely, solution is unique.[]

5. problem with constant value at point of infinity

In previous sections, problems with solution \mathbf{u} with $\mathbf{0}$ value at point of infinity as of following view are analized.

$$\|\mathbf{u}\|_{\mathbf{L}^q(\mathbf{R}^n)}^q = \sum_{k=0}^{\infty} \int_{|\mathbf{x}| \in [k, k+1]} dV(\mathbf{x}) |\mathbf{u}(\mathbf{x})|^q < \infty, q \in [2, \infty), t \in (0, \infty)$$

In this section, problem with solution $\mathbf{u}^\#$ with constant value $\mathbf{c} \in \mathbf{R}^n$ at point of infinity as of following view is analized.

$$\|\mathbf{u}^\# - \mathbf{c}\|_{\mathbf{L}^q(\mathbf{R}^n)}^q = \sum_{k=0}^{\infty} \int_{|\mathbf{x}| \in [k, k+1]} dV(\mathbf{x}) |\mathbf{u}^\#(\mathbf{x}) - \mathbf{c}|^q < \infty, q \in [2, \infty), t \in (0, \infty)$$

Namely, initial value problem of which initial value $\mathbf{a}^\#$ and solution $\mathbf{u}^\#$ satisfy following conditions(5.1)(5.2) will be analized.

$$(5.1) \quad \mathbf{a}^\# \in \mathbf{c} + \mathcal{P}\mathbf{L}^{[2, q]}, \partial^\alpha \mathbf{a}^\# \in \mathbf{L}^{[2, q]}, |\alpha| = 1, \dots, m$$

$$(5.2) \quad \mathbf{u}^\# \in \mathbf{c} + \mathcal{P}\mathbf{L}_{(0, \infty)}^{[2, q]}$$

$$\partial_t \mathbf{u}^\# - \nu \Delta \mathbf{u}^\# + (\mathbf{u}^\# \cdot \partial) \mathbf{u}^\# + \frac{1}{\rho} \partial p = \mathbf{0}, (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

$$\mathbf{u}^\#(0, \mathbf{x}) = \mathbf{a}^\#(\mathbf{x}), \mathbf{x} \in \mathbf{R}^n$$

With regard to existance of global in time solution and its characteristic of this problem, argument regards to problem which is already analized depends on transformation of equation is effective. Namely, by setting $\tilde{\mathbf{u}} = \mathbf{u}^\# - \mathbf{c}$, $\tilde{\mathbf{a}} = \mathbf{a}^\# - \mathbf{c}$, this problem is transformed to initial value problem of which initial value $\tilde{\mathbf{a}}$ and solution $\tilde{\mathbf{u}}$ satisfy following conditions (5.1)~(5.2)~.

$$(5.1) \quad \tilde{\mathbf{a}} \in \mathcal{P}\mathbf{L}^{[2, q]}, \partial^\alpha \tilde{\mathbf{a}} \in \mathbf{L}^{[2, q]}, |\alpha| \leq m$$

$$(5.2) \quad \tilde{\mathbf{u}} \in \mathcal{P}\mathbf{L}_{(0, \infty)}^{[2, q]}$$

$$\partial_t \tilde{\mathbf{u}} - \nu \Delta \tilde{\mathbf{u}} + (\mathbf{c} \cdot \partial) \tilde{\mathbf{u}} + (\tilde{\mathbf{u}} \cdot \partial) \tilde{\mathbf{u}} + \frac{1}{\rho} \partial p = \mathbf{0}, (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

$$\tilde{\mathbf{u}}(0, \mathbf{x}) = \tilde{\mathbf{a}}(\mathbf{x}), \mathbf{x} \in \mathbf{R}^n$$

These equations above differ from NS equations at point that these have linear term $(\mathbf{c} \cdot \partial) \tilde{\mathbf{u}}$. Correspond to this diffrent point, following kernel $\tilde{K}^{(\nu)}$ that correspond to linear equation $\partial_t f - \nu \Delta f + (\mathbf{c} \cdot \partial) f = 0$ over whole space \mathbf{R}^n , instead of heat kernel $K^{(\nu)}$, will be used.

$$\tilde{K}^{(\nu)}(t, \mathbf{x}) = \frac{1}{\sqrt{4\pi\nu t^n}} \exp\left(-\frac{(\mathbf{x} - \mathbf{c}t)^2}{4\nu t}\right)$$

In case of there exist derivatives of solutions, following equation (5.3) is equivarent to initial value problem (5.1)~(5.2)~.

$$(5.3) \quad \tilde{\mathbf{u}}_t = \tilde{K}_t * \mathbf{a} - \int_0^t d\tau \mathcal{P}(\partial \tilde{K}_{t-\tau} * \tilde{\mathbf{u}}_\tau \tilde{\mathbf{u}}_\tau) \quad , (t, \mathbf{x}) \in (0, \infty) \times \mathbf{R}^n$$

Kernel \tilde{K} has similar characteristic as heat kernel K , especially expressions $\|\tilde{K}\|_q = \|K\|_q, \|\partial^\alpha \tilde{K}\|_q = \|\partial^\alpha K\|_q$ are confirmed. Based on these expressions, almost similar results in previous section are confirmed and it is confirmed that this problem has global in time solution with following form.

$$\mathbf{u}^\# = \mathbf{c} + \tilde{\mathbf{u}}^{(L)} + \tilde{\mathbf{u}}^{(NL)}$$

On the other hand, solution \mathbf{u} of problem in previous sections and solution $\mathbf{u}^\#$ of problem in this section have following relation.

$$(5.4) \quad \mathbf{u}^\#(t, \mathbf{x}) = \mathbf{c} + \mathbf{u}(t, \mathbf{x} - \mathbf{c}t)$$

This is proved as follows.

$$\begin{aligned} (\partial_t - \nu \Delta) \mathbf{u}^\#(t, \mathbf{x}) &= (\partial_t - \nu \Delta) \mathbf{u}(t, \mathbf{x} - \mathbf{c}t) - (\mathbf{c} \cdot \partial) \mathbf{u}(t, \mathbf{x} - \mathbf{c}t) \\ &= -((\mathbf{u}(t, \mathbf{x} - \mathbf{c}t) + \mathbf{c}) \cdot \partial) \mathbf{u}(t, \mathbf{x} - \mathbf{c}t) - \frac{1}{\rho} \partial p \\ &= -(\mathbf{u}^\#(t, \mathbf{x}) \cdot \partial) \mathbf{u}^\#(t, \mathbf{x}) - \frac{1}{\rho} \partial p \end{aligned}$$

Using relation (5.4), based on existence of solution of problem in previous sections, existence of solution of problem in this section is automatically confirmed.

note (relation to CMI problem)

Relation between results of this paper and CMI (Clay Mathematical Institute) problem^[8] is as follows.

In the CMI problem, 4 candidate propositions (A)(B)(C)(D) are given, 2 of which (A)(C) correspond to initial value problem over 3 dimensional whole space \mathbf{R}^3 . In these 2 propositions (A)(C), regard to initial value, conditions that are consisted from (1) nondivergence, and (2) existence of arbitrary times derivatives on spatial variables and its spatial decreasing.

$$|\partial^\alpha \mathbf{a}| \leq C_{\alpha, K} (1 + x)^{-K} \quad , \mathbf{x} \in \mathbf{R}^3, \alpha \in (\{0\} \cup \mathbf{N})^3, K \geq 0$$

These 2 conditions are sufficient condition for initial value condition $\mathbf{a} \in \mathcal{P}\mathbf{L}^2 \cap \mathbf{L}^\infty$ in the present paper, therefore, results of the present paper conclude for the CMI problem, viz. proposition (A) is confirmed and proposition (C) is denied.

Results of the present paper give not only existence of global in time solutions, but also various characteristic regard to solutions like uniqueness, partial derivatibility, equable continuous on initial value, etc. Also results of the present paper weaken conditions of propositions from these 2 propositions (A)(B), and give similar results for comprehensive initial values.

On the other hand, results related to 2 propositions (B)(D) are given in other paper^[9], which are like as results related to 2 propositions (A)(C). Although, as for uniqueness, attention should be required.

bibliography

- [1] C.L.M.H.Navier, Memoiré sur les lois du mouvement des fluides, *Mémoires Acad. Roy. Sci. Inst. France* **6**(1823), 389-440
- [2] G.G.Stokes, On the Theories of the Internal Friction of Fluids in Motion, and of the Equilibrium and Motion of Elastic Solids, *Trans. Camb. Phil. Soc.*, **8**(1845), 287-319
- [3] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Mathematica*. **63**(1934), 193-248

- [4] E. Hopf, Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen, *Math. Nachr* **4** (1950), 213-231
- [5] A. A. Kiselev and O.A. Ladyzenskaya, On the existence and uniqueness of the solution of the nonstationary problem for a viscous incompressible fluid, *Izv. Akad. Nauk SSSR, Ser. Mat.*, **21**(1957), 655-680
- [6] S. Ito, The existence and the uniqueness of regular solution of non-stationary Navier-Stokes equation, *J. Fac. Sci. Univ. Tokyo Sect. I* **9** (1961), 103-140
- [7] T. Kato and H. Fujita, On the nonstationary Navier-Stokes system *Rend. Sem. Mat. Univ. Padova* **9** (1962), 243-260
- [8] T. Kato, Strong L^p -solution of the Navier-Stokes equation in \mathbf{R}^m , with applications to weak solutions in \mathbf{R}^m , with applications to weak solutions, *Math. Z.* **187** (1984), 471-480
- [9] Y. Giga and T. Miyakawa, Solution in L_r of the Navier-Stokes initial value problem, *Arch. Ration. Mech. Anal.* **89** (1985), 267-281
- [10] C. Fefferman, Existence & Smoothness of the Navier-Stokes Equations, Millennium Prize Problems, Clay Mathematical Institute, 2000,
<http://www.claymath.org/prizeproblems/navierstokes.htm>.
- [11] N.Isobe, Global in Time Solvability of Incompressive NSIVP in Periodic Space,-