

Blind Quantum Computing

For the first time, physicists have demonstrated that clients who possess only classical computers—and no quantum devices—can outsource computing tasks to quantum servers that perform blind quantum computing. [15]

Experiments at Space Scale project, which involves making use of the Micius satellite—the first sent aloft to conduct quantum networking experiments. [14]

Just two weeks ago, China demonstrated its prowess in the field of quantum technology by becoming the first to teleport information from Earth to a satellite in space using the simple mechanics of quantum entanglement. [13]

The researchers showed that the combination of these two properties can be used to transfer an encoded digital signal without information loss, which has potential applications for realizing highly efficient optical communication systems. [12]

Physicists from the University of Würzburg have designed a light source that emits photon pairs, which are particularly well suited for tap-proof data encryption. The experiment's key ingredients: a semiconductor crystal and some sticky tape. [11]

Quantum cryptography involves two parties sharing a secret key that is created using the states of quantum particles such as photons. The communicating parties can then exchange messages by conventional means, in principle with complete security, by encrypting them using the secret key. Any eavesdropper trying to intercept the key automatically reveals their presence by destroying the quantum states. [10]

Optical photons would be ideal carriers to transfer quantum information over large distances. Researchers envisage a network where information is processed in certain nodes and transferred between them via photons. [9]

While physicists are continually looking for ways to unify the theory of relativity, which describes large-scale phenomena, with quantum theory, which describes small-scale phenomena, computer scientists are searching for technologies to build the quantum computer using Quantum Information.

In August 2013, the achievement of "fully deterministic" quantum teleportation, using a hybrid technique, was reported. On 29 May 2014, scientists announced a reliable way of transferring data by quantum teleportation. Quantum teleportation of data had been done before but with highly unreliable methods.

The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the Wave-Particle Duality and the electron's spin also, building the Bridge between the Classical and Quantum Theories.

The Planck Distribution Law of the electromagnetic oscillators explains the electron/proton mass rate and the Weak and Strong Interactions by the diffraction patterns. The Weak Interaction changes the diffraction patterns by moving the electric charge from one side to the other side of the diffraction pattern, which violates the CP and Time reversal symmetry.

The diffraction patterns and the locality of the self-maintaining electromagnetic potential explains also the Quantum Entanglement, giving it as a natural part of the Relativistic Quantum Theory and making possible to build the Quantum Computer with the help of Quantum Information.

Contents

Preface.....	3
Blind quantum computing for everyone	4
Chinese team sends quantum keys to ground stations and teleports ground to satellite signals	5
China will launch an "unhackable" quantum communication network in its Jinan city next month	6
Optical lens can transfer digital information without loss	7
Novel light sources made of 2-D materials.....	8
Experiments started with sticky tape.....	8
Monolayers enable novel lasers.....	9
How to Win at Bridge Using Quantum Physics	9
Quantum Information	9
Heralded Qubit Transfer.....	10
Quantum Teleportation	11
Quantum cryptography set for lift-off	11
Losses and curvature	11
Polarization is preserved.....	12
Rotation on the fly	12
Quantum Computing	13
Quantum Entanglement	13
The Bridge	13
Accelerating charges	14
Relativistic effect	14

Heisenberg Uncertainty Relation	14
Wave – Particle Duality	14
Atomic model	14
The Relativistic Bridge	15
The weak interaction	15
The General Weak Interaction	16
Fermions and Bosons	16
Van Der Waals force	17
Electromagnetic inertia and mass	17
Electromagnetic Induction	17
Relativistic change of mass.....	17
The frequency dependence of mass	17
Electron – Proton mass rate	17
Gravity from the point of view of quantum physics	18
The Gravitational force	18
The Higgs boson	18
Higgs mechanism and Quantum Gravity	19
What is the Spin?	19
The Graviton	19
Conclusions	20
References	20

Author: George Rajna

Preface

While physicists are continually looking for ways to unify the theory of relativity, which describes large-scale phenomena, with quantum theory, which describes small-scale phenomena, computer scientists are searching for technologies to build the quantum computer.

Australian engineers detect in real-time the quantum spin properties of a pair of atoms inside a silicon chip, and disclose new method to perform quantum logic operations between two atoms. [5]

Quantum entanglement is a physical phenomenon that occurs when pairs or groups of particles are generated or interact in ways such that the quantum state of each particle cannot be described independently – instead, a quantum state may be given for the system as a whole. [4]

I think that we have a simple bridge between the classical and quantum mechanics by understanding the Heisenberg Uncertainty Relations. It makes clear that the particles are not point like but have a

dx and dp uncertainty.

Blind quantum computing for everyone

For the first time, physicists have demonstrated that clients who possess only classical computers—and no quantum devices—can outsource computing tasks to quantum servers that perform blind quantum computing. "Blind" means the quantum servers do not have full information about the tasks they are computing, which ensures that the clients' computing tasks are kept secure. Until now, all blind quantum computing demonstrations have required that clients have their own quantum devices in order to delegate tasks for blind quantum computing.

The team of physicists, led by Jian-Wei Pan and Chao-Yang Lu at the University of Science and Technology of China, have published a paper on the demonstration of blind quantum computing for classical clients in a recent issue of Physical Review Letters.

"We have demonstrated for the first time that a fully classical client can delegate a quantum computation to untrusted quantum servers while maintaining full privacy," Lu told Phys.org.

The idea behind blind quantum computing is that, while there are certain computing tasks that quantum computers can perform exponentially better than classical computers, quantum computing still involves expensive, complex hardware that will make it inaccessible for most clients. So instead of everyone owning their own quantum computing devices, blind quantum computing makes it possible for clients to outsource their computing tasks to quantum servers that do the job for them. Ensuring that the quantum computing is performed blindly is important, since many of the potential applications of quantum computing will likely require a high degree of security.

Although several blind quantum computing protocols have been performed in the past few years, they have all required that the clients have the ability to perform certain quantum tasks, such as prepare or measure qubit states. Eliminating this requirement will provide greater access to blind quantum computing, since most clients only have classical computing systems.

In the new study, the physicists experimentally demonstrated that a classical client can outsource a simple problem (factoring the number 15) to two quantum servers that do not fully know what problem they are solving. This is because each server completes part of the task, and it is physically impossible for the servers to communicate with each other. To ensure that the quantum servers are performing their tasks honestly, the client can give them "dummy tasks" that are indistinguishable from the real task to test their honesty and correctness.

The researchers expect that the new method can be scaled up for realizing secure, outsourced quantum computing, which could one day be implemented on quantum cloud servers and make the power of quantum computing widely available.

"Blind quantum computing protocol is an important privacy-preserving technique for future secure quantum cloud computing and secure quantum networks," Lu said. "Applying our implemented blind quantum computing protocol, classical clients could delegate computation tasks to servers 'in the cloud' blindly and correctly without directly owning quantum devices. It saves resources and makes scalable quantum computing possible."

In the future, the physicists want to make blind quantum computing even easier for clients by further reducing the requirements.

"We plan to study more robust blind quantum computing protocols with fewer required resources and fewer constraints theoretically and experimentally," Lu said. "We will also explore blind quantum computing for more application scenarios, such as multi-user blind quantum computing, publicly verifiable quantum computing, and secure multi-party quantum computing." [15]

Chinese team sends quantum keys to ground stations and teleports ground to satellite signals

Two Chinese teams working with quantum encryption and entanglement have achieved two more goals toward building a quantum space-based communication network. In the first experiment, one team succeeded in sending quantum keys from a satellite to two ground stations. In the second, another team sent entangled photons from the ground to a satellite. Both teams were made up of researchers from several institutions in China and both have published their results in the journal *Nature*.

Quantum-based networks have been proposed as the next innovation for both speeding up and sending more information through communications networks—to that end, teams in countries such as Japan, the U.S. and China have been working hard to better understand how to actually create them. Most in the field agree that national, international and global quantum networks will require data to be sent at least partially via satellites because traditional media such as fiber cable results in too much interference and data loss. Quantum networks are also theorized to be hack-proof because observing the keys that are used to unlock the data would destroy them due to their quantum nature. In this new effort, the two research teams have fulfilled the second and third goals (the first was to break the distance record for sending entangled particles and was achieved this past June) outlined by officials directing that country's Quantum Experiments at Space Scale project, which involves making use of the Micius satellite—the first sent aloft to conduct quantum networking experiments.

The first team reports that they have successfully transmitted multiple quantum keys from the satellite to two ground-based stations in China—one in Xinglong the other in Nanshan. The distance involved was between 645 and 1,200 kilometers. The main result of the experiment was proving that it could be done.

The second team reported that they successfully sent entangled particles from ground stations in China to the satellite, complementing their experiments in June, in which the process was reversed.

In achieving their goals, the Chinese researchers have moved to the forefront of quantum-based network communications and have demonstrated that the country's leaders are serious about implementing the first global quantum system over the course of the next decade. [14]

China will launch an "unhackable" quantum communication network in its Jinan city next month

Just two weeks ago, China demonstrated its prowess in the field of quantum technology by becoming the first to teleport information from Earth to a satellite in space using the simple mechanics of quantum entanglement.

It's time for another first; in August, China plans to launch the world's largest quantum communications network, spanning 2,000 km and linking Beijing with Shanghai, passing through a message hub in China's Jinan city.

Jinan will use this network extensively for communication between government offices within the city, with 200 users spread across the government, finance, military, and electricity sectors; this will make it the first city in the world to use a quantum communications network for its government's functions.

The link between Beijing and Shanghai has been in testing since 2013, but the Jinan network's trials began two months ago. As reported by China's Xinhua News Agency, the groundbreaking network has exchanged data more than 51,000 times so far, with a success rate of 99 percent; this data includes things like secure phone calls, files and documents, as well as faxes – because we still need fax on a quantum communications network.

Quantum communication works in a unique manner; the technology still relies on the tried-and-tested fiber optic cables, but the key difference is in how the data is sent.

To begin a transfer of information, a key is first generated and sent over the network via traditional means; once the key is received, a single quantum-entangled photon is sent over the same fiber network, which can then be used to teleport information and decrypt it using the key that was received beforehand. However, due to the nature of quantum-entangled particles, any tampering (or intercept) of the photon results in either an alteration or destruction of it. This makes interception a physical impossibility, and thus China's claim of an "unhackable" communications network may stand true, at least with current technology.

To be clear: China isn't the first to build a quantum communication network – such networks have been operating in the US and Europe for the past few years, but only as research projects. China is, however, the first to use a network like this for a practical application.

China's – and the world's – need for a quantum communications network stems from the recent progress made with quantum computers. The traditional ways of encrypting data rely on simply hiding it behind an extremely difficult math problem; it's easy to decrypt the data if you have the key to this problem, but next to impossible if you don't. That is, for a traditional computer, but a quantum computer is much better at crunching numbers simultaneously, making it capable of brute forcing through some of the hardest encryption methods within a reasonable time.

Once matured, the technology will most likely be used by everyone – governments, banks, militaries, and everything in-between – but China may just be the first to have it for itself. [13]

Optical lens can transfer digital information without loss

Researchers have designed an optical lens that exhibits two properties that so far have not been demonstrated together: self-focusing and an optical effect called the Talbot effect that creates repeating patterns of light. The researchers showed that the combination of these two properties can be used to transfer an encoded digital signal without information loss, which has potential applications for realizing highly efficient optical communication systems.

The scientists, Xiangyang Wang and Hui Liu at Nanjing University, Huanyang Chen at Xiamen University, along with their coauthors, have published a paper on the new lens, called a "conformal lens," in a recent issue of Physical Review Letters.

This type of a conformal lens, which is also known as a Mikaelian lens, arose from the field of transformation optics, which is based on the idea that lenses can direct light in analogy with how the curved geometry of spacetime bends light in general relativity.

The main goal of the study was to design a conformal lens that works simultaneously in two different regimes: the geometry optics regime, in which light is treated as a particle, and the wave optics regime, which also accounts for the wave-like properties of light.

Working in both regimes is challenging because the two regimes have two seemingly opposing requirements for the size of the working wavelengths. On one hand, the working wavelengths must be much smaller than the size of the lens, but at the same time they must be larger than the basic units that make up the lens.

To address this challenge, the researchers started with a Maxwell's fish-eye lens, which dates back to the 1850s, as the basis for the conformal lens. They explained that trying to realize a lens with the desired properties using conventional transformation optics is very challenging, in part due its demands on a three-dimensional medium. On the other hand, conformal transformation optics places demands on a two-dimensional medium, which eases the fabrication requirements.

"Although transformation optics can be used to design many novel optical devices, it is usually very difficult to use in practical systems, especially in the visible regime," Liu told Phys.org. "In our work, we have established a feasible experiment platform to obtain conformation transformation optical devices."

After constructing the conformal lens, the researchers demonstrated that the lens exhibits both self-focusing, which is a property of geometric optics, and the Talbot effect, which is a property of wave optics. In this way, the device connects the two distinct realms of geometry optics and wave optics.

Most interesting for potential applications is that the conformal Talbot effect displayed here is very different from the ordinary Talbot effect in other media due to the additional self-focusing property. One of the biggest differences is that, unlike the ordinary Talbot effect which experiences boundary diffraction, the conformal Talbot effect does not.

As a result of its lack of diffraction, the conformal Talbot effect can be used to transfer encoded optical patterns over long distances with a very small amount of distortion. The researchers expect that this ability could lead to a highly efficient method of transferring digital information in future high-speed optical communication systems with no information loss.

"We can send a stream of optical digits '0' and '1' by parallel communication, which is much faster than the serial communication used in regular optical waveguides or optical fibers," Liu said. "The conformal Talbot effect can help reduce transmission errors because of its non-diffractive properties and good self-focusing of the field patterns."

In the future, the researchers plan to explore various potential applications of conformal transformation optics, such as designing novel integrated photonic chips that can transport and process information in micro-optical circuits. These "conformal photonic chips" may one day be used in future quantum computers.

"We hope conformal transformation optics can be used in quantum simulators and quantum computers in the future," Chen said. "We also plan to mimic the quantum effects in the curved space of general relativity using conformal transformation optics, such as the horizon of a black hole and Hawking radiation." [12]

Novel light sources made of 2-D materials

Physicists from the University of Würzburg have designed a light source that emits photon pairs, which are particularly well suited for tap-proof data encryption. The experiment's key ingredients: a semiconductor crystal and some sticky tape.

So-called monolayers are at the heart of the research activities. These so-called "super materials" have been surrounded by hype over the past decade. This is because they show great promise to revolutionise many areas of physics.

In physics, the term "monolayer" refers to solid materials of minimum thickness. Occasionally, it is only a single layer of atoms thick; in crystals, monolayers can be three or more layers. Experts also speak of two-dimensional materials. In this form, monolayers can exhibit unexpected properties that make them interesting for research. The so-called transition metal dichalcogenides (TMDC) are particularly promising. They behave like semiconductors and can be used to manufacture ultra-small and energy-efficient chips, for example.

Moreover, TMDCs are capable of generating light when supplied with energy. Dr. Christian Schneider, Professor Sven Höfling and their research team from the Chair of Technical Physics of the Julius-Maximilians-Universität Würzburg (JMU) in Bavaria, Germany, have harnessed exactly this effect for their experiments.

Experiments started with sticky tape

First, a monolayer was produced using a simple method. The researchers used a piece of sticky tape to peel a multi-layer film from a TMDC crystal. Using the same procedure, they stripped increasingly thin layers from the film, repeating the process until the material on the tape was only one layer thick.

The researchers then cooled this monolayer to a temperature of just above absolute zero and excited it with a laser. This caused the monolayer to emit single photons under specific conditions. "We were now able to show that a specific type of excitement produces not one but exactly two photons," Schneider explains.

"The light particles are generated in pairs, so to speak."

Such two-photon sources can be used to transfer 100 percent tap-proof information. For this purpose, the light particles are entangled. The state of the first photon then has a direct impact on that of the second photon, regardless of the distance between the two. This state can be used to encrypt communication channels.

Monolayers enable novel lasers

In a second study, the JMU scientists demonstrated another application of exotic monolayers. They mounted a monolayer between two mirrors and again stimulated it with a laser. The radiation excited the TMDC plate itself to emit photons. These were reflected back to the plate by the mirrors, where they excited atoms to create new photons.

"We call this process strong coupling," Schneider explains. The light particles are cloned during this process, in a manner of speaking. "Light and matter hybridise, forming new quasi particles in the process: exciton polaritons," the physicist says. For the first time, it is possible to detect these polaritons at room temperature in atomic monolayers.

In the short term, this will open up interesting new applications. The "cloned" photons have properties similar to laser light. But they are manufactured in completely different ways. Ideally, the production of new light particles is self-sustaining after the initial excitation without requiring any additional energy supply. In a laser, however, the light-producing material has to be excited energetically from the outside on a permanent basis. This makes the new light source highly energy efficient.

Moreover, it is well suited to study certain quantum effects. [11]

How to Win at Bridge Using Quantum Physics

Contract bridge is the chess of card games. You might know it as some stuffy old game your grandparents play, but it requires major brainpower, and preferably an obsession with rules and strategy. So how to make it even geekier? Throw in some quantum mechanics to try to gain a competitive advantage. The idea here is to use the quantum magic of entangled photons—which are essentially twins, sharing every property—to transmit two bits of information to your bridge partner for the price of one. Understanding how to do this is not an easy task, but it will help elucidate some basic building blocks of quantum information theory. It's also kind of fun to consider whether or not such tactics could ever be allowed in professional sports. [6]

Quantum Information

In quantum mechanics, quantum information is physical information that is held in the "state" of a quantum system. The most popular unit of quantum information is the qubit, a two-level quantum system. However, unlike classical digital states (which are discrete), a two-state quantum system can actually be in a superposition of the two states at any given time.

Quantum information differs from classical information in several respects, among which we note the following:

However, despite this, the amount of information that can be retrieved in a single qubit is equal to one bit. It is in the processing of information (quantum computation) that a difference occurs.

The ability to manipulate quantum information enables us to perform tasks that would be unachievable in a classical context, such as unconditionally secure transmission of information. Quantum information processing is the most general field that is concerned with quantum information. There are certain tasks which classical computers cannot perform "efficiently" (that is, in polynomial time) according to any known algorithm. However, a quantum computer can compute the answer to some of these problems in polynomial time; one well-known example of this is Shor's factoring algorithm. Other algorithms can speed up a task less dramatically - for example, Grover's search algorithm which gives a quadratic speed-up over the best possible classical algorithm.

Quantum information, and changes in quantum information, can be quantitatively measured by using an analogue of Shannon entropy. Given a statistical ensemble of quantum mechanical systems with the density matrix S , it is given by.

Many of the same entropy measures in classical information theory can also be generalized to the quantum case, such as the conditional quantum entropy. [7]

Heralded Qubit Transfer

Optical photons would be ideal carriers to transfer quantum information over large distances. Researchers envisage a network where information is processed in certain nodes and transferred between them via photons. However, inherent losses in long-distance networks mean that the information transfer is subject to probabilistic errors, making it hard to know whether the transfer of a qubit of information has been successful. Now Gerhard Rempe and colleagues from the Max Planck Institute for Quantum Optics in Germany have developed a new protocol that solves this problem through a strategy that "heralds" the accurate transfer of quantum information at a network node.

The method developed by the researchers involves transferring a photonic qubit to an atomic qubit trapped inside an optical cavity. The photon-atom quantum information transfer is initiated via a quantum "logic-gate" operation, performed by reflecting the photon from the atom-cavity system, which creates an entangled atom-photon state. The detection of the reflected photon then collapses the atom into a definite state. This state can be one of two possibilities, depending on the photonic state detected: Either the atom is in the initial qubit state encoded in the photon and the transfer process is complete, or the atom is in a rotated version of this state. The authors were able to show that the roles of the atom and photon could be reversed. Their method could thus be used as a quantum memory that stores (photon-to-atom state transfer) and recreates (atom-to-photon state transfer) a single-photon polarization qubit. [9]

Quantum Teleportation

Quantum teleportation is a process by which quantum information (e.g. the exact state of an atom or photon) can be transmitted (exactly, in principle) from one location to another, with the help of classical communication and previously shared quantum entanglement between the sending and receiving location. Because it depends on classical communication, which can proceed no faster than the speed of light, it cannot be used for superluminal transport or communication of classical bits. It also cannot be used to make copies of a system, as this violates the no-cloning theorem. Although the name is inspired by the teleportation commonly used in fiction, current technology provides no possibility of anything resembling the fictional form of teleportation. While it is possible to teleport one or more qubits of information between two (entangled) atoms, this has not yet been achieved between molecules or anything larger. One may think of teleportation either as a kind of transportation, or as a kind of communication; it provides a way of transporting a qubit from one location to another, without having to move a physical particle along with it.

The seminal paper first expounding the idea was published by C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres and W. K. Wootters in 1993. Since then, quantum teleportation has been realized in various physical systems. Presently, the record distance for quantum teleportation is 143 km (89 mi) with photons, and 21 m with material systems. In August 2013, the achievement of "fully deterministic" quantum teleportation, using a hybrid technique, was reported. On 29 May 2014, scientists announced a reliable way of transferring data by quantum teleportation. Quantum teleportation of data had been done before but with highly unreliable methods. [8]

Quantum cryptography set for lift-off

Exchanging messages with almost complete security by exploiting the strange laws of quantum mechanics should in future be possible on a global scale. That is the conclusion of physicists in Italy, who have found that the delicate states needed for quantum cryptography can be transmitted via laser beam from an orbiting satellite to a receiver on the surface of the Earth. The researchers say that the relatively simple technology needed for such encryption could be incorporated into conventional communications satellites.

Quantum cryptography involves two parties sharing a secret key that is created using the states of quantum particles such as photons. The communicating parties can then exchange messages by conventional means, in principle with complete security, by encrypting them using the secret key. Any eavesdropper trying to intercept the key automatically reveals their presence by destroying the quantum states.

Losses and curvature

Such cryptographic systems are already produced commercially, but they use fibre-optic cables. Losses in the cables limit the distance over which quantum keys can be sent to about 100 km, and that distance cannot be increased using repeaters, as is the case with classical data, because it is impossible to carry out the necessary amplification. Alternatively, quantum bits, or qubits, can be transmitted through the atmosphere, but this approach has a similar distance limit imposed by the curvature of the Earth.

This is where satellites could help. A single satellite, for example, could be used to send quantum data to two people on the Earth's surface to enable those people to share a secret key. To date, however, no device capable of generating or detecting quantum states – such as single photons – has been placed in orbit.

Paolo Villoresi of the University of Padua and colleagues have taken a creative approach to this problem by using the Matera Laser Ranging Observatory in southern Italy. This facility usually directs laser pulses at passing satellites and then measures the reflected pulses in order to measure tiny variations in the Earth's gravitational field. In 2008 Villoresi's team worked with a group of physicists at the University of Vienna to bounce very weak laser pulses from a satellite and then show that less than one photon per pulse could be detected on the ground (see "Single photons make the trek from space").

Polarization is preserved

In this latest research, the Italian group has gone one better, showing that it is possible to preserve the polarization state of those photons. Doing so is essential to quantum cryptography because it is the property of polarization – the orientation of a wave's oscillation – that is used to define the value of the qubits that make up a quantum key.

The researchers prepared the observatory's laser photons in one of four polarization states – horizontal, vertical, left-circular or right-circular – and beamed each of the states in 10-second bursts towards five satellites (including NASA's Jason-2) in orbits up to 2000 km above the Earth's surface. Their aim was to establish whether or not they could limit the fraction of qubits in each burst that had the wrong polarization after reflection to less than 11%. Above this figure, information theory dictates that no secret key can be established.

Villoresi and colleagues found, as hoped, that the error rates from four of the satellites were in single-figure percentages. These satellites employ corner-cube retroreflectors with metallic coatings, which are needed to preserve polarization states. The fifth satellite has uncoated retroreflectors and generated error rates of about 40%.

"Our results prove that quantum-key distribution from an orbiting terminal and a base station is not only a promising idea but nowadays is realizable," the researchers write.

Rotation on the fly

The tests did not involve the satellites transmitting qubits that could be used to make actual quantum keys, since the polarizations of the qubits were determined on the ground. But the researchers say that a straightforward modification of existing retroreflectors could make quantum-key generation a reality. All that is needed, they say, is to add a device known as a Faraday rotator and a random-number generator to each retroreflector in order to rotate the polarization of incoming photons on the fly.

Scientists in China have already developed a satellite that will generate quantum keys, and plan to launch it next year. This mission will create entangled pairs of photons in space and then send the two halves of each pair simultaneously to two communicating parties on the ground. The retroreflector-based scheme, on the other hand, involves transmitting the key to each user

separately. According to Villoresi, the latter approach will be much cheaper and easier to implement and, he says, could "piggyback" on satellites due to be launched anyway. [10]

Quantum Computing

A team of electrical engineers at UNSW Australia has observed the unique quantum behavior of a pair of spins in silicon and designed a new method to use them for "2-bit" quantum logic operations.

These milestones bring researchers a step closer to building a quantum computer, which promises dramatic data processing improvements.

Quantum bits, or qubits, are the building blocks of quantum computers. While many ways to create a qubits exist, the Australian team has focused on the use of single atoms of phosphorus, embedded inside a silicon chip similar to those used in normal computers.

The first author on the experimental work, PhD student Juan Pablo Dehollain, recalls the first time he realized what he was looking at.

"We clearly saw these two distinct quantum states, but they behaved very differently from what we were used to with a single atom. We had a real 'Eureka!' moment when we realized what was happening – we were seeing in real time the 'entangled' quantum states of a pair of atoms." [5]

Quantum Entanglement

Measurements of physical properties such as position, momentum, spin, polarization, etc. performed on entangled particles are found to be appropriately correlated. For example, if a pair of particles is generated in such a way that their total spin is known to be zero, and one particle is found to have clockwise spin on a certain axis, then the spin of the other particle, measured on the same axis, will be found to be counterclockwise. Because of the nature of quantum measurement, however, this behavior gives rise to effects that can appear paradoxical: any measurement of a property of a particle can be seen as acting on that particle (e.g. by collapsing a number of superimposed states); and in the case of entangled particles, such action must be on the entangled system as a whole. It thus appears that one particle of an entangled pair "knows" what measurement has been performed on the other, and with what outcome, even though there is no known means for such information to be communicated between the particles, which at the time of measurement may be separated by arbitrarily large distances. [4]

The Bridge

The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Quantum Theories. [1]

Accelerating charges

The moving charges are self maintain the electromagnetic field locally, causing their movement and this is the result of their acceleration under the force of this field. In the classical physics the charges will distributed along the electric current so that the electric potential lowering along the current, by linearly increasing the way they take every next time period because this accelerated motion.

The same thing happens on the atomic scale giving a dp impulse difference and a dx way difference between the different part of the not point like particles.

Relativistic effect

Another bridge between the classical and quantum mechanics in the realm of relativity is that the charge distribution is lowering in the reference frame of the accelerating charges linearly: $ds/dt = at$ (time coordinate), but in the reference frame of the current it is parabolic: $s = a/2 t^2$ (geometric coordinate).

Heisenberg Uncertainty Relation

In the atomic scale the Heisenberg uncertainty relation gives the same result, since the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that they product is about the half Planck reduced constant. For the proton this Δx much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greater proton mass.

This means that the electron and proton are not point like particles, but has a real charge distribution.

Wave – Particle Duality

The accelerating electrons explains the wave – particle duality of the electrons and photons, since the elementary charges are distributed on Δx position with Δp impulse and creating a wave packet of the electron. The photon gives the electromagnetic particle of the mediating force of the electrons electromagnetic field with the same distribution of wavelengths.

Atomic model

The constantly accelerating electron in the Hydrogen atom is moving on the equipotential line of the proton and it's kinetic and potential energy will be constant. Its energy will change only when it is changing its way to another equipotential line with another value of potential energy or getting free with enough kinetic energy. This means that the Rutherford-Bohr atomic model is right and only that changing acceleration of the electric charge causes radiation, not the steady acceleration. The steady acceleration of the charges only creates a centric parabolic steady electric field around the charge, the magnetic field. This gives the magnetic moment of the atoms, summing up the proton and electron magnetic moments caused by their circular motions and spins.

The Relativistic Bridge

Commonly accepted idea that the relativistic effect on the particle physics is the fermions' spin - another unresolved problem in the classical concepts. If the electric charges can move only with accelerated motions in the self maintaining electromagnetic field, once upon a time they would reach the velocity of the electromagnetic field. The resolution of this problem is the spinning particle, constantly accelerating and not reaching the velocity of light because the acceleration is radial. One origin of the Quantum Physics is the Planck Distribution Law of the electromagnetic oscillators, giving equal intensity for 2 different wavelengths on any temperature. Any of these two wavelengths will give equal intensity diffraction patterns, building different asymmetric constructions, for example proton - electron structures (atoms), molecules, etc. Since the particles are centers of diffraction patterns they also have particle - wave duality as the electromagnetic waves have. [2]

The weak interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry. The Electroweak Interaction shows that the Weak Interaction is basically electromagnetic in nature. The arrow of time shows the entropy grows by changing the temperature dependent diffraction patterns of the electromagnetic oscillators.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a 1/2 spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $\frac{1}{2}$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell-Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking!!! This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating; it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change. There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

Fermions and Bosons

The fermions are the diffraction patterns of the bosons such a way that they are both sides of the same thing.

Van Der Waals force

Named after the Dutch scientist Johannes Diderik van der Waals – who first proposed it in 1873 to explain the behaviour of gases – it is a very weak force that only becomes relevant when atoms and molecules are very close together. Fluctuations in the electronic cloud of an atom mean that it will have an instantaneous dipole moment. This can induce a dipole moment in a nearby atom, the result being an attractive dipole–dipole interaction.

Electromagnetic inertia and mass

Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

Relativistic change of mass

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

The frequency dependence of mass

Since $E = h\nu$ and $E = mc^2$, $m = h\nu / c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

Electron – Proton mass rate

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force. [2]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

Gravity from the point of view of quantum physics

The Gravitational force

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass ratio $M_p=1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass.

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Higgs boson

By March 2013, the particle had been proven to behave, interact and decay in many of the expected ways predicted by the Standard Model, and was also tentatively confirmed to have + parity and zero spin, two fundamental criteria of a Higgs boson, making it also the first known scalar particle to be discovered in nature, although a number of other properties were not fully proven and some partial results do not yet precisely match those expected; in some cases data is also still awaited or being analyzed.

Since the Higgs boson is necessary to the W and Z bosons, the dipole change of the Weak interaction and the change in the magnetic effect caused gravitation must be conducted. The Wien law is also important to explain the Weak interaction, since it describes the T_{\max} change and the diffraction patterns change. [2]

Higgs mechanism and Quantum Gravity

The magnetic induction creates a negative electric field, causing an electromagnetic inertia. Probably it is the mysterious Higgs field giving mass to the charged particles? We can think about the photon as an electron-positron pair, they have mass. The neutral particles are built from negative and positive charges, for example the neutron, decaying to proton and electron. The wave – particle duality makes sure that the particles are oscillating and creating magnetic induction as an inertial mass, explaining also the relativistic mass change. Higher frequency creates stronger magnetic induction, smaller frequency results lesser magnetic induction. It seems to me that the magnetic induction is the secret of the Higgs field.

In particle physics, the Higgs mechanism is a kind of mass generation mechanism, a process that gives mass to elementary particles. According to this theory, particles gain mass by interacting with the Higgs field that permeates all space. More precisely, the Higgs mechanism endows gauge bosons in a gauge theory with mass through absorption of Nambu–Goldstone bosons arising in spontaneous symmetry breaking.

The simplest implementation of the mechanism adds an extra Higgs field to the gauge theory. The spontaneous symmetry breaking of the underlying local symmetry triggers conversion of components of this Higgs field to Goldstone bosons which interact with (at least some of) the other fields in the theory, so as to produce mass terms for (at least some of) the gauge bosons. This mechanism may also leave behind elementary scalar (spin-0) particles, known as Higgs bosons.

In the Standard Model, the phrase "Higgs mechanism" refers specifically to the generation of masses for the W^\pm , and Z weak gauge bosons through electroweak symmetry breaking. The Large Hadron Collider at CERN announced results consistent with the Higgs particle on July 4, 2012 but stressed that further testing is needed to confirm the Standard Model.

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor

(compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

Conclusions

The tests did not involve the satellites transmitting qubits that could be used to make actual quantum keys, since the polarizations of the qubits were determined on the ground. But the researchers say that a straightforward modification of existing retroreflectors could make quantum-key generation a reality. All that is needed, they say, is to add a device known as a Faraday rotator and a random-number generator to each retroreflector in order to rotate the polarization of incoming photons on the fly. [10]

The method developed by the researchers involves transferring a photonic qubit to an atomic qubit trapped inside an optical cavity. The photon-atom quantum information transfer is initiated via a quantum "logic-gate" operation, performed by reflecting the photon from the atom-cavity system, which creates an entangled atom-photon state. [9]

In August 2013, the achievement of "fully deterministic" quantum teleportation, using a hybrid technique, was reported. On 29 May 2014, scientists announced a reliable way of transferring data by quantum teleportation. Quantum teleportation of data had been done before but with highly unreliable methods. [8]

One of the most important conclusions is that the electric charges are moving in an accelerated way and even if their velocity is constant, they have an intrinsic acceleration anyway, the so called spin, since they need at least an intrinsic acceleration to make possible their movement .

The accelerated charges self-maintaining potential shows the locality of the relativity, working on the quantum level also. [1]

The bridge between the classical and quantum theory is based on this intrinsic acceleration of the spin, explaining also the Heisenberg Uncertainty Principle. The particle – wave duality of the electric charges and the photon makes certain that they are both sides of the same thing.

The Secret of Quantum Entanglement that the particles are diffraction patterns of the electromagnetic waves and this way their quantum states every time is the result of the quantum state of the intermediate electromagnetic waves. [2]

The key breakthrough to arrive at this new idea to build qubits was to exploit the ability to control the nuclear spin of each atom. With that insight, the team has now conceived a unique way to use the nuclei as facilitators for the quantum logic operation between the electrons. [5]

Basing the gravitational force on the accelerating Universe caused magnetic force and the Planck Distribution Law of the electromagnetic waves caused diffraction gives us the basis to build a Unified Theory of the physical interactions also.

References

[1] The Magnetic field of the Electric current and the Magnetic induction

http://academia.edu/3833335/The_Magnetic_field_of_the_Electric_current

[2] 3 Dimensional String Theory

http://academia.edu/3834454/3_Dimensional_String_Theory

[3] Graviton Production By Two Photon and Electron-Photon Processes In Kaluza-Klein Theories With Large Extra Dimensions

<http://arxiv.org/abs/hep-ph/9909392>

[4] Quantum Entanglement

http://en.wikipedia.org/wiki/Quantum_entanglement

[5] Pairing up single atoms in silicon for quantum computing

<http://phys.org/news/2014-06-pairing-atoms-silicon-quantum.html#nwlT>

[6] How to Win at Bridge Using Quantum Physics

<http://www.wired.com/2014/06/bridge-quantum-mechanics/>

[7] Information Entropy-Theory of Physics

https://www.academia.edu/3836084/Information_-_Entropy_Theory_of_Physics

[8] Quantum Teleportation

http://en.wikipedia.org/wiki/Quantum_teleportation

[9] Synopsis: Heralded Qubit Transfer

<http://physics.aps.org/synopsis-for/10.1103/PhysRevLett.114.220501>

[10] Quantum cryptography set for lift-off

<http://physicsworld.com/cws/article/news/2015/jun/29/quantum-cryptography-set-for-lift-off>

[11] Novel light sources made of 2-D materials

<http://phys.org/news/2016-10-sources-d-materials.html>

[12] Optical lens can transfer digital information without loss

<https://phys.org/news/2017-07-optical-lens-digital-loss.html>

[13] China will launch an "unhackable" quantum communication network in its Jinan city next month

<https://www.neowin.net/news/china-will-launch-an-unhackable-quantum-communication-network-in-its-jinan-city-next-month>

[14] Chinese team sends quantum keys to ground stations and teleports ground to satellite signals

<https://phys.org/news/2017-08-chinese-team-quantum-keys-ground.html>

[15] Blind quantum computing for everyone

<https://phys.org/news/2017-08-quantum.html>

