

Recursive Future Average Of A Time Series Data Based On Cosine Similarity- RF

ISSN 1751-3030

Author:

Ramesh Chandra Bagadi

Data Scientist

INSOFE (International School Of Engineering),

Hyderabad, India.

rameshcbagadi@uwalumni.com

+91 9440032711

Technical Note

Abstract

In this research Technical Note the author have presented a Recursive Future Average Of A Time Series Data Based on Cosine Similarity.

Theory

The Recursive Future Average Of A Time Series Data Based on Cosine Similarity can be given by the following methods:

Method 1:

$$y_{n+1} = \frac{\sum_{i=1}^n (y_i) \{CS(y_i, y_{n+1})\}}{\left\{ \sum_{i=1}^n (\{CS(y_i, y_{n+1})\}^2) \right\}^{1/2}}$$

$$\text{where } CS(y_i, y_{n+1}) = \left\{ \frac{\text{Smaller of } (y_i, y_{n+1})}{\text{Larger of } (y_i, y_{n+1})} \right\}$$

when the Time Series Data is of the kind

$$S = \{y_1, y_2, y_3, \dots, y_{n-1}, y_n\}$$

Method 2:

$$y_{n+1} = \frac{\sum_{i=1}^n (y_i) \{CS(y_i, y_{n+1})\} \{CS(y_i, y_{n+1})\}}{\sum_{i=1}^n \{CS(y_i, y_{n+1})\}}$$

$$\text{where } CS(y_i, y_{n+1}) = \left\{ \frac{\text{Smaller of } (y_i, y_{n+1})}{\text{Larger of } (y_i, y_{n+1})} \right\}$$

when the Time Series Data is of the kind

$$S = \{y_1, y_2, y_3, \dots, y_{n-1}, y_n\}$$

Deriving motivation from this concept, we further extend this formula using [1] as

$$y_{n+1} = \frac{\left\{ \sum_{i=1}^n (y_i) \{CS(y_i, y_{n+1})\} + \sum_{i=1}^n ({}^1 y_i) \{CS({}^1 y_i, y_{n+1})\} + \sum_{i=1}^n ({}^2 y_i) \{CS({}^2 y_i, y_{n+1})\} + \dots + \sum_{i=1}^n ({}^r y_i) \{CS({}^r y_i, y_{n+1})\} \right\}}{\left\{ \sum_{i=1}^n \left(\{CS(y_i, y_{n+1})\}^2 \right) + \sum_{i=1}^n \left(\{CS({}^1 y_i, y_{n+1})\}^2 \right) + \sum_{i=1}^n \left(\{CS({}^2 y_i, y_{n+1})\}^2 \right) + \dots + \sum_{i=1}^n \left(\{CS({}^r y_i, y_{n+1})\}^2 \right) \right\}^{1/2}}$$

where ${}^1 y_i = \frac{\left\{ y_i y_{n+1} - \frac{(\text{Smaller of } (y_i, y_{n+1}))^2}{y_i y_{n+1}} \right\}}{y_{n+1}}$ and

${}^2 y_i = \frac{\left\{ {}^1 y_i y_{n+1} - \frac{(\text{Smaller of } ({}^1 y_i, y_{n+1}))^2}{{}^1 y_i y_{n+1}} \right\}}{y_{n+1}}$,, i.e., and so on, so forth

${}^k y_i = \frac{\left\{ {}^{k-1} y_i y_{n+1} - \frac{(\text{Smaller of } ({}^{k-1} y_i, y_{n+1}))^2}{{}^{k-1} y_i y_{n+1}} \right\}}{y_{n+1}}$

upto

${}^p y_i = \frac{\left\{ {}^{p-1} y_i y_{n+1} - \frac{(\text{Smaller of } ({}^{p-1} y_i, y_{n+1}))^2}{{}^{p-1} y_i y_{n+1}} \right\}}{y_{n+1}}$ such that we can write

$$y_{n+1} = \frac{\left\{ \sum_{i=1}^n (y_i) \{CS(y_i, y_{n+1})\} + \sum_{k=1}^p \sum_{i=1}^n ({}^k y_i) \{CS({}^k y_i, y_{n+1})\} \right\}}{\left\{ \sum_{i=1}^n \left(\{CS(y_i, y_{n+1})\}^2 \right) + \sum_{k=1}^p \sum_{i=1}^n \left(\{CS({}^k y_i, y_{n+1})\}^2 \right) \right\}^{1/2}}$$

where p is a number such that ${}^p y_i \rightarrow 0$.

References

1. Bagadi, R. (2016). Proof Of As To Why The Euclidean Inner Product Is A Good Measure Of Similarity Of Two Vectors. *PHILICA.COM Article number 626*.
http://philica.com/display_article.php?article_id=626
2. <http://www.philica.com/advancedsearch.php?author=12897>
3. http://www.vixra.org/author/ramesh_chandra_bagadi