

# Where Standard Physics Runs into Infinite Challenges, Atomism Predicts Exact Limits

Espen Gaarder Haug\*  
Norwegian University of Life Sciences

November 27, 2017

## Abstract

Where standard physics runs into infinite challenges, atomism predicts exact limits. We summarize the mathematical results briefly in a table in this note and are also revisiting the energy momentum relationship based on this view.

**Key words:** Rest-mass energy, rest-mass, relativistic energy, relativistic mass, proper velocity, momentum, kinetic energy, acceleration, rapidity, temperature.

## 1 Mathematical Summary of Upper Limits Predicted by Atomism

These are some of the main findings in a series of papers I have posted on this topic (see [1, 2, 3, 4, 5, 7, 8, 9, 10]). This is first draft, and I will update this paper later on. Comments are welcome.

In standard physics there is no limit below infinity for kinetic energy, for momentum, for relativistic mass, for relativistic energy, for temperature, or for proper velocity. Further, there is no good explanation for what the Planck acceleration truly represents. Many physicists assume that Planck time is the shortest possible unit of time, but in fact, no rest-mass in standard theory can be accelerated for even one Planck second, as this would mean that the mass was traveling at the speed of light and had infinite kinetic energy.

Under mathematical atomism (based on just two postulates), we get all the equations of special relativity when using Einstein-Poincaré synchronized clocks, and we also get a series of new predictions. We obtain the exact upper boundary conditions on a series of relativistic formulas that are linked to the entities found by Max Planck in 1906 [11].

Atomism leads to a quantized relativity theory that helps us to understand the Planck mass in a new way. The Planck mass is the very key to understanding physics, and in our view it can only be grasped fully through atomism. Still atomism leads to breaks in Lorentz symmetry at the Planck scale, this is consistent with what is predicted from several quantum gravity theories.

## 2 Energy Momentum Relationship

Here we show that the energy momentum relationship has a limit equal to the rest- mass energy of the Planck mass for any fundamental particle

---

\*e-mail [espenhaug@mac.com](mailto:espenhaug@mac.com). Thanks to Victoria Terces for assisting with manuscript editing.

	For non-Planck mass particles	For Planck mass particle
Rest-mass	$m$	$m_p$ (last one Planck second)
Rest-mass energy	$E = mc^2$	$E = m_p c^2$
Maximum relativistic energy	$E = m_p c^2$	$E = m_p c^2$
Maximum relativistic mass	$m_p$ then becomes light	$m_p$ (last one Planck second)
Maximum velocity	$v_{max} = c \sqrt{1 - \frac{l_p^2}{\lambda^2}}$	$v_{max} = 0$ (and $c$ when dissolved)
Maximum proper velocity	$W_{max} = c \frac{\bar{\lambda}}{l_p} \sqrt{1 - \frac{l_p^2}{\lambda^2}}$	$W_{max} = 0$
Maximum momentum	$p_{max} = \hbar \sqrt{\frac{1}{l_p^2} - \frac{1}{\lambda^2}}$	$p_{max} = 0$
Maximum kinetic energy	$E_{k,max} = \hbar c \left( \frac{1}{l_p} - \frac{1}{\lambda} \right)$	$E_{k,max} = 0$ (at rest)
Maximum acceleration	$a_{max} = \left( \frac{c^2}{l_p} - \frac{c^2}{\lambda} \right) \sqrt{1 - \frac{l_p^2}{\lambda^2}}$	0 and $a_p = \frac{c^2}{l_p}$ becomes light
Maximum force	$F_{max} = \left( \frac{\hbar c}{l_p^2} - \frac{\hbar c}{\lambda l_p} \right) \sqrt{1 - \frac{l_p^2}{\lambda^2}}$	$F_{max} = \frac{\hbar c}{l_p^2}$ (One Planck second)
Maximum rapidity	$w_{max} \approx \ln \left( 2 \frac{\bar{\lambda}}{l_p} \right)$	$w_{max} = 0$ (at rest)
Maximum Doppler shift	$f_2 = f_1 \sqrt{\frac{4 - \frac{l_p^2}{\lambda^2}}{\frac{l_p^2}{\lambda^2}}} \approx 2 \frac{c}{l_p}$	None (at rest)
Maximum length contraction	$\bar{\lambda} \sqrt{1 - \frac{v_{max}^2}{c^2}} = l_p$	$l_p$ (at rest, no contraction) Reduced Compton wavelength Planck particle
Maximum time dilation	$\frac{\bar{\lambda}}{c} \sqrt{1 - \frac{v_{max}^2}{c^2}} = \frac{l_p}{c}$ (Planck time) $\frac{\frac{\bar{\lambda}}{c}}{\sqrt{1 - \frac{v_{max}^2}{c^2}}} = \frac{\bar{\lambda}^2}{l_p c}$	0 (at rest, no time dilation)
Maximum temperature	$T_{max} = \frac{\hbar c \left( \frac{1}{l_p} - \frac{1}{\lambda} \right)}{k_b}$	$T_p = \frac{m_p c^2}{k_b} = \frac{\hbar c}{l_p k_b}$ (for one Planck second)
Fuel needed for maximum velocity for each fundamental particle.	$\geq 2m_p$	0 (Planck mass particle at rest)

**Table 1:** The table shows a series of new boundary conditions that are given by atomism.

$$\begin{aligned}
E_{max}^2 &= p_{max}^2 c^2 + m^2 c^4 \\
E_{max} &= \sqrt{p_{max}^2 c^2 + m^2 c^4} \\
E_{max} &= \sqrt{\left( \hbar \sqrt{\frac{1}{l_p^2} - \frac{1}{\lambda^2}} \right)^2 c^2 + \left( \frac{\hbar}{\bar{\lambda} c} \right)^2 c^4} \\
E_{max} &= \sqrt{\hbar^2 \left( \frac{1}{l_p^2} - \frac{1}{\lambda^2} \right) c^2 + \frac{\hbar^2}{\lambda^2} \frac{1}{c^2} c^4} \\
E_{max} &= \hbar c \sqrt{\frac{1}{l_p^2} - \frac{1}{\lambda^2} + \frac{1}{\lambda^2}} \\
E_{max} &= \frac{\hbar}{l_p} c = m_p c^2
\end{aligned} \tag{1}$$

In the special case of a Planck mass particle (see also [6]), we have that  $\bar{\lambda} = l_p$  and this gives

$$\begin{aligned}
E_{max} &= \sqrt{\left(\hbar\sqrt{\frac{1}{l_p^2} - \frac{1}{l_p^2}}\right)^2 c^2 + \left(\frac{\hbar}{l_p} \frac{1}{c}\right)^2 c^4} \\
E_{max} &= \sqrt{\hbar^2 \left(\frac{1}{l_p^2} - \frac{1}{l_p^2}\right) c^2 + \frac{\hbar^2}{l_p^2} \frac{1}{c^2} c^4} \\
E_{max} &= \hbar c \sqrt{\frac{1}{l_p^2} - \frac{1}{l_p^2} + \frac{1}{l_p^2}} \\
E_{max} &= \frac{\hbar}{l_p} c = m_p c^2
\end{aligned} \tag{2}$$

That is the same end result as for any other particle. Note, however, that the momentum for a Planck mass particle is zero. This is because a Planck mass particle only can exist when it is at absolute rest. The Planck mass particle is the same as observed across reference frames; the same is true for the Planck length and Planck time. Other particles have less rest-mass, but the maximum momentum ensures that they have total maximal energy equal to the Planck mass particle.

### 3 Maximum Acceleration Force

I have not calculated the maximum acceleration force before in any of my papers. The maximum relativistic acceleration force is given by<sup>1</sup>

$$\begin{aligned}
F_{max} &= \frac{m}{\sqrt{1 - \frac{v_{max}^2}{c^2}}} a_{max} \\
F_{max} &= m_p \left(\frac{c^2}{l_p} - \frac{c^2}{\lambda}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}} \\
F_{max} &= \frac{\hbar}{l_p} \frac{1}{c} \left(\frac{c^2}{l_p} - \frac{c^2}{\lambda}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}} \\
F_{max} &= \left(\frac{\hbar c}{l_p^2} - \frac{\hbar c}{\lambda l_p}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}}
\end{aligned} \tag{4}$$

For a Planck particle it is

$$F_{max} = m_p a_p = \frac{\hbar}{l_p} \frac{1}{c} \frac{c^2}{l_p} = \frac{\hbar c}{l_p^2} \tag{5}$$

Again the Planck mass particle is very unique.

---

<sup>1</sup>One could mistakenly forget to take the relativistic mass., this gives

$$\begin{aligned}
F_{max} &= m a_{max} \\
F_{max} &= m \left(\frac{c^2}{l_p} - \frac{c^2}{\lambda}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}} \\
F_{max} &= \frac{\hbar}{\lambda} \frac{1}{c} \left(\frac{c^2}{l_p} - \frac{c^2}{\lambda}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}} \\
F_{max} &= \left(\frac{\hbar c}{l_p \lambda} - \frac{\hbar c}{\lambda^2}\right) \sqrt{1 - \frac{l_p^2}{\lambda^2}}
\end{aligned} \tag{3}$$

That we clearly think is the incorrect way of doing this.

## References

- [1] E. G. Haug. *Unified Revolution: New Fundamental Physics*. Oslo, E.G.H. Publishing, 2014.
- [2] E. G. Haug. The Planck mass particle finally discovered! The true God particle! Good bye to the point particle hypothesis! <http://vixra.org/abs/1607.0496>, 2016.
- [3] E. G. Haug. Deriving the maximum velocity of matter from the Planck length limit on length contraction. <http://vixra.org/abs/1612.0358>, 2016.
- [4] E. G. Haug. A new solution to Einstein's relativistic mass challenge based on maximum frequency. <http://vixra.org/abs/1609.0083>, 2016.
- [5] E. G. Haug. Modern physics' incomplete absurd relativistic mass interpretation. And the simple solution that saves Einstein's formula. <http://vixra.org/abs/1612.0249>, 2016.
- [6] E. G. Haug. The Planck mass must always have zero momentum: Relativistic energy-momentum relationship for the Planck mass. <http://vixra.org/abs/1611.0230>, 2016.
- [7] E. G. Haug. The Lorentz transformation at the maximum velocity for a mass. <http://vixra.org/abs/1609.0151>, 2016.
- [8] E. G. Haug. The ultimate limits of the relativistic rocket equation: The Planck photon rocket. *Acta Astronautica*, 136, 2017.
- [9] E. G. Haug. The incompatibility of the Planck acceleration and modern physics? <http://vixra.org/abs/1710.0340>, 2017.
- [10] E. G. Haug. A maximum limit on proper velocity. <http://vixra.org/abs/1702.0230>, 2017.
- [11] M. Planck. *The Theory of Radiation*. Dover 1959 translation, 1906.