

# Research Project Primus

Predrag Terzić

Podgorica , Montenegro

e-mail: pedja.terzic@hotmail.com

May 1 , 2019

## 1 Theorems and Conjectures

**Theorem 1.1.** A natural number  $n > 2$  is a prime iff  $\prod_{k=1}^{n-1} k \equiv n - 1 \pmod{\sum_{k=1}^{n-1} k}$ .

**Theorem 1.2.** Let  $p \equiv 5 \pmod{6}$  be prime then ,  $2p + 1$  is prime iff  $2p + 1 \mid 3^p - 1$ .

**Theorem 1.3.** Let  $p_n$  be the  $n$ th prime , then

$$p_n = 1 + \sum_{k=1}^{2 \cdot (\lfloor n \ln(n) \rfloor + 1)} \left( 1 - \left[ \frac{1}{n} \cdot \sum_{j=2}^k \left[ \frac{3 - \sum_{i=1}^j \left[ \frac{\lfloor \frac{j}{i} \rfloor}{\lfloor \frac{j}{i} \rfloor} \right]}{j} \right] \right] \right)$$

**Theorem 1.4.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$  , where  $j$  and  $x$  are nonnegative integers . Let  $N = k \cdot 2^m - 1$  such that  $m > 2$  ,  $3 \mid k$  ,  $0 < k < 2^m$  and

$$\begin{cases} k \equiv 1 \pmod{10} \text{ with } m \equiv 2, 3 \pmod{4} \\ k \equiv 3 \pmod{10} \text{ with } m \equiv 0, 3 \pmod{4} \\ k \equiv 7 \pmod{10} \text{ with } m \equiv 1, 2 \pmod{4} \\ k \equiv 9 \pmod{10} \text{ with } m \equiv 0, 1 \pmod{4} \end{cases}$$

Let  $S_i = S_{i-1}^2 - 2$  with  $S_0 = P_k(3)$  , then  $N$  is prime iff  $S_{m-2} \equiv 0 \pmod{N}$

**Theorem 1.5.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$  , where  $j$  and  $x$  are nonnegative integers . Let  $N = k \cdot 2^m - 1$  such that  $m > 2$  ,  $3 \mid k$  ,  $0 < k < 2^m$  and

$$\begin{cases} k \equiv 3 \pmod{42} \text{ with } m \equiv 0, 2 \pmod{3} \\ k \equiv 9 \pmod{42} \text{ with } m \equiv 0 \pmod{3} \\ k \equiv 15 \pmod{42} \text{ with } m \equiv 1 \pmod{3} \\ k \equiv 27 \pmod{42} \text{ with } m \equiv 1, 2 \pmod{3} \\ k \equiv 33 \pmod{42} \text{ with } m \equiv 0, 1 \pmod{3} \\ k \equiv 39 \pmod{42} \text{ with } m \equiv 2 \pmod{3} \end{cases}$$

Let  $S_i = S_{i-1}^2 - 2$  with  $S_0 = P_k(5)$  , then  $N$  is prime iff  $S_{m-2} \equiv 0 \pmod{N}$

**Theorem 1.6.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$ , where  $j$  and  $x$  are nonnegative integers. Let  $N = k \cdot 2^m + 1$  such that  $m > 2$ ,  $0 < k < 2^m$  and

$$\left\{ \begin{array}{l} k \equiv 1 \pmod{42} \text{ with } m \equiv 2, 4 \pmod{6} \\ k \equiv 5 \pmod{42} \text{ with } m \equiv 3 \pmod{6} \\ k \equiv 11 \pmod{42} \text{ with } m \equiv 3, 5 \pmod{6} \\ k \equiv 13 \pmod{42} \text{ with } m \equiv 4 \pmod{6} \\ k \equiv 17 \pmod{42} \text{ with } m \equiv 5 \pmod{6} \\ k \equiv 19 \pmod{42} \text{ with } m \equiv 0 \pmod{6} \\ k \equiv 23 \pmod{42} \text{ with } m \equiv 1, 3 \pmod{6} \\ k \equiv 25 \pmod{42} \text{ with } m \equiv 0, 2 \pmod{6} \\ k \equiv 29 \pmod{42} \text{ with } m \equiv 1, 5 \pmod{6} \\ k \equiv 31 \pmod{42} \text{ with } m \equiv 2 \pmod{6} \\ k \equiv 37 \pmod{42} \text{ with } m \equiv 0, 4 \pmod{6} \\ k \equiv 41 \pmod{42} \text{ with } m \equiv 1 \pmod{6} \end{array} \right.$$

Let  $S_i = S_{i-1}^2 - 2$  with  $S_0 = P_k(5)$ , then  $N$  is prime iff  $S_{m-2} \equiv 0 \pmod{N}$ .

**Theorem 1.7.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$ , where  $j$  and  $x$  are nonnegative integers. Let  $N = k \cdot 2^m + 1$  such that  $m > 2$ ,  $0 < k < 2^m$  and

$$\left\{ \begin{array}{l} k \equiv 1 \pmod{6} \text{ and } k \equiv 1, 7 \pmod{10} \text{ with } m \equiv 0 \pmod{4} \\ k \equiv 5 \pmod{6} \text{ and } k \equiv 1, 3 \pmod{10} \text{ with } m \equiv 1 \pmod{4} \\ k \equiv 1 \pmod{6} \text{ and } k \equiv 3, 9 \pmod{10} \text{ with } m \equiv 2 \pmod{4} \\ k \equiv 5 \pmod{6} \text{ and } k \equiv 7, 9 \pmod{10} \text{ with } m \equiv 3 \pmod{4} \end{array} \right.$$

Let  $S_i = S_{i-1}^2 - 2$  with  $S_0 = P_k(8)$ , then  $N$  is prime iff  $S_{m-2} \equiv 0 \pmod{N}$ .

**Theorem 1.8.** Let  $N = k \cdot 2^n + 1$  with  $n > 1$ ,  $k$  is odd,  $0 < k < 2^n$ ,  $3 \mid k$  and

$$\left\{ \begin{array}{l} k \equiv 3 \pmod{30}, \quad \text{with } n \equiv 1, 2 \pmod{4} \\ k \equiv 9 \pmod{30}, \quad \text{with } n \equiv 2, 3 \pmod{4} \\ k \equiv 21 \pmod{30}, \quad \text{with } n \equiv 0, 1 \pmod{4} \\ k \equiv 27 \pmod{30}, \quad \text{with } n \equiv 0, 3 \pmod{4} \end{array} \right.$$

then  $N$  is prime iff  $5^{\frac{N-1}{2}} \equiv -1 \pmod{N}$ .

**Theorem 1.9.** Let  $N = k \cdot 2^n + 1$  with  $n > 1$ ,  $k$  is odd,  $0 < k < 2^n$ ,  $3 \mid k$  and

$$\left\{ \begin{array}{l} k \equiv 3 \pmod{42}, \quad \text{with } n \equiv 2 \pmod{3} \\ k \equiv 9 \pmod{42}, \quad \text{with } n \equiv 0, 1 \pmod{3} \\ k \equiv 15 \pmod{42}, \quad \text{with } n \equiv 1, 2 \pmod{3} \\ k \equiv 27 \pmod{42}, \quad \text{with } n \equiv 1 \pmod{3} \\ k \equiv 33 \pmod{42}, \quad \text{with } n \equiv 0 \pmod{3} \\ k \equiv 39 \pmod{42}, \quad \text{with } n \equiv 0, 2 \pmod{3} \end{array} \right.$$

then  $N$  is prime iff  $7^{\frac{N-1}{2}} \equiv -1 \pmod{N}$ .

**Theorem 1.10.** Let  $N = k \cdot 2^n + 1$  with  $n > 1$ ,  $k$  is odd,  $0 < k < 2^n$ ,  $3 \mid k$  and

$$\left\{ \begin{array}{l} k \equiv 3 \pmod{66}, \quad \text{with } n \equiv 1, 2, 6, 8, 9 \pmod{10} \\ k \equiv 9 \pmod{66}, \quad \text{with } n \equiv 0, 1, 3, 4, 8 \pmod{10} \\ k \equiv 15 \pmod{66}, \quad \text{with } n \equiv 2, 4, 5, 7, 8 \pmod{10} \\ k \equiv 21 \pmod{66}, \quad \text{with } n \equiv 1, 2, 4, 5, 9 \pmod{10} \\ k \equiv 27 \pmod{66}, \quad \text{with } n \equiv 0, 2, 3, 5, 6 \pmod{10} \\ k \equiv 39 \pmod{66}, \quad \text{with } n \equiv 0, 1, 5, 7, 8 \pmod{10} \\ k \equiv 45 \pmod{66}, \quad \text{with } n \equiv 0, 4, 6, 7, 9 \pmod{10} \\ k \equiv 51 \pmod{66}, \quad \text{with } n \equiv 0, 2, 3, 7, 9 \pmod{10} \\ k \equiv 57 \pmod{66}, \quad \text{with } n \equiv 3, 5, 6, 8, 9 \pmod{10} \\ k \equiv 63 \pmod{66}, \quad \text{with } n \equiv 1, 3, 4, 6, 7 \pmod{10} \end{array} \right.$$

then  $N$  is prime iff  $11^{\frac{N-1}{2}} \equiv -1 \pmod{N}$ .

**Theorem 1.11.** A positive integer  $n$  is prime iff  $\varphi(n)! \equiv -1 \pmod{n}$

**Theorem 1.12.** For  $m \geq 1$  number  $n$  greater than one is prime iff

$$(n^m - 1)! \equiv (n - 1)!^{\left\lceil \frac{(-1)^{m+1}}{2} \right\rceil} \cdot n^{\frac{n^m - mn + m - 1}{n-1}} \pmod{n^{\frac{n^m - mn + m + n - 2}{n-1}}}$$

**Theorem 1.13.** Sequence  $S_i$  is defined as  $S_i = \begin{cases} 8 & \text{if } i = 0; \\ (S_{i-1}^2 - 2)^2 - 2 & \text{otherwise.} \end{cases}$  then,  $F_n = 2^{2^n} + 1$ , ( $n \geq 2$ ) is a prime if and only if  $F_n$  divides  $S_{2^{n-1}-1}$ .

**Theorem 1.14.** Let  $p \equiv 1 \pmod{6}$  be prime and let  $5 \nmid 4p + 1$ , then  $4p + 1$  is prime iff  $4p + 1 \mid 2^{2^p} + 1$ .

**Theorem 1.15.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $F_n(b) = b^{2^n} + 1$  such that  $n \geq 2$  and  $b$  is even number. Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_b(6)$ , thus If  $F_n(b)$  is prime, then  $S_{2^n-1} \equiv 2 \pmod{F_n(b)}$ .

**Theorem 1.16.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $E_n(b) = \frac{b^{2^n} + 1}{2}$  such that  $n > 1$ ,  $b$  is odd number greater than one. Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_b(6)$ , thus If  $E_n(b)$  is prime, then  $S_{2^n-1} \equiv 6 \pmod{E_n(b)}$ .

**Theorem 1.17.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ .

Let  $N_p(b) = \frac{b^p + 1}{b + 1}$ , where  $p$  is an odd prime and  $b$  is an odd natural number greater than one.

CASE(1).  $b \equiv 1, 9 \pmod{12}$ , or  $b \equiv 3, 7 \pmod{12}$  and  $p \equiv 1 \pmod{4}$ , or  $b \equiv 5 \pmod{12}$  and  $p \equiv 1, 7 \pmod{12}$ , or  $b \equiv 11 \pmod{12}$  and  $p \equiv 1, 11 \pmod{12}$ .

CASE(2).  $b \equiv 3, 7 \pmod{12}$  and  $p \equiv 3 \pmod{4}$ , or  $b \equiv 5 \pmod{12}$  and  $p \equiv 5, 11 \pmod{12}$ , or  $b \equiv 11 \pmod{12}$  and  $p \equiv 5, 7 \pmod{12}$ .

Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_b(4)$ . Suppose  $N_p(b)$  is prime, then :

- $S_{p-1} \equiv P_b(4) \pmod{N_p(b)}$  if Case(1) holds ;
- $S_{p-1} \equiv P_{b+2}(4) \pmod{N_p(b)}$  if Case(2) holds ;

**Theorem 1.18.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ .

Let  $M_p(a) = \frac{a^p - 1}{a - 1}$ , where  $p$  is an odd prime and  $a$  is an odd natural number greater than one

CASE(1).  $a \equiv 3, 11 \pmod{12}$ , or  $a \equiv 5, 9 \pmod{12}$  and  $p \equiv 1 \pmod{4}$ , or  $a \equiv 7 \pmod{12}$  and  $p \equiv 1, 7 \pmod{12}$ , or  $a \equiv 1 \pmod{12}$  and  $p \equiv 1, 11 \pmod{12}$ .

CASE(2).  $a \equiv 5, 9 \pmod{12}$  and  $p \equiv 3 \pmod{4}$ , or  $a \equiv 7 \pmod{12}$  and  $p \equiv 5, 11 \pmod{12}$ , or  $a \equiv 1 \pmod{12}$  and  $p \equiv 5, 7 \pmod{12}$ .

Let  $S_i = P_a(S_{i-1})$  with  $S_0 = P_a(4)$ . Suppose  $M_p(a)$  is prime, then :

- $S_{p-1} \equiv P_a(4) \pmod{M_p(a)}$  if Case(1) holds ;
- $S_{p-1} \equiv P_{a-2}(4) \pmod{M_p(a)}$  if Case(2) holds ;

**Conjecture 1.1.** Let  $b_n = b_{n-2} + \text{lcm}(n-1, b_{n-2})$  with  $b_1 = 2$ ,  $b_2 = 2$  and  $n > 2$ . Let  $a_n = b_{n+2}/b_n - 1$ , then

1. Every term of this sequence  $a_i$  is either prime or 1. 2. Every odd prime number is member of this sequence. 3. Every new prime in sequence is a next prime from the largest prime already listed.

**Conjecture 1.2.** Let  $b_n = b_{n-1} + \text{lcm}(\lfloor \sqrt{n^3} \rfloor, b_{n-1})$  with  $b_1 = 2$  and  $n > 1$ . Let  $a_n = b_{n+1}/b_n - 1$ , then

1. Every term of this sequence  $a_i$  is either prime or 1. 2. Every odd prime of the form  $\lfloor \sqrt{n^3} \rfloor$  is member of this sequence. 3. Every new prime of the form  $\lfloor \sqrt{n^3} \rfloor$  in sequence is a next prime from the largest prime already listed.

**Conjecture 1.3.** Let  $b_n = b_{n-1} + \text{lcm}(\lfloor \sqrt{2} \cdot n \rfloor, b_{n-1})$  with  $b_1 = 2$  and  $n > 1$ . Let  $a_n = b_{n+1}/b_n - 1$ , then

1. Every term of this sequence  $a_i$  is either prime or 1. 2. Every prime of the form  $\lfloor \sqrt{2} \cdot n \rfloor$  is member of this sequence. 3. Every new prime of the form  $\lfloor \sqrt{2} \cdot n \rfloor$  in sequence is a next prime from the largest prime already listed.

**Conjecture 1.4.** Let  $b_n = b_{n-1} + \text{lcm}(\lfloor \sqrt{3} \cdot n \rfloor, b_{n-1})$  with  $b_1 = 3$  and  $n > 1$ . Let  $a_n = b_{n+1}/b_n - 1$ , then

1. Every term of this sequence  $a_i$  is either prime or 1. 2. Every prime of the form  $\lfloor \sqrt{3} \cdot n \rfloor$  is member of this sequence. 3. Every new prime of the form  $\lfloor \sqrt{3} \cdot n \rfloor$  in sequence is a next prime from the largest prime already listed.

**Conjecture 1.5.** Let  $b$  and  $n$  be a natural numbers,  $b \geq 2$ ,  $n > 2$  and  $n \neq 9$ . Then  $n$  is prime if

and only if  $\sum_{k=1}^{n-1} (b^k - 1)^{n-1} \equiv n \pmod{\frac{b^n - 1}{b - 1}}$

**Conjecture 1.6.** Let  $a$ ,  $b$  and  $n$  be a natural numbers,  $b > a > 1$ ,  $n > 2$  and  $n \notin \{4, 9, 25\}$ .

Then  $n$  is prime iff  $\prod_{k=1}^{n-1} (b^k - a) \equiv \frac{a^n - 1}{a - 1} \pmod{\frac{b^n - 1}{b - 1}}$

**Conjecture 1.7.** Let  $a$ ,  $b$  and  $n$  be a natural numbers,  $b > a > 0$ ,  $n > 2$  and  $n \notin \{4, 9, 25\}$ .

Then  $n$  is prime iff  $\prod_{k=1}^{n-1} (b^k + a) \equiv \frac{a^n + 1}{a + 1} \pmod{\frac{b^n - 1}{b - 1}}$

**Conjecture 1.8.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - 1$  such that  $k > 0$ ,  $3 \nmid k$ ,  $k < 2^n$ ,  $b > 0$ ,  $b$  is even number,  $3 \nmid b$  and  $n > 2$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{kb/2}(P_{b/2}(4))$ , then  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$ .

**Conjecture 1.9.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$ , where  $j$  and  $x$  are nonnegative integers. Let  $N = k \cdot 2^m + 1$  with  $k$  odd,  $0 < k < 2^m$  and  $m > 2$ . Let  $F_n$  be the  $n$ th Fibonacci number and let  $S_i = S_{i-1}^2 - 2$  with  $S_0 = P_k(F_n)$ , then  $N$  is prime iff there exists  $F_n$  for which  $S_{m-2} \equiv 0 \pmod{N}$ .

**Conjecture 1.10.** Let  $P_j(x) = 2^{-j} \cdot \left( (x - \sqrt{x^2 - 4})^j + (x + \sqrt{x^2 - 4})^j \right)$ , where  $j$  and  $x$  are nonnegative integers. Let  $F_m(b) = b^{2^m} + 1$  with  $b$  even,  $b > 0$  and  $m \geq 2$ . Let  $F_n$  be the  $n$ th Fibonacci number and let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(P_{b/2}(F_n))$ , then  $F_m(b)$  is prime iff there exists  $F_n$  for which  $S_{m-2} \equiv 0 \pmod{F_m(b)}$ .

**Conjecture 1.11.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n - b - 1$  such that  $n > 2$ ,  $b \equiv 0, 6 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv P_{(b+2)/2}(6) \pmod{N}$ .

**Conjecture 1.12.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n - b - 1$  such that  $n > 2$ ,  $b \equiv 2, 4 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv -P_{b/2}(6) \pmod{N}$ .

**Conjecture 1.13.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n + b + 1$  such that  $n > 2$ ,  $b \equiv 0, 6 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv P_{b/2}(6) \pmod{N}$ .

**Conjecture 1.14.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n + b + 1$  such that  $n > 2$ ,  $b \equiv 2, 4 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv -P_{(b+2)/2}(6) \pmod{N}$ .

**Conjecture 1.15.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n - b + 1$  such that  $n > 3$ ,  $b \equiv 0, 2 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv P_{b/2}(6) \pmod{N}$ .

**Conjecture 1.16.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n - b + 1$  such that  $n > 3$ ,  $b \equiv 4, 6 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv -P_{(b-2)/2}(6) \pmod{N}$ .

**Conjecture 1.17.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n + b - 1$  such that  $n > 3$ ,  $b \equiv 0, 2 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv P_{(b-2)/2}(6) \pmod{N}$ .

**Conjecture 1.18.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = b^n + b - 1$  such that  $n > 3$ ,  $b \equiv 4, 6 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(6)$ , thus if  $N$  is prime, then  $S_{n-1} \equiv -P_{b/2}(6) \pmod{N}$ .

**Conjecture 1.19.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$

Let  $N = k \cdot 3^n - 2$  such that  $n > 3$ ,  $k \equiv 1, 3 \pmod{8}$  and  $k > 0$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_{3k}(6)$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_3(6) \pmod{N}$

**Conjecture 1.20.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$

Let  $N = k \cdot 3^n - 2$  such that  $n > 3$ ,  $k \equiv 5, 7 \pmod{8}$  and  $k > 0$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_{3k}(6)$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_1(6) \pmod{N}$

**Conjecture 1.21.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot 3^n + 2$  such that  $n > 2$ ,  $k \equiv 1, 3 \pmod{8}$  and  $k > 0$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_{3k}(6)$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_3(6) \pmod{N}$

**Conjecture 1.22.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot 3^n + 2$  such that  $n > 2$ ,  $k \equiv 5, 7 \pmod{8}$  and  $k > 0$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_{3k}(6)$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_1(6) \pmod{N}$

**Conjecture 1.23.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - c$  such that  $b \equiv 0 \pmod{2}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 1, 7 \pmod{8}$  Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_{(b/2) \cdot \lceil c/2 \rceil}(6) \pmod{N}$

**Conjecture 1.24.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - c$  such that  $b \equiv 0, 4, 8 \pmod{12}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 3, 5 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_{(b/2) \cdot \lfloor c/2 \rfloor}(6) \pmod{N}$

**Conjecture 1.25.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - c$  such that  $b \equiv 2, 6, 10 \pmod{12}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 3, 5 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv -P_{(b/2) \cdot \lfloor c/2 \rfloor}(6) \pmod{N}$

**Conjecture 1.26.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n + c$  such that  $b \equiv 0 \pmod{2}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 1, 7 \pmod{8}$  Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_{(b/2) \cdot \lfloor c/2 \rfloor}(6) \pmod{N}$

**Conjecture 1.27.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n + c$  such that  $b \equiv 0, 4, 8 \pmod{12}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 3, 5 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv P_{(b/2) \cdot \lceil c/2 \rceil}(6) \pmod{N}$

**Conjecture 1.28.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n + c$  such that  $b \equiv 2, 6, 10 \pmod{12}$ ,  $n > bc$ ,  $k > 0$ ,  $c > 0$  and  $c \equiv 3, 5 \pmod{8}$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(6))$ , thus If  $N$  is prime then  $S_{n-1} \equiv -P_{(b/2) \cdot \lceil c/2 \rceil}(6) \pmod{N}$

**Conjecture 1.29.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = 2 \cdot 3^n - 1$  such that  $n > 1$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_3(a)$ , where  $a = \begin{cases} 6, & \text{if } n \equiv 0 \pmod{2} \\ 8, & \text{if } n \equiv 1 \pmod{2} \end{cases}$  thus,  $N$  is prime iff  $S_{n-1} \equiv a \pmod{N}$

**Conjecture 1.30.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = 8 \cdot 3^n - 1$  such that  $n > 1$ . Let  $S_i = P_3(S_{i-1})$  with  $S_0 = P_{12}(4)$  thus,  $N$  is prime iff  $S_{n-1} \equiv 4 \pmod{N}$

**Conjecture 1.31.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot 6^n - 1$  such that  $n > 2$ ,  $k > 0$ ,  $k \equiv 2, 5 \pmod{7}$  and  $k < 6^n$  Let  $S_i = P_6(S_{i-1})$  with  $S_0 = P_{3k}(P_3(5))$ , thus  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.32.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot 6^n - 1$  such that  $n > 2$ ,  $k > 0$ ,  $k \equiv 3, 4 \pmod{5}$  and  $k < 6^n$  Let  $S_i = P_6(S_{i-1})$  with  $S_0 = P_{3k}(P_3(3))$ , thus  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.33.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - 1$  such that  $n > 2$ ,  $k < 2^n$  and

$$\begin{cases} k \equiv 3 \pmod{30} \text{ with } b \equiv 2 \pmod{10} \text{ and } n \equiv 0, 3 \pmod{4} \\ k \equiv 3 \pmod{30} \text{ with } b \equiv 4 \pmod{10} \text{ and } n \equiv 0, 2 \pmod{4} \\ k \equiv 3 \pmod{30} \text{ with } b \equiv 6 \pmod{10} \text{ and } n \equiv 0, 1, 2, 3 \pmod{4} \\ k \equiv 3 \pmod{30} \text{ with } b \equiv 8 \pmod{10} \text{ and } n \equiv 0, 1 \pmod{4} \end{cases}$$

Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(18))$ , then  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.34.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - 1$  such that  $n > 2$ ,  $k < 2^n$  and

$$\begin{cases} k \equiv 9 \pmod{30} \text{ with } b \equiv 2 \pmod{10} \text{ and } n \equiv 0, 1 \pmod{4} \\ k \equiv 9 \pmod{30} \text{ with } b \equiv 4 \pmod{10} \text{ and } n \equiv 0, 2 \pmod{4} \\ k \equiv 9 \pmod{30} \text{ with } b \equiv 6 \pmod{10} \text{ and } n \equiv 0, 1, 2, 3 \pmod{4} \\ k \equiv 9 \pmod{30} \text{ with } b \equiv 8 \pmod{10} \text{ and } n \equiv 0, 3 \pmod{4} \end{cases}$$

Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(18))$ , then  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.35.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$ , where  $m$  and  $x$  are nonnegative integers. Let  $N = k \cdot b^n - 1$  such that  $n > 2$ ,  $k < 2^n$  and

$$\begin{cases} k \equiv 21 \pmod{30} \text{ with } b \equiv 2 \pmod{10} \text{ and } n \equiv 2, 3 \pmod{4} \\ k \equiv 21 \pmod{30} \text{ with } b \equiv 4 \pmod{10} \text{ and } n \equiv 1, 3 \pmod{4} \\ k \equiv 21 \pmod{30} \text{ with } b \equiv 8 \pmod{10} \text{ and } n \equiv 1, 2 \pmod{4} \end{cases}$$

Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{bk/2}(P_{b/2}(3))$ , then  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.36.** Let  $F_p$  be the  $p$ th Fibonacci number. If  $p$  is prime, not 5, and  $M \geq 2$  then  $M^{F_p} \equiv M^{(p-1)(1-\frac{p}{5})/2} \pmod{\frac{M^p-1}{M-1}}$

**Conjecture 1.37.** Let  $b$  and  $n$  be a natural numbers,  $b \geq 2$ ,  $n > 1$  and  $n \notin \{4, 8, 9\}$ . Then  $n$  is prime if and only if  $\sum_{k=1}^n (b^k + 1)^{n-1} \equiv n \pmod{\frac{b^n - 1}{b - 1}}$

**Conjecture 1.38.** If  $q$  is the smallest prime greater than  $\prod_{i=1}^n C_i + 1$ , where  $\prod_{i=1}^n C_i$  is the product of the first  $n$  composite numbers, then  $q - \prod_{i=1}^n C_i$  is prime.

**Conjecture 1.39.** If  $q$  is the greatest prime less than  $\prod_{i=1}^n C_i - 1$ , where  $\prod_{i=1}^n C_i$  is the product of the first  $n$  composite numbers, then  $\prod_{i=1}^n C_i - q$  is prime.

**Conjecture 1.40.** Let  $n$  be an odd number and  $n > 1$ . Let  $T_n(x)$  be Chebyshev polynomial of the first kind and let  $P_n(x)$  be Legendre polynomial, then  $n$  is a prime number if and only if the following congruences hold simultaneously •  $T_n(3) \equiv 3 \pmod{n}$  •  $P_n(3) \equiv 3 \pmod{n}$

**Conjecture 1.41.** Let  $n$  be a natural number greater than two. Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$ . Let  $T_n(x)$  be Chebyshev polynomial of the first kind, then  $n$  is a prime number if and only if  $T_n(x) \equiv x^n \pmod{x^r - 1, n}$ .

**Conjecture 1.42.** Let  $n$  be a natural number greater than two and  $n \neq 5$ . Let  $T_n(x)$  be Chebyshev polynomial of the first kind. If there exists an integer  $a$ ,  $1 < a < n$ , such that  $T_{n-1}(a) \equiv 1 \pmod{n}$  and for every prime factor  $q$  of  $n - 1$ ,  $T_{(n-1)/q}(a) \not\equiv 1 \pmod{n}$  then  $n$  is prime. If no such number  $a$  exists then  $n$  is composite.

**Conjecture 1.43.** Let  $P_a(x) = 2^{-a} \cdot \left( (x - \sqrt{x^2 - 4})^a + (x + \sqrt{x^2 - 4})^a \right)$ . Let  $N = k \cdot b^m \pm 1$  with  $b$  an even positive integer,  $0 < k < b^m$  and  $m > 2$ . Let  $F_n$  be the  $n$ th Fibonacci number and let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{kb/2}(P_{b/2}(F_n))$ , then  $N$  is prime iff there exists  $F_n$  for which  $S_{m-2} \equiv 0 \pmod{N}$ .

**Conjecture 1.44.** Let  $n$  be a natural number greater than one. Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$ . Let  $L_n(x)$  be Lucas polynomial, then  $n$  is a prime number if and only if  $L_n(x) \equiv x^n \pmod{x^r - 1, n}$ .

**Conjecture 1.45.** Let  $b$  and  $n$  be a natural numbers,  $b \geq 2$ , then  $\frac{b^n - 1}{b - 1} \cdot \frac{b^{\sigma(n)} - 1}{b - 1} \equiv b + 1 \pmod{\frac{b^{\varphi(n)} - 1}{b - 1}}$  for all primes and no composite with the exception of 4 and 6.

**Conjecture 1.46.** Let  $b$  and  $n$  be a natural numbers,  $b \geq 2$ , then  $\frac{b^{\varphi(n)} - 1}{b - 1} (b^{\tau(n)} - 1) + b \equiv b^{n-1} \pmod{\frac{b^n - 1}{b - 1}}$  for all primes and no composite with the exception of 4.

**Conjecture 1.47.** Let  $p$  be prime number greater than three and let  $T_n(x)$  be Chebyshev polynomial of the first kind, then  $T_{p-1}(2) \equiv 1 \pmod{p}$  if and only if  $p \equiv 1, 11 \pmod{12}$ .

**Conjecture 1.48.** Let  $p$  be prime number greater than two and let  $T_n(x)$  be Chebyshev polynomial of the first kind, then  $T_{p-1}(3) \equiv 1 \pmod{p}$  if and only if  $p \equiv 1, 7 \pmod{8}$ .

**Conjecture 1.49.** Let  $p$  be prime number greater than three and let  $T_n(x)$  be Chebyshev polynomial of the first kind , then  $T_{p-1}(5) \equiv 1 \pmod{p}$  if and only if  $p \equiv 1, 5, 19, 23 \pmod{24}$

**Conjecture 1.50.** Let  $n$  be an odd natural number greater than one , let  $k$  be a natural number such that  $k \leq n$  , then  $n$  is prime if and only if : 
$$\sum_{i=0}^{k-1} i^{n-1} + \sum_{j=0}^{n-k} j^{n-1} \equiv -1 \pmod{n}$$

**Conjecture 1.51.** Let  $n$  be a natural number greater than one and let  $T_n(x)$  be Chebyshev polynomial of the first kind , then  $n$  is prime if and only if : 
$$\sum_{k=0}^{n-1} 2T_{n-1}\left(\frac{k}{2}\right) \equiv -1 \pmod{n} .$$

**Conjecture 1.52.** Let  $n$  be a natural number greater than one and let  $L_n(x)$  be Lucas polynomial , then  $n$  is prime if and only if : 
$$\sum_{k=0}^{n-1} L_{n-1}(k) \equiv -1 \pmod{n} .$$

**Conjecture 1.53.** Let  $p$  be an odd prime number , let  $R_p(3) = \frac{3^p-1}{2}$  and let  $S_i = S_{i-1}^3 + 3S_{i-1}$  with  $S_0 = 36$  , then  $R_p(3)$  is prime number iff  $S_{p-1} \equiv 36 \pmod{R_p(3)}$  .

**Conjecture 1.54.** Let  $p$  be an odd prime number greater than three , let  $R_p(-3) = \frac{3^p+1}{4}$  and let  $S_i = S_{i-1}^3 + 3S_{i-1}$  with  $S_0 = 36$  , then  $R_p(-3)$  is prime number iff  $S_{p-1} \equiv 36 \pmod{R_p(-3)}$  .

**Conjecture 1.55.** Let  $P_n^{(a)}(x) = \left(\frac{1}{2}\right) \cdot \left( (x - \sqrt{x^2 + a})^n + (x + \sqrt{x^2 + a})^n \right)$  . Given an odd integer  $n (\geq 3)$  and integer  $a$  coprime to  $n$  ,  $n$  is prime if and only if  $P_n^{(a)}(x) \equiv x^n \pmod{n}$  holds .

**Conjecture 1.56.** Let  $n$  be an odd natural number greater than one . Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$  . Let  $P_n(x)$  be Legendre polynomial , then  $n$  is a prime number if and only if  $P_n(x) \equiv x^n \pmod{x^r - 1, n}$  .

**Conjecture 1.57.** Let  $n$  be a natural number greater than one and let  $F_n(x)$  be Fibonacci polynomial , then  $n$  is prime if and only if : 
$$\sum_{k=0}^{n-1} F_n(k) \equiv -1 \pmod{n} .$$

**Conjecture 1.58.** Let  $a_n$  be the least unused prime greater than 3 such that  $(a_n + a_{n-1})/2$  is prime, with  $a_0 = 13$  , then :

1. Every term of this sequence  $a_i$  is prime of the form  $12k + 1$  .
2. Every prime of the form  $12k + 1$  is a member of this sequence .

**Conjecture 1.59.** Let  $m$  and  $n$  be a natural numbers ,  $m \geq 1$  ,  $n > 2$  ,  $n \neq 9$  and  $\gcd(m, n) = 1$  . Then  $n$  is prime if and only if 
$$\sum_{k=1}^{n-1} (2^{mk} - 1)^{n-1} \equiv n \pmod{2^n - 1}$$

**Conjecture 1.60.** Let  $p, q, r$  be three consecutive prime numbers such that  $p \geq 11$  and  $p < q < r$  , then  $\frac{1}{p^2} < \frac{1}{q^2} + \frac{1}{r^2}$  .

**Conjecture 1.61.** Let  $p$  and  $q$  be consecutive prime numbers such that  $p \geq 5$  and  $p < q$ , then  $\left\lfloor \frac{q}{p} - \frac{p}{q} \right\rfloor = 0$ .

**Conjecture 1.62.** Let  $a, n, k$  be natural numbers greater than 0. If  $n$  is a prime number then  $\sum_{d|n} (\sigma_k(d) \cdot a^{n/d}) \equiv 2a \pmod{n}$

**Conjecture 1.63.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $F_n(b) = b^{2^n} + 1$  where  $b$  is an even integer,  $3 \nmid b, 5 \nmid b$  and  $n \geq 2$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(P_{b/2}(8))$ , then  $F_n(b)$  is prime iff  $S_{2^n-2} \equiv 0 \pmod{F_n(b)}$ .

**Conjecture 1.64.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $F_n(b) = b^{2^n} + 1$  where  $b$  is an even integer,  $3 \nmid b, b \equiv 2, 4, 10, 12 \pmod{14}$  and  $n \geq 2$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(P_{b/2}(5))$ , then  $F_n(b)$  is prime iff  $S_{2^n-2} \equiv 0 \pmod{F_n(b)}$ .

**Conjecture 1.65.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $F_n(b) = b^{2^n} + 1$  where  $b$  is an even integer,  $5 \nmid b, b \equiv 2, 4, 10, 12 \pmod{14}$  and  $n \geq 2$ . Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(P_{b/2}(12))$ , then  $F_n(b)$  is prime iff  $S_{2^n-2} \equiv 0 \pmod{F_n(b)}$ .

**Conjecture 1.66.** Let  $a_n = 62a_{n-1} - a_{n-2}$  with  $a_1 = 8$  and  $a_2 = 488$ , let  $b_n = 482b_{n-1} - b_{n-2}$  with  $b_1 = 22$  and  $b_2 = 10582$ , then each member of the sequences  $\{a_n\}$  and  $\{b_n\}$  can be used as an initial value for Inkeri's primality test for Fermat numbers.

**Conjecture 1.67.** Let  $P_m(x) = 2^{-m} \cdot \left( (x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m \right)$

Let  $N = k \cdot b^n - 1$  such that  $n > 2, k < 2^n$  and

$$\begin{cases} k \equiv 27 \pmod{30} \text{ with } b \equiv 2 \pmod{10} \text{ and } n \equiv 1, 2 \pmod{4} \\ k \equiv 27 \pmod{30} \text{ with } b \equiv 4 \pmod{10} \text{ and } n \equiv 1, 3 \pmod{4} \\ k \equiv 27 \pmod{30} \text{ with } b \equiv 8 \pmod{10} \text{ and } n \equiv 2, 3 \pmod{4} \end{cases}$$

Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_{b/2}(P_{b/2}(3))$ , then  $N$  is prime iff  $S_{n-2} \equiv 0 \pmod{N}$

**Conjecture 1.68.** Let  $n$  be a natural number greater than two. Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$ . Let  $H_n(x)$  be Hermite polynomial, then  $n$  is either a prime number or Fermat pseudoprime to base 2 if and only if  $H_n(x) \equiv 2x^n \pmod{x^r - 1, n}$ .

**Conjecture 1.69.** Let  $n$  be an odd natural number greater than one. Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$ . Let  $P_n^{(\alpha, \beta)}(x)$  be Jacobi polynomial such that  $\alpha, \beta$  are natural numbers and  $\alpha + \beta < n$ , then  $n$  is a prime number if and only if  $P_n^{(\alpha, \beta)}(x) \equiv x^n \pmod{x^r - 1, n}$ .

**Conjecture 1.70.** Let  $n$  be an odd natural number greater than one. Let  $r$  be the smallest odd prime number such that  $r \nmid n$  and  $n^2 \not\equiv 1 \pmod{r}$ . Let  $F_n(x)$  be Fibonacci polynomial, then  $n$  is prime if and only if  $F_n(2x) \equiv (1 + x^2)^{\frac{n-1}{2}} \pmod{x^r - 1, n}$ .

**Conjecture 1.71.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $F_n(b) = b^{2^n} + 1$  where  $b$  is an even natural number and  $n \geq 2$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{F_n(b)}\right) = -1$  and  $\left(\frac{a+2}{F_n(b)}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{b/2}(P_{b/2}(a)) \pmod{F_n(b)}$ . Then  $F_n(b)$  is prime if and only if  $S_{2^n-2} \equiv 0 \pmod{F_n(b)}$ .

**Conjecture 1.72.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = k \cdot b^n + 1$  where  $k$  is positive natural number,  $k < 2^n$ ,  $b$  is an even positive natural number and  $n \geq 3$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{N}\right) = -1$  and  $\left(\frac{a+2}{N}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb/2}(P_{b/2}(a)) \pmod N$ . Then  $N$  is prime if and only if  $S_{n-2} \equiv 0 \pmod N$ .

**Conjecture 1.73.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M = k \cdot b^n - 1$  where  $k$  is positive natural number,  $k < 2^n$ ,  $b$  is an even positive natural number and  $n \geq 3$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{M}\right) = 1$  and  $\left(\frac{a+2}{M}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb/2}(P_{b/2}(a)) \pmod M$ . Then  $M$  is prime if and only if  $S_{n-2} \equiv 0 \pmod M$ .

**Conjecture 1.74.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = k \cdot b^n + 1$  where  $k$  is an even positive natural number,  $k < 2^n$ ,  $b$  is an odd positive natural number greater than one and  $n \geq 2$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{N}\right) = -1$  and  $\left(\frac{a+2}{N}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb/2}(a) \pmod N$ . Then, if  $N$  is prime then  $S_{n-1} \equiv a \pmod N$ .

**Conjecture 1.75.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M = k \cdot b^n - 1$  where  $k$  is an even positive natural number,  $k < 2^n$ ,  $b$  is an odd positive natural number greater than one and  $n \geq 2$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{M}\right) = 1$  and  $\left(\frac{a+2}{M}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb/2}(a) \pmod M$ . Then, if  $M$  is prime then  $S_{n-1} \equiv a \pmod M$ .

**Conjecture 1.76.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M_p(a) = \frac{a^p - 1}{a - 1}$  where  $a$  is a natural number greater than one and  $p$  is an odd prime number. Let  $c$  be a natural number greater than two such that  $\left(\frac{c-2}{M_p(a)}\right) = \left(\frac{c+2}{M_p(a)}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_a(S_{i-1})$  with  $S_0 = P_a(c)$ . Then, if  $M_p(a)$  is prime then  $S_{p-1} \equiv P_a(c) \pmod{M_p(a)}$ .

**Conjecture 1.77.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N_p(b) = \frac{b^p + 1}{b + 1}$  where  $b$  is a natural number greater than one and  $p$  is an odd prime number. Let  $c$  be a natural number greater than two such that  $\left(\frac{c-2}{N_p(b)}\right) = \left(\frac{c+2}{N_p(b)}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_b(c)$ . Then, if  $N_p(b)$  is prime then  $S_{p-1} \equiv P_b(c) \pmod{N_p(b)}$ .

**Conjecture 1.78.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M = k \cdot b^n - c$  where  $k, b, n, c$  are natural numbers such that  $k > 0$ ,  $b > 1$ ,  $n > 1$  and  $c > 0$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{M}\right) = -1$  and  $\left(\frac{a+2}{M}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb}(a) \pmod M$ . Then, if  $M$  is prime then  $S_{n-1} \equiv P_{c-1}(a) \pmod M$ .

**Conjecture 1.79.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = k \cdot b^n + c$  where  $k, b, n, c$  are natural numbers such that  $k > 0$ ,  $b > 1$ ,  $n > 1$  and  $c > 0$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{N}\right) = 1$  and  $\left(\frac{a+2}{N}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb}(a) \pmod N$ . Then, if  $N$  is prime then  $S_{n-1} \equiv P_{c-1}(a) \pmod N$ .

**Conjecture 1.80.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M_p(a) = \frac{a^p - 1}{a - 1}$  where  $a$  is a natural number greater than one and  $p$  is an odd prime number. Let  $c$  be a natural number greater than two such that  $\left(\frac{c-2}{M_p(a)}\right) = -1$  and  $\left(\frac{c+2}{M_p(a)}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_a(S_{i-1})$  with  $S_0 = P_a(c)$ . Then, if  $M_p(a)$  is prime then  $S_{p-1} \equiv P_{a-2}(c) \pmod{M_p(a)}$ .

**Conjecture 1.81.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N_p(b) = \frac{b^p + 1}{b + 1}$  where  $b$  is a natural number greater than one and  $p$  is an odd prime number. Let  $c$  be a natural number greater than two such that  $\left(\frac{c-2}{N_p(b)}\right) = -1$  and  $\left(\frac{c+2}{N_p(b)}\right) = 1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0 = P_b(c)$ . Then, if  $N_p(b)$  is prime then  $S_{p-1} \equiv P_{b+2}(c) \pmod{N_p(b)}$ .

**Conjecture 1.82.** Let  $n_1, n_2, \dots, n_k$  be a sequence of  $k$  consecutive odd composite numbers. Let  $\text{gpf}(n_i)$  be the greatest prime factor of  $n_i$ . Then, all  $\text{gpf}(n_i)$ ,  $1 \leq i \leq k$  are mutually different.

**Conjecture 1.83.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = k \cdot b^n + 1$  where  $k$  is positive natural number,  $k < 2^n$ ,  $b$  is a positive natural number greater than one and  $n \geq 3$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{N}\right) = -1$  and  $\left(\frac{a+2}{N}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb^2/2}(a) \pmod{N}$ . Then  $N$  is prime if and only if  $S_{n-2} \equiv -2 \pmod{N}$ .

**Conjecture 1.84.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $M = k \cdot b^n - 1$  where  $k$  is positive natural number,  $k < 2^n$ ,  $b$  is a positive natural number greater than one and  $n \geq 3$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{a-2}{M}\right) = 1$  and  $\left(\frac{a+2}{M}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb^2/2}(a) \pmod{M}$ . Then  $M$  is prime if and only if  $S_{n-2} \equiv -2 \pmod{M}$ .

**Conjecture 1.85.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = 4 \cdot 3^n - 1$  where  $n \geq 3$ . Let  $S_i = S_{i-1}^3 - 3S_{i-1}$  with  $S_0 = P_9(6)$ . Then  $N$  is prime if and only if  $S_{n-2} \equiv 0 \pmod{N}$ .

**Conjecture 1.86.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = 4 \cdot 3^n + 1$  where  $n \geq 3$ . Let  $S_i = S_{i-1}^3 - 3S_{i-1}$  with  $S_0 = P_9(4)$ . Then  $N$  is prime if and only if  $S_{n-2} \equiv 0 \pmod{N}$ .

**Conjecture 1.87.** Let  $P_m(x) = 2^{-m} \cdot ((x - \sqrt{x^2 - 4})^m + (x + \sqrt{x^2 - 4})^m)$ . Let  $N = k \cdot b^n \pm 1$  where  $k$  is positive natural number,  $4 \mid k$ ,  $k < 2^n$ ,  $b$  is an odd positive natural number greater than one and  $n \geq 3$ . Let  $a$  be a natural number greater than two such that  $\left(\frac{2-a}{N}\right) = \left(\frac{a+2}{N}\right) = -1$  where  $(\ )$  denotes Jacobi symbol. Let  $S_i = P_b(S_{i-1})$  with  $S_0$  equal to the modular  $P_{kb^2/4}(a) \pmod{N}$ . Then  $N$  is prime if and only if  $S_{n-2} \equiv 0 \pmod{N}$ .