

Gravity and Cosmic Wavelength Graviton Action Densities

Julian Williams

April 2018

jjawilliams@xtra.co.nz

Abstract

This paper builds on ideas developed in an earlier paper [7] which looked at Standard Model particles built from infinite superpositions borrowing mass from a Higgs type scalar field, and energy from zero point fields. At cosmic wavelengths, zero point energy densities are infinitesimal. To make available and required zero point energies equal, space expands exponentially with time. This balance occurs at a minimum graviton wavenumber k_{\min} (or maximum wavelength). The density $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$, of k_{\min} gravitons has an invariant $K_{Gk_{\min}}$ in all coordinates, for all spacetime. The value of $k_{\min} \approx R_{Horizon}^{-1}$ decreases with cosmic time T , but increases around mass concentrations, inversely with the clock rate $\sqrt{g_{00}}$ in the local metric. It also increases with peculiar velocities relative to comoving coordinates. This paper proposes that this relates with what we define as an “Invariant Four Volume Spherically Symmetric Action Density” at that maximum wavelength k_{\min} which itself varies in direct proportion to the measurement of the averaged local CMB temperature increase. The same proportionality is true for the increase of k_{\min} , and the average CMB temperature in any metric. Borrowed cosmic wavelength quanta are Planck scale zero point action modes, redshifted from a holographic spherical horizon receding at virtually light velocity. This fits an infinitesimally modified General Relativity. We also extend these arguments to include angular momentum and the Kerr Metric. The earlier paper, for simplicity, included only the vast majority ($\psi_{\text{Universe}} * \psi_m + \psi_m * \psi_{\text{Universe}}$) of k_{\min} gravitons around a mass concentration m , We now include the relatively smaller number $\psi_m * \psi_m$ of k_{\min} gravitons emitted by mass m , adding a dimensionless m^2 / r^2 term to $2m / r$ in the metric, which becomes $g_{00} = 1 - m / r - 1.4m^2 / r^3 = g_{rr}^{-1}$ in the non rotating case, and is equivalent to ≈ 700 metres extra distance to the centre of the sun for all the planets, with no change in their orbital periods. The effect of m^2 / r^2 is significant close to Black Holes. The radius of a non-rotating Black Hole increases $\approx 27.5\%$ from $r = 2m$ to $r \approx 2.55m$, but maximum spin Black Holes remain at $r = m$. Only the last cycle or so of black hole mergers would be significantly affected. The extra acceleration due to m^2 / r^2 could slightly speed up mergers for any total angular momentum and mass. This may allow spins to be aligned with their mutual orbits; as thought more probable in some recent mergers. It also increases their apparent mass slightly. The change in the Riemannian tensor due to m^2 / r^2 is of same form, but opposite sign, when compared with the r_Q^2 / r^2 term in electrically charged, Reissner-Nordstrom and rotating Kerr-Newman, metrics. A negative energy massless particle in the Energy-Momentum tensor can generate this term in the metric, just as massless particles in the electromagnetic field do with diagonal stress tensor terms contracting to zero.

1	Introduction.....	4
2	The Expanding Universe and General Relativity	8
2.1	Zero point energy densities are limited	8
2.1.1	Virtual Particles and Infinite Superpositions	8
2.1.2	Virtual graviton density at wavenumber k in a causally connected Universe	9
2.2	Can we relate this to General Relativity?	12
2.2.1	Restricting the range of General Relativity to well inside the horizon.....	12
2.2.2	The Schwarzschild metric near large masses	16
2.3	Angular Momentum and the Kerr Metric	18
2.3.1	Stress tensor sources for spin 2 gravitons & 4 current sources for spin 1	24
2.3.2	Circularly polarized gravitons from corotating space.....	24
2.3.3	Why there are no transverse polarized gravitons in co-rotating coordinates? ...	26
2.3.4	Does our time polarized k_{\min} value in co-rotating coordinates make sense?	26
2.4	The Expanding Universe	29
2.4.1	Holographic horizons and red shifted Planck scale zero point modes.....	30
2.4.2	Plotting available quanta densities, and required quanta densities	33
2.4.3	A possible exponential expansion in a flat matter dominated universe.....	35
2.4.4	The radiation dominated era up to approximately 47,000 years.....	38
2.4.5	Peculiar velocities and non-commoving coordinates	39
2.4.6	Invariant 4 volume cosmic wavelength graviton densities	39
2.4.7	Cosmic wavelength graviton action densities.....	40
2.4.8	If Gravity is due to metric changes then what about exchanged momentum? ..	41
2.5	An Infinitesimal change to General Relativity at Cosmic Scale	42
2.5.1	Is inflation in this proposed scenario really necessary?.....	43
2.5.2	Why do we think virtual particle pairs do not matter?.....	43
2.6	Messing up what was starting to look promising, or maybe not?	45
2.6.1	The k_{\min} virtual gravitons emitted by the mass interacting with itself.....	45
2.6.2	What does this extra term mean for non rotating black holes?.....	46

2.6.3	What does it mean for rotating black holes?.....	46
2.6.4	Determinant of the metric and the k_{\min} graviton constant $K_{Gk_{\min}}$	49
2.6.5	The Reissner-Nordstrom Metric and $\mathbf{m}^2 / \mathbf{r}^2$ terms.....	49
2.6.6	The Kerr-Newman Metric and $\mathbf{m}^2 / \mathbf{r}^2$ terms	51
2.6.7	What is the effect of this term in the solar system?	52
2.6.8	Can we measure this difference?	53
2.6.9	What about the Hulse Taylor binary pulsar, can it show this change?	54
2.6.10	Gravitational Wave observations of Black Hole mergers.....	55
2.7	Revisiting some other aspects of the first paper	56
2.7.1	Infinitesimal rest masses	56
2.7.2	Redshifted zero point energy from the horizon behaves differently to local	56
2.7.3	Revisiting the building of infinite superpositions.....	57
2.8	Gravitational Waves	57
2.8.1	Constant transverse areas in low energy waves.....	58
2.8.2	What might happen in high energy waves?	58
2.8.3	No connection between wave frequency and radiated quanta energy	59
2.9	Four Vectors and Four Volume Action Densities	59
2.9.1	Graviton densities represented as invariant 4 velocities	59
2.9.2	Four volumes in changing metrics	60
3	Some Loose Ends.....	61
3.1.1	Preferred Frames	61
3.1.2	Solar System Constraints and do our proposed changes fit?	61
3.1.3	Action Principles and the Einstein Field Equations.....	61
4	Conclusions.....	62
5	References.....	64

1 Introduction

The universe we live in is currently described by two models: “The Standard Model of Particle Physics” and “The Standard Model of Cosmology”. While the Standard Model of Particle Physics is remarkably accurate in its predictions, is mathematically elegant and apparently complete in many respects; it is also at the same time incomplete. Supersymmetry, proposed to solve some of its issues, is at this date not panning out as expected, with some physicists questioning whether supersymmetry is the hoped for answer. Neutrinos have a small mass, which the Standard Model does not predict. Gravity is not included and there is no force unification without supersymmetry.

The Standard Model of Cosmology or “The Lambda-CDM Model” requires Dark Energy to explain the accelerating expansion of space, with no good understanding of what causes it. Without initial inflation, there is no good explanation of why space is Euclidean, or flat on average, or why regions initially causally separated are so homogeneous; but there is still no widely accepted understanding of what causes this inflation.

In the first paper [7] we attempted to show that the fundamental particles of the Standard Model can be built from infinite superpositions apart from infinitesimal, but important differences. They all had mass which naturally divided into two sets. Spin 2 gravitons, spin 1 photons and gluons, all had infinitesimal mass approximately the inverse (always) of the causally connected horizon radius *of the observable universe* $R_{ov} \approx 46 \times 10^9$ light years or $\approx 10^{-33} eV$. (This value is close to some recent proposals [8] giving gravitons a mass of $< 10^{-33} eV$ to explain the accelerating expansion of the universe.) The rest had finite masses of micro electron volts upwards. Infinite superpositions are always built in some rest frame in which they had no net momentum \mathbf{p} , but only \mathbf{p}^2 terms. In the “infinitesimal” mass set this rest frame can be, and usually is, travelling at virtually light velocity, as seen from our usual (nearly) comoving frame. We also divided the world of all interactions into two sets.

(a) **Primary Interactions** are purely virtual. They build all the fundamental particles in the form of infinite superpositions. We can not see any direct signs of primary interactions.

(b) **Secondary Interactions** are all the others that occur between fundamental particles, both virtual and real. They are the real world of experiments that the Standard Model is all about.

The rules for borrowing energy from zero point fields can be different for both (a) & (b). Primary interactions are between spin zero particles borrowed from a Higgs type scalar field and the zero point fields.

In the 1970's models were proposed with preons as common building blocks of leptons and quarks [10] [11] [12] [13] In contrast with the spin zero particles in this paper, most of these

earlier models used *real spin 1/2* building blocks. As in these earlier models, this paper also calls the common building blocks preons; *but here the preons are both virtual, and spin zero bosons*. There are only three preons; red, green and blue, all with positive electric charge. There are also only three anti preons; antired, antigreen and antiblue, all negatively charged. As preons are spin zero, there can be no weak charge involved in primary interactions. This is all explained more fully in the first paper. These preons build all spin 1/2 leptons and quarks, spin 1 gluons, photons, W & Z particles, plus spin 2 gravitons. This is in contrast to only leptons and quarks in earlier preon models. We found that the fundamental forces do not unite at the Planck energy cutoff of superpositions. They relate with each other in a manner that meshes nicely with the Standard Model, but do not relate with versions including supersymmetry. In the final third of this first paper we tried to fit infinite superpositions with General Relativity and The Standard Model of Cosmology. Because these infinite superpositions borrow energy from zero point fields, which have virtually zero density at cosmic wavelengths; it only works if space expands exponentially with time, and if space is flat on average. The equations we derived looked the same for all comoving observers. Regardless of an observer's position in the universe this expansion looked the same apart from the effect of initial quantum fluctuations at the start. This may remove one of the key reasons for inflation. The universe in this proposed scenario should look the same, and be flat on average, for all observers with or without inflation. Even to observers near the horizon or outside it. The properties and equations controlling distant universes should be identical to ours and there would be no multiverses which are a natural endpoint of inflation. We found that all particles have a maximum wavelength that is approximately the same as the size of the causally connected universe at any cosmic time T . At this maximum wavelength there is a minimum wavenumber we called k_{\min} . We found that the density of k_{\min} gravitons at this maximum wavelength was always proportional to a universal invariant which we labelled $K_{Gk_{\min}}$. The same invariance applies to action densities@ k_{\min} . We connected this with infinitesimally modified GR equations locally, but significant implications at cosmic scale.

Solutions to $R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}[T_{\mu\nu} - T_{\mu\nu}(\text{Background})]$ are consistent with $\rho_{Gk_{\min}} = K_{Gk_{\min}}dk_{\min}$

where $K_{Gk_{\min}}$ is invariant in all coordinates through all Spacetime, and $\rho_{Gk_{\min}}$ is the density of maximum wavelength k_{\min} gravitons, but wavenumber k_{\min} depends on local clocks or $\sqrt{g_{00}}$.

These solutions are equivalent to what we defined as a

"Spherically Symmetric Invariant Four Volume k_{\min} Graviton Action Density"

In comoving coordinates $T_{\mu\nu}(\text{Background})$ has just one component $T_{00} = \rho_U$ the average density of the universe, or only a few hydrogen atoms per cubic metre. This modification limits the range of GR to scales smaller than the radius of the universe and guarantees flatness on average regardless of the value of Ω . The overall exponential expansion of space

is controlled by equations balancing the zero point action available at cosmic wavelenths to that borrowed by infinite superpositions. Dark Energy is not required for this accelerating expansion, but Dark Matter is still required inside galaxies because of centrifugal forces due to their fast rotation. We found a spin 2 (massive graviton type) infinite superposition as a possible dark matter candidate, that won't show up in current weak interaction type searches. Spacetime has to warp in accord with GR around mass concentrations to make available the zero point energy required by their extra cosmic wavelength gravitons. The first paper only looked at long range gravitons emitted by a mass interacting with the rest of the mass in the universe $(\psi_{\text{Universe}} * \psi_m + \psi_m * \psi_{\text{Universe}})$, it ignored the relatively smaller number of cosmic wavelength gravitons emitted by the mass interacting with itself $(\psi_m * \psi_m)$. This paper looks at the $(\psi_m * \psi_m)$ term, which is most significant near black holes. It unfortunately messes up a nice agreement with the Schwarzschild solution by adding a dimensionless m^2 / r^2 term to the usual $2m / r$ in the metric. The effect is equivalent to increasing the distance to the centre of the sun by about 700 metres for all the planets, assuming no change in their orbital periods. This may well be measurable in the foreseeable future. But a non rotating black hole radius increases approximately 27.5% from $r = 2m$ to $r \approx 2.55m$. With angular momentum this becomes the modified ergosphere maximum diameter, but the radius of a maximum spin black hole is unchanged at $r = m$.

These changes, mainly close to Black holes, initially appear to introduce a tension with the field equations of General Relativity. However in Section 2.6 we look at the similarities between this m^2 / r^2 term and the dimensionless r_Q^2 / r^2 terms of both the Reissner-Nordstrom and Kerr-Newman charged black hole metrics. Their effects on the Riemannian curvature tensor are of the same form, but opposite sign to that from an m^2 / r^2 term. There are no covariance problems with electromagnetic field massless particles. If we include in the Stress-Energy tensor a massless negative energy particle, covariance is similarly maintained. It is a bit like, but not the same as, including negative gravitational field energy; which Einstein specifically excluded because of covariance problems. This m^2 / r^2 term increases the merging energies for a fixed mass at any radius. The same applies to the gravitational wave radiated energies. It increases the apparent masses of merging black holes above those derived with only an m / r term. It introduces extra radial acceleration, and may speed up final mergers of black holes for any total mass and spin. This may relate with the merger [26], where if General Relativity holds to the horizon, spins were found unlikely to be aligned with their mutual orbits, as current astrophysics theory had expected. The accuracy of these observations will almost certainly increase with time, either confirming this or not.

In the *rest frame* in which the particles are built from infinite superpositions the spin zero preons are born with *zero momentum*. This means they are *born with infinite wavelength*

allowing the possibility that they *can borrow zero point energy from an infinite distance*. We proposed that they borrow redshifted Planck energy zero point quanta from a holographic horizon receding at light like velocities relative to comoving coordinates instantaneously on that horizon. This is necessary because at cosmic wavelengths of $\approx R_{OU}$ the density of zero point modes is almost zero, and insufficient to build all the fundamental particles; gravitons in particular. The Riemannian spacetime curvature tensor is controlled by the need to keep both “**The Graviton $K_{Gk_{min}}$, & Action Density @ k_{min} or, $K(k_{min} \text{ Action})$ ” invariant. For the sake of clarity, this paper repeats a heavily revised portion of the final third of that first paper, but now includes cosmic wavelength gravitons emitted by the mass interacting with itself ($\psi_m * \psi_m$), the effects of angular momentum, and gravitational waves.**

Einstein published his General Theory of Relativity [1] 100 years ago. There have been many attempts over the intervening years to modify it with different goals in mind. A dissertation by Germanis [2] discusses some of these modifications [3] [4] [5] [6]. On its initial publication it was criticized for not including gravitational field energy, but over the last century, many physicists have tried unsuccessfully to covariantly do this. The modifications proposed in these papers, are the extra m^2 / r^2 term in the metric with its large effect close to black holes, and our equations being consistent with $T_{\mu\nu}$ changing to $T_{\mu\nu} - T_{\mu\nu}(\text{Background})$, which has an infinitesimal effect locally, but significant implications at cosmic scale. The Λ -CDM Model of Cosmology is based on General Relativity as it is currently interpreted. It requires Dark energy to accelerate the expansion, it requires $\Omega \approx 1$, it requires “Inflation” so that regions initially out of causal contact can have (almost) uniform properties, and to produce the observed average flatness. The ideas proposed in these two papers, may well eliminate the need for these requirements. If both $K_{Gk_{min}}$ & $K(k_{min} \text{ Action})$ are invariant at all points in spacetime, the equations controlling the expansion of space and the warping of spacetime around mass concentrations are the same for all observers in this universe and should also be for those far away. There should be no multiverses and no need for anthropic arguments. The original arguments behind the Cosmological Model, of uniformity on average everywhere, should be absolutely true. While the arguments proposed in these two papers are radical, and no doubt contain many errors, the principles behind them may well suggest a possible different path forward. But much tidying up, and putting these ideas on a more rigorous foundation, would be required. It is almost certainly to our evolutionary advantage that what we call established, or collective knowledge, or paradigms particularly in science, change slowly; and only after evidence for change builds to a tipping point. In the end however, science, as it always has in the past, slowly but surely progresses towards the simplest explanations. Finally, so that these ideas are accessible to the widest possible audience, many more details than required by experts in the field are included, with the simplest possible explanations.

2 The Expanding Universe and General Relativity

2.1 Zero point energy densities are limited

If fundamental particles can be built from energy borrowed from the spatial component of zero point fields, and this energy source is limited, (particularly at cosmic wavelengths) there must be implications for the maximum possible densities of these particles. In section 2.2.3 in [7] we discussed how preons build massive fundamental particles, and are born from a Higg's type scalar field with zero momentum in the laboratory rest frame. Infinitesimal mass particles such as gravitons borrow their mass from the time component of the same zero point fields. In this frame they have infinite wavelength and can borrow from anywhere in the universe. This suggests there should be little effect on localized densities, but possibly on overall average densities in any universe. So which fundamental particle is there likely to be most of? Working in Planck, or natural units with $G=1$ and a graviton coupling between Planck masses of one, there are approximately $M \approx 10^{61}$ Planck masses within the causally connected observable universe. Their average distance apart is approximately the radius R_{OH} of this region. There should be approximately $M^2 \approx 10^{122}$ virtual gravitons with wavelengths of the order of radius R_{OH} within this same volume. No other fundamental particle is likely to approach these values, for example the number of virtual photons *of this extreme wavelength* is much smaller. (Virtual particles emerging from the vacuum are covered in section 2.5.2) If this density of virtual gravitons needs to borrow more energy from the zero point fields than what is available at these extreme wavelengths does this somehow control the maximum possible density of a causally connected universe?

2.1.1 Virtual Particles and Infinite Superpositions

Looking carefully at section 3.3 in [7] we showed there that, for all interactions between fundamental particles represented as infinite superpositions, the actual interaction is between only single wavenumber k superpositions of each particle. *We are going to conjecture that a virtual particle of wavenumber k for example is just such a single wavenumber k member.* Only if we actually measure the properties of real particles do we observe the properties of the full infinite superposition. The full properties do not exist until measurement, just as in so many other examples in quantum mechanics. We will use this conjectured virtual property from here on when looking at the probability density of virtual gravitons of the maximum cutoff wavelength. These virtual gravitons would be a superposition of the three modes $n=3,4,5$ of a single wavenumber k , as in Table 4.3.1 in [7]. Time polarized, or spherically symmetric, versions we conjecture (See section 2.3.1) are a further equal $(1/\sqrt{3})$ superposition of $m=-2,0,+2$ states of the above $n=3,4,5$ mode superpositions. A spin 2 graviton in an $m=+2$ state is simply a superposition of the three modes $n=3,4,5$ as above but all in an $m=+2$ state. This is explained in the first paper section 3.2.2 page 30 [7].


2.1.2 Virtual graviton density at wavenumber k in a causally connected Universe

From here on we will use Planck units $\hbar = c = G = 1$. When we looked at scalar potentials between electric charges in [7] we used time polarized virtual photons in a simple example that works for both photons and gravitons, using field energy densities rather than exchanged 4 momentum. However in quantum field theory, scalar or coulomb forces are due to exchanged 3 momentum with time polarized photons (or gravitons), but gravitational forces appear to be due to changes in the metric, and not exchanged 3 momentum. However, in the meantime we will continue to use time polarized graviton densities as we did in [7] and discuss exchanged momentum in section 2.4.8. Also if observers, at the centre of their universe, are moving at a peculiar velocity β_p relative to commoving coordinates, the average velocity of all mass in the universe is moving with the opposite velocity $-\beta_p$. Over all thin spherical shells of matter at the same radius, they can choose pairs of small areas in opposite directions. The spatially polarized vectors due to their velocity exactly cancel for all pairs. Central observers see only time polarized gravitons regardless of peculiar velocity. This is the same as magnetic vectors cancelling at the centre of long current conductors. Spin 2 gravitons couple to the stress tensor in contrast to 4 currents for spin 1. Because of the above the only important term is the mass/energy density T_{00} (or simply ρ) or its transformed value in any other coordinates, as flow of momentum density terms cancel out. We can thus use the same wavefunctions as we did in [7] Eq's. (3.4.1) for time polarized gravitons and photons.

$$\text{Using } (\psi_1 + \psi_2) * (\psi_1 + \psi_2) = (\psi_1 * \psi_1) + (\psi_1 * \psi_2 + \psi_2 * \psi_1) + (\psi_2 * \psi_2)$$

$$\text{The interaction term is } \psi_1 * \psi_2 + \psi_2 * \psi_1 = \frac{4k}{4\pi r_1 r_2} e^{-k(r_1+r_2)} \cos k(r_1 - r_2) \quad (2.1. 1)$$

Figure 2.1. 1



Where r_1 & r_2 are the distances to some point P from two charges or masses 1 & 2, and we are looking at the interaction at point P as in Figure 2.1. 1. Equation (2.1. 1) is strictly true *only in flat space* but it is still approximately true in small curvatures when $2m/r \ll \ll 1$, which we will assume applies almost everywhere throughout the universe except in the infinitesimal fraction of space close to black holes. In both sections 3.4 & 3.5 in [7] for simplicity and clarity, we delayed using coupling constants and emission probabilities in the wavefunctions until necessary. We do the same here. There is also some minimum wavenumber k which we call k_{\min} where for all $k < k_{\min}$ there is insufficient zero point energy available. At this maximum wavelength $k_{\min} \approx 1/R_{OU}$ ($= 1/R_{\text{ObsevableUniverse}}$), for all $k < k_{\min}$, Eq. (2.1. 1) cuts off exponentially. Section 6 in [7] shows gravitons have infinitesimal rest

mass m_0 , of the same order as this minimum wavenumber k_{\min} . At these extreme k values this rest mass must be included in the wavefunction negative exponential term. It is normally irrelevant for infinitesimal masses. Section 6.2 in [7] looks at $N = 2$ infinitesimal rest masses finding $\langle K_{k_{\min}} \rangle^2 \approx 1$. Using Equ's. (3.1.11) & (3.2.10) in [7] with $\hbar = c = 1$

$$\langle K_{k_{\min}} \rangle^2 = \frac{s \langle n \rangle^2 k_{\min}^2}{2m_0^2} \approx 1 \text{ \& for spin 2 gravitons } \langle K_{k_{\min}} \rangle^2 = \frac{\langle n \rangle^2 k_{\min}^2}{m_0^2} \approx 1 \text{ or } m_0 = \langle n \rangle k_{\min} \quad (2.1. 2)$$

From Table 4.3.1 in [7] we find

$$\text{For } N = 2 \text{ spin 2 gravitons } \langle n \rangle \approx 3.33 \text{ so that } m_0 \approx 3.33k_{\min} \quad (2.1. 3)$$

This virtual mass m_0 increases the ΔE term in $\Delta E \cdot \Delta T \approx \hbar / 2$ for the virtual graviton from $\Delta E = k$ to $\Delta E = \sqrt{k^2 + m_0^2}$. This reduces the region $r \approx \Delta T \approx \Delta E^{-1}$ over which it can be found, which is controlled by the exponential decay term e^{-kr} in its wavefunction. This term becomes $e^{-r\sqrt{k^2+m_0^2}}$ as we approach k_{\min} . Using Eq. (2.1. 3) we can define a k' such that

$$k' = \sqrt{k^2 + m_0^2} \approx \sqrt{k^2 + 3.33^2 k_{\min}^2} \text{ and } k'_{\min} \approx \sqrt{k_{\min}^2 + 11.09k_{\min}^2} \approx 3.477k_{\min} \quad (2.1. 4)$$

A normalized virtual massless graviton wavefunction is $\psi = \sqrt{\frac{2k}{4\pi}} \frac{e^{-kr+ikr}}{r}$ see Eq. (3.4.1) in [7] and for infinitesimal mass gravitons this becomes using Eq. (2.1. 4)

$$\text{A massless } \psi = \sqrt{\frac{2k}{4\pi}} \frac{e^{-kr+ikr}}{r} \text{ becomes with infinitesimal mass } \sqrt{\frac{2k'}{4\pi}} \frac{e^{-k'r+ikr}}{r} \quad (2.1. 5)$$

Thus the massless interaction term in Eq. (2.1. 1) becomes with this infinitesimal mass m_0

$$\psi_1^* \psi_2 + \psi_2^* \psi_1 = \frac{4k'}{4\pi r_1 r_2} e^{-k'(r_1+r_2)} \cos k(r_1 - r_2) \quad (2.1. 6)$$




Figure 2.1. 2

Let point P be anywhere in the interior region of a universe as in Figure 2.1. 2. Let the average density (or its equivalent transformed value) be ρ_U (subscript U for universe) Planck masses/energy density per unit volume. Consider two spherical shells initially in commoving coordinates around the central point P of radii r_1 & r_2 and thicknesses dr_1 & dr_2 with masses $dm_1 = \rho_U dv_1 = 4\pi r_1^2 dr_1 \rho_U$ & $dm_2 = \rho_U dv_2 = 4\pi r_2^2 dr_2 \rho_U$ Now we expect graviton coupling α_G to be 1 between Planck masses, but we will assume we don't know this and solve it later to find it appears to be true.

Temporarily define Secondary graviton coupling between Planck masses as α_G ? (2.1. 7)

In section 3.4.1 in [7] Eq. (3.4.3) used a scalar emission probability $(2\alpha/\pi)(dk/k)$ which becomes $(2\alpha_G/\pi)(dk/k)$ between Planck masses. But we must include an exponential cutoff $(1 - \text{Exp}[-0.61k^2/k_{\min}^2])$ near k_{\min} (See section 6.2 in [7]). Now distant galaxies recede at light like and greater velocities, but quantum interactions are instantaneous over all space. Thus, as we integrate over radii r_1 & $r_2 = 0 \rightarrow \infty$, we will assume we can use the same equations as if space is not expanding. Using coupling probability $(1 - \text{Exp}[-0.61k^2/k_{\min}^2])(2\alpha_G/\pi)(dk/k)$ between Planck masses we can integrate over both radii r_1 & r_2 ; but to avoid counting all pairs of masses dm_1 & dm_2 twice, we divide the integral by two. The total probability density of virtual gravitons at any point P in the universe at wavenumber k is using Eq. (2.1. 6)

$$\begin{aligned} \rho_{Gk} &= \frac{\rho_U^2}{2} \alpha_G (1 - e^{-0.61k^2/k_{\min}^2}) \frac{2}{\pi} \frac{dk}{k} \iint_{0 \rightarrow \infty} 4\pi r_1^2 dr_1 \cdot 4\pi r_2^2 dr_2 \cdot \frac{4k'}{4\pi r_1 r_2} e^{-k'(r_1+r_2)} \cos k(r_1 - r_2) \\ &= 16\alpha_G (1 - e^{-0.61k^2/k_{\min}^2}) \rho_U^2 \frac{k'}{k} dk \iint_{0 \rightarrow \infty} r_1 r_2 e^{-k'(r_1+r_2)} \cos k(r_1 - r_2) \cdot dr_1 \cdot dr_2 \end{aligned}$$

Expanding $\cos k(r_1 - r_2) = \cos kr_1 \cos kr_2 + \sin kr_1 \sin kr_2$, and then using:

$$\int_{r=0}^{r=\infty} r \text{Exp}(-k'r) \cos(kr) dr = \frac{k'^2 - k^2}{(k'^2 + k^2)^2} \quad \text{and} \quad \int_{r=0}^{r=\infty} r \text{Exp}(-k'r) \sin(kr) dr = \frac{2k'k}{(k'^2 + k^2)^2}$$

$$\begin{aligned} \rho_{Gk} &= 16\alpha_G (1 - e^{-0.61k^2/k_{\min}^2}) \rho_U^2 \frac{k'}{k} dk \frac{(k'^2 + k^2)^2}{(k'^2 + k^2)^4} \\ &= 16\alpha_G (1 - e^{-0.61k^2/k_{\min}^2}) \rho_U^2 \frac{k'}{k} dk \frac{1}{(k'^2 + k^2)^2} \end{aligned} \quad (2.1. 8)$$

From Eq.(2.1. 4) $k' = \sqrt{k^2 + m_0^2} \approx \sqrt{k^2 + 11.09k_{\min}^2}$ and we can write Eq. (2.1. 8) as

$$\begin{aligned} \rho_{Gk} &= 16\alpha_G \rho_U^2 (1 - e^{-0.61k^2/k_{\min}^2}) \frac{\sqrt{k^2 + 11.09k_{\min}^2}}{(2k^2 + 11.09k_{\min}^2)^2} \frac{dk}{k} \\ &= 16\alpha_G \frac{\rho_U^2}{k_{\min}^4} dk \frac{(1 - e^{-0.61x^2}) \sqrt{x^2 + 11.09}}{x(2x^2 + 11.09)^2} \quad \text{where} \quad x = \frac{k}{k_{\min}} \end{aligned}$$

$$\text{Wavelength } k \text{ Probability Density } \rho_{Gk} \approx \frac{0.149\alpha_G\rho_U^2}{k_{\min}^4} dk \left[\frac{108 (1 - e^{-0.61k^2/k_{\min}^2}) \sqrt{x^2 + 11.09}}{x (2x^2 + 11.09)^2} \right] \quad (2.1.9)$$

Where the blue square bracket is 1 when $k / k_{\min} = x = 1$

$$\text{Cutoff wavelength Probability Density } \rho_{Gk_{\min}} \approx \frac{0.149\alpha_G\rho_U^2}{k_{\min}^4} dk_{\min} \text{ when } \frac{k}{k_{\min}} = x = 1$$

As we think $K_{G_{\min}}$ will prove to be a spacetime invariant, we will write this as follows.

$$\text{Cutoff wavelength Probability Density } \rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min} \text{ where } K_{Gk_{\min}} \approx \frac{0.149\alpha_G\rho_U^2}{k_{\min}^4} \quad (2.1.10)$$

2.2 Can we relate this to General Relativity?

The above assumes a homogeneous universe that is essentially flat on average. At any cosmic time T it assumes there is always some value k_{\min} where the borrowed energy density $E_{Gk_{\min}} = E_{ZP_{\min}}$ the available zero point energy density @ k_{\min} . We have initially assumed commoving coordinates, but at peculiar velocities our spherical shells become ellipses and Eq. (2.1. 10) or $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ should remain true at any peculiar velocity, also in all coordinates as we hope to show later. So what happens if we put a small mass concentration $+m_1$ at some point? The gravitons it emits must surely increase the local density of k_{\min} gravitons upsetting the balance between borrowed energy and that available. However General Relativity tells us that near mass concentrations the metric changes, radial rulers shrink and local observers measure larger radial lengths. This expands locally measured volumes, lowering their measurement of the background $\rho_{Gk_{\min}}$. But clocks also slow down, increasing the locally measured value of k_{\min} . Let us look at whether we can relate these changes in rulers and clocks with the $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ of Eq. (2.1. 10).

2.2.1 Restricting the range of General Relativity to well inside the horizon

Let us refer back to Eq. (3.4.2) in [7] and the steps we took to derive it; but now including $k' = \sqrt{k + m_0} \approx \sqrt{k^2 + 11.09k_{\min}^2}$ as in Eq. (2.1. 4)

$$\psi_1 * \psi_2 + \psi_2 * \psi_1 = \frac{4k'}{4\pi r_1 r_2} e^{-k'(r_1+r_2)} \cos[k(r_1 - r_2)] \quad (2.2. 1)$$

And assume that *space has to be approximately flat* with errors $\propto 1 - (1 - 2m/r)^{1/2} \approx m/r$. If we now focus on Figure 2.1. 1, equation (2.2. 1) is the probability that an infinitesimal mass virtual graviton of wavenumber k is at the point P if all other factors are one. Let us now put

a mass of m_1 Planck masses as in Figure 2.2. 1. Also assume that the *point P* is reasonably close to mass m_1 (in relation to the horizon radius) at distance r_1 as in Figure 2.2. 1 and the vast majority of the rest of the mass inside the causally connected or observable horizon R_{OH} is at various radii r , equal to r_2 of Eq.(2.2. 1) where $r_2 = r \gg r_1$ and thus $\cos[k(r_1 - r)] \approx \cos(-kr)$ & $e^{-k'(r_1+r_2)} \approx e^{-kr}$. This is equivalent to localizing General Relativity to much smaller than horizon radii. *Only under these conditions can we approximate Eq. (2.2. 1) as*

$$\psi_1 * \psi_2 + \psi_2 * \psi_1 \approx \frac{4k'}{4\pi r_1 r} e^{-kr} \cos(-kr) \quad (2.2. 2)$$




Figure 2.2. 1

The background gravitons are time polarized and we are effectively looking at the scalar potential of this central mass relative to the rest of the universe, so this is a time polarized or scalar interaction with no directional effects due to spatial polarization. We can consider simple spherical shells (again initially in commoving coordinates) of thickness dr and radius r around a central observer at the point P which have mass $dm = \rho_U 4\pi r^2 dr$. At each radius r the coupling factor including an exponential cutoff is $(1 - e^{-0.61k^2/k_{min}^2})(2\alpha_G / \pi)(dk / k)$ between Planck masses. Again assuming instantaneous quantum coupling as if space is not expanding:

$$\text{Coupling factor} = (1 - e^{-0.61k^2/k_{min}^2}) \frac{2\alpha_G m_1}{\pi} dm \frac{dk}{k} = (1 - e^{-0.61k^2/k_{min}^2}) \frac{2\alpha_G m_1}{\pi} \rho_U 4\pi r^2 dr \frac{dk}{k} \quad (2.2. 3)$$

$$\begin{aligned} \text{Including this coupling factor } & (1 - e^{-0.61k^2/k_{min}^2}) \left(\frac{2\alpha_G m_1}{\pi} \frac{dk}{k} \rho_U 4\pi r^2 dr \right) (\psi_1 * \psi_2 + \psi_2 * \psi_1) \\ & \approx (1 - e^{-0.61k^2/k_{min}^2}) \left(\frac{2\alpha_G m_1}{\pi} \frac{dk}{k} \rho_U 4\pi r^2 dr \right) \left(\frac{4k'}{4\pi r_1 r} e^{-kr} \cos(-kr) \right) \\ & \approx (1 - e^{-0.61k^2/k_{min}^2}) \alpha_G \frac{m_1}{r_1} \frac{8\rho_U}{\pi} \frac{k' dk}{k} r e^{-kr} \cos(-kr) dr \end{aligned} \quad (2.2. 4)$$

This is virtual graviton density at point P due to each spherical shell. (ignoring the relatively *small number of particularly* k_{\min} gravitons emitted by mass m_1 itself ($\psi_{m_1} * \psi_{m_1}$) see section 2.6.1). Integrating over radius $r = 0 \rightarrow \infty$ the virtual graviton density at wavenumber k using Eq's. (2.1. 4 & (2.2. 4)

$$\begin{aligned}\Delta\rho_G &= (1 - e^{-0.61k^2/k_{\min}^2})\alpha_G \frac{m_1}{r_1} \frac{8\rho_U}{\pi} \frac{k'dk}{k} \int_0^\infty r e^{-kr} \cos(-kr) dr \\ &= (1 - e^{-0.61k^2/k_{\min}^2})\alpha_G \frac{m_1}{r_1} \frac{8\rho_U}{\pi} \frac{k'dk}{k} \left[\frac{(k'^2 - k^2)}{(k'^2 + k^2)^2} \right]\end{aligned}\quad (2.2. 5)$$

Now $k'^2 = k^2 + m_0^2 \approx k^2 + 11.09k_{\min}^2$ and if $k = k_{\min}$ then $k_{\min}'^2 \approx 12.09k_{\min}^2$ & so when $k = k_{\min}$:

$$\Delta\rho_{Gk_{\min}} \approx (1 - e^{-0.61k^2/k_{\min}^2})\alpha_G \frac{m_1}{r_1} \frac{8\rho_U}{\pi} \frac{\sqrt{12.09k_{\min}^2} dk_{\min}}{k_{\min}} \left[\frac{(12.09k_{\min}^2 - k_{\min}^2)}{(12.09k_{\min}^2 + k_{\min}^2)^2} \right]\quad (2.2. 6)$$

$$\Delta\rho_{Gk_{\min}} = (\psi_{\text{Universe}} * \psi_{m_1} + \psi_{m_1} * \psi_{\text{Universe}}) \approx (1 - e^{-0.61k^2/k_{\min}^2}) \cdot \alpha_G \frac{m_1}{r_1} 0.573 \frac{\rho_U}{k_{\min}^2} dk_{\min}$$

Equation (2.1. 10) hypothesizes $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$. In a metric far from masses where $g_{\mu\nu} = \eta_{\mu\nu}$, k_{\min} has its lowest value. As we approach any mass k_{\min} increases to k_{\min}'' where we use green double primes when $g_{\mu\nu} \neq \eta_{\mu\nu}$ to avoid confusion with the k' & k'_{\min} of Eq. (2.1. 4). At a radius r from mass m the Schwarzschild metric is $(1 - 2m/r)^{\pm 1/2}$ for the time and radial terms. Radial rulers shrink and clocks slow, measured local volume V & frequency k_{\min} both increase as $\approx 1 + \frac{m}{r}$.

$$\text{Thus both } \frac{V + \Delta V}{V} \approx 1 + \frac{m}{r} \text{ and also } \frac{k_{\min}''}{k_{\min}} \approx 1 + \frac{m}{r} \text{ if } r \gg \gg m$$

$$\text{Then using } \rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min} \text{ \& } \rho_{Gk_{\min}}'' = K_{Gk_{\min}} dk_{\min}''$$

$$1 + \frac{m}{r} \approx \frac{V + \Delta V}{V} = 1 + \frac{\Delta V}{V} \approx \frac{k_{\min}''}{k_{\min}} = \frac{dk_{\min}''}{dk_{\min}} = \frac{\rho_{k_{\min}}''}{\rho_{k_{\min}}}\quad (2.2. 7)$$

So in this metric the total number of k_{\min} gravitons is the original ($g_{\mu\nu} = \eta_{\mu\nu}$) $\rho_{Gk_{\min}}$ of Eq. (2.1. 10) plus the extra due to a local mass of Eq. (2.2. 6) but we have to divide this number by the increased volume to get the new density $\rho_{Gk_{\min}}'' \approx (1 + \frac{m}{r})\rho_{Gk_{\min}}$. Thus using Eq. (2.2. 7)

$$\text{The new } \rho_{Gk_{\min}}'' \approx \frac{\rho_{Gk_{\min}} + \Delta\rho_{Gk_{\min}}}{1 + \Delta V/V} \approx \frac{\rho_{Gk_{\min}} + \Delta\rho_{Gk_{\min}}}{(1 + m/r)} \approx (1 + m/r)\rho_{Gk_{\min}}$$

$$\rho_{Gk_{\min}} + \Delta\rho_{Gk_{\min}} = (1 + m/r)^2 \rho_{Gk_{\min}} \approx (1 + 2m/r)\rho_{Gk_{\min}} \quad (\text{if } r \gg \gg m)$$

$$\frac{\rho_{Gk \min} + \Delta\rho_{Gk \min}}{\rho_{Gk \min}} \approx 1 + \frac{2m}{r}$$

$$\frac{\Delta\rho_{Gk \min}}{\rho_{Gk \min}} \approx 2\frac{m}{r} \quad (2.2. 8)$$

We can now put Eq's.(2.2. 6) into Eq. (2.2. 8), and dropping the now unnecessary subscripts, both graviton coupling constant α_G and the exponential cutoff $(1 - e^{-0.61k^2/k_{\min}^2})$ cancel:

$$\frac{\Delta\rho_{Gk \min}}{\rho_{Gk \min}} \approx \frac{(1 - e^{-0.61k^2/k_{\min}^2})\alpha_G \left[\frac{m}{r} \right] 0.573 \frac{\rho_U}{k_{\min}^2} dk_{\min}}{(1 - e^{-0.61k^2/k_{\min}^2})\alpha_G 0.3247 \frac{\rho_U^2}{k_{\min}^4} dk_{\min}} \approx \left[\frac{m}{r} \right] \left[\frac{1.765k_{\min}^2}{\rho_U} \right] \approx 2\frac{m}{r} \quad (2.2. 9)$$

(Strictly speaking we should be using dk_{\min}'' in the top line of this equation but the error is second order as we are approximating with $r \gg \gg m$. We will do this more accurately below for large masses.) In any metric both ρ_U & k_{\min} transform their values but k_{\min}^2 / ρ_U is invariant. For the above to be consistent with General Relativity this suggests that:

“At all points inside the horizon, and at any cosmic time T , the red highlighted part of Eq.(2.2.9) is ≈ 2 in Planck units. This is simply equivalent to putting $G/c^2 = 1 = G = c$ ”.

Thus we conjecture

$$\text{The local measured density of the universe } \rho_U \approx (0.8823)k_{\min}^2 \approx 0.8823 \frac{\Upsilon^2}{R_{OU}^2} \quad (2.2. 10)$$

The parameter $\Upsilon = k_{\min} R_{OH}$ in radians is close to 1.

Putting Eq. (2.2. 10) the average density ρ_U into Eq.(2.1. 10) gives $\rho_{Gk \min}$ & $K_{Gk \min}$.

$$\text{Cutoff Wavelength Graviton Probability Density } \rho_{Gk \min} \approx \frac{0.149\alpha_G \rho_U^2}{k_{\min}^4} dk_{\min}$$

$$\rho_{Gk \min} \approx \frac{0.149\alpha_G (0.8823k_{\min}^2)^2}{k_{\min}^4} dk_{\min} \approx 0.115\alpha_G dk_{\min} = K_{Gk \min} dk_{\min} \quad (2.2. 11)$$

Where we label $K_{Gk \min} \approx 0.115\alpha_G$ as "The k_{\min} Graviton Invariant".

If our conjectures are true, this is the number density of k_{\min} cutoff wavelength gravitons excluding possible effects of virtual particles emerging from the vacuum. In section 2.5.2 we argue that these do not change the $K_{Gk \min}$ of Eq. (2.2. 11). However $K_{Gk \min}$ does depend on the graviton coupling constant α_G between Planck masses, but α_G cancels out in Eq.(2.2. 9) It does not affect the allowed universe average density ρ_U in Eq. (2.2. 10).

2.2.2 The Schwarzschild metric near large masses

At a radius r from a mass m (dropping the now unnecessary subscripts) the Schwarzschild metric is $(1 - 2m/r)^{\pm 1/2}$ for the time and radial terms which can be written as

$$\sqrt{g_{rr}} = \frac{1}{\sqrt{1 - 2m/r}} = \frac{1}{\sqrt{g_{tt}}} = \frac{1}{\sqrt{1 - \beta_M^2}} = \gamma_M \quad (2.2. 12)$$

Velocity β_M ($c = 1$) is that reached by a small mass falling from infinity and $\gamma_M^{\pm 1}$ is the metric change in clocks and rulers due to mass m . We are using green symbols with the subscript M for metrics $g_{\mu\nu} \neq \eta_{\mu\nu}$ as we did for k_{\min}'' above. The symbols $\gamma_M^{\pm 1}$ help clarity in what follows.

$$\beta_M^2 = \frac{2m}{r}$$

$$\gamma_M^2 = \frac{1}{1 - 2m/r} = g_{rr} = \frac{1}{g_{tt}}$$

Using these symbols $k_{\min}'' = \gamma_M k_{\min} \rightarrow dk_{\min}'' = \gamma_M dk_{\min} \rightarrow \rho_{Gk_{\min}}'' = \gamma_M^2 \rho_{Gk_{\min}}$ (2.2. 13)

In sections 2.1.2 & 2.2.2 we approximated in flat space. The wavelength of k_{\min} gravitons span approximately to the horizon. They fill all of space. We can think of the non flat space around even a large black hole as an infinitesimal bubble on the scale of the observable universe. The normalizing constant of a k_{\min} wavefunction emitted from a localized mass is only altered very close to this mass. Over the vast majority of space it is unaltered. Only close to this mass will local observers measure $k_{\min}'' = \gamma_M k_{\min}$ due to the change in clocks. There is also a local dilution of the normalizing constant due to changing radial rulers. We will consider both these changes in two steps to help illustrate our argument. Now repeat the derivation of $\Delta\rho_{Gk_{\min}}$ as in section 2.2.1 but with a large central mass as in Figure 2.2. 1.

At the point P consider Eq.(2.2. 2) @ k_{\min} : $\psi_1 * \psi_2 + \psi_2 * \psi_1 \approx \frac{4k_{\min}'}{4\pi r_1 r} e^{-k_{\min}' r} \cos(-k_{\min}' r)$.

The red part is the normalizing factor discussed above where we will *initially ignore the dilution* due to the local increase in volume. In deriving Eq.(2.2. 2) we ignored the exponential decay term and phase angle term from the local mass as $e^{-k_{\min}' r_1}$ & $\cos(-k_{\min}' r_1) \approx 1$, even in the space around large black holes. The green $k'r$ & kr terms are phase angles only applying to the vastly distant masses that are virtually fixed by the time they approach even large black holes; increasing only infinitesimally in any local metric. So treating them as fixed and *ignoring the dilution factor* this equation is unaltered. As the exponential cutoff is unchanged we are left with the coupling factor

$$\frac{2\alpha_G}{\pi} \frac{dk_{\min}}{k_{\min}} \text{ which is the same as } \frac{2\alpha_G}{\pi} \frac{dk_{\min}''}{k_{\min}''} = \frac{2\alpha_G}{\pi} \frac{\gamma_M dk_{\min}}{\gamma_M k_{\min}}$$

Dropping the now unnecessary subscripts and temporarily ignoring dilution factors and clock changes we can rederive Eq's (2.2. 4), (2.2. 5) & (2.2. 6) to get with large masses:

$$\Delta\rho_{Gk\min} \approx (1 - e^{-0.61k^2/k_{\min}^2})\alpha_G \frac{m}{r} (0.573) \frac{\rho_U}{k_{\min}^2} dk_{\min} \approx \frac{m}{r} 0.261\alpha_G \frac{\rho_U}{k_{\min}^2} dk_{\min} @ k_{\min}$$

But $\frac{\rho_U}{k_{\min}^2} \approx 0.8823$ from Eq. (2.2. 10) so $\Delta\rho_{Gk\min} \approx \frac{2m}{r} 0.115\alpha_G dk_{\min} @ k_{\min} .$

Using Eq. (2.2. 11) $K_{Gk\min} \approx 0.115\alpha_G$ and $\Delta\rho_{Gk\min} \approx \frac{2m}{r} K_{Gk\min} dk_{\min} @ k_{\min} .$

Using $\beta_M^2 = \frac{2m}{r}$ and to help illustrate metric changes we temporarily include the factor γ_M^2

$$\text{Before metric changes } \Delta\rho_{Gk\min} \approx \beta_M^2 \gamma_M^2 K_{Gk\min} dk_{\min} \quad (2.2. 14)$$

So the total k_{\min} graviton density before metric changes is the original $\rho_{Gk\min} \approx K_{Gk\min} dk_{\min}$ plus the extra $\Delta\rho_{Gk\min} \approx \beta_M^2 \gamma_M^2 K_{Gk\min} dk_{\min} .$ Thus before metric changes

$$\begin{aligned} \rho_{Gk\min}(\text{Total}) &= K_{Gk\min} dk_{\min} + \beta_M^2 \gamma_M^2 K_{Gk\min} dk_{\min} = (1 + \beta_M^2 \gamma_M^2) K_{Gk\min} dk_{\min} \\ \text{But } (1 + \beta_M^2 \gamma_M^2) &= 1 + \frac{\beta_M^2}{1 - \beta_M^2} = \gamma_M^2 \end{aligned} \quad (2.2. 15)$$

$$\text{Before changes } \rho_{Gk\min}(\text{Total}) = K_{Gk\min} dk_{\min} + \beta_M^2 \gamma_M^2 K_{Gk\min} dk_{\min} = \gamma_M^2 K_{Gk\min} dk_{\min}$$

If we now increase the volume to that in the new metric, the new volume is $\sqrt{g_{rr}} = \gamma_M$ times the original volume. So in the new metric we must divide this value by γ_M to get

$$\text{Diluted } \rho_{Gk\min}(\text{Total}) = \frac{K_{Gk\min} dk_{\min}}{\gamma_M} + \beta_M^2 \gamma_M K_{Gk\min} dk_{\min} = \gamma_M K_{Gk\min} dk_{\min} \text{ but in the new metric time changes make } k_{\min}'' = \gamma_M k_{\min} \text{ and } dk_{\min}'' = \gamma_M dk_{\min} \text{ or } dk_{\min} = dk_{\min}'' / \gamma_M$$

$$\text{In the new metric } \rho_{3Gk\min}(\text{Total}) = K_{Gk\min} \frac{dk_{\min}''}{\gamma_M^2} + \beta_M^2 K_{Gk\min} dk_{\min}'' = K_{Gk\min} dk_{\min}'' \quad (2.2. 16)$$

If for example $\gamma_M = 2$, frequencies are doubled so $k_{\min}'' = 2k_{\min}$, the number density of gravitons ($\rho_{Gk\min}'' = 2\rho_{Gk\min}$) is doubled, but so is the measurement of a local small volume element, which is now $V = 2$. The above equations tell us that the original $\rho_{Gk\min}$ background gravitons which occupied one unit of volume is now compressed into 1/2 a unit of volume and the remaining 3/2 units of volume is taken up by extra gravitons due to the central mass. Figure 2.2. 2 illustrates this. The metric adjusts itself so that $K_{Gk\min}$ (*the cutoff wavelength graviton probability constant*) is an invariant number, and this should be true in all metrics at any peculiar velocity (See Figure 2.5. 1 also.) What we have done in this section is only true if the increase in measured volume is equal to the increase in measured frequency. In the Schwarzschild metric this is equivalent to saying that $g_{rr} \cdot g_{tt} = 1$ or $|g| = 1$. But what happens in the Kerr metric with angular momentum?




Figure 2.2. 2 An infinitesimal local volume in a Schwarzschild metric where $\sqrt{g_{rr}} = \gamma_M = 2$.

2.3 Angular Momentum and the Kerr Metric

In the Schwarzschild metric the increase in volume is the same as the frequency increase as $g_{rr} \cdot g_{tt} = 1$ and $g_{\theta\theta} \cdot g_{\phi\phi} = r^4 \sin^2 \theta$ is invariant if there is no angular momentum. With angular momentum both $g_{\theta\theta}$ & $g_{\phi\phi}$ change. The volume ratio of $g_{\mu\nu} \neq \eta_{\mu\nu}$ space, to $g_{\mu\nu} = \eta_{\mu\nu}$ space in

$$\text{any metric at fixed } r \text{ \& } \theta \text{ is } \frac{V'}{V} = \sqrt{\frac{(g'_{rr} \cdot g'_{\theta\theta} \cdot g'_{\phi\phi})(g_{\mu\nu} \neq \eta_{\mu\nu})}{(g_{rr} \cdot g_{\theta\theta} \cdot g_{\phi\phi})(g_{\mu\nu} = \eta_{\mu\nu})}} = \sqrt{\frac{(g'_{rr} \cdot g'_{\theta\theta} \cdot g'_{\phi\phi})(g_{\mu\nu} \neq \eta_{\mu\nu})}{r^4 \sin^2 \theta}} \quad (2.3. 1)$$

The Kerr metric can be written in Boyer-Lindquist coordinates as

$$\left[\begin{array}{l} g_{\theta\theta} = r^2 + \alpha^2 \cos^2 \theta \\ g_{\phi\phi} = (r^2 + \alpha^2 + \frac{r_s r}{\rho^2} \alpha^2 \sin^2 \theta) \sin^2 \theta \\ g_{t\phi} = \frac{r_s r}{g_{\theta\theta}} \alpha \sin^2 \theta \\ g_{rr} = \frac{g_{\theta\theta}}{\Delta} \quad \& \quad g_{tt} = 1 - \frac{r_s r}{g_{\theta\theta}} \end{array} \right.$$

Where $\Delta = r^2 + r_s r + \alpha^2$ and $\alpha = \frac{J}{mc}$ and $r_s = \frac{2Gm}{c} = 2m$ is the Schwarzschild radius in

Planck units where $G = c = 1$. Everything is in units of length or (length)², except g_{rr} & g_{tt} which are dimensionless. Because we want volume ratios as in Eq. (2.3. 1) we can write the above version of the Kerr metric in a dimensionless form, leaving the length squared, and length terms $r^2, r^2 \sin^2 \theta$ & $r \sin \theta$ in $r^2 d\theta^2, r^2 \sin^2 \theta d\phi^2$ & $r \sin \theta d\phi$ etc outside the metric tensor. This effectively gives us the denominator $r^4 \sin^2 \theta$ we want in Eq. (2.3. 1) as we will see. We must also remember that angular momentum parameter α is a length dimension.

Writing the above in dimensionless form as follows, using $-+++$ for the line element ds^2 :

A Dimensionless form of the Kerr Metric where

$$\Delta = 1 + \frac{\alpha^2}{r^2} - A \quad \text{and} \quad A = \frac{2m}{r} \quad \text{but we will add an also dimensionless } \frac{m^2}{r^2} \text{ later. See section 2.6}$$

(We assume silent $G = c^2 = 1$ Planck value constants in $A = \frac{2m}{r} +$ a dimensionless term)

$$\left[\begin{array}{l} g_{\varphi\varphi} = 1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta \\ g_{\theta\theta} = 1 + \frac{\alpha^2}{r^2} \cos^2 \theta \\ g_{rr} = \frac{g_{\theta\theta}}{\Delta} \\ g_{t\varphi} = \frac{A}{g_{\theta\theta}} \frac{\alpha}{r} \sin \theta \\ g_{tt} = 1 - \frac{A}{g_{\theta\theta}} \end{array} \right. \quad (2.3. 2)$$

The space surrounding a rotating mass corotates with it. If we move in this corotating reference frame there is a new metric time component, which after some rearranging of plus

and minus signs just for convenience, we can write as: $g'_{tt} = g_{tt} + \frac{g_{t\varphi}^2}{g_{\phi\phi}}$.

Thus using Eq. (2.3. 2)

$$\begin{aligned} g'_{tt} &= g_{tt} + \frac{g_{t\varphi}^2}{g_{\phi\phi}} = \left(1 - \frac{A}{g_{\theta\theta}}\right) + \frac{\frac{A^2}{g_{\theta\theta}^2} \frac{\alpha^2}{r^2} \sin^2 \theta}{\left[1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta\right]} \\ &= \left(1 - \frac{A}{g_{\theta\theta}}\right) + \frac{\frac{A^2}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta}{g_{\theta\theta} \left[1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta\right]} \\ &= \frac{g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta\right) - A \left(1 + \frac{\alpha^2}{r^2}\right) - \frac{A^2}{g_{\theta\theta}^2} \frac{\alpha^2}{r^2} \sin^2 \theta + \frac{A^2}{g_{\theta\theta}^2} \frac{\alpha^2}{r^2} \sin^2 \theta}{g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta\right)} \\ &= \frac{-A \left(1 + \frac{\alpha^2}{r^2} - \frac{\alpha^2}{r^2} \sin^2 \theta\right) + g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2}\right)}{g_{\theta\theta} g_{\phi\phi}} \\ &= \frac{-A \left(1 + \frac{\alpha^2}{r^2} \cos^2 \theta\right) + g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2}\right)}{g_{\theta\theta} g_{\phi\phi}} \\ g'_{tt} &= g_{\theta\theta} \frac{-A g_{\theta\theta} + g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2}\right)}{g_{\phi\phi}} = \frac{g_{\theta\theta} \left(1 + \frac{\alpha^2}{r^2} - A\right)}{g_{\theta\theta} g_{\phi\phi}} = \frac{\Delta}{g_{\phi\phi}} \end{aligned} \quad (2.3. 3)$$

We have explicitly gone through this to show that if the parameter $A = 2m/r$ is dimensionless, there is potentially freedom to change it without changing Eq. (2.3. 3).

(See Section 2.6.6 as this is similar to what happens in the Kerr-Newman metric, where instead of a dimensionless m^2/r^2 term, a dimensionless r_0^2/r^2 or equivalently a dimensionless Q^2/r^2 is included in term A . See for example Table 2.6. 2 and Table 2.6. 3.)

We will work in corotating frames. Space is swirling around the black hole effectively at rest in these frames, simplifying our calculations and equations. (Section 2.9 puts this into a four vector form, invariant in all frames.) If a small mass, at rest at infinity in the same rest frame as the rotating black hole, falls inwards, it will have the same circumferential velocity as the corotating rest frames at all radii. It will be falling radially through these corotating frames. As in section 2.2.2 we call this radial velocity β_M where as in the non-rotating case

$$\frac{1}{\sqrt{1-\beta_M^2}} = \gamma_M \quad \text{but now} \quad \frac{1}{\sqrt{1-\beta_M^2}} = \gamma_M = \frac{1}{\sqrt{g'_{tt}}} \quad \text{the new inverse rate of clocks.}$$

$$\begin{aligned} \text{In corotating frames} \quad \frac{1}{\gamma_M^2} &= g'_{tt} = \frac{\Delta}{g_{\phi\phi}} \\ \gamma_M^2 &= \frac{g_{\phi\phi}}{\Delta} \end{aligned} \quad (2.3. 4)$$

Frequencies measured in corotating frames increase as γ_M . Similarly using Eq's. (2.3. 1) & (2.3. 4) we can get the (three) volume element ratio in this corotating reference frame.

$$\text{The 3 volume element ratio } V = \sqrt{(g_{rr} \cdot g_{\theta\theta} \cdot g_{\phi\phi})} = \sqrt{\frac{g_{\theta\theta}}{\Delta} g_{\theta\theta} g_{\phi\phi}} = g_{\theta\theta} \sqrt{\frac{g_{\phi\phi}}{\Delta}} = g_{\theta\theta} \gamma_M \quad (2.3. 5)$$

With angular momentum we no longer have the same increase in frequency as volume as in the Schwarzschild case. With no angular momentum we found that the probability density of time polarized k_{\min} gravitons Eq. (2.2. 14) $\Delta\rho_{Gk_{\min}} \approx \gamma_M^2 \beta_M^2 K_{Gk_{\min}} dk_{\min} = \gamma_M^2 \frac{2m}{r} K_{Gk_{\min}} dk_{\min}$.

(Again temporarily adding γ_M^2). With rotation we will find a circularly polarized $\cos^2 \theta$ type distribution of gravitons around the axis. These add to the time polarized dimensionless number $\frac{2m}{r}$ to get an as yet unknown number we simply label as X where $X > \frac{2m}{r}$

$$(2.3. 6)$$

Let us rewrite Eq.(2.2. 14) as $\Delta\rho_{Gk_{\min}} \approx \gamma_M^2 X K_{Gk_{\min}} dk_{\min}$ with rotation

Where the factor γ_M^2 is for clarity only. Repeating the derivation of Eq.(2.2. 15)

$$\rho_{Gk_{\min}} (\text{Undiluted Total}) = K_{Gk_{\min}} dk_{\min} + \gamma_M^2 X K_{Gk_{\min}} dk_{\min} = (1 + \gamma_M^2 X) K_{Gk_{\min}} dk_{\min}$$

As in Eq.(2.2. 16) we need to divide this undiluted total by the new volume $V = g_{\theta\theta}\gamma_M$ in Eq. (2.3. 5) to get the new k_{\min} graviton density $\rho''_{Gk\min}$. If our conjectures are correct $\rho''_{Gk\min} = K_{Gk\min} dk''_{\min}$ is always true, and as our measurement of k_{\min} increases to $k''_{\min} = \gamma_M k_{\min}$ in the new metric, $\rho''_{Gk\min} = K_{Gk\min}\gamma_M dk_{\min}$. So rewriting Eq.(2.2. 16) as follows

$$\rho''_{Gk\min} = \frac{(1 + \gamma_M^2 X) K_{Gk\min} dk_{\min}}{V} = \frac{(1 + \gamma_M^2 X) K_{Gk\min} dk_{\min}}{g_{\theta\theta}\gamma_M} = \gamma_M K_{Gk\min} dk_{\min} = K_{Gk\min} dk''_{\min}$$

$$(1 + \gamma_M^2 X) K_{Gk\min} dk_{\min} = g_{\theta\theta}\gamma_M^2 K_{Gk\min} dk_{\min}$$

$$1 + \gamma_M^2 X = g_{\theta\theta}\gamma_M^2$$

$$X = g_{\theta\theta} - \frac{1}{\gamma_M^2} = \left(1 + \frac{\alpha^2}{r^2} \cos^2 \theta\right) - \frac{1}{\gamma_M^2}$$

$$X = \left(1 + \frac{\alpha^2}{r^2} \cos^2 \theta\right) - \frac{\Delta}{g_{\phi\phi}} \text{ using Eq. (2.3. 4)}$$

$$X = 1 + \frac{\alpha^2}{r^2} \cos^2 \theta - \frac{1 + \frac{\alpha^2}{r^2} - A}{1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta} \text{ using Eq's.(2.3. 2)}$$

We can write this as

$$X = \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{\left[1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta\right] - \left[1 + \frac{\alpha^2}{r^2} - A\right]}{1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta}$$

$$X = \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A \left[1 + \frac{\alpha^2}{r^2} \frac{\sin^2 \theta}{g_{\theta\theta}}\right]}{1 + \frac{\alpha^2}{r^2} + \frac{A}{g_{\theta\theta}} \frac{\alpha^2}{r^2} \sin^2 \theta}$$

$$X = \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A \left[1 + \frac{\alpha^2}{r^2} \frac{\sin^2 \theta}{g_{\theta\theta}}\right]}{g_{\phi\phi}} \text{ using Eq's.(2.3. 2)}$$

$$X = \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A}{g_{\phi\phi}} + A \frac{\alpha^2}{r^2} \frac{\sin^2 \theta}{g_{\phi\phi} g_{\theta\theta}}$$

Section 2.3.3 discusses why there is no separate term in $A \frac{\alpha^2}{r^2} \frac{\sin^2 \theta}{g_{\phi\phi} g_{\theta\theta}}$ so we will write this as

$$\begin{aligned}
\mathbf{X} &= \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A g_{\theta\theta}}{g_{\phi\phi} g_{\theta\theta}} + A \frac{\alpha^2 \sin^2 \theta}{r^2 g_{\phi\phi} g_{\theta\theta}} \\
\mathbf{X} &= \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A(1 + \frac{\alpha^2}{r^2} \cos^2 \theta)}{g_{\phi\phi} g_{\theta\theta}} + A \frac{\alpha^2 \sin^2 \theta}{r^2 g_{\phi\phi} g_{\theta\theta}} \\
\text{Which we finally write as } \mathbf{X} &= \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A(1 + \frac{\alpha^2}{r^2})}{g_{\phi\phi} g_{\theta\theta}}
\end{aligned} \tag{2.3. 7}$$

Putting $A = \frac{2m}{r}$, the *extra* k_{\min} virtual gravitons $\gamma_M^2 \mathbf{X}$ (due to a mass m rotating with angular parameter α that has dimensions of length) are the following two polarization groups (The background k_{\min} virtual gravitons have been normalized to one when $\gamma_M = 1$)

Circularly polarized spin2: $\left[\frac{\alpha^2}{r^2} \cos^2 \theta \right] \times (m = \pm 2)$ & Time polarized spin 2: $\left[\frac{\frac{2m}{r} (1 + \frac{\alpha^2}{r^2})}{g_{\phi\phi} g_{\theta\theta}} \right]$

We can rewrite Eq. (2.2. 16) using $1 + \gamma_M^2 \mathbf{X} = g_{\theta\theta} \gamma_M^2$ or $\frac{1}{\gamma_M^2} + \mathbf{X} = g_{\theta\theta} = 1 + \frac{\alpha^2}{r^2} \cos^2 \theta$

$$\left[\frac{1}{\gamma_M^2} + \frac{\alpha^2}{r^2} \cos^2 \theta + \frac{A(1 + \frac{\alpha^2}{r^2})}{g_{\phi\phi} g_{\theta\theta}} = 1 + \frac{\alpha^2}{r^2} \cos^2 \theta \right] \text{ or } \left[\frac{1}{\gamma_M^2} + \frac{A(1 + \frac{\alpha^2}{r^2})}{g_{\phi\phi} g_{\theta\theta}} = 1 \right] \text{ where } \left[\frac{A(1 + \frac{\alpha^2}{r^2})}{g_{\phi\phi} g_{\theta\theta}} = \beta_M^2 \right]$$

We have a 4 vector equation again as circular polarization cancels on both sides. The main thing to notice here is that the circularly polarized k_{\min} gravitons are independent of the central mass, suggesting they are due to the effect of the rotation of space, or frame dragging, on the k_{\min} graviton background. We will discuss this in section 2.3.2. The extra k_{\min} gravitons due to the central mass have a $(1 + \alpha^2 / r^2) / (g_{\phi\phi} g_{\theta\theta})$ factor, distorting them from spherical symmetry. Figure 2.3.1 & Figure 2.3. 2 compare the above with spinning charged spheres in electromagnetism. The electrostatic energy density surrounding a charged sphere however, reduces with radius as r^{-4} , and magnetic energy as r^{-6} , or two more powers of radius. With gravity however we have been looking at the probability density of minimum wavenumber k_{\min} gravitons surrounding a mass. With no angular momentum there are only time polarized k_{\min} gravitons, and their extra probability density drops as r^{-1} , as so far we have only focussed on those k_{\min} gravitons (the vast majority), that interact with the rest of the mass in the universe. If a charged sphere rotates, there is a radial magnetic field of circularly polarized $m = \pm 1$ photons varying in intensity as $\cos^2 \theta$ and a transverse magnetic field (of transversely polarized $m = \pm 1$ photons) varying as $\sin^2 \theta$ as in Figure 2.3.1.




Figure 2.3.1 Spinning electrically charged sphere. At same radius $B_R @(\theta=0) = 2B_T @(\theta = \pi / 2)$.




Figure 2.3. 2 Spinning mass m with angular momentum length parameter α as viewed in a corotating frame. There are circularly polarized $(m = \pm 2) \times k_{\min}$ gravitons due to the effect of frame dragging on the background time polarized k_{\min} gravitons. There are no transversely polarized $(m = \pm 2) \times k_{\min}$ gravitons due to a rotating mass m as seen in a corotating frame. Radially polarized extra k_{\min} gravitons due to mass m are distorted from spherical symmetry as $(1 + \alpha^2 / r^2) / (g_{\varphi\varphi} g_{\theta\theta})$. For $r \gg r_{Sw}$ we can ignore the effects of $g_{\theta\theta}, g_{\varphi\varphi}$. as they rapidly tend to one, with the metric written in dimensionless form as in Equ's.(2.3. 2).

2.3.1 Stress tensor sources for spin 2 gravitons & 4 current sources for spin 1

Spin 1 particles behave like a 4 vector as they come from a 4 current source, transforming with velocity as in the Special Relativity transformations of Minkowski spacetime. Spin 2 gravitons in contrast come from mass/energy density sources. There are two factors in their transformations with velocity. One from the mass increase per source particle, and the second from the increase in particles per unit length due to length contraction. Thus Spin 2 particles transform as a 4x4 rank 2 tensor, which Einstein connected with spacetime curvature.

The rules of quantum mechanics tell us that spherically symmetric spin 2 particles should be equal $1/\sqrt{5}$ superpositions of $m = -2, -1, 0, +1, +2$ states. But the shape of gravitational waves behaves like transversely polarized $m = \pm 2$ particles, suggesting the k_{\min} gravitons surrounding mass concentrations may only consist of time polarized, plus $m = \pm 2$ circularly polarized, spin 2 particles. In [7] we showed that a spherically symmetric spin one state is built from equal $1/\sqrt{3}$ superpositions of $m = -2, 0, +2$ states, and if this is true then we will conjecture that the same superposition of $m = -2, 0, +2$ states must be able to build a spherically symmetric spin 2 state. When we looked at non rotating spherical masses it appeared that, even close to black holes, the spherical symmetry of the Schwarzschild metric suggested similarly spherically symmetric, time polarized, extra k_{\min} gravitons down to the horizon; with space expanding only radially. Thus before we considered angular momentum we could treat all k_{\min} gravitons as only time polarized. A stress tensor source with no angular momentum has spherically symmetric spacetime curvature with time polarized k_{\min} gravitons. But angular momentum in the source produces cylindrically symmetric spacetime curvature. We still have radially polarized k_{\min} gravitons (in co-rotating coordinates) due to the central mass, but distorted from spherical symmetry as $(1 + \alpha^2 / r^2) / (g_{\varphi\varphi} g_{\theta\theta})$ which only affects the close in region, disappearing as $\alpha \rightarrow 0$. But there are also circularly polarized $m = \pm 2$ k_{\min} gravitons only related to angular momentum. These circularly polarized k_{\min} gravitons do not have the $2m/r$ factor and must be very different. As we will discuss below it appears that they are generated from the background time polarized k_{\min} gravitons by the swirling velocity of corotating space.

2.3.2 Circularly polarized gravitons from corotating space

The circularly polarized gravitons do not have a $2m/r$ factor. The Kerr metric is an exact solution to Einstein's field equations, which we conjecture (in an infinitesimally modified form as in Eq. (2.5. 1) are consistent with the k_{\min} Graviton constant being invariant at all points in spacetime, or that Eq. (2.2. 11) is always true. If this is so then Eq. (2.3. 7) should be true also. We can perhaps just accept that it must be true, but at the same time we can look at whether it makes sense?

The angular momentum parameter α has dimensions of length, and is defined as $\alpha = \frac{J}{mc}$.

Because angular momentum is the cross product of momentum by radius or $m\mathbf{v} \times \mathbf{r}$, we can think of this length parameter as a vector of length α , pointing along the axis of spin, with components $\alpha \cos \theta$ at any polar angle θ to the spin axis. Space corotates around spinning masses with angular velocity $\Omega = \frac{g_{t\phi}}{g_{\phi\phi}}$ which in the plane of the equator simplifies to

$$\Omega = \frac{r_s \alpha c}{r^3 + r \alpha^2 + r_s \alpha^2} \approx \frac{r_s \alpha c}{r^3} \text{ when } r \gg r_s \text{ \& } \alpha.$$

$$\text{At large radii the corotating velocity } V = \Omega \times \mathbf{r} \approx \frac{r_s \alpha c}{r^2} \quad (2.3. 8)$$

Because r_s & α have dimensions of length this equation has dimensions of velocity, and if we divide it by c it is dimensionless. We will call it $\beta_{\text{Corotating}} = \beta_C$

$$\text{At large radii } \beta_{\text{Corotating}} = \beta_C = \frac{V}{c} = \frac{\Omega \times r}{c} \approx \frac{r_s \alpha}{r^2} \text{ a dimensionless number.} \quad (2.3. 9)$$

If we now think of $\alpha = \frac{J}{mc}$ as $\alpha = \frac{m \mathbf{v} \times \mathbf{r}}{m c} = \frac{\mathbf{v} \times \mathbf{r}}{c}$ we can consider a similar vector along the spin axis consisting of the cross product of the corotating velocity of space $\frac{V}{c} \approx \frac{r_s \alpha}{r^2}$ by the radius r . The length along the spin axis of this cross product vector $\frac{\mathbf{V} \times \mathbf{r}}{c}$ is simply $\frac{r_s \alpha}{r}$.

$$\text{At the equator: Length of vector } \frac{\mathbf{V} \times \mathbf{r}}{c} \text{ along the spin axis is } \approx \frac{r_s \alpha}{r} \text{ for } r \gg r_s \quad (2.3. 10)$$

We need this vector length to be a dimensionless number representing the amplitude that a background time polarized k_{\min} graviton generates a circularly polarized k_{\min} graviton around the spin axis. If we divide Eq. (2.3. 10) by the Schwarzschild radius r_s , all rotating black holes with the same percentage of maximum spin look identical, and we get a dimensionless magnitude as required

$$\text{Magnitude of normalized dimensionless vector } \left| \frac{\mathbf{V} \times \mathbf{r}}{r_s c} \right| \approx \frac{r_s \alpha}{r_s r} \approx \frac{\alpha}{r} \quad (2.3. 11)$$

The whirling velocity of space is a maximum out from the equator, but circularly polarized gravitons generated in this region have to be distributed on this shell around the spin axis as the square of the component of angular momentum. We thus conjecture that the probability of *background* time polarized k_{\min} gravitons, on a corotating thin spherical shell at large radius, generating circularly polarized k_{\min} gravitons around the spin axis on the same shell is

$$\text{Probability of } \frac{\text{Extra circularly polarized } m = \pm 2 \times k_{\min} \text{ gravitons}}{\text{Background time polarized } k_{\min} \text{ gravitons}} \approx \frac{\alpha^2 \cos^2 \theta}{r^2} \quad (2.3. 12)$$

There is a background density of time polarized k_{\min} gravitons on each corotating spherical shell. The swirling velocity of these k_{\min} gravitons generates extra circularly polarized k_{\min} gravitons around the spin axis with a $\cos^2 \theta$ distribution around the spin axis on the same shell, in agreement with Figure 2.3. 2. For simple explanatory purposes, we approximated at

large radii only. At small radii we must use $\Omega = \frac{g_{t\phi}}{g_{\phi\phi}} = \frac{r_s \alpha c}{r^3 g_{\phi\phi} g_{\theta\theta}}$. On the equator $g_{\theta\theta} = 1$, the

co-rotation velocity $V = \Omega \times \mathbf{r} \sqrt{g_{\phi\phi}}$. The circumferential volume generating these circularly polarized gravitons also expands as $\sqrt{g_{\phi\phi}}$. Rederiving Eq's. (2.3. 9) and those following, the

effective angular momentum term becomes $\Omega \times \mathbf{r} \times \mathbf{r} \cdot \sqrt{g_{\phi\phi}} \cdot \sqrt{g_{\phi\phi}} = \left[\frac{r_s \alpha c}{r^3 g_{\phi\phi}} \right] \cdot r^2 g_{\phi\phi} = \frac{r_s \alpha c}{r}$

just as before; our derivation applies down to the equatorial horizon. This circular polarization appears to be the result of the swirling or corotating velocity of space as it has no mass term, only angular momentum terms.

2.3.3 Why there are no transverse polarized gravitons in co-rotating coordinates?

In Section 2.1 we showed that the majority of k_{\min} gravitons around any non rotating mass m is due to the interaction between that mass and the rest of the mass in the universe $\Delta \rho_{Gk \min} \propto (\psi_{\text{Universe}} * \psi_m + \psi_m * \psi_{\text{Universe}})$; and these were all time polarized k_{\min} gravitons. Let us imagine that a rotating mass emits transversely polarized k_{\min} gravitons, there will only be a small number unless there are also transversely polarized k_{\min} gravitons from the rest of the universe for their amplitudes to interact with. But from what we have just done above there appears to be only circularly polarized k_{\min} gravitons due to the corotation of space. Also if a rotating mass emits its own circularly polarized k_{\min} gravitons, these would interact with the circularly polarized k_{\min} gravitons due to the corotation of space. It thus appears that, when observed in corotating coordinates, a rotating mass does not itself emit either tranverse or circularly polarized k_{\min} gravitons. This perhaps makes sense, as in corotating frames, we are effectively at rest above the horizon which is rotating in sync with us.

2.3.4 Does our time polarized k_{\min} value in co-rotating coordinates make sense?

The inner horizon radius R is defined when $\Delta = 1 + \frac{\alpha^2}{r^2} - A = 1 + \frac{\alpha^2}{r^2} - \frac{2m}{r} = 0$ where we initially define the dimensionless number $A = \frac{2m}{r}$. Using $A = \frac{2m}{r}$ the inner horizon is

where $r^2 - 2mr + \alpha^2 = 0$. So horizon radius $R = r = m + \sqrt{m^2 + \alpha^2} = 2m$ when $\alpha = 0$, and at maximum spin $r = R = m$ when $\alpha = m$. But we will, for generality, revert to the dimensionless A and look at what happens near the horizon for various spins.

Let A_H be the value of A at the horizon where $1 + \frac{\alpha^2}{R^2} - A_H = 0$ is always true

$$\text{So } \frac{\alpha^2}{R^2} = A_H - 1 \quad \text{and} \quad A_H = 1 + \frac{\alpha^2}{R^2} \quad (2.3. 13)$$

At the horizon $g_{\theta\theta} g_{\phi\phi} = (1 + \frac{\alpha^2}{R^2} \cos^2 \theta)(1 + \frac{\alpha^2}{R^2} + A_H \frac{\alpha^2}{R^2} \frac{\sin^2 \theta}{g_{\theta\theta}}) = g_{\theta\theta} (A_H + A_H \frac{\alpha^2}{R^2} \frac{\sin^2 \theta}{g_{\theta\theta}})$

$$= g_{\theta\theta} A_H (1 + \frac{\alpha^2}{R^2} \frac{\sin^2 \theta}{g_{\theta\theta}}) = A_H (g_{\theta\theta} + \frac{\alpha^2}{R^2} \sin^2 \theta)$$

At the horizon $g_{\phi\phi} g_{\theta\theta} = A_H (1 + \frac{\alpha^2}{R^2} \cos^2 \theta + \frac{\alpha^2}{R^2} \sin^2 \theta) = A_H (1 + \frac{\alpha^2}{R^2}) = A_H^2$

$$\text{and independent of angle } \theta \text{ near the horizon only. This is true regardless of} \quad (2.3. 14)$$

spin from zero to maximum as the radius shrinks.

Near the horizon the black hole surface area is $4\pi R^2 \sqrt{g_{\phi\phi} g_{\theta\theta}} = 4\pi R^2 A_H$ (2.3. 15)

We have shown in co-rotating frames the extra time polarized k_{\min} graviton density due to a

central mass is $\Delta\rho_{Gk_{\min}} = A \cdot \frac{1 + \frac{\alpha^2}{R^2}}{g_{\theta\theta} g_{\phi\phi}} K_{Gk_{\min}} dk_{\min}$ where $A = \frac{2m}{r}$ so far, and dimensionless.

Using Eq's. (2.3. 13) & (2.3. 14) above, near the horizon in a corotating frame, this becomes

Extra time polarized k_{\min} graviton density near horizon for all Black holes

$$\Delta\rho_{Gk_{\min}} = A_H \cdot \frac{1 + \frac{\alpha^2}{R^2}}{g_{\theta\theta} g_{\phi\phi}} K_{Gk_{\min}} dk_{\min} = \frac{A_H^2}{A_H^2} K_{Gk_{\min}} dk_{\min} = K_{Gk_{\min}} dk_{\min} \quad (2.3. 16)$$

Ignoring the factor $K_{Gk_{\min}} dk_{\min}$ the extra radially polarized k_{\min} graviton probability density is *always one in Planck units* regardless of spin and is spherically symmetric, but only near the horizon where the background density $K_{Gk_{\min}} dk_{\min} / \gamma_M^2 \rightarrow 0$ as $\beta_M^2 \approx 1$ & $\gamma_M^2 \rightarrow \infty$, providing it is observed (somehow) in a corotating reference frame. It can also be shown that near the

horizon of a black hole $\gamma_M^2 \approx \frac{4R^2}{s^2}$ and $\gamma_M \approx \frac{2R}{s}$ is always true regardless of the degree of

spin, and the value of the dimensionless number A_H , where R is the horizon radius and s the proper distance from it, providing it is all measured in co-rotating coordinates. The region well above the horizon is not spherically symmetric until several Schwarzschild radii away where spherical symmetry is gradually retained as in the non-rotating case.

Also near the horizon this density due to a central mass is so great that we can effectively ignore the background value, but the rotation of space generates circularly polarized k_{\min} gravitons, of probability density $\frac{\alpha^2}{r^2} \cos^2 \theta$ from this background. The extra radially, plus circularly, polarized k_{\min} graviton probability density near the horizon ignoring the factor $K_{Gk_{\min}} dk_{\min}$ is

$$\text{Time polarized } 1 + \text{Circularly polarized } \frac{\alpha^2}{R^2} \cos^2 \theta = (1 + \frac{\alpha^2}{R^2} \cos^2 \theta) \quad (2.3. 17)$$

= $g_{\theta\theta}$ as in our original derivation, but ignoring the background, which is infinitesimal near the horizon.

As the Kerr metric is an exact solution for rotating black holes we can say that if the extra k_{\min} gravitons due to a rotating mass are consistent with X as in Eq. (2.3. 7) then it is also consistent with keeping the Graviton constant $K_{Gk_{\min}}$ as in Eq. (2.2. 11) invariant in the spacetime surrounding it. (We come back to this, and potential changes to the dimensionless term $A = 2m / r$ in section 2.6.) When we looked at non rotating black holes in section 2.2.2 we used simple first principles to show that the warping of spacetime around them is consistent with an invariant Graviton constant $K_{Gk_{\min}}$. With rotating black holes we turned the argument around and assumed this invariance to derive the extra probability densities of time, and circular polarized k_{\min} gravitons, before the density dilution from the expansion of space around the rotating mass. Equations (2.3. 16) & (2.3. 17) can perhaps increase our confidence that our hypothesis is possibly correct. If it is correct on the horizon, and also far from a rotating black hole, we will conjecture that it is also correct in all regions between, even if it might not initially appear to be so. It is important to remember that the Kerr metric is an exact solution for rotating black holes, not for rotating masses in general. We have only considered here the exact solution. We can thus perhaps summarize section 2.3 as follows:

Spherically symmetric spacetime curvature generates only time polarized k_{\min} gravitons.

Cylindrically symmetric spacetime curvature, due to angular momentum, generates time polarized k_{\min} gravitons and circularly polarized $m = \pm 2 k_{\min}$ gravitons in corotating coordinates.

We have not yet included the relatively small number of k_{\min} gravitons emitted by the mass itself ($\psi_m * \psi_m$), which has effect close to black holes; but we will first look at the expanding universe. This is a much revised version of section 5.3 in [7] with Figure 2.4. 1 & Equ's.(2.4. 12) helping to make clear why the k_{\min} graviton constant $K_{Gk_{\min}}$ is invariant throughout spacetime. It is the cutoff wavenumber where densities for action available equals action required, by k_{\min} graviton superpositions. The value of k_{\min} reduces with cosmic time T but increases around mass concentrations with the local metric clock rates. See Figure 2.5. 1.

2.4 The Expanding Universe

Section 2.1.1 describes virtual gravitons as superpositions of the three modes $n=3,4,5$ at a single wavenumber k as in Table 4.3.1 in [7] which also tells us $\langle n \rangle = 3.33$. Equation (3.2.1) in that paper tells us $\langle \mathbf{p}_k(\text{debt}) \rangle = -\langle \beta_k \rangle^2 \langle n \rangle \hbar \mathbf{k}$ is the debt of $\hbar \mathbf{k}$ spatially polarized quanta they borrow. Equation (2.1. 2) tells us $m_0 = \langle n \rangle k_{\min}$ & Eq. (3.1.6) in [7] says they borrow from time polarized quanta a mass $m_0 / (\gamma \sqrt{2s}) = m_0 / (2\gamma) = \langle n \rangle k_{\min} / (2\gamma)$ for spin 2 gravitons. Equation 6.2.2 in [7] tells us that @ k_{\min} for gravitons $\gamma_{\min}^2 = 2$ & $\beta_{\min}^2 = 1/2$. From this we can see that @ k_{\min} , the spatially polarized debt is $\sqrt{2}$ larger than the time polarized debt. So we only need to consider the spatially polarized debt when equating quanta available to quanta required, as when these are equal there is a small surplus of time polarized quanta. However to plot these curves near $k = k_{\min}$ we need the number density of gravitons at any wavenumber k , so rewriting Eq. (2.1. 9) using Eq.(2.2. 10) for ρ_U^2 / k_{\min}^4 & Eq.(2.2. 11)

$$\rho_{Gk} \approx \frac{0.149 \alpha_G \rho_U^2}{k_{\min}^4} dk \left[\frac{108 (1-e^{-0.61x^2}) \sqrt{x^2+11.09}}{x (2x^2+11.09)^2} \right] \approx 0.115 \alpha_G dk \left[\frac{108 (1-e^{-0.61x^2}) \sqrt{x^2+11.09}}{x (2x^2+11.09)^2} \right] \quad (2.4. 1)$$

Both blue boxes are one when $k / k_{\min} = x = 1$. Using Eq's. (3.1.11), (3.1.12) & (3.2.10) in [7]

$$\langle \beta_k \rangle^2 = \frac{\langle K_k \rangle^2}{1 + \langle K_k \rangle^2}. \quad \text{For } N = 2 \text{ spin 2; } \langle K_k \rangle = \frac{\langle n \rangle k}{m_0} \quad \text{and using } m_0 \approx 3.33 k_{\min};$$

$$\langle \beta_k \rangle^2 = \frac{k^2}{k^2 + k_{\min}^2} = \frac{x^2}{x^2 + 1} \quad \text{where } x = \frac{k}{k_{\min}} \text{ so wavefunction } \psi_k \text{ borrows } \langle \beta_k \rangle^2 \langle n \rangle \approx (3.33) \frac{x^2}{x^2 + 1}$$

wavenumber k quanta. But wavenumber k virtual quanta last for time $\Delta T \approx \hbar / 2k$, whereas the time the superposition lasts is $\Delta T' \approx \hbar / 2E$. We are borrowing Energy \times Time or Action, and the superposition energy $E = k' = \sqrt{k^2 + 11.09 k_{\min}^2}$ as in Eq. (2.1. 4) lasts for a shorter time when k is near k_{\min} . So the Action Quanta of Energy \times Time required, reduces as

$$\frac{k}{k'} = \frac{k}{\sqrt{k^2 + 11.09 k_{\min}^2}} = \frac{x}{\sqrt{x^2 + 11.09}} \quad \text{and the quanta density required @ } k \text{ by gravitons is thus}$$

$$\rho_{Quanta @ k} \approx \frac{0.115 \alpha_G}{2.09} dk \left[2.09 \cdot \frac{3.33 \times x^2}{x^2 + 1} \cdot \frac{x}{\sqrt{x^2 + 11.09}} \right] \cdot \left[\frac{108 (1-e^{-0.61x^2}) \sqrt{x^2 + 11.09}}{x (2x^2 + 11.09)^2} \right]$$

$$\rho_{Quanta @ k} \approx 0.0555 \alpha_G dk \left[\frac{751 x^2 (1-e^{-0.61x^2})}{(x^2 + 1) (2x^2 + 11.09)^2} \right] \quad \text{All blue boxes are}$$

$$\text{one when } \frac{k}{k_{\min}} = x = 1 \quad \& \quad \rho_{Quanta @ k_{\min}} = \rho_{Qk_{\min}} \approx 0.0555 \alpha_G dk_{\min} \quad @ \quad k_{\min} \quad (2.4. 2)$$

But the density of zero point modes available @ k_{\min} is $k_{\min}^2 dk / \pi^2$ (ignoring some small factors). Even if $\alpha_G \ll 1$ this is too small by about $k_{\min}^2 \approx 1/R_{OH}^2$. However the area of the causally connected horizon $4\pi R_{OH}^2$ suggests possible connections with Holographic horizons and the AdS/CFT correspondence [24] but in a very different way.

2.4.1 Holographic horizons and red shifted Planck scale zero point modes

Maldacena [24] proposed AntiDesitter or Hyperbolic spacetime where Planck modes on a 2D horizon are (almost) infinitely redshifted at the origin by an (almost) infinite change in the metric. *In contrast we have assumed flat is space on average to the horizon.* In section 2.2.3 in [7] we defined a rest frame, in which preons are born with zero momentum and infinite wavelength, forming infinite superpositions. If we also have a spherical horizon with Planck scale modes, but receding locally at the velocity of light, these Planck modes can be absorbed by infinite wavelength preons (from that receding horizon) and red shifted in a radially focussed manner inwards. We will argue in what follows, that at the centre where the infinite superpositions are built, approximately 1/6 of these Planck modes can be absorbed from that horizon with wavelengths of the order of the horizon radius. *This potential possibility only exists because zero momentum preons have an infinite wavelength.* If any source of radiation recedes at velocity $\beta = v/c$ the frequency/wavenumber reduces as $k_{observer} = k_{source} [\gamma(1-\beta)]$ where $\gamma = (1-\beta^2)^{-1/2}$. In the extreme relativistic limit $\beta \rightarrow 1$ & we can put $1-\beta = \Delta\beta = \varepsilon$.

Putting $1-\beta = \Delta\beta = \varepsilon$ implies $\beta = 1-\varepsilon$ and $\beta^2 \approx 1-2\varepsilon$

$$1-\beta^2 = \gamma^{-2} \approx 2\varepsilon \text{ and } \gamma \approx 1/\sqrt{2\varepsilon}$$

$$\text{Thus } \frac{k_{Observer}}{k_{Source}} = [\gamma(1-\beta)] \approx \frac{\varepsilon}{\sqrt{2\varepsilon}} \approx \sqrt{\frac{\varepsilon}{2}} \approx \frac{1}{2\gamma} \approx 10^{-61} \text{ at current time} \quad (2.4. 3)$$

There is always some rest frame travelling at nearly light velocity that can redshift Planck energy modes into a $k_{\min} \approx 1/R_{OH}$ mode and also many other frames travelling at various lower velocities that can redshift Planck energy modes into any $k > k_{\min}$ mode . This is special relativity applying locally. But in sections 2.1.2 & 2.2.1 we used the fact that clocks in comoving coordinates tick at the same rate. So how does Eq.(2.4. 3) help? Space between comoving galaxies expands with cosmic or proper time t and is called the scale factor $a(t)$. It is normally expressed as $a(t) \propto t^p$ and we will start at time $t = T_0$ with time T now.

$$\text{Thus } \dot{a}(t) \propto pt^{p-1} \text{ and the Hubble parameter } H(t) = \frac{\dot{a}(t)}{a(t)} = \frac{p}{t} \quad (2.4. 4)$$

We have been assuming to here that space is flat on average and will use the properties that in flat space at the current time the coordinate, proper and comoving distances are all equal. Writing the present scale factor normalized to one so that $a(T) = 1$ implies $a(t) = t^p / T^p$, we can get the causally connected horizon radius and the horizon velocity V . Using Eq. (2.4. 4)

The horizon radius $R_{OH} = \int_{T_0}^T \frac{dt}{a(t)} = T^p \int_{T_0}^T \frac{dt}{t^p} = \frac{T - T_0}{1 - p} \approx \frac{T}{1 - p}$ if $T \gg T_0$ & p constant. (2.4. 5)

In flat space horizon velocity $V = \frac{dR_{OH}}{dT} = \frac{d}{dT} \left[T^p \int_{T_0}^T \frac{dt}{t^p} \right]$ then using $d(u \cdot v) = u \cdot dv + v \cdot du$

for $(T \gg T_0)$: $\frac{dR_{OH}}{dT} = \frac{T^p}{T^p} + \frac{R_{OH}}{T^p} (pT^{p-1}) = 1 + \frac{p}{T} R_{OH}$ but $\frac{p}{T}$ is the Hubble constant (2.4. 6)

at time T so the horizon velocity $V = 1 + H(T)R_{OH}$ regardless of how p behaves.

The *Hubble flow velocity of a comoving galaxy on the horizon* is $V' = H(T)R_{OH}$ and thus from Eq. (2.4. 6) the horizon velocity is always $V = 1 + V'$. In other words the horizon is moving at light velocity relative to comoving coordinates instantaneously on the horizon as measured by a central observer. Now clocks tick at the same rate in all comoving galaxies but clocks moving at almost the horizon light velocity (relative to comoving coordinates instantaneously on the horizon) will tick extremely slowly or as $1/\gamma \approx 10^{-61}$ from Eq.(2.4. 3) as special relativity applies locally in this case. Thus Planck modes on the receding horizon will obey Eq's.(2.4. 3) as seen in all comoving coordinates. Let us now imagine an infinity of frames all travelling at various relativistic velocities relative to comoving coordinates instantaneously on the horizon, and radially as seen by central observers. We can think of these as spherical shells on the horizon *all of one Planck length thickness as measured by observers moving radially with them*. Transverse dimensions do not change for all radially moving observers and the effective surface area of all these shells is $4\pi R_{OH}^2$. The internal volume of all these shells, as measured in rest frames by observers moving radially with them, and each of these observers measures their thickness ΔR as one Planck length; is

$$\text{Rest frame internal shell volume } V = 4\pi R_{OH}^2 \Delta R = 4\pi R_{OH}^2 \quad (2.4. 7)$$

We want the zero point quanta available where *these quanta have Planck energy ΔE , lasting for Planck time ΔT such that $\Delta E \times \Delta T \approx \hbar / 2$* . Before redshifting, a single zero point quanta thus has Planck energy (temporarily using a single primed k' that is not the k' of Eq.(2.1. 4)) where $k' = 1$ before, and k after the frequency change. The density of modes in this shell is $k'^2 dk' / \pi^2$ (where each quanta & the superposition it builds both last for time $\Delta T \approx \hbar / (2\Delta E)$)

$$\frac{k'^2 dk'}{\pi^2} \text{ quanta, which we will write as mode quanta density } \frac{k'^3}{\pi^2} \frac{dk'}{k'}. \quad (2.4. 8)$$

At Planck scale $k' = 1$ and redshifting to k then using Eq. (2.4. 3) $k = k' / \gamma$ & $dk = dk' / \gamma$. Thus $dk' / k' = dk / k$. As $k = 1$ Eq.(2.4. 8) becomes

$$\text{Planck Scale Quanta Density before redshifting} = \frac{1^3}{\pi^2} \frac{dk'}{k'} = \frac{1}{\pi^2} \frac{dk}{k} \quad (2.4. 9)$$

Multiplying density by volume ie. Eq's. (2.4. 7) & (2.4. 9) gives the total Planck scale quanta inside the rest frame shell $4\pi R_{OH}^2 \cdot \frac{1}{\pi^2} \frac{dk}{k}$. Two thirds of these modes are transverse and one third radial. Only the inward $1/6$ of these modes can be radially redshifted inwards. After redshifting to wavenumber k , in flat on average space in a thought experiment, we can imagine them forming spherical standing waves, with a central spherical first node at radius $R = \lambda/4 = \pi/2k$, where λ is the De Broglie wavelength of momentum k particles or waves. The polarization directions are spherically symmetric (as required to build infinite superpositions in their rest frame, see section 2.4.7), forming virtual spin 1 quanta with a radial probability of $\psi * \psi \propto 2k \cos^2 kr$. Inside this sphere the expectation value of the radius that a quantum is at is $\langle r \rangle = \pi/4k$ as $\langle \cos^2 kr \rangle = 1/2$, so the expectation value of the probability density is

$$\frac{2k \langle \cos^2 kr \rangle}{4\pi \langle r \rangle^2} = \frac{2k \cos^2(\pi/4)}{4\pi(\pi/4k)^2} = \frac{k}{4\pi} \frac{16k^2}{\pi^2} = \frac{1.62k^3}{4\pi} = \frac{1.62\Upsilon^3}{4\pi R_{OH}^3} \left[\frac{k}{k_{\min}} \right]^3 \text{ where we have used } \Upsilon = k_{\min} R_{OH}$$

This is the average probability density of a single quantum. So the total density is this single quantum probability density, times the number from the horizon; but we also need to divide by 2 as we are only considering the spatially polarized or vector half. Again using $\Upsilon = k_{\min} R_{OH}$ the total quanta density becomes, after dividing by the two factors of 2 & 6

$$\rho_{Quanta@k} \approx \frac{1}{2} \cdot \frac{1}{6} \cdot \left[4\pi R_{OH}^2 \frac{1}{\pi^2} \frac{dk}{k} \right] \frac{1.62\Upsilon^3}{4\pi R_{OH}^3} \left[\frac{k}{k_{\min}} \right]^3 \approx \frac{\Upsilon^2}{7.4\pi^2} dk \left[\frac{k}{k_{\min}} \right]^2 \approx \frac{\Upsilon^2}{7.4\pi^2} dk \cdot x^2 \text{ where } x = \frac{k}{k_{\min}}$$

$$\text{Density of quanta available after redshifting } \rho_{Quanta@k} \approx \frac{\Upsilon^2}{7.4\pi^2} x^2 dk \quad (2.4. 10)$$

Now an observer at the centre of all this sees flat space being added inside the horizon at the rate of the horizon velocity $V = 1 + H(T)R_{OH}$ as in Eq.(2.4. 6). We will conjecture that the space added in one unit of Planck time inside the expanding horizon also creates the source of these zero point action quanta that we can borrow. Thus Eq. (2.4. 10) becomes

$$\text{Density of } k_{\min} \text{ quanta available } \rho'_k \approx \frac{\Upsilon^2 V}{7.4\pi^2} \left[\frac{k}{k_{\min}} \right]^2 dk \approx \frac{\Upsilon^2 V}{7.4\pi^2} x^2 dk$$

$$\approx \frac{\Upsilon^2 (1 + H \cdot R_{OH})}{7.4\pi^2} x^2 dk \text{ where } x = \frac{k}{k_{\min}} \quad (2.4. 11)$$

2.4.2 Plotting available quanta densities, and required quanta densities




Figure 2.4. 1 plots Eq's. (2.4. 2) & (2.4. 11)) as a function of $x = k / k_{min}$. Going through similar procedures for the time mode quanta as for space modes, we have plotted both time and space. An exponential $e^{-0.61x^2}$ cutoff fits available and required reasonably for $k < k_{min}$ in both cases; also showing there should be an adequate supply of time mode quanta from the horizon for all infinitesimal mass particles. The spatial mode crosses @ $k = k_{min}$. Both plots always look the same at all cosmic times T . And, in any metric only the value of k_{min} changes. It only works in an expanding flat universe. Equating required and available spatial modes @ $k = k_{min}$

$$\text{Quanta available} \approx \frac{\Upsilon^2 V}{7.4\pi^2} dk_{min} = \text{Quanta required} \approx 0.055\alpha_G dk_{min} = K_{Qk_{min}} dk_{min} \quad (2.4. 12)$$

$$\text{Where } K_{Qk_{min}} = 0.055\alpha_G \text{ is the "Quanta required @ } k_{min} \text{ Constant " \& } \alpha_G \approx \frac{\Upsilon^2 V}{4}$$

Equation (2.2. 10) the average density of the universe $\rho_U \approx 0.8823 \frac{\Upsilon^2}{R_{OH}^2}$ allows us to solve the present value of $\Upsilon = k_{min} R_{OH}$. Using WMAP data for Baryonic and Dark Matter density $\rho_U \approx 5.6 \times 10^{-124}$ in Planck units & the radius $R_{OH} \approx 2.7 \times 10^{61}$ Planck lengths ($\approx 46.5 \times 10^9$ light years) puts $\rho_U \times R_{OH}^2 \approx 0.42$ in Planck units so $\rho_U \times R_{OH}^2 \approx 0.8823 \Upsilon^2 \approx 0.42$ now.

$$\text{Using } \Lambda\text{-CDM model \& WMAP data } \Upsilon = k_{\min} R_{OH} \approx 0.69 \text{ now} \quad (2.4. 13)$$




Figure 2.4. 2 plots $\rho_U \times R_{OH}^2 \approx 0.8823 \Upsilon^2$

The Λ -CDM model & WMAP data Horizon velocity $V = 1 + H(T)R_{OH} \approx 1 + 3.37 \approx 4.37$ if $H(T) \approx 1/T$ now, and putting this and $\Upsilon \approx 0.69$ into Eq.(2.4. 12) we get an approximate value for the graviton coupling constant α_G

$$\text{Using } \Lambda\text{-CDM model \& WMAP data } \alpha_G \approx \frac{\Upsilon^2 V}{4} \approx \frac{(0.69)^2 \times 4.37}{4} \approx \frac{2.08}{4} \approx 0.52 \quad (2.4. 14)$$

Or if $\alpha_G = 1$ as expected, we should expect $\Upsilon^2 V \approx 4$

Using the current Λ -CDM model and WMAP data with $\Upsilon^2 V \approx 2.08$, puts α_G in the right ball park, suggesting that our approximate analyses is not too far off the mark, and that we can perhaps turn it around to show we should expect $\Upsilon^2 V \approx 4$. The Λ -CDM model has to be fine tuned for flatness requiring a critical density. It also has to have a fixed ratio of Dark Energy to total matter to get the observed accelerated expansion. The current figures are $\approx 5\%$ Baryonic $\approx 23\%$ Dark matter and the rest Dark energy. This puts the ratio of Dark Matter to Baryonic as ≈ 4.5 to 1 whereas it can be as much as 9 or even 10 to 1 in some galaxies. If for example it was approaching 10 to 1 then $\Upsilon^2 V \approx 4$ at the current horizon velocity and horizon radius we used above. However the most important part of the above is that $\Upsilon^2 V$ has to be constant, and as we will see this naturally leads to exponential expansion. In the next section we will find slightly different values for both the horizon radius and velocity, which combined with a smaller increase in Dark matter of about 6.5 to 1, can give $\Upsilon^2 V \approx 4$. And it all only works in flat on average space.

2.4.3 A possible exponential expansion in a flat matter dominated universe

We have been assuming space is flat on average, and this allows very simple calculations. Using Euclidean space properties, of equal coordinate, proper and comoving distances at the current time, let the scale factor be a with density $\rho \propto \frac{1}{a^3}$ in this era. Eq. (2.2. 10) tells us the

average density of the universe $\rho_U \approx 0.8823 \frac{\Upsilon^2}{R_{OH}^2}$ or $\rho_U = K \frac{\Upsilon^2}{R_{OH}^2} = \frac{1}{a^3}$ where

$$K = 0.8823 \text{ is constant and } a^3 = KR^2\Upsilon^{-2} \rightarrow a = K'R^{2/3}\Upsilon^{-2/3} \text{ where } R = R_{OH} \quad (2.4. 15)$$

The Hubble parameter H is

$$H = \frac{\dot{a}}{a} = \frac{(2/3)K'R^{-1/3}\Upsilon^{-2/3} dR/dt}{K'R^{2/3}\Upsilon^{-2/3}} - \frac{(2/3)K'R^{2/3}\Upsilon^{-5/3} d\Upsilon/dt}{K'R^{2/3}\Upsilon^{-2/3}} = \frac{2}{3} \left[\frac{1}{R} \frac{dR}{dt} - \frac{1}{\Upsilon} \frac{d\Upsilon}{dt} \right]$$

Thus the Hubble Horizon flow velocity @ R_{OH} is $V' = H \cdot R = \frac{2}{3} \left[\frac{dR}{dt} - \frac{R}{\Upsilon} \frac{d\Upsilon}{dt} \right]$ (2.4. 16)

We can also write either of Eq's.(2.4. 14) as $\Upsilon^2 V = \text{a constant } K$, with $\Upsilon^2 dV + 2\Upsilon d\Upsilon V = 0$.

Thus $\frac{1}{2V} \frac{dV}{dT} = -\frac{1}{\Upsilon} \frac{d\Upsilon}{dT}$ and Eq. (2.4. 6) tells us that the Horizon velocity $V = \frac{dR_{OH}}{dt} = \frac{dR}{dt}$.

Equation (2.4. 6) also tells us that $V' = H \cdot R = V - 1$ so we can write Eq. (2.4. 16) as

$$\left[3(V-1) - 2V = -\frac{2R}{\Upsilon} \frac{d\Upsilon}{dt} = \frac{2R}{2V} \frac{dV}{dt} \right] \rightarrow V-3 = \frac{R}{V} \frac{dV}{dt} \rightarrow \frac{dV}{dt} = \frac{V}{R} (V-3) \quad (2.4. 17)$$

We will look for an exponential increase of the horizon velocity so $dV/dt > 0$ and $3 \leq V \leq \infty$. Let us simply try $V = 3\text{Exp}(bt)$ with $V > 3$ for all values of b & $t > 0$. Assume a starting time after transition of $t_0 \approx 0$ initially, and only consider times $t \gg t_0$.

If space is flat we can simply put $R = \int_{t_0}^t V dt = \int_{t_0}^t 3\text{Exp}(bt) dt$ & $R \approx \frac{3[\text{Exp}(bt)-1]}{b}$ if $(t \gg t_0)$

Putting this value for R plus $V = 3\text{Exp}(bt)$ & $V-3 = 3[\text{Exp}(bt)-1]$ into Eq. (2.4. 17)

$$\frac{dV}{dt} = \frac{V}{R} (V-3) = 3\text{Exp}(bt) \cdot \frac{b}{3[\text{Exp}(bt)-1]} \cdot 3[\text{Exp}(bt)-1] = 3b\text{Exp}(bt).$$

But $V = 3\text{Exp}(bt)$ and again $\frac{dV}{dt} = \frac{d}{dt} [3\text{Exp}(bt)] = 3b\text{Exp}(bt)$. Thus Eq's. (2.2. 10) & (2.4. 14)

are consistent with $V = 3\text{Exp}(bt)$ for positive b regardless of the value of graviton coupling α_G

$$\text{A possible expansion solution is } V = 3\text{Exp}(bt) \text{ \& } R = \frac{3[\text{Exp}(bt)-1]}{b}, b > 0. \quad (2.4. 18)$$

But is this consistent with the local special relativity requirement for R_{OH} ? In other words does R @ time $T = a(T) \int_0^T \frac{dt}{a(t)} = \frac{3[Exp(bT) - 1]}{b}$? Now Eq. (2.4. 15) tells us the scale factor $a^3 = KR^2\Upsilon^{-2} \rightarrow a = K'R^{2/3}\Upsilon^{-2/3}$ but Eq.(2.4. 14) says $\Upsilon^2 \propto 1/V$ so the scale factor $a \propto V^{1/3}R^{2/3}$. From Eq. (2.4. 18) ignoring the constant factors 3 & b, $V \propto Exp(bt)$ & $R \propto [Exp(bt) - 1]$

$$\begin{aligned} \text{The scale factor } a(t) &\propto Exp(bt)^{1/3}[Exp(bt) - 1]^{2/3} \text{ and } R = a(T) \int_0^T \frac{dt}{a(t)} \\ &= Exp(bT)^{1/3}[Exp(bT) - 1]^{2/3} \int_0^T \frac{dt}{Exp(bt)^{1/3}[Exp(bt) - 1]^{2/3}} \\ &= \frac{3[Exp(bT) - 1]}{b} \end{aligned} \quad (2.4.19)$$

And Eq. (2.4. 18) appears to be a consistent exponential expansion for both V and R .

From Eq. (2.4. 14) we showed $\frac{1}{2V} \frac{dV}{dT} = -\frac{1}{\Upsilon} \frac{d\Upsilon}{dT}$. Using Eq. (2.4. 18) $V = 3Exp(bt)$ & $\frac{dV}{dt} = 3bExp(bt)$ implies $\Upsilon = K \cdot Exp(-bt/2)$. The current Λ -CDM/WMAP value $\Upsilon \approx 0.69$ from Eq. (2.4. 13), and our best guess of $b \approx 0.48$ from Figure 2.4. 3 yields

$$\Upsilon = k_{\min} R_{OH} \approx 0.88Exp(-0.24t) \text{ in radians} \quad (2.4. 20)$$

Several of the above formulae only apply in Euclidean or flat on average space.

This simple exponential expansion starts after the transition, and is very different to current Λ -CDM models, which keep the Hubble parameter $H = \dot{a}/a \approx 2/3t$ constant (if $\Omega = 1$) until Dark Energy starts to take effect. Current Λ -CDM models put the Hubble parameter as $H = \dot{a}/a \approx 1/T$ at present (based on $T \approx 13.8 \times 10^9$ years). In the plots below we put $T \approx 13.8 \times 10^9 \text{ years} = 1$, with R_{OH} or radius R becoming multiples of $T = 1$. Using Eq. (2.4. 6) $V = 1 + H(T)R$, Figure 2.4. 3 plots the Hubble parameter by time ($T = 1$) now, as a function of the exponential time coefficient b , showing if $b = 0$ that $H(t)$ always $= 2/(3t)$ as in current cosmology at critical density with no dark energy. Also if $H \approx 1/T$ now, the best guess is $b \approx 0.48$. This yields $R \approx 3.85T$ or $\approx 15\%$ greater than current cosmology. Figure 2.4. 4 plots horizon velocity which @ $V \approx 4.85$ now is also $\approx 11\%$ greater. The current Λ -CDM model puts Baryonic matter at $\approx 5\%$ and Dark matter at $\approx 23\%$ but if we make this ratio say about 6.4 to 1, the total matter density of the universe increases from $\approx 28\%$ to $\approx 37\%$, and ρ_U increases as $37/28 \approx 1.32$. Now $\rho_U \times R_{OH}^2 \approx 0.8823\Upsilon^2$ and if R_{OH} is 15% greater, then $\Upsilon'^2 \approx 1.32 \times 1.15^2 \times \Upsilon^2 \approx 1.75 \times 0.47 \approx 0.825$ as $\Upsilon^2 \approx 0.47$ in current Λ -CDM models. If $V' \approx 4.85$ then $\Upsilon'^2 V' \approx 4$ which fits our model. Figure 2.4. 5 plots the scale factor based on $b \approx 0.48$, but of course the actual value of b or rate of change with time must be in agreement with the redshifts currently observed when looking back in time. These could well change b and radius R . Figure 2.4. 6 plots the transition to positive acceleration of the scale factor showing the effect of changing the value of b .




Figure 2.4. 3




Figure 2.4. 4




Figure 2.4. 5




Figure 2.4. 6

These plots show an $\approx 15\%$ increase in Horizon Radius, an $\approx 11\%$ increase in Horizon Velocity, which if combined with an increased Dark Matter to Baryonic Matter ratio of ≈ 6.5 (versus the Λ -CDM ratio ≈ 4.5) gives the required $\Upsilon^2 V \approx (0.91)^2 \times 4.85 \approx 4$




Figure 2.4. 7 Plots $\Upsilon = k_{\min} R_{OH} \approx 1.155 \text{Exp}(-0.24t)$ out to 10 times the age of the universe.

2.4.4 The radiation dominated era up to approximately 47,000 years




Figure 2.4. 8 An example of a possible radiation dominated exponential expansion.

In the mass dominated era the density $\rho \propto 1/a^3$, and in the preceding radiation dominant era $\rho \propto 1/a^4$. We can repeat section 2.4.3 with horizon velocity $V = 2\text{Exp}(ct)$, & horizon radius $R = \int_0^t V dt = \int_0^t 2\text{Exp}(ct) dt = 2[\text{Exp}(ct) - 1]/c$. The horizon velocity starts @ $V = 2$ and horizon radius @ $R = 2t$ with a scale factor $a \propto t^{1/2}$ and is the value used in current models of this era before transition, which predict results in close agreement with the current measurements of normal matter in the universe. At the start of the matter dominated era, $V = 3$, and one possibility is $V \approx 2\text{Exp}(0.4t)$, where time is normalized to one at the end of this era. This exponential expansion can, with some smooth transition, continue on into the $V = 3\text{Exp}(0.48t)$ of the matter era, with the normalization of time then changed to the current

cosmos age. If $\alpha_G = 1$ as in Eq.(2.4. 14) with $V = 2$, Υ has to be ≈ 1.4 initially to get $\Upsilon^2 V \approx 4$. By the end of this era when the horizon velocity has increased to $V \approx 3$, Υ has exponentially reduced to $\Upsilon \approx 1.155$ to keep $\Upsilon^2 V \approx 4$. Because the transition time is so small in relation to the $\approx 10^{10}$ year age of the universe, this era has insignificant affect on all our above graphs.

2.4.5 Peculiar velocities and non-commoving coordinates

We have up to here been focussing more on commoving coordinates for simplicity. Velocities relative to comoving coordinates are usually referred to as peculiar velocities, so, does all our previous work still apply in non-comoving coordinates? The average momentum of the universe is zero in comoving coordinates and all background gravitons are time polarized. As explained in section 2.1.2, if we move at a peculiar velocity, equal and opposite gravitomagnetic vectors all sum to zero, resulting in zero spatially polarized gravitons, just as the magnetic field is zero at the centre of long circular conductors with uniform currents. Thus the background, at the centre of any observer's universe, in non-rotating spacetime, always contains only time polarized or spherically symmetric k_{\min} gravitons, regardless of peculiar velocities.

We can think of a box of these k_{\min} gravitons fixed in commoving coordinates. It will have a 3 volume density $\rho_{Gk_{\min}}^{3V} = K_{Gk_{\min}} dk_{\min}$ as we have previously calculated. (The superscripts here are for convenience only, and nothing to do with tensors.) If we now move relative to it at peculiar velocity β_P it will shrink in size as $\gamma_P^{-1} = \sqrt{1 - \beta_P^2}$ so that it's new 3 volume density $\rho_{Gk_{\min}}^{3V} = K_{Gk_{\min}} dk_{\min}''$, where $dk_{\min}'' / dk_{\min} = \gamma_P$ is the local increase in wavenumber k_{\min} . If we repeat our derivations of the background 3 volume density, and the extra emitted by local mass concentrations, we find they also both increase by $dk_{\min}'' / dk_{\min} = \gamma_P$ with no change in the ratio $\Delta\rho / \rho$, so all our logic is unchanged at any peculiar velocity. But all this, is the same as saying that at any peculiar velocity, and in any metric, the 4 volume density of k_{\min} gravitons is Invariant at any cosmic time T.

2.4.6 Invariant 4 volume cosmic wavelength graviton densities

$$\text{Define } \rho_{Gk_{\min}}^{3V} = \frac{k_{\min} \text{ Gravitons}}{3 \text{ Volume}} = \frac{k_{\min} \text{ Gravitons}}{\Delta x \Delta y \Delta z} \text{ and as 4 volume } \Delta x \Delta y \Delta z \Delta t = \Delta x' \Delta y' \Delta z' \Delta t'$$

$$\rho_{Gk_{\min}}^{4V} = \frac{k_{\min} \text{ Gravitons}}{4 \text{ Volume}} = \frac{k_{\min} \text{ Gravitons}}{\Delta x \Delta y \Delta z \Delta t} = \frac{k_{\min} \text{ Gravitons}}{\Delta x' \Delta y' \Delta z' \Delta t'} \text{ is an invariant.}$$

We will define 4 volume k_{\min} graviton density at any point in space-time as

$$\begin{aligned} & \text{4 Volume Density } \rho_{Gk_{\min}}^{4V} = \text{3 Volume Density } \rho_{Gk_{\min}}^{3V} \\ & \text{but only in commoving flat space coordinates, however} \\ & \rho_{Gk_{\min}}^{4V} \text{ is invariant in all coordinates and in any metric.} \end{aligned} \tag{2.4. 21}$$

This is equivalent to dividing k''_{\min} , in any metric, at any peculiar velocity, by $\gamma_p \gamma_M$ thus returning it to flat space commoving value k_{\min} at any cosmic time T . Similarly Eq.(2.2. 16)

$$\rho_{Gk_{\min}}^{3V}(\text{Total}) = K_{Gk_{\min}} \frac{dk''_{\min}}{\gamma_M^2} + \beta_M^2 K_{Gk_{\min}} dk''_{\min} = K_{Gk_{\min}} dk''_{\min} \text{ in commoving coordinates,}$$

which can be written as

$$\rho_{Gk_{\min}}^{3V}(\text{Total}) = K_{Gk_{\min}} \frac{dk''_{\min}}{\gamma_M^2} + \Delta\rho_{Gk_{\min}}^{3V}, \text{ becomes using invariant 4 volume notation}$$

$$\rho_{Gk_{\min}}^{4V}(\text{Total}) = K_{Gk_{\min}} \frac{dk_{\min}}{\gamma_M^2} + \beta_M^2 K_{Gk_{\min}} dk_{\min} = K_{Gk_{\min}} dk_{\min} \text{ which can be written as}$$

$$\rho_{Gk_{\min}}^{4V}(\text{Total}) = K_{Gk_{\min}} \frac{dk_{\min}}{\gamma_M^2} + \Delta\rho_{Gk_{\min}}^{4V}$$

Where $\Delta\rho_{Gk_{\min}}^{4V} = \beta_M^2 K_{Gk_{\min}} dk_{\min}$; $\Delta\rho_{Gk_{\min}}^{3V} = \beta_M^2 K_{Gk_{\min}} dk''_{\min}$; $\frac{dk''_{\min}}{dk_{\min}} = \frac{\Delta x \Delta y \Delta z}{\Delta x' \Delta y' \Delta z'} = \frac{\Delta t'}{\Delta t}$

As both $\rho_{Gk_{\min}}^{4V}$ & $\Delta\rho_{Gk_{\min}}^{4V}$ are invariant, their ratio is also invariant in any coordinates, and at any peculiar velocity at any particular cosmic time. But the flat space commoving value of k_{\min} decreases with cosmic time. We also know that $\beta_M^2 = 2m / r$ is always true.

2.4.7 Cosmic wavelength graviton action densities

In deriving Eq. (2.4. 2) we said that each k_{\min} graviton always borrows a fixed amount of action, where $Action = \Delta E \cdot \Delta T$ per graviton is constant but $\Delta E \propto k_{\min}$. So if four volume density $\rho_{4V_{Gk_{\min}}}$ is invariant the four volume action density required by k_{\min} gravitons must also be invariant. In Eq.(2.4. 11) we calculated the k_{\min} action or quanta density available from the horizon in commoving coordinates. But if we move at a peculiar velocity β_p , both energy ΔE and time ΔT increase as γ_p , so the four volume action density available from this source should increase by γ_p^2 and not be invariant, appearing to destroy our logic. (If there is more action available than what is required by k_{\min} gravitons there is nothing to keep their density controlled.) If we go back to the building of superpositions (page 13 in [7]), we see that in their rest frame, they have a wave equation generated from a vector potential squared, or A^2 term. This can only come from a spherically symmetric source of spatially polarized quanta. At high frequencies this spherically symmetric source is the invariant vector potential squared portion of the local zero point fields. At low frequencies this source is from the receding horizon, but it must also be spherically symmetric (see sections 2.4.1, 2.7.2 & 2.7.3). We derived Eq. ((2.4. 11) in commoving coordinates where the spatially polarized source from the horizon is spherically symmetric as required to build superpositions in their rest frame. But it is not spherically symmetric at peculiar velocities. We showed (page 22 in [7]) that if we move at a peculiar velocity β_p relative to a spherically symmetric spin one source, its probability of being spherically symmetric is $1/\gamma_p^2$. But the 4 volume action density increases as γ_p^2 so these two factors cancel. Exactly the same happens in any metric.

The 4 volume k_{\min} spin one quanta action density from the horizon increases as γ_M^2 but the spherical symmetry probability drops as $1/\gamma_M^2$, and both these effects cancel again.

Spherically symmetric 4 volume k_{\min} action density available from the receding horizon is always invariant at any particular cosmic time in a region of space. (At any cosmic time it depends on the value of k_{\min} in commoving flat space but decreases with cosmic time.)

So our hypothesis is that at any point in spacetime: Gravity is consistent with the spherically symmetric 4 volume action density available from a receding horizon always being equal to the spherically symmetric 4 volume k_{\min} action density required by gravitons; with both remaining invariant in any coordinates.

We will use the superscript ss for spherically symmetric invariant 4 volume densities.

Define Invariant Spherically Symmetric 4 Volume Action Density as $\rho_{Qk_{\min}}^{4VSS}$

Where $\rho_{Qk_{\min}}^{4VSS}$ action available $\approx 0.48(\rho_{Gk_{\min}}^{4VSS})$ required by k_{\min} gravitons. (2.4. 22)

This equation is true in any coordinates, and at any point in spacetime.

2.4.8 If Gravity is due to metric changes then what about exchanged momentum?

Let us now consider exchanging k_{\min} time polarized gravitons between Planck masses (or in fact between any masses) instead of simply considering k_{\min} graviton densities. As we have noted many times, by far the vast majority of gravitons in the universe are near k_{\min} , so we will only look at the effect of k_{\min} exchanges. If the 4 volume densities of both spherically symmetric k_{\min} action available, and k_{\min} gravitons are invariant everywhere, each mass is interacting with the rest of the universe in a spherically symmetric manner. Quantum field theory tells us that coulomb or scalar forces are due to the exchange of virtual photon 3 momentum. Assuming virtual gravitons are no different, and this exchange happening in a spherically symmetric manner, there can be no nett force in any direction due to these exchanges, but only momentum squared terms. If two masses are orbiting each other they will also be exchanging higher frequency gravitons, but to a first approximation the effect of these is small in relation to the spherically symmetric exchanges of k_{\min} gravitons with the rest of the universe. But perhaps spin 2 gravitons do not exchange 3 momentum in such interactions. On Page 34 in [7], at the bottom, we mention a possible such interaction between superpositions with no 3 momentum exchanged, but only time will tell. Regardless, the warping of spacetime does guarantee spherical symmetry of k_{\min} exchanges.


Einstein always felt that gravitational forces were fictitious, that gravity was due to metric changes only, and not exchanged gravitons. If in time, our hypothesis proves to be true, he may well prove to have been correct.

2.5 An Infinitesimal change to General Relativity at Cosmic Scale

We started everything here in flat space with no mass concentrations. So, uniform densities don't curve space. We introduced mass concentrations and space has to curve around them so as to keep our spherically symmetric 4 volume k_{\min} action densities, required and available, invariant. If we think of the mass in the universe as a dust of density ρ_U essentially at rest in comoving coordinates we can define a tensor $T_{\mu\nu}$ (Background). In comoving coordinates $T_{\mu\nu}$ (Background) has only one non zero term T_{00} (Background) = ρ_U . In any other coordinates this same $T_{\mu\nu}$ (Background) tensor is transformed by the usual tensor transformations that apply in GR. If these coordinates move at peculiar velocity β_p then T_{00}' (Background) = $\gamma_p^2 \rho_U = \gamma_p^2 T_{00}$ (Background). The same transformation happens in any metric but with $\gamma_M^2 \rho_U$. We argue that Eq.(2.4. 22) is consistent with the infinitesimally modified Einstein field equations

$$G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi G}{c^4} [T_{\mu\nu} - T_{\mu\nu} \text{ (Background)}] \quad (2.5. 1)$$

This infinitesimal modification is only relevant in the extreme case as $T_{\mu\nu}$ approaches $T_{\mu\nu}$ (Background). Far from mass concentrations $T_{\mu\nu} \leq T_{\mu\nu}$ (Background). Space curvature, in these remote voids, is in general somewhere between slightly negative and zero; but the causally connected universe is always flat on average regardless of the value of Ω . Equation (2.5. 1) is also consistent with Eq. (2.1. 10) $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ at all cosmic times. If there is no inflation, in flat comoving coordinates, at the Big Bang or slightly after, k_{\min} starts at just under one and is always close to the inverse of the causally connected horizon radius. It is also close to the inverse of cosmic time T . It is always at its minimum far from mass concentrations, but increases with the slower clock rates in the local metric around mass concentrations as in Figure 2.5. 1



2.5.1 Is inflation in this proposed scenario really necessary?

There are two main reasons, usually given, for why inflation is necessary:

(a) The average flatness of space.

(b) The almost uniform temperature of the cosmic microwave background from regions that were initially out of causal contact.

If we put $T_{\mu\nu}(\text{Local}) = T_{\mu\nu}(\text{Background})$ in Eq.(2.5. 1), the right hand side is identically zero,

and $G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \equiv 0$ on average throughout all space. The average curvature of all

space must be zero and space is compelled to be flat on average. In section 2.4.3 we found that space naturally expands exponentially as in Eq. (2.4. 18) and plotted in Figure 2.4. 4. The value of the constant b in $V = 3Exp(bt)$ has to fit experimental observations. But if it is some fundamental constant, which does not seem unreasonable, it must be the same for all comoving observers. If this is so, the physics is identical for all such observers regardless of whether they are in causal contact. Provided we can assume identical starting points everywhere, of say the Planck temperature at cosmic time $T = 0$, then apart from quantum fluctuations, the average background temperature should be some function of cosmic time T for all comoving observers, or at least up to the time the universe became transparent. The physics controlling this should be identical in each comoving frame. Causal contact should not be essential for this. Inflation only guarantees that the starting temperature is uniform everywhere when it stops at approximately $T = 0$. It also has to assume identical physics everywhere from $T \approx 0$ for about the first 375,000 years, or until the universe is transparent. What we are proposing in this paper should produce results not too different from this.

2.5.2 Why do we think virtual particle pairs do not matter?

For almost a century it has been a puzzle why spacetime is not massively curved by Planck scale zero point energy densities. However space appears to be flat on average regardless of this massive Planck scale zero point energy density so **something must be different and what is it?** In the first paper [7] we conjectured that virtual particles are just single wavenumber k superposition members, whereas real, or observable particles are full infinite superpositions of all wavenumbers k from k_{\min} to k_{Planck} . Only full infinite superpositions have real properties that can be measured (such as measured mass/energy) rather than implied. Because k_{\min} virtual gravitons are such single members they can couple to k_{\min} members of full infinite superpositions. On the other hand virtual particles out of the vacuum are mainly short lived high k single value members that will not couple to k_{\min} ; provided these virtual particles are single wavenumber k members only, and of a higher k value than k_{\min} . The density of k_{\min}

virtual pairs from the vacuum is virtually zero, as it is based on the Lorentz invariant supply of local zero point fields, not from the receding horizon (sections 2.7.2 & 2.7.3 clarify this). But this is not the full story. The virtual particles that dress electrons and quarks for example add mass to the real particles. In fact the majority of proton and neutron mass is due to the virtual gluons interacting between quarks. If short lived virtual particles somehow contribute to the mass of full infinite superpositions, then these high k value virtual particles indirectly contribute to the k_{\min} virtual graviton coupling, which is based on the actual mass of the infinite superposition as in Eq. (3.2.3) in [7]. The conservation of energy, or more correctly 4 momentum, says that what we call “real matter or energy” can last for close to the age of the universe. It will have mass and by definition it can be weighed. It can move around, even close to the speed of light, but it is conserved. Gravitons that last this long we have called k_{\min} gravitons and they can only couple to real, or long lasting energy/matter that can be weighed in whatever manner. The rotating dark matter in galaxies we cannot weigh directly, but it contributes to the theoretical weight of a galaxy. We have to allow for this mass when studying galaxy dynamics.

The particle beams in accelerators have real energy which can be temporarily converted into virtual particles. The total energy or 4 momentum is always conserved, but can fluctuate for time $\Delta T \approx 1 / 2\Delta E$. The long term average is what counts. In this sense the mass of short lived virtual particles can contribute to k_{\min} virtual graviton coupling, just as it does in the virtual particle dressing of real charged particles as above. If it can be somehow weighed, it will couple to k_{\min} virtual gravitons. But we cannot weigh the zero point background.

2.6 Messing up what was starting to look promising, or maybe not?

2.6.1 The k_{\min} virtual gravitons emitted by the mass interacting with itself

In section 2 we started out by finding the average k_{\min} graviton probability density in a uniform universe. We then placed a mass concentration in it, and calculated the extra probability density of k_{\min} gravitons (before the dilution due to local space expansion) due to the amplitude of this mass multiplied by the amplitude of the rest of the mass in the universe. This ended up being proportional to $2m/r$ in Planck units. (In this section we revert to simple 3 volume k_{\min} graviton probability densities, with no need for 4 volume superscripts.)

$$\Delta\rho_{Gk_{\min}} = (\psi_{\text{Universe}} * \psi_m) + (\psi_m * \psi_{\text{Universe}}) \propto 2m/r \text{ as in Eq.(2.2. 6)}$$

And this is true in weak field metrics, except as we start approaching the Schwarzschild radius because of the extra k_{\min} gravitons from the mass interacting with itself: $\psi_m * \psi_m$.

$$\text{Using Eq. (2.1. 5) and coupling probability: } (1 - e^{-0.61k^2/k_{\min}^2}) \left[\frac{2\alpha_G}{\pi} m^2 \frac{dk}{k} \right]$$

$$\psi_m * \psi_m = (1 - e^{-0.61k^2/k_{\min}^2}) \left[\frac{2\alpha_G}{\pi} m^2 \frac{dk}{k} \right] \cdot \left[\frac{2k'}{4\pi} \frac{e^{-2k'r}}{r^2} \right] = (1 - e^{-0.61k^2/k_{\min}^2}) \alpha_G \frac{m^2}{r^2} \frac{k'e^{-2k'r}}{\pi^2} \frac{dk}{k}$$

$$\text{Also using Eq. (2.1. 4) } k' = \sqrt{k^2 + 11.09k_{\min}^2} \approx 3.477k_{\min} \text{ when } k = k_{\min}$$

$$\begin{aligned} \psi_m * \psi_m &= (1 - e^{-0.61}) \alpha_G \frac{m^2}{r^2} \frac{3.477k_{\min} e^{-2(3.477k_{\min}r)}}{\pi^2} \frac{dk_{\min}}{k_{\min}} \\ &= \alpha_G \frac{m^2}{r^2} \frac{1.588e^{-6.95k_{\min}r}}{\pi^2} dk_{\min} \text{ when } k = k_{\min} \end{aligned}$$

The radial exponential decay term $e^{-6.9k_{\min}r} \approx 1$ as we are only interested in radii r that are small in relation to the observable radius of the Universe $R_{OU} \approx k_{\min}^{-1}$, just as in the assumptions we made in section 2.2.1. Thus in these regions we can approximate this equation with good accuracy as

$$\psi_m * \psi_m \approx \alpha_G \frac{m^2}{r^2} \frac{1.588}{\pi^2} dk_{\min}$$

$$\Delta\rho_{Gk_{\min}} \text{ due to self emission } \psi_m * \psi_m \approx \alpha_G \frac{m^2}{r^2} 0.161 dk_{\min}$$

$$\approx 1.4 \frac{m^2}{r^2} 0.115 \alpha_G dk_{\min} \text{ when } k = k_{\min}$$

$$\Delta\rho_{Gk\min} \text{ due to } \psi_m * \psi_m \approx 1.4 \frac{m^2}{r^2} K_{Gk\min} dk_{\min} \text{ using Eq.(2.2. 11)} \quad (2.6. 1)$$

2.6.2 What does this extra term mean for non rotating black holes?

When deriving Eq.(2.2. 14) we found (about two equations previous) that due to interactions with the rest of the Universe $\Delta\rho_{Gk\min} \approx \frac{2m}{r} 0.115\alpha_G dk_{\min} \approx 2 \frac{m}{r} K_{Gk\min} dk_{\min}$

$$\text{Thus } \Delta\rho_{Gk\min} \text{ total} \approx \left[2 \frac{m}{r} + 1.4 \frac{m^2}{r^2} \right] K_{Gk\min} dk_{\min} \text{ in Planck units.} \quad (2.6. 2)$$

Staying on our current path appears to contradict General Relativity, but temporarily ignoring this, let us repeat section 2.2.2 which modifies a non rotating black hole metric to

$$g'_{00} = 1 - \frac{2m}{r} - 1.4 \frac{m^2}{r^2} = \frac{1}{g'_{rr}}; \quad \gamma_M^2 = \frac{1}{1 - 2m/r - 1.4m^2/r^2} \quad (2.6. 3)$$

$$\beta_M^2 = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$$

Where β_M is the velocity reached by a small test mass falling in from infinity in the same rest frame. Applying the same procedures as in section 2.2.2 we can use Equ's. (2.6. 3) to show that $\rho_{Gk\min} = K_{Gk\min} dk_{\min}$ in this new metric, and we will discuss how this relates with General Relativity in sections 2.6.5 & 2.6.6. The modified non rotating horizon radius occurs when $r^2 - 2mr - 1.4m^2 = 0$ or the:

$$\text{Modified non rotating horizon radius } r \approx 2.55m \quad (2.6. 4)$$

or $\approx 27.5\%$ larger than the Schwarzschild value.

2.6.3 What does it mean for rotating black holes?

In section 2.3 when we looked at the Kerr Metric we used a dimensionless form of the metric in Equ's.(2.3. 2). We also used a dimensionless parameter A where we initially put $A = 2m/r$. We also showed that we could change A without changing $g'_t = \Delta / g_{\varphi\varphi}$ the time component in the corotating frame, provided there is a modified $\Delta = 1 + \frac{\alpha^2}{r^2} - A$. So again temporarily ignoring potential conflicts with General Relativity let us change $A = \frac{2m}{r}$ to

$A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$ and look at the consequences. Firstly from Equ's. (2.6. 3) we can see that $A = \beta_M^2$ where β_M is the radial inward velocity, in a corotating rest frame, of a small test mass

falling from infinity (in the rest frame of the rotating black hole centre). The inner event horizon is the radius where $g_{rr} \rightarrow \infty$ so using Equ's.(2.3. 2) let $g_{rr} = \frac{g_{\theta\theta}}{\Delta} \rightarrow \infty$ or put

$$\begin{aligned}\Delta &= 1 + \frac{\alpha^2}{r^2} - A = 0 \\ &= 1 + \frac{\alpha^2}{r^2} - \frac{2m}{r} - 1.4 \frac{m^2}{r^2} = 0 \\ \text{or } r^2 + \alpha^2 - 2mr - 1.4m^2 &= 0\end{aligned}$$

$$r = \frac{2m \pm \sqrt{4m^2 + 5.6m^2 - 4\alpha^2}}{2}$$

Event Horizon radius $r = \frac{2m \pm \sqrt{9.6m^2 - 4\alpha^2}}{2}$ (2.6. 5)

When $\alpha = 0$ $r = \frac{2m \pm \sqrt{9.6m^2}}{2} \approx \frac{2m + 3.1m}{2} \approx 2.55m$ as in the non rotating case.

Maximum spin is when $4\alpha^2 = 9.6m^2$ or $\alpha_{\max} \approx 1.55m$

At this maximum spin $r = m$ as in the usual Kerr Metric.

The outer horizon occurs when $g_{tt} = 1 - \frac{A}{g_{\theta\theta}} = 0$ or $g_{\theta\theta} - A = 0$ and using Equ's.(2.3. 2)

$$\begin{aligned}1 + \frac{\alpha^2}{r^2} \cos^2 \theta - A &= 1 + \frac{\alpha^2}{r^2} \cos^2 \theta - \frac{2m}{r} - 1.4 \frac{m^2}{r^2} = 0 \\ r^2 - 2mr - 1.4m^2 + \alpha^2 \cos^2 \theta &= 0\end{aligned}$$

$$r = \frac{2m + \sqrt{4m^2 + 5.6m^2 - 4\alpha^2 \cos^2 \theta}}{2}$$

$$r = \frac{2m + \sqrt{9.6m^2 - 4\alpha^2 \cos^2 \theta}}{2}$$

Ergosphere radius $r = \frac{2m + \sqrt{9.6m^2 - 4\alpha^2}}{2}$ @ $\theta = 0$ & π (2.6. 6)

$$= \frac{2m + \sqrt{9.6m^2}}{2} \approx 2.55m \text{ @ } \theta = \frac{\pi}{2}$$

Figure 2.6.1 illustrates these changes from the Kerr Metric. The main effect from changing A is to allow an increase in maximum spin from $\alpha = m$ to $\alpha \approx 1.55m$, and $\approx 27.5\%$ increase in the maximum ergosphere radius from $r = 2m$ to $2.55m$. It appears to contradict General Relativity which we discuss in sections 2.6.5 & 2.6.6, but provided the extra densities of time polarized and $m = \pm 2$ circular gravitons are as in Eq.(2.3. 7) with $A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$ then

$\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ is still true in rotating space outside black holes.




Figure 2.6.1 Modified Kerr Metric with the dimensionless parameter A changed from $A = \frac{2m}{r} \rightarrow A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$. It initially appears to clash with GR near the horizon.




Figure 2.6. 2 Spinning black hole mass m with angular momentum length parameter α , but with the dimensionless parameter A changed from $A = \frac{2m}{r} \rightarrow A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$. The determinant of the metric is independent of A . The denominator terms $g_{\theta\theta}$ & $g_{\phi\phi}$, in dimensionless form as in Equ's. (2.3. 2), rapidly tend to one for radii $r \gg r_{sw}$, and can then be ignored. It shows the probability densities of time polarized, and circularly polarized $m = \pm 2$ k_{\min} gravitons as in Eq.(2.3. 7) in this modified metric which keeps the k_{\min} graviton constant $K_{Gk_{\min}}$ invariant outside the black hole. This is as observed in corotating coordinates.

2.6.4 Determinant of the metric and the k_{\min} graviton constant $K_{Gk_{\min}}$

Working in dimensionless form as in Equ's.(2.3. 2), using Eq. (2.3. 3) $g'_{tt} = \frac{\Delta}{g_{\phi\phi}}$ and the steps used in its derivation; the determinant of the metric is

$$|g_{\mu\nu}| = (g_{tt}g_{\phi\phi} - g_{t\phi}^2)g_{\theta\theta}g_{rr} = g'_{tt}g_{\phi\phi}g_{\theta\theta}g_{rr} = \frac{\Delta}{g_{\phi\phi}}g_{\phi\phi}g_{\theta\theta} \frac{g_{\theta\theta}}{\Delta} = g_{\theta\theta}^2 = (1 + \frac{\alpha^2}{r^2} \cos^2 \theta)^2$$

As 4 volumes are invariant in relativity and $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ is true in corotating frames

$$\text{If } |g_{\mu\nu}| = g_{\theta\theta}^2 = (1 + \frac{\alpha^2}{r^2} \cos^2 \theta)^2 \text{ then } \rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min} \text{ is true in all frames, and is independent of the dimensionless parameter } A. \quad (2.6. 7)$$

Despite what initially appears to be a conflict with General Relativity (which we discuss below), if the metric determinant Eq. (2.6. 7) is $g_{\theta\theta}^2$ then the k_{\min} graviton probability density is always $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ in all frames outside the black hole, and this is also true if there is no rotation, regardless of the value of the dimensionless parameter A . (See section 2.9)

2.6.5 The Reissner-Nordstrom Metric and m^2/r^2 terms

Reissner [27][28] solved the metric surrounding an electrically charged non-rotating mass not long after Schwarzschild had solved the metric around a static mass. He added the electromagnetic stress tensor surrounding a charge to the usual Einstein Energy-momentum tensor, in the region where the mass density term had previously been zero as in the Schwarzschild case. As before we will put $G = c = 1$ so we can work in Planck masses. The Schwarzschild radius $r_s = 2Gm/c^2$ has length dimension and thus $2Gm/rc^2$ becomes $2m/r$, and both $2m/r$ and m^2/r^2 are effectively dimensionless as, in these units, mass effectively has a length dimension.

Reissner similarly used the characteristic length r_Q where $r_Q^2 = \frac{Q^2 G}{4\pi\epsilon_0 c^4}$

Working in length units of charge with the Coulomb force constant $\frac{1}{4\pi\epsilon_0} = 1$ (2.6. 8)

If $G = c = 1$ & these units of charge $\frac{r_Q^2}{r^2} \equiv \frac{Q^2}{r^2}$ are both dimensionless numbers.

Table 2.6. 1. Both parameters, mass m and charge Q , effectively have dimensions of length.

Metric	Schwarzschild	Modified Schwarzschild	Reissner-Nordstrom
$g_{00} = g_{rr}^{-1}$	$1 - \frac{2m}{r}$	$1 - \frac{2m}{r} - 1.4 \frac{m^2}{r^2}$	$1 - \frac{2m}{r} + \frac{Q^2}{r^2}$

Using our modified Schwarzschild metric from Eq (2.6. 3) we can see the similarities to the Reissner-Nordstrom metric for a charged mass, providing we measure charge parameter Q in a similar manner to measuring mass in Planck units. The signs are reversed however.

We can crudely think of this another way as follows: In the units we have been working in, the electrostatic field energy outside any radius r is $Q^2/2r$. This mass/energy must be subtracted from the original central charged mass as work is done bringing these charged particles together to establish the field energy.

So we can very crudely say the original central mass m becomes $m' = m - \frac{Q^2}{2r}$ at radius r .

$$\text{Thus } \frac{2m'}{r} = \frac{2m}{r} - \frac{Q^2}{r^2} \text{ and the new } g_{00} = 1 - \frac{2m'}{r} = 1 - \frac{2m}{r} + \frac{Q^2}{r^2}.$$

It is very tempting from this, to think of our modified Schwarzschild metric, as somehow including the negative gravitational field energy; which in Planck units is $-m^2/2r$ outside radius r . Using the same logic as the electrostatic case, but reversing signs, as gravitational

field energy is negative, the original central mass m becomes $m' = m + \frac{m^2}{2r}$ at radius r .

$$\text{Thus } \frac{2m'}{r} = \frac{2m}{r} + \frac{m^2}{r^2} \text{ and the new } g_{00} = 1 - \frac{2m'}{r} = 1 - \frac{2m}{r} - \frac{m^2}{r^2}.$$

Of course our coefficient of 1.4 for m^2/r^2 does not fit this scenario, but our analysis is full of approximations and we could have it wrong. Roger Penrose in Chapter 19 of his “Road to Reality” gives a very good discussion on the concerns of many eminent physicists early last century when General Relativity was first published. They worried that gravitational energy was not explicitly included in the stress tensor. But Einstein could not do this and maintain covariance. In the century since, many eminent physicists have tried unsuccessfully to include gravitational energy in a covariant manner. So we must conclude that it is probably not related to gravitational energy; and as we have shown in this section, it is really due to the small number of k_{\min} gravitons (except close in) emitted by the mass itself.

The Maxwell stress tensor tells us in the the electrostatic case, that if the field is in the z direction, there is a tension or negative pressure $P_z = -E^2/2$ along the z axis and transverse positive pressures $P_x = P_y = +E^2/2$ such that $P_x + P_y + P_z = E^2/2$ and the mass/energy density $\rho = E^2/2$ if they are all in appropriate units. The stress tensor contracts to $\rho - P_x - P_y - P_z = 0$ and this is a property of massless particles. Thus the presence of an electromagnetic field does not alter field equation covariance. So if we simply reverse all these signs with a negative mass energy density of $\rho = -1.4m^2/2$ with transverse tensions $P_x = P_y = -1.4m^2/2$ and in the field direction positive pressure $P_z = 1.4m^2/2$ such that the stress tensor contracts again contracts to $\rho - P_x - P_y - P_z = 0$. We can thus include a negative energy massless particle in the stress tensor in the same way as in the positive energy electrostatic case, and similarly maintain covariance.

2.6.6 The Kerr-Newman Metric and m^2/r^2 terms

In 1965 Newman [29][30] solved the charged version of the axisymmetric rotating black hole solved earlier by Kerr [31] in 1962. In section 2.3 and Equ's (2.3. 2) we introduced the dimensionless parameter $A = 2m/r$ where as above we have assumed a silent $G = 1$ in the numerator and a silent $c^2 = 1$ in the denominator and in section 2.6 modified this to get a dimensionless $A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$. We showed in section 2.3 that provided this A is dimensionless it does not change Equ's (2.3. 3). If we look carefully at the Kerr-Newman metric we can see that it fits Equ's (2.3. 2) provided we put $A = \frac{2m}{r} - \frac{r_Q^2}{r^2}$ which is equivalent to putting $A = \frac{2m}{r} - \frac{Q^2}{r^2}$ where $r_Q^2 = \frac{Q^2 G}{4\pi\epsilon_0 c^4}$ and we have again measured charge Q in length units as in Equ's (2.6. 8).

Thus our modified Kerr metric where $A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$ is again similar to:

The Kerr-Newman metric where $A = \frac{2m}{r} - \frac{Q^2}{r^2}$ but with opposite signs.

These two metrics are the rotating versions of our modified Schwarzschild metric and the Reissner-Nordstrom metrics. We can perhaps summarize this in the following two tables.

Table 2.6. 2 The non rotating metrics where dimensionless parameter A is as in Eq. (2.3. 2) The modified Schwarzschild and Reissner-Nordstrom metrics both have the same form of changes to the Reimannian curvature tensor but of opposite sign.

Schwarzschild	Modified Schwarzschild	Reissner-Nordstrom
$A = \frac{2m}{r}$	$A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$	$A = \frac{2m}{r} - \frac{Q^2}{r^2}$

Table 2.6. 3 The rotating versions of the above. Again the modified Kerr and Kerr-Newman metrics both have the same form of changes to the Reimannian tensor but of opposite sign.

Kerr	Modified Kerr	Kerr-Newman
$A = \frac{2m}{r}$	$A = \frac{2m}{r} + 1.4 \frac{m^2}{r^2}$	$A = \frac{2m}{r} - \frac{Q^2}{r^2}$

Again massless particles in the electromagnetic field apply equally in the Reissner-Nordstrom and Kerr-Newmann metrics. The arguments we used above in the non rotating case using massless negative energy particles in our modified stress tensor apply equally in the rotating case. The small changes in the Riemannian curvature tensor, due to this m^2/r^2 term, are of opposite sign for both our modified Kerr and Schwarzschild metrics, when compared to the Kerr-Newman and Reissner-Nordstrom metrics, but of exactly the same form.

So, provided we include such an appropriate negative energy massless particle in the stress tensor, solutions to $R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi G}{c^4} [T_{\mu\nu} - T_{\mu\nu}(\text{Background})]$ are consistent with k_{\min} graviton probability density $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$ where $K_{Gk_{\min}}$ is invariant for all observers; whether they are near the horizon of black holes, or if they are at our current cosmic horizon. Also for any observers outside it, who are looking at their own cosmic horizons; and for all cosmic time since the big bang. But wavenumber k_{\min} depends on the local metric and cosmic time. It is approximately the inverse of the causally connected radius at any cosmic time.

Einstein based his remarkable equation on the “Equivalence Principle”, or the same physics in all free falling frames as in empty space; with covariant tensor equations that apply in all coordinates throughout all spacetime. He wanted it to be similar to Gauss’s law and Poisson’s equation $\nabla^2 \phi = \rho$ ignoring constants, but in curved spacetime. This naturally leads to inverse square force laws with inverse potentials where masses are concerned, but the inclusion of an m^2 / r^2 potential term in the metric due to $\psi_m * \psi_m$ seems to mess all this up. But does it really? Could it be trying to tell us something that we need to know, but did not want to know? Quantum mechanics in the form of QED tells us that, close to the Compton wavelength, the normally simple inverse square force law starts to change, close in shielding makes fundamental electric charge appear to increase, and QED takes over with incredible accuracy. Simple inverse square electric force laws had ruled with remarkable accuracy for over a century before QED arrived on the scene. In fact it was the announcement of the Lamb Shift at the Long Island conference in 1947 that started the big breakthroughs in QED. World War II developments in radar had enabled these remarkably accurate experiments. Is it possible that similar developments today will allow improvements in Gravitational Wave observation accuracy? Developments that may see effects in gravity close to black holes with some parallels to QED changes inside the Compton wavelength of electric charges?

2.6.7 What is the effect of this term in the solar system?

The distance to Mars can be measured very precisely as we have instruments on the surface that can reflect radar from Earth at known locations. On the other hand we don’t know the exact diameter of the Sun. If we look at the outer rim it will be deflected outwards by $\approx 1.75/2$ arc seconds (half that of the gravitational bending of starlight because it is coming from the rim). At the distance of the sun, $\approx 150 \times 10^6$ KM this is roughly 640 KM in radius. Even if we optically measure the diameter precisely with no error the actual sun diameter will be about 1275 KM smaller so we only know the true diameter approximately. We also do not know the exact surface of radar reflection. The Astronomical unit is quoted as 149,597,870,700 metres, but this is really based on knowing interplanetary measurements accurately and then using Kepler’s laws modified by the Schwarzschild metric to give us this

level of accuracy. So, let us do a crude first order approximation of what happens if we include an m^2/r^2 term in the metric. Using low velocity (compared with light) Christoffel symbol approximations and circular orbits for simple comparisons the accelerations are:

$$\omega^2 r \approx \frac{1}{2} \frac{d}{dr} g_{00} \approx \frac{1}{2} \frac{d}{dr} \left(1 - \frac{2m}{r}\right) \approx \frac{m}{r^2} \text{ in the usual Schwarzschild case if } g_{00} \approx 1 \text{ and}$$

$$\omega'^2 r \approx \frac{1}{2} \frac{d}{dr} g'_{00} \approx \frac{1}{2} \frac{d}{dr} \left(1 - \frac{2m}{r} - \frac{1.4m^2}{r^2}\right) \approx \frac{m}{r^2} + \frac{1.4m^2}{r^3} \approx \frac{m}{r^2} \left(1 + \frac{1.4m}{r}\right) \text{ in the modified metric}$$

case. So $\omega^2 \approx \frac{m}{r^3}$ in the usual Schwarzschild case and $\omega'^2 \approx \frac{m}{r^3} \left(1 + \frac{1.4m}{r}\right)$ in the modified metric case. In weak gravitational field accelerations we can replace mass m with a new effective mass $m' = m \left(1 + \frac{1.4m}{r}\right)$ but orbital periods and angular velocities ω cannot change as we know them very precisely. So we will try the following modification to all planetary radii

$$\omega^2 = \frac{m}{r^3} = \frac{m}{(r + \Delta r)^3} \left(1 + \frac{1.4m}{r + \Delta r}\right) = \frac{m}{r^3 \left(1 + \frac{\Delta r}{r}\right)^3} \left(1 + \frac{1.4m}{r + \Delta r}\right) \approx \frac{m}{r^3 \left(1 + \frac{3\Delta r}{r}\right)} \left(1 + \frac{1.4m}{r + \Delta r}\right)$$

$$\omega^2 = \frac{m}{r^3} \approx \frac{m}{r^3} \left(1 - \frac{3\Delta r}{r}\right) \left(1 + \frac{1.4m}{r + \Delta r}\right) \approx \frac{m}{r^3} \left(1 - \frac{3\Delta r}{r} + \frac{1.4m}{r + \Delta r}\right) \text{ and if } \omega \text{ is unchanged}$$

$$1 - \frac{3\Delta r}{r} + \frac{1.4m}{r + \Delta r} \approx 1 \quad \text{and} \quad \frac{3\Delta r}{r} \approx \frac{1.4m}{r + \Delta r} \approx \frac{1.4m}{r} \quad \text{thus} \quad \Delta r \approx \frac{1.4m}{3}$$

The Schwarzschild radius $R_{sw} = 2m$ and the extra distance to the sun $\Delta r \approx \frac{1.4m}{3} \approx \frac{1.4R_{sw}}{6}$

The Schwarzschild radius of the Sun is $R_{sw} \approx 3 \text{ km}$ so $\Delta r \approx \frac{1.4R_{sw}}{6} \approx 0.7 \text{ km}$.

The change Δr for our solar system is about 700 metres. But all interplanetary radial separations do not change. So we can still use the old metric and the astronomical unit unchanged with Kepler's laws to a first approximation, or the new metric and just add 700 metres to all the planetary radii from the sun. Orbital periods are identical to a very high accuracy. The gravitational constant does not change in both cases.

What we have done here is a bit like dipoles with the electrostatic field dropping as $\propto 1/r^2$ and the resultant field as $\propto \Delta r/r^3$ where $\frac{1}{r^2} - \frac{1}{(r + \Delta r)^2} \approx \frac{2\Delta r}{r^3}$. However, in the non-spinning gravity case there is spherical symmetry but not in an electric dipole.

2.6.8 Can we measure this difference?

We used circular orbits for a simple crude calculation but the same arguments apply in a slightly more complicated way for eccentric orbits; in a similar manner as Kepler's original arguments with elliptical orbits that sweep out equal angular segment areas with time. The orbit of Mars in particular is highly eccentric and Earth much less so. If the eccentricity of

both Earth and Mars orbits were known, to better than say a hundred metres or so between max and min, we should be able to check this difference by measuring the distance (also to around a 100 metre or so accuracy) between Mars and Earth at various points around their orbits. It would seem however that this would be pushing at the very border of current technology, as radar measurements to the sun are inherently a little blurry due to surface variability. We need these to get a very precise value for the eccentricity of Earth's orbit. Even if we can measure Earth-Mars with complete accuracy we have to add in errors due to lack of accuracy in Earth's eccentricity. Also when Mars is on the opposite side of the sun there will a Shapiro type delay that is equivalent to roughly a 15 km error that reduces logarithmically with the minimum radial distance of the signal from the sun. Even if the beam passes through a half Earth-Sun radius there is still a few km error. All these effects introduce possible errors that make it difficult to measure a 700 metre difference in all planetary radii.




Figure 2.6. 3 Scales are grossly exaggerated for clarity. We have also assumed circular orbits here for simplicity, and ignored errors due to the centre of mass of the solar system not being at centre of sun. We have also assumed infinitesimal planet masses so we can simply ignore the effect they have on each other.

2.6.9 What about the Hulse Taylor binary pulsar, can it show this change?

The timing of this pulsar is accurate to 14 significant figures and it would initially seem that this accuracy would show up such differences. However, the semi-major orbit of this binary is $\approx 2 \times 10^9$ metres, with a decay rate of ≈ 3.5 metres per orbit, or a change $\Delta r \approx 100$ metres over 30 years due to gravitational radiated energy. If we totally ignore this change in the radius, treating it as effectively zero, the accumulated time delay is parabolic; or proportional to elapsed time squared. If we include the small effect of the change in radius,

$2m/r + 1.4m^2/r^2$ increases minutely to $2m/(r - \Delta r) + m^2/(r - \Delta r)^2$ adding two minute cubic terms, both proportional to elapsed time cubed, where the m^2/r^2 contribution is about 10^{-7} of that due to the $2m/r$ term. Even the cubic effect of a $\Delta r \approx 100$ metres change in the usual m/r term (which is currently $m/r \approx 5 \times 10^{-7}$) on the parabola over 30 years, is miniscule. The chances of measuring either the m/r , or the m^2/r^2 cubic terms are very small in the foreseeable future; let alone distinguish between them. The best chance of measuring any difference will almost certainly turn out to be gravitational wave observations.

2.6.10 Gravitational Wave observations of Black Hole mergers

Some of the mergers observed so far have been suggesting relatively larger Black Hole masses than current astrophysics theory had expected. If we look at our new metric term we

$$\text{can write } g_{00} = 1 - \frac{2m}{r} - 1.4 \frac{m^2}{r^2} = g_{00} \approx 1 - \frac{2m}{r} \left(1 + 0.7 \frac{m}{r}\right) = 1 - \frac{2m'}{r} \quad \text{where } m' \approx 1 + 0.7 \frac{m}{r}$$

For a maximum spin black hole when $r = m$ we can say the effective mass at merger is $m' \approx 1.7m$ or about 70% greater. The addition of an m^2/r^2 term in our modified metrics increases the total merging energy, and hence that in the resulting gravitational waves. Inward radial accelerations would appear to be greater also. However computer simulations with these changed metrics would be required to model all this in detail, but our rough analysis above suggests that the masses of the black holes before merging could well be less than what they have so far seemed. In other words, a pair of smaller black holes merging might create the gravitational waves current theory predicts from the mergers of two, up to maybe 70% larger black holes. Spins had also been expected to be roughly perpendicular to their orbiting plane, but their merging speeds don't tie up with this. Is it possible that this unexpected behaviour is trying to tell us something is different; something different in the metric as we get close to Black Hole Horizons?

Finally in this section, does this extra m^2/r^2 term alter what we said in Eq.(2.3. 16)? We first used $A = 2m/r$, before we introduced the self emission term $1.4m^2/r^2$ we found that the extra time polarized k_{\min} graviton density near the horizon, for all black holes is

$$A_H \cdot \frac{1 + \frac{\alpha^2}{R^2}}{g_{\theta\theta} g_{\phi\phi}} = \frac{A_H^2}{A_H^2} = 1 \quad \text{and this is still true, but now } A_H = \frac{2m}{R} + 1.4 \frac{m^2}{R^2} = 1 + \frac{\alpha'^2}{R^2}$$

Where α' is the increased spin parameter due to the extra $1.4m^2/R^2$ term and we have also reused Eq's. (2.3. 13) & (2.3. 14). Everything we did there is not affected by this extra term.

2.7 Revisiting some other aspects of the first paper

2.7.1 Infinitesimal rest masses

In section 6.2 in [7] we showed infinitesimal rest mass $N = 2$ infinite superpositions have $\langle K_{k_{\min}} \rangle^2 = 1$. From Table 4.3.1 in [7]

$N = 2$ infinitesimal rest mass spin 1 superpositions have $\langle n \rangle \approx 3.98$

$N = 2$ infinitesimal rest mass spin 2 superpositions have $\langle n \rangle \approx 3.33$

Using Eq's. (3.1.11) in [7] and Eq. (2.2. 10).

$$\begin{aligned} \langle K_{k_{\min}} \rangle^2 &= \frac{\langle n \rangle^2 s}{2} \tilde{\lambda}_c^2 k_{\min}^2 \approx \frac{15.82}{2} \tilde{\lambda}_c^2 k_{\min}^2 = 1 \text{ or } \tilde{\lambda}_c \approx 0.355 \frac{R_{OH}}{\Upsilon} \text{ for Spin 1} \\ &\approx \frac{11.09 \times 2}{2} \tilde{\lambda}_c^2 k_{\min}^2 = 1 \text{ or } \tilde{\lambda}_c \approx 0.300 \frac{R_{OH}}{\Upsilon} \text{ for Spin 2} \end{aligned}$$

Using the value for $\Upsilon \approx 0.69$ from Eq. (2.4. 13) based on WMAP data which also puts the horizon radius at $\approx 46 \times 10^9$ light years and $R_{OH} \approx 2.7 \times 10^{61}$ in Planck lengths.

Spin	$\langle n \rangle$	Compton Wavelength λ_c	Infinitesimal Rest Mass
1	3.98	$\approx 0.52 R_{OH}$	$\approx 8.8 \times 10^{-34} eV$.
2	3.33	$\approx 0.43 R_{OH}$	$\approx 1.04 \times 10^{-33} eV$.

Table 2.6. 4 Infinitesimal rest masses of $N = 2$ photons, gluons & gravitons.

These Compton wavelengths and rest masses are the present values, reducing slowly but exponentially with cosmic time T . They are based on WMAP data where $\Omega = 1$ and could be slightly greater if Ω does not need to be one as we have discussed. They also depend on the actual value of b in the exponential expansion $V = 3Exp(bt)$. These infinitesimal rest masses limit the range of virtual photons, gluons and gravitons to approximately the horizon. The graviton rest masses above are close to recent proposals explaining the accelerating expansion of the cosmos [8].

2.7.2 Redshifted zero point energy from the horizon behaves differently to local

Local zero point energies are Lorentz invariant. At high frequencies there is no shortage locally to build the high frequency components of superpositions. If a massive $N = 1$ virtual pair emerges from the vacuum its life is short and it places little demand on long range quanta. If there were no redshifted supply from the horizon there would be only a few modes of the local supply of $k_{\min} \approx 1/R_{OU}$ quanta inside the horizon. Because preons are born with zero momentum and infinite wavelength they can however absorb a different supply of redshifted $k_{\min} \approx 1/R_{OU}$ quanta from the receding horizon as we have discussed.

This k_{\min} quanta redshifted supply behaves differently to normal Lorentz invariant zero point local fields. It behaves as $K_{Qk_{\min}} = 0.055\alpha_G$ "The Quanta required @ k_{\min} Invariant" of Eq. (2.4. 12) Where $K_{Qk_{\min}} \approx 0.48 \times K_{Gk_{\min}}$ "The k_{\min} Graviton Invariant" of Eq. (2.2. 11). This redshifted supply is only available in a continuously expanding universe to preons born with zero momentum, or infinite wavelength, in the rest frame in which infinite superpositions are built.

2.7.3 Revisiting the building of infinite superpositions

In section 2 of the first paper we developed equations to determine the probability of each mode of a superposition using local zero point fields and when we found the cosmic wavelength supply inadequate we switched to a different redshifted supply for long range quanta. So how do we justify our use of the local zero point fields to determine mode probabilities and behaviours? There is simply a plentiful supply of high frequency local zero point fields. This local supply is adequate for high densities of superpositions for all modes from the Planck energy $k=1$ high energy mode cutoffs to somewhere around $k \approx 10^{-20}$ or near nuclear wavelengths. Thus, until we reach somewhere near nuclear densities, there is a sufficient supply of local high frequency zero point fields to build infinite superpositions. The coupling to local zero point fields in this high frequency region determines the primary coupling behaviour (see page 5 Part I) of all the standard model particles. There is however a gradual transition to absorbing quanta from the redshifted horizon supply as the wavelength increases. Because the redshifted supply of k_{\min} quanta behaves as the invariants $K_{Qk_{\min}}$ or $K_{Gk_{\min}}$ above and entirely differently to Lorentz invariant local zero point fields, spacetime has to warp around mass concentrations and the universe has to expand.

2.8 Gravitational Waves

Our hypothesis has been throughout, that the warping of spacetime is directly related to maintaining the maximum wavelength, or k_{\min} graviton density $\rho_{Gk_{\min}} = G_{Gk_{\min}} dk_{\min}$ invariant throughout all spacetime. Around non-rotating (spherically symmetric) mass concentrations this warping decreases inversely with radius (at least well away from black holes) but always in a spherically symmetric manner as the extra k_{\min} gravitons due to this mass are distributed in the same spherical way. Likewise we get cylindrical symmetry for rotating mass concentrations. Both these types of symmetries are the lowest action/energy stable state of the metric. Disturbances to this stable state will travel as waves at the speed of light.

2.8.1 Constant transverse areas in low energy waves

If these mass concentrations accelerate, then just like accelerating electric charges they will radiate gravitational energy in the form of real transversely polarized $m = \pm 2$, gravitons. This energy is a disturbance or oscillation in this lowest energy state k_{\min} graviton background.

Perhaps $\rho_{Gk_{\min}} = G_{Gk_{\min}} dk_{\min}$ can oscillate about its mean, but if it cannot change during these disturbances, what might be going on? Let us imagine a region of spacetime far from mass concentrations where the metric $g_{\mu\nu} = \eta_{\mu\nu}$ and using t, x, y, z coordinates let $g_{00} = -1, g_{xx} = +1, g_{yy} = +1, g_{zz} = +1$. Ignoring signs the determinant of the metric $|g_{\mu\nu}| = 1$. Let a gravitational wave pass through in the z direction with a transverse wave in the x, y plane. We know that a circular transverse ring of particles will oscillate into, and out of, ellipses perpendicular to each other, in such a manner that the enclosed area does not change, or that $g_{xx} \cdot g_{yy} = 1$ during this oscillation. Thus the measured volume of space does not change as the wave passes through and $\rho_{Gk_{\min}} = G_{Gk_{\min}} dk_{\min}$ does not change. The determinant of the metric $|g_{\mu\nu}| = 1$ also does not change. This is only approximately true as there are extra real transversely polarized $m = \pm 2, k_{\min}$ gravitons passing through due to the energy in the wave, but the error is second order unless the amplitude of the wave is quite large.

2.8.2 What might happen in high energy waves?

We can imagine the extra gravitons around a mass concentration and the background gravitons as in section 2.2 (if they are undergoing an acceleration as in binary pairs) generating real transversely polarized $m = \pm 2$, gravitons. This has some parallels to what we found in the Kerr metric, but now with real gravitons. But the intensity, or probability density, of these real gravitons will drop as the inverse radius squared, at least when far away. We can also show from Equ's. (2.1. 9) & (2.2. 5) that most of these gravitons are close to the locally measured value of the k_{\min} wavenumber. About 66% are between k_{\min} & $2k_{\min}$ and about 96% are between k_{\min} & $5k_{\min}$. Thus most of this radiated energy is near k_{\min} . Just as measured volumes around mass concentrations had to increase to accommodate extra k_{\min} gravitons, the transverse area of the wave has to increase in relation to the oscillating constant area. Ignoring signs again, if $g_{xx} \cdot g_{yy} = 1 + \varepsilon$ then $g_{00} = (1 - \varepsilon)^{-1}$ to keep the metric determinant $|g_{\mu\nu}| = 1$. The energy density in the wave increases the local measurement of k_{\min} , but $\rho_{Gk_{\min}} = G_{Gk_{\min}} dk_{\min}$ can remain invariant if it can't oscillate. Close to orbiting binary black holes or neutron stars this radiated energy intensity is huge and the changes in $g_{xx} \cdot g_{yy}$ & g_{00} become large in relation to the oscillating changes. Transverse areas and hence measured volumes change significantly. This is in complete contrast to what happens at extremely large distances, such as when we observe gravitational waves here on Earth, where the transverse areas are virtually constant during these oscillations.

2.8.3 No connection between wave frequency and radiated quanta energy

The frequency of the radiated wave is twice the orbital frequency of the binary pair source. As most of the energy in the wave is in quanta near k_{\min} there is no connection with the frequency of the radiated wave as in spin 1 photons in electromagnetism. In the recently observed gravitational waves the wave frequency was ≈ 250 cycles per second just before merger with wavelengths ≈ 1200 kilometres or approximately 10^{41} Planck lengths, whereas the wavelength of k_{\min} gravitons is $1/k_{\min} \approx R_{OU} \approx 10^{62}$ Planck lengths. The ratio between them is $\approx 10^{21}$. This ratio is inverse to the binary pair orbital frequency. It could only approach one if the orbital period is approximately twice the age of the universe.

2.9 Four Vectors and Four Volume Action Densities

2.9.1 Graviton densities represented as invariant 4 velocities

Four velocity vectors have the property that $U_0^2 - U_1^2 = 1$ is invariant under local Lorentz transformations; where U_0 is the *time component* of the four velocity, and U_1 the *spatial component*. We will, as previously, use the notation

$$U_0^2 = \gamma_M^2 \quad \text{and} \quad U_1^2 = \gamma_M^2 \beta_M^2 \quad \text{where} \quad \gamma_M^2 = \frac{1}{1 - \beta_M^2}$$

We can think of the spatial component U_1 as the four velocity $\gamma_M \beta_M$ of a free falling mass that came from rest at infinity, in the same coordinate frame as the black hole, and pointing radially inwards. We can also write

$$U_0^2 - U_1^2 = 1 \quad \text{as} \quad 1 + U_1^2 = U_0^2 \quad \text{or} \quad 1 + \gamma_M^2 \beta_M^2 = \gamma_M^2.$$

This was what we did for the Schwarzschild metric when we had temporarily multiplied both sides by γ_M^2 and normalized the background k_{\min} graviton three volume probability density to 1 with $\gamma_M^2 \beta_M^2$ the extra k_{\min} graviton density due to a central mass, and γ_M^2 the total; this equation only applies before we have expanded the volume and changed time in the new metric. Because this is a 4 vector relationship it is true in all coordinates. Multiplying both sides temporarily by γ_M^2 does not change its validity.

We can also add a term $\gamma_M^2 X^2$ to both sides to get $1 + \gamma_M^2 \beta_M^2 + \gamma_M^2 X^2 = \gamma_M^2 + \gamma_M^2 X^2$ and still maintain covariance as $(\gamma_M^2 + \gamma_M^2 X^2) - (\gamma_M^2 \beta_M^2 + \gamma_M^2 X^2) = 1$, and we can put $X^2 = \frac{\alpha^2}{r^2} \cos^2 \theta$

so that:

$$1 + \gamma_M^2 \beta_M^2 + \gamma_M^2 \frac{\alpha^2}{r^2} \cos^2 \theta = \gamma_M^2 + \gamma_M^2 \frac{\alpha^2}{r^2} \cos^2 \theta = g_{\theta\theta} \gamma_M^2.$$

We are not adding another 4 vector here; we are simply adding squared terms, which are equal on each side, so that Lorentz invariance is not affected. This is still an invariant equation in any coordinates. In the above the local metric clock rate is always $1/\gamma_M$.

The three volume probability density of circularly polarized k_{\min} gravitons due to rotation, before volume expansion and time changes in the new metric, always obeys $\gamma_M^2 X^2 = \gamma_M^2 \frac{\alpha^2}{r^2}$ and the remaining k_{\min} graviton three volume probability density is $\gamma_M^2 \beta_M^2$.

This section on invariant 4 vectors, is really just saying the same thing as our invariant 4 volume densities of k_{\min} gravitons.

2.9.2 Four volumes in changing metrics

Using our dimensionless form of the metric tensor, the nonrotating space metric determinant has magnitude $|\text{Det } g| = |g| = |g_{tt}g_{rr}g_{\theta\theta}g_{\phi\phi}| = 1$, but we want the square root of this $\sqrt{|g|} = 1$.

However in rotating space this becomes $\sqrt{|g|} = g_{\theta\theta} = 1 + \cos^2 \frac{\alpha^2}{r^2}$ which reverts to $\sqrt{|g|} = g_{\theta\theta} = 1$ when the angular momentum length parameter $\alpha = 0$. At a large radius from any mass concentration let us start with a unit four volume such that $\Delta^4 x = \Delta t \Delta x \Delta y \Delta z = 1$ when $g_{\mu\nu} = \eta_{\mu\nu}$, where for simplicity we use x, y & z for the space components. As we approach the central mass in the new metric, this four volume becomes

$$\Delta^4 x' = \sqrt{|g|} \Delta t \Delta x \Delta y \Delta z = g_{\theta\theta} = 1 + \frac{\alpha^2}{r^2} \cos^2 \theta = \Delta t' \Delta x' \Delta y' \Delta z'$$

Four volumes at a fixed point in spacetime are invariant as coordinates change, and also as the metric changes if in nonrotating space. In rotating space however it increases as $g_{\theta\theta}$.

$$\frac{\text{Curved spacetime 4 volume}}{\text{Flat spacetime 4 volume}} = \frac{\Delta^4 x'}{\Delta^4 x} = \frac{\Delta t' \Delta x' \Delta y' \Delta z'}{\Delta t \Delta x \Delta y \Delta z} = g_{\theta\theta} = 1 \text{ when angular momentum is zero.}$$

We also know that clocks change as $\Delta t' = \sqrt{g_{tt}} \Delta t = \frac{\Delta t}{\gamma_M}$ in curved spacetime so that

$$\frac{\Delta t' \Delta x' \Delta y' \Delta z'}{\Delta t \Delta x \Delta y \Delta z} = \frac{\Delta t \cdot \Delta x' \Delta y' \Delta z'}{\gamma_M \cdot \Delta t \Delta x \Delta y \Delta z} = g_{\theta\theta} = \frac{\Delta x' \Delta y' \Delta z'}{\gamma_M \cdot \Delta x \Delta y \Delta z}$$

The expanded spatial volume in the new metric $\Delta^3 x' = \Delta x' \Delta y' \Delta z' = \gamma_M g_{\theta\theta} \Delta x \Delta y \Delta z = \gamma_M g_{\theta\theta} \Delta^3 x$.

Spatial volume in any metric expands as $\frac{V'}{V} = \frac{\Delta^3 x'}{\Delta^3 x} = \frac{d^3 x'}{d^3 x} = \gamma_M g_{\theta\theta} = (1 + \frac{\alpha^2}{r^2} \cos^2 \theta) \gamma_M$

Where as above we have defined $\gamma_M = g_{tt}^{-1/2}$ as the local metric clock rate.

The 4 volume density invariance of k_{\min} graviton still applies, as the extra circularly polarized gravitons due to angular momentum, occupy this expanded volume.

3 Some Loose Ends

3.1.1 Preferred Frames

It might seem that we have been arguing for a preferred frame. But there is really no difference in what we are proposing compared to current physics. In commoving frames the cosmic microwave background is isotropic. At peculiar velocity β_p it is no longer isotropic, and the average background temperature increases by γ_p , exactly the same increase as k_{\min} to $k'_{\min} = \gamma_p k_{\min}$, and that is if we could measure it, which is most unlikely. We have frequently talked in this paper about local observers measuring k_{\min} , but only as a thought experiment, and the average (over all directions) background temperature can be used to measure either γ_{peculiar} or γ_{metric} . There are no other changes in physics in this commoving frame; it is exactly as Einstein originally postulated, an important experimentally verified feature of General Relativity [33]. However it does make everything we did here much simpler if we work in commoving coordinates. All the mass moving at peculiar velocities in random directions does not affect the average universe density of either k_{\min} gravitons or the k_{\min} action density that they require. We calculated the average density of k_{\min} action from the horizon in these commoving coordinates. But if we think in terms of spherically symmetric four volume k_{\min} action density invariance, then whether we are in a non commoving frame, or in a non-flat metric, it makes no difference; and is why we can use 4 vector notation for the extra k_{\min} gravitons around mass concentrations.

3.1.2 Solar System Constraints and do our proposed changes fit?

See “The Confrontation between General Relativity and Experiment” Clifford M. Will. [33] Probably the most important constraint mentioned in this review is the Cassini Time Delay data that gives a fit with GR of $\approx 10^{-5}$ for signals passing close to the solar horizon, where our extra $1.4m^2/r^2$ term is $\approx 3 \times 10^{-6}$. So it should be within the Cassini Constraint and also within the light deflection constraint. The remaining changes are discussed in section 2.6.7.

3.1.3 Action Principles and the Einstein Field Equations

The field equations of GR can be derived from an invariant action principle $\delta I = 0$ where $I = \frac{1}{16\pi G} \int R \sqrt{-g} \cdot d^4x + I_m(\psi_m, g_{\mu\nu})$ and R is the Ricci scalar, with I_m the matter action which depends on matter fields ψ_m coupled to the metric g . Varying the action with respect to $g_{\mu\nu}$ we obtain Einstein’s field equations $G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi G T_{\mu\nu}$. This paper suggests however, that an “Invariant spherically symmetric 4 volume cosmic wavelength graviton action density” applies to the solutions of infinitesimally modified stress tensor.

At any fixed cosmic time T in comoving coordinates, a homogenous flat universe with no mass concentrations has both 3 volume & 4 volume spherically symmetric action density invariance for cosmic wavelength gravitons, but only 4 volume invariance with local mass concentrations. We argue the changing metric due to this relates with gravity. This 4 volume invariance is true in all frames. However an infinitesimal change to Einstein's Energy Momentum tensor is required with almost zero local effects, but significant implications at cosmic scale. The extra $\approx 1.4m^2 / r^2$ term we had to include in the metric (mainly significant around Black Holes) is irrelevant in these invariant properties, but must have some effect on Black Hole mergers, with possibly faster supermassive Black Hole formation than had been expected.

4 Conclusions

If the fundamental particles can be formed from infinite superpositions as outlined in [7], our hypothesis is that the warping of spacetime is consistent with a maximum wavelength, or k_{\min} graviton probability density $\rho_{Gk_{\min}} = K_{Gk_{\min}} dk_{\min}$, or equivalently, an invariant "Four Volume Spherically Symmetric Action Density". But the value of k_{\min} decreases with cosmic time and is roughly inverse to the horizon radius, it also depends on the local clockrate $\sqrt{g_{00}}$

Thinking in a simple way and using the proportionality symbol \propto as follows:

In a universe with no mass concentrations $\rho_{Gk_{\min}} \propto (\Psi_{\text{Universe}} * \Psi_{\text{Universe}})$. With a concentration of mass m , $\rho'_{Gk_{\min}} \propto (\Psi_{\text{Universe}} * \Psi_{\text{Universe}}) + (\Psi_{\text{Universe}} * \Psi_m + \Psi_m * \Psi_{\text{Universe}}) + (\Psi_m * \Psi_m)$ but space expands locally to restore $\rho'_{Gk_{\min}}$ back to $\rho_{Gk_{\min}} = G_{Gk_{\min}} dk_{\min}$. The green term $(\Psi_{\text{Universe}} * \Psi_m + \Psi_m * \Psi_{\text{Universe}})$ requires $2m/r$ in the metric, and meshes well with an infinitesimally modified General Relativity. This modification changes the T_{00} component from $T_{00} = \rho$, where ρ is the local mass density, to $T_{00} = \rho - \rho_U$, where ρ_U is the average density of the Universe (only a few hydrogen atoms per cubic metre). It matches the Schwarzschild metric, and fits the Kerr metric. In the earlier paper we focused only on this term to illustrate a possible connection with quantum mechanics, provided the fundamental particles can be made from infinite superpositions borrowing action/energy from zero point fields. This paper messes up that nice connection by introducing the troublesome blue term $(\Psi_m * \Psi_m)$ with its associated m^2 / r^2 in the metric. This requires a (possibly problematic?) negative energy massless particle to be added to Einstein's stress tensor, similar to the positive energy massless particles in the electromagnetic field. This m^2 / r^2 term is of opposite sign to the r_Q^2 / r^2 , or equivalently dimensionless Q^2 / r^2 term, of both the Reissner-Nordstrom and Kerr-Newman metrics. The effect on the Riemannian curvature tensor is of identical form, but opposite sign to these metrics. It does not, however, alter the event horizon radius of a maximum spin black hole, but allows about 55% more spin.

These values as with other findings are only approximate however. A more rigorous analysis of everything we have done will almost certainly change these findings, and the coefficient of m^2 / r^2 in the metric due to $(\psi_m * \psi_m)$, and the extra maximum spin. But if our conjecture is true, this m^2 / r^2 term will not go away. The extra radial acceleration that it introduces could alter the merging rate of black holes. The third merger observed [26] suggests that General Relativity holds to the horizon, but the spins appear not to align with their mutual orbit, possibly challenging conventional astrophysics. If the spins are aligned, as had been thought more probable, the merging rate is slightly too fast. Could we turn this around and say:

“If the spins are in fact aligned, does General Relativity need modifying near the horizon?”

This paper requires two such changes, one that is significant, mainly near black holes, and the other only significant at cosmic scale. Testing these two changes has to await future accuracy improvements in gravitational wave detectors and further refinements in our observations of the accelerating expansion of space.

Finally, supermassive black holes are appearing much earlier in cosmic time than expected. Could this extra m^2 / r^2 term, alter the inflow rate of the surrounding swirling matter; above that of current models? Also the Hubble parameter, predicted by Λ -CDM models based on Cosmic Microwave data, is $\approx 9\%$ less than the recent, or more current, Hubble parameter measurements [32]. The expansion velocity in our exponential expansion model predicts expansion velocities $\approx 11\%$ greater than Λ -CDM models. Is there a connection here? The future refinements expected in these measurements over the coming years, mentioned above, will no doubt clarify this.

5 References

- [1] A. Einstein, “*The Foundation of the General theory of Relativity*” (1916) *Annalen Phys.*, 49:769-822
- [2] D. Germanis, “*General Relativity and Modified Gravity*,” September 2011 and references therein. <https://workspace.imperial.ac.uk>
- [3] R. Jackiw, S.Y.Pi, “*Chern-Simons Modification of General Relativity*” (2003) [arXiv:gr-qc/0308071v2](https://arxiv.org/abs/gr-qc/0308071v2)
- [4] U.Ertem, “*Generalized Chern-Symons Modified Gravity in First-Order Formalism*” (2009), [arXiv:0912.1433v1](https://arxiv.org/abs/0912.1433v1)
- [5] C. Brans and R.H. Dicke, “*Mach’s Principle and a Relativistic Theory of Gravitation*”, *Phys. Rev.* 124, 925 (1961).
- [6] Carl Brans, “*Invariant Approach to the Geometry of Spaces in General Relativity*,” *Jour. Math. Phys.*, 6 94 (1965)
- [7] J.Williams, “*Exploring the Possibility of New Physics Part I: Quantum Mechanics and the Warping of Spacetime*”, [viXra:1303.0029v6](https://arxiv.org/abs/1303.0029v6)
- [8] De Rham, C., Gabadadze, G. & Tolley, A.J. *Phys. Rev. Lett.* 106, 231101 (2011)
- [9] De Rham, C., Gabadadze, G., Heisenberg, L. & Pirtskhalava, D. *Phys. Rev.D*83, 103516 (2011)
- [10] J.C. Pati, A. Salam, *Phys. Rev.* D10, 275 (1974).
- [11] See for example H. Terazawa, K. Akama, *Phys. Lett.* B96, 276 (1980) and references therein.
- [12] H. Harari, *Phys. Lett.* B86, 83 (1979).
- [13] M. Shupe, *Phys. Lett.* B86, 87 (1979).
- [14] A. Rae, “*Quantum Mechanics 3rd edition*”, (Institute of Physics Publishing, Bristol, 1992).
- [15] R. Feynman, “*The Feynman Lectures on Physics.*” Volume III. (Addison Wesley, Reading, Massachusetts, 1964).
- [16] I. Aitchison, A. Hey, “*Gauge Theories in Particle Physics, 2nd edition*”, (Institute of Physics Publishing, Bristol, 1989).
- [17] Cosmological Constraints from Measurements of Type 1a Supernovae discovered during the first 1.5 years of the Pan-STARRS 1 Survey. [arXiv:1310.3828](https://arxiv.org/abs/1310.3828)
- [18] W.G. Unruh, “*Notes on Black Hole Evaporation*,” *Phys. Rev.D*14 (4):870892 (1976)
- [19] D. Marolf and J. Polchinski, *Phys. Rev. Lett.* 111, 171301 (2013).

- [20] A. Almheiri, D. Marolf, J. Polchinski, and J. Sully, *Journal of High Energy Physics*. 2013, No. 2, 063 (2013).
- [21] K. Papadodimas and S. Raju, “An infalling Observer in AdS/CFT” [arXiv:1211.6767](https://arxiv.org/abs/1211.6767) (2012)
- [22] R. Bousso, “Frozen Vacuum” [arXiv:1308.3697](https://arxiv.org/abs/1308.3697) (2013)
- [23] J. Maldacena and L. Susskind, “Cool Horizons for Entangled Black Holes” [arXiv:1306.0533](https://arxiv.org/abs/1306.0533) (2013)
- [24] Maldacena, Juan (1998). "The Large N limit of superconformal field theories and supergravity". *Advances in Theoretical and Mathematical Physics*. 2: 231–252. [arXiv:hep-th/9711200](https://arxiv.org/abs/hep-th/9711200).
- [25] Lense, J.; Thirring, H. (1918). “On the Influence of the Proper Rotation of Central Bodies on the Motions of Planets and Moons According to Einstein's Theory of Gravitation” *Physikalische Zeitschrift*. 19: 156–163. [Bibcode:1918PhyZ...19..156L](https://ui.adsabs.harvard.edu/abs/1918PhyZ...19..156L).
- [26] B.P.Abbott *et al.* (Ligo Scientific and Virgo Collaboration) GW170104: “Observation of a 50-Solar- Mass Binary Black Hole Coalescence of Redshift 0.2” *Phys. Rev. Lett.* **118**, 221101 <https://doi.org/10.1103/PhysRevLett.118.221101>
- [27] Reissner, H. (1916). “Über die Eigengravitation des elektrischen Feldes nach der Einsteinschen Theorie”. *Annalen der Physik (in German)*. 50: 106–120. [Bibcode:1916AnP...355..106R](https://ui.adsabs.harvard.edu/abs/1916AnP...355..106R). [doi:10.1002/andp.19163550905](https://doi.org/10.1002/andp.19163550905).
- [28] Nordström, G. (1918). “On the Energy of the Gravitational Field in Einstein's Theory”. *Verhandl. Koninkl. Ned. Akad. Wetenschap., Afdel. Natuurk., Amsterdam*. 26: 1201–1208.
- [29] Newman, Ezra; Janis, Allen (1965). “Note on the Kerr Spinning-Particle Metric”. *Journal of Mathematical Physics*. 6 (6): 915–917. [Bibcode:1965JMP....6..915N](https://ui.adsabs.harvard.edu/abs/1965JMP....6..915N). [doi:10.1063/1.1704350](https://doi.org/10.1063/1.1704350).
- [30] Newman, Ezra; Couch, E.; Chinnapared, K.; Exton, A.; Prakash, A.; Torrence, R. (1965). “Metric of a Rotating, Charged Mass”. *Journal of Mathematical Physics*. 6 (6): 918–919. [Bibcode:1965JMP....6..918N](https://ui.adsabs.harvard.edu/abs/1965JMP....6..918N). [doi:10.1063/1.1704351](https://doi.org/10.1063/1.1704351).
- [31] Kerr, RP (1963). “Gravitational field of a spinning mass as an example of algebraically special metrics”. *Physical Review Letters*. 11: 237–238. [Bibcode:1963PhRvL..11..237K](https://ui.adsabs.harvard.edu/abs/1963PhRvL..11..237K). [doi:10.1103/PhysRevLett.11.237](https://doi.org/10.1103/PhysRevLett.11.237).
- [32] AG Reiss (2018). “New Parallaxes of Galactic Cepheids from Spatially Scanning the Hubble Space Telescope” arxiv.org/abs/1801.01120
- [33] Clifford M. Will. (2006) “The Confrontation between General Relativity and Experiment” [arxiv:gr-qc/0519972v2](https://arxiv.org/abs/gr-qc/0519972v2)