

# Boundary Matrices and the Marcus-de Oliveira Determinantal Conjecture

Ameet Sharma

## ARTICLE HISTORY

Compiled April 22, 2019

## ABSTRACT

We present notes on the Marcus-de Oliveira conjecture. The conjecture concerns the region in the complex plane covered by the determinants of the sums of two normal matrices with prescribed eigenvalues. Call this region  $\Delta$ . This paper focuses on boundary matrices of  $\Delta$ . We prove 2 theorems regarding these boundary matrices. This paper uses ideas from [1].

## KEYWORDS

determinantal conjecture, Marcus-de Oliveira, determinants, normal matrices, convex-hull

## 1. Introduction

Marcus [2] and de Oliveira [3] made the following conjecture. Given two normal matrices  $A$  and  $B$  with prescribed eigenvalues  $a_1, a_2 \dots a_n$  and  $b_1, b_2 \dots b_n$  respectively,  $\det(A + B)$  lies within the region:

$$co\left\{ \prod (a_i + b_{\sigma(i)}) \right\}$$

where  $\sigma \in S_n$ .  $co$  denotes the convex hull of the  $n!$  points in the complex plane. As described in [1], the problem can be restated as follows. Given two diagonal matrices,  $A_0 = \text{diag}(a_1, a_2 \dots a_n)$  and  $B_0 = \text{diag}(b_1, b_2 \dots b_n)$ , let:

$$\Delta = \{ \det(A_0 + UB_0U^*) : U \in U(n) \} \quad (1)$$

where  $U(n)$  is the set of  $n \times n$  unitary matrices. Then we can write the conjecture as:

**Conjecture 1.1** (Marcus-de Oliveira Conjecture).

$$\Delta \subseteq \text{co}\left\{\prod (a_i + b_{\sigma(i)})\right\} \quad (2)$$

Let

$$M(U) = \det(A_0 + UB_0U^*). \quad (3)$$

The paper is organized as follows. In section 2 we define terms and functions that will be used in the rest of the paper. These definitions are necessary to state our results. In section 3, we state 3 lemmas and 2 theorems that form the bulk of the paper. We state them in the order they are proved.

## 2. Preparatory definitions

### 2.1. Terms

For the purposes of this paper we define an **ordinary point** of a curve as a point on the curve that is not any kind of singular point. ie, an ordinary point has the following properties:

- It is not a cusp.
- It is not a point of self-intersection.
- It is not an isolated point.
- The curve has a unique tangent at that point.

Given a unitary matrix  $U$  and square, diagonal matrices  $A_0$  and  $B_0$  all of dimension  $n \times n$ ,

- If  $M(U)$  is a point on  $\partial\Delta$  (the boundary of  $\Delta$ ), we call  $M(U)$  a boundary point of  $\Delta$  and we call  $U$  a **boundary matrix** of  $\Delta$ . See eq. (1) and eq. (3).
- We define the **B-matrix** of  $U$  as  $UB_0U^*$ .
- We define the **C-matrix** of  $U$  as  $A_0 + UB_0U^*$ .
- We define the **F-matrix** of  $U$  as  $C^{-1}A_0 - A_0C^{-1}$  where  $C$  is the C-matrix of  $U$ . Note that the F-matrix is only defined when  $C$  is invertible, or equivalently when  $\det(C) = M(U) \neq 0$ . See eq. (3). Also note that since  $A_0$  is diagonal, the F-matrix is a zero-diagonal matrix. The idea for using the F-matrix comes from [1], Theorem 4, p.27.

Throughout the rest of the paper, we'll assume  $A_0$  and  $B_0$  are defined, even if we don't explicitly mention them.

## 2.2. Functions given a unitary matrix $U$

Given a unitary matrix  $U$  with B-matrix  $B$ , C-matrix  $C$  and F-matrix  $F$ . For every skew-hermitian matrix  $Z$ , we define the following functions

let

$$U_Z(t) = (e^{Zt})U \quad (4)$$

where  $t$  is any real number.

Since the exponential of a skew-hermitian matrix is unitary,  $U_Z(t)$  is a function of unitary matrices.

let

$$B_Z(t) = U_Z(t)B_0U_Z^*(t) \quad (5)$$

let  $C_Z(t) = A_0 + B_Z(t)$

We note that  $B_Z(0) = B$  and  $C_Z(0) = C$ .

let

$$R_Z(t) = \det(C_Z(t)) \quad (6)$$

We can see by eq. (1) that  $R_Z(t) \subseteq \Delta$ .

$$R_Z(0) = A_0 + UB_0U^*$$

So by eq. (3) we see that  $R_Z(0) = M(U)$ .

So all the  $R_Z(t)$  functions go through  $M(U)$  at  $t = 0$ .

We shall refer to these functions in the rest of the paper with the same notation (for example  $R_Z(t)$  for a skew-hermitian matrix  $Z$ .  $R_{Z_1}(t)$  for a skew-hermitian matrix  $Z_1$ ). Note that  $R_Z(t)$  requires  $A_0, B_0, U$  and  $Z$  in order to be defined. But we won't explicitly mention  $A_0$  and  $B_0$ . All the results in this paper assume there are two diagonal matrices  $A_0$  and  $B_0$  defined in the background.

## 2.3. Skew-Hermitian matrices $Z^{ab}$ and $Z^{ab,i}$

Given two integers  $a, b$  where  $1 \leq a, b \leq n$  and  $a \neq b$ .

We define the  $n \times n$  skew-hermitian matrix  $Z^{ab}$  as follows.  $Z_{ab}^{ab} = -1$  (the element at the  $a$ th row and  $b$ th column is -1.)  $Z_{ba}^{ab} = 1$  (the element at the  $b$ th row and  $a$ th column is 1.) And all other elements are 0. Note that  $Z^{ab} = -Z^{ba}$ .

We define the  $n \times n$  skew-hermitian matrix  $Z^{ab,i}$  as follows.  $Z_{ab}^{ab,i} = i$  and  $Z_{ba}^{ab,i} = i$ .

All other elements are zero. Note that  $Z^{ab,i} = Z^{ba,i}$ .

It is straightforward to verify that  $Z^{ab}$  and  $Z^{ab,i}$  are skew-hermitian.

### 3. Main Results

**Lemma 3.1.** *Given a unitary matrix  $U$  with  $M(U) \neq 0$ . Let  $F$  be its  $F$ -matrix. Then  $R'_Z(0) = M(U)tr(ZF)$  for any skew-hermitian matrix  $Z$ .*

**Lemma 3.2.** *Given an  $n \times n$  zero-diagonal matrix  $W$ . Given  $tr(Z^{ab}W) = 0$  and  $tr(Z^{ab,i}W) = 0$  for all pairs  $(a,b)$  where  $1 \leq a,b \leq n$  and  $a \neq b$ . Then  $W$  is the zero-matrix.*

**Lemma 3.3.** *Given a boundary matrix  $U$  with  $M(U) \neq 0$  and with  $F$ -matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ . Let  $L$  be the tangent line to  $\partial\Delta$  at  $M(U)$  with direction vector  $v$ . Then for every skew-hermitian matrix  $Z$ ,  $tr(ZF) = cv$  where  $c$  is some real number.*

**Theorem 3.4.** *Given a boundary matrix  $U$  with  $M(U) \neq 0$  and with  $F$ -matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ . Then  $F$  can be written uniquely in the form  $F = e^{i\theta}H$  where  $H$  is a zero-diagonal hermitian matrix and  $0 \leq \theta < \pi$ .*

**Theorem 3.5.** *Given a boundary matrix  $U$  with  $M(U) \neq 0$  and with  $F$ -matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ . Let  $L$  be the tangent line to  $\Delta$  at  $M(U)$ . By the previous theorem we know that  $F = e^{i\theta}H$  for some real  $0 \leq \theta < \pi$ . Then  $L$  makes an angle  $\arg(M(U)) + \theta + \pi/2$  with the positive real axis.*

### 4. Proof of lemma 3.1

The proof given here uses ideas from [1], Theorem 4, p.26-27. But the proof given here is complete on its own.

**Proof.** We're given a unitary matrix  $U$  where  $M(U) \neq 0$ . So its  $F$ -matrix is well-defined and we call it  $F$ . Let  $B$  be its  $B$ -matrix, and  $C$  be its  $C$ -matrix. Given an arbitrary skew-hermitian matrix  $Z$ .

We can use Jacobi's formula [4] on eq. (6) to find  $R'_Z(t)$

$$R'_Z(t) = tr(\det(C_Z(t))C_Z^{-1}(t)C'_Z(t)) \quad (7)$$

$$R'_Z(0) = tr(\det(C_Z(0))C_Z^{-1}(0)C'_Z(0))$$

We can substitute  $C$  for  $C_Z(0)$ .

$$R'_Z(0) = tr(\det(C)C^{-1}C'_Z(0))$$

$$R'_Z(0) = \det(C) \text{tr}(C^{-1}C'_Z(0))$$

We know that  $C'_Z(t) = B'_Z(t)$  so

$$R'_Z(0) = \det(C) \text{tr}(C^{-1}B'_Z(0))$$

By section 2.1 and eq. (3) we know that  $\det(C) = M(U)$

$$R'_Z(0) = M(U) \text{tr}(C^{-1}B'_Z(0)) \quad (8)$$

Using eq. (5),

$$B'_Z(t) = \frac{dU_Z(t)}{dt} B_0 U_Z^*(t) + U_Z(t) B_0 \frac{dU_Z^*(t)}{dt} \quad (9)$$

Using eq. (4),

$$\frac{dU_Z(t)}{dt} = Z e^{Zt} U$$

$$U_Z^*(t) = (U^*) e^{-Zt}$$

$$\frac{dU_Z^*(t)}{dt} = -(U^*) Z e^{-Zt}$$

Substitute these and eq. (4) into eq. (9)

$$B'_Z(t) = Z e^{Zt} U B_0 (U^*) e^{-Zt} - (e^{Zt} U B_0 (U^*) Z e^{-Zt}$$

$$B'_Z(0) = Z U B_0 U^* - U B_0 (U^*) Z$$

Using the definition of the C-matrix in section 2.1

$$B'_Z(0) = Z(C - A_0) - (C - A_0)Z$$

$$C^{-1}B'_Z(0) = C^{-1}ZC - C^{-1}ZA_0 - Z + C^{-1}A_0Z$$

$$\text{tr}(C^{-1}B'_Z(0)) = \text{tr}(C^{-1}ZC) - \text{tr}(C^{-1}ZA_0) - \text{tr}(Z) + \text{tr}(C^{-1}A_0Z)$$

The first and third terms cancel since similar matrices have the same trace.

$$\text{tr}(C^{-1}B'_Z(0)) = -\text{tr}(C^{-1}ZA_0) + \text{tr}(C^{-1}A_0Z).$$

Using the idea that  $\text{tr}(XY) = \text{tr}(YX)$

$$\text{tr}(C^{-1}B'_Z(0)) = -\text{tr}(ZA_0C^{-1}) + \text{tr}(ZC^{-1}A_0)$$

$$\text{tr}(C^{-1}B'_Z(0)) = \text{tr}(Z(C^{-1}A_0 - A_0C^{-1}))$$

$$\text{tr}(C^{-1}B'_Z(0)) = \text{tr}(ZF)$$

Substitute this into eq. (8) to get

$$R'_Z(0) = M(U) \text{tr}(ZF) \quad (10)$$

This proves lemma 3.1. □

### 5. Proof of lemma 3.2

**Proof.** Given an  $n \times n$  zero-diagonal matrix  $W$ . Given that for every pair  $(a,b)$  where  $1 \leq a, b \leq n$  and  $a \neq b$ ,

$$\text{tr}(Z^{ab}W) = 0.$$

$$\text{tr}(Z^{ab,i}W) = 0$$

(See section 2.3 for definitions of  $Z^{ab}$  and  $Z^{ab,i}$ ).

by direct computation we see that

$$\text{tr}(Z^{ab}W) = W_{ab} - W_{ba} = 0$$

$$\text{tr}(Z^{ab,i}W) = (W_{ab} + W_{ba})i = 0$$

Solving these, we get that  $W_{ab} = 0$  and  $W_{ba} = 0$ . So all the off-diagonal elements of  $W$  are zero. Hence  $W$  is the zero-matrix. □

### 6. Proof of lemma 3.3

**Proof.** Given a boundary matrix  $U$  with  $M(U) \neq 0$  and with F-matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ . Let  $L$  be the tangent line to  $\partial\Delta$  at  $M(U)$ . Let  $v$  be the direction vector of the line  $L$ . Note that  $v$  is just a non-zero complex number.

Let  $Z$  be a skew-hermitian matrix. By lemma 3.1 we know that  $R'_Z(0) = M(U)\text{tr}(ZF)$ .

Since  $R_Z(t) \subseteq \Delta$  and  $R_Z(0) = M(U)$ , we know that  $R'_Z(0) = kv$  for some real number  $k$ . (since  $L$  is the unique tangent to  $\partial\Delta$  at  $M(U)$ , then it must be the tangent to every curve that lies in  $\Delta$ , goes through  $M(U)$  and has a well-defined derivative at  $M(U)$ ).

$$\text{So, } M(U)\text{tr}(ZF) = kv$$

$$\text{tr}(ZF) = \left(\frac{k}{M(U)}\right)v \quad \square$$

### 7. Proof of theorem 3.4

**Proof.** Given a boundary matrix  $U$  with  $M(U) \neq 0$  and with F-matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ .

We pick an arbitrary pair  $(a,b)$  such that  $1 \leq a, b \leq n$  and  $a \neq b$

We have two skew-hermitian matrices  $Z^{ab}$  and  $Z^{ab,i}$  defined as per section 2.3.

By direct computation we see that

$$\text{tr}(Z^{ab}F) = F_{ab} - F_{ba}$$

$$\text{tr}(Z^{ab,i}F) = (F_{ab} + F_{ba})i$$

Suppose  $F_{ab} = F_{ab,r} + iF_{ab,i}$ . (note that these are not tensors.  $F_{ab,r}$  is just the real component of  $F_{ab}$  and  $F_{ab,i}$  is just the imaginary component.) We can substitute this in to get

$$\text{tr}(Z^{ab}F) = (F_{ab,r} - F_{ba,r}) + i(F_{ab,i} - F_{ba,i}) \quad (11)$$

$$\text{tr}(Z^{ab,i}F) = (-F_{ab,i} - F_{ba,i}) + i(F_{ab,r} + F_{ba,r}) \quad (12)$$

We know by lemma 3.3 that these are collinear vectors in the complex plane.

So we know that

$$(F_{ab,i} - F_{ba,i})(-F_{ab,i} - F_{ba,i}) = (F_{ab,r} + F_{ba,r})(F_{ab,r} - F_{ba,r})$$

We can simplify this to get:

$$F_{ab,r}^2 + F_{ab,i}^2 = F_{ba,r}^2 + F_{ba,i}^2$$

$$|F_{ab}| = |F_{ba}|$$

We can write:

$$F_{ab} = |F_{ab}|\angle\theta_{ab}$$

$$F_{ba} = |F_{ab}|\angle\theta_{ba}$$

Note that we are given that  $F \neq 0$ . Note that we already know by section 2.1 that  $F$  is zero-diagonal.

We will divide the possible values of  $F$  into multiple cases. First we split all cases into two. The first is when only one pair of elements of the  $F$ -matrix,  $F_{ab}$  and  $F_{ba}$  is nonzero. The second case is when multiple pairs of elements of the  $F$ -matrix are nonzero. We shall further subdivide the second case using the fact that all  $\text{tr}(ZF)$  values are collinear. We can divide these cases into 3 possibilities: 1. All nonzero  $\text{tr}(ZF)$  values are imaginary. 2. All nonzero  $\text{tr}(ZF)$  values are real. 3. All nonzero  $\text{tr}(ZF)$  values are not real or imaginary. (note that since  $F$  is nonzero, we don't have to deal with the possibility that  $\text{tr}(ZF)$  is 0 for all skew-hermitian matrices  $Z$ . see lemma 3.2).

So we have 4 cases to deal with.

**Case 1:  $|F_{ab}|$  is non-zero for only one pair  $\{a, b\}$  where  $a \neq b$**

In this case,

$$H = e^{-(\theta_{ab} + \theta_{ba})/2}F \text{ is a hermitian matrix, and we're finished.}$$

**Case 2:**  $|F_{ab}|$  is non-zero for multiple pairs  $\{a, b\}$  where  $a \neq b$ . For any skew-hermitian  $\mathbf{Z}$ , when  $\text{tr}(\mathbf{ZF})$  is non-zero, it is imaginary.

If  $|F_{ab}| \neq 0$ , then by eq. (11) and eq. (12),  $\theta_{ab} = -\theta_{ba}$ . This holds for all distinct pairs  $\{a, b\}$ , so our  $F$ -matrix is already hermitian, and we're done.

**Case 3:**  $|F_{ab}|$  is non-zero for multiple pairs  $\{a, b\}$  where  $a \neq b$ . For any skew-hermitian  $\mathbf{Z}$ , when  $\text{tr}(\mathbf{ZF})$  is non-zero, it is real.

If  $|F_{ab}| \neq 0$ , then by eq. (11) and eq. (12),  $\theta_{ab} = \pi - \theta_{ba}$ . This holds for all distinct pairs  $\{a, b\}$

$H = e^{-i(\frac{\pi}{2})}F$  is hermitian and we're done.

**Case 4:**  $|F_{ab}|$  is non-zero for multiple pairs  $\{a, b\}$  where  $a \neq b$ . For any skew-hermitian matrix  $\mathbf{Z}$ , when  $\text{tr}(\mathbf{ZF})$  is non-zero, it isn't real or imaginary.

Suppose  $|F_{ab}| \neq 0$  and  $|F_{cd}| \neq 0$

if  $\text{tr}(Z^{ab}F) \neq 0$ , then

$$\text{slope of } \text{tr}(Z^{ab}F) = \frac{\sin(\theta_{ab}) - \sin(\theta_{ba})}{\cos(\theta_{ab}) - \cos(\theta_{ba})} = -\cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

if  $\text{tr}(Z^{ab,i}F) \neq 0$ :

$$\text{slope of } \text{tr}(Z^{ab,i}F) = \frac{\cos(\theta_{ab}) + \cos(\theta_{ba})}{-\sin(\theta_{ab}) - \sin(\theta_{ba})} = -\cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

We know that since  $|F_{ab}| \neq 0$ , at least one of  $\text{tr}(Z^{ab}F)$  or  $\text{tr}(Z^{ab,i}F)$  is non-zero.

similarly,

if  $\text{tr}(Z^{cd}F) \neq 0$ , then

$$\text{slope of } \text{tr}(Z^{cd}F) = -\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right)$$

if  $\text{tr}(Z^{cd,i}F) \neq 0$ :

$$\text{slope of } \text{tr}(Z^{cd,i}F) = -\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right)$$

We know that since  $|F_{cd}| \neq 0$ , at least one of  $\text{tr}(Z^{cd}F)$  or  $\text{tr}(Z^{cd,i}F)$  is non-zero.

So we have:

$$\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right) = \cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right) \text{ (lemma 3.3)}$$

therefore:

$$\frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2} + n\pi \text{ for some integer } n.$$

We can freely adjust  $\theta_{cd}$  by  $-2n\pi$ . It makes no difference since  $|F_{cd}| \angle \theta_{cd} = |F_{cd}| \angle (\theta_{cd} - 2n\pi)$

So after the adjustment we have:

$$\frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2}.$$

We make the same adjustment for any pair  $\{c, d\} \neq \{a, b\}$  where  $|F_{cd}| \neq 0$

We set  $\beta = \frac{\theta_{ab} + \theta_{ba}}{2}$

let  $H = e^{-i\beta} F$

For some pair (x,y) where  $x \neq y$  and  $|H_{xy}| \neq 0$ ,

$$H_{xy} = |H_{xy}| \angle \alpha_{xy}$$

$$\alpha_{xy} = -\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right) + \theta_{xy}$$

$$\alpha_{yx} = -\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right) + \theta_{yx}$$

But because of our adjustments,

$$\frac{\theta_{ab} + \theta_{ba}}{2} = \frac{\theta_{xy} + \theta_{yx}}{2}$$

Plugging this into the above two formulas we have

$$\alpha_{xy} = \frac{\theta_{xy} - \theta_{yx}}{2}$$

$$\alpha_{yx} = -\left(\frac{\theta_{xy} - \theta_{yx}}{2}\right)$$

Therefore H is zero-diagonal, with transpositional elements of equal magnitude and opposite arguments. Therefore H is hermitian.

So in all 4 cases we can write  $F = e^{i\beta} H$  for some hermitian matrix H and some real  $\beta$ . But we've not arrived at a unique representation for F yet.

Suppose

$$F = e^{i\beta_1} H_1 = e^{i\beta_2} H_2$$

$$e^{i(\beta_1 - \beta_2)} H_1 = H_2$$

$$e^{i(\beta_1 - \beta_2)} H_1 = H_2 = H_2^* = e^{i(\beta_2 - \beta_1)} H_1^* = e^{i(\beta_2 - \beta_1)} H_1$$

So

$$(e^{i(\beta_1 - \beta_2)} - e^{i(\beta_2 - \beta_1)}) H_1 = 0$$

Since  $F \neq 0$ , we know  $H_1 \neq 0$  so

$$e^{i(\beta_1 - \beta_2)} - e^{i(\beta_2 - \beta_1)} = 0$$

$$e^{i(\beta_1 - \beta_2)} = e^{i(\beta_2 - \beta_1)}$$

Then

$$\beta_1 - \beta_2 = \beta_2 - \beta_1 + 2k\pi, \text{ for any integer } k$$

$$\beta_1 = \beta_2 + k\pi$$

So if we restrict all  $\beta$  to  $0 \leq \beta < \pi$ , we have a unique representation since k is forced to 0.

This completes our proof of theorem 3.4. □

## 8. Proof of theorem 3.5

Given a boundary matrix  $U$  with  $M(U) \neq 0$  and F-matrix  $F \neq 0$ . Given  $M(U)$  is an ordinary point of  $\partial\Delta$ . Let  $L$  be the tangent line to  $\partial\Delta$  at  $M(U)$ .

**Proof.** By theorem 3.4 we know that

$$F = e^{i\theta} H \quad (13)$$

for some real  $0 \leq \theta < \pi$  and some zero-diagonal hermitian matrix  $H$ .

We can substitute eq. (13) into eq. (11) and eq. (12) and simplify to get:

$$\text{tr}(Z^{ab} F) = 2H_{ab,i} e^{i(\theta+\pi/2)} \quad (14)$$

$$\text{tr}(Z^{ab,i} F) = 2H_{ab,r} e^{i(\theta+\pi/2)} \quad (15)$$

By lemma 3.2 we know that at least one of the above equations is nonzero for some pair  $(a,b)$ . So then using lemma 3.1 we know that  $R'_Z(0) = M(U)\text{tr}(ZF) \neq 0$  for some skew-hermitian matrix  $Z$ .

So by eq. (14) and eq. (15) we see that for some skew-hermitian matrix  $Z$ ,  $\text{tr}(ZF)$  forms an angle of  $(\theta + \pi/2)$  or  $(\theta + 3\pi/2)$  with the positive real axis (depending on whether the coefficient is negative or not). Therefore  $R'_Z(0)$  forms an angle  $\text{arg}(M(U)) + \theta + \pi/2$  or  $\text{arg}(M(U)) + \theta + 3\pi/2$  with the positive real axis.

Therefore the line  $L$  forms an angle  $\text{arg}(M(U)) + \theta + \pi/2$  with the positive real axis (since this is a line as opposed to a vector, a rotation of  $\pi$  makes no difference).

This completes our proof of theorem 3.5. □

## References

- [1] Bebiano N, Querió J. The determinant of the sum of two normal matrices with prescribed eigenvalues. *Linear Algebra and its Applications*. 1985;71:23–28.
- [2] Marcus M. Derivations, plücker relations and the numerical range. *Indiana University Math Journal*. 1973;22:1137–1149.
- [3] de Oliveira GN. Research problem: Normal matrices. *Linear and Multilinear Algebra*. 1982; 12:153–154.
- [4] Wikipedia contributors. Jacobi's formula — Wikipedia, the free encyclopedia ; 2019. [Online; accessed 13-February-2019]; Available from: [https://en.wikipedia.org/w/index.php?title=Jacobi%27s\\_formula&oldid=880845059](https://en.wikipedia.org/w/index.php?title=Jacobi%27s_formula&oldid=880845059).