

# Epstein Barr Virus and atrial fibrillation – A causal link?

Ilija Barukčić<sup>1</sup>

<sup>1</sup> Internist, Horandstrasse, DE-26441 Jever, Germany

Correspondence: Ilija Barukčić, Horandstrasse, DE-26441 Jever, Germany. Tel: 0049-(0)4466-333. E-mail: Barukcic@t-online.de

Received: October 6, 2018; Accepted: October 6, 2018; Published: October 6, 2018

## Abstract

**Objective:** Atrial fibrillation (AF) is very frequent and clinically significant arrhythmia. The incidence of atrial fibrillation is continuously rising. Meanwhile several risk factors for AF development have been identified but the etiology is not cleared.

**Methods:** A systematic review and re-analysis of studies which investigated the relationship between AF and some risk factors was conducted. The method of the *conditio sine qua non* relationship, the method of the *conditio per quam* relationship, the method of the exclusion relationship and the mathematical formula of the causal relationship *k* were used to proof the hypothesis. Significance was indicated by a p-value of less than 0.05.

**Results:** The studies analysed were able to provide direct and indirect evidence that AF is caused by a process of inflammation while a direct identification of the factor causing AF was not possible. Still, it appears to be very probable that Epstein-Barr virus (EBV) is the cause of AF.

**Conclusion:** Atrial fibrillation (AF) is caused by an inflammatory process.

**Keywords:** Epstein-Barr virus, atrial fibrillation, causal relationship

## 1. Introduction

In general, atrial fibrillation (AF) is regarded as a disease of the elderly, with a substantially increase (Chugh et al., 2014) of incidence and prevalence with age but asymptomatic atrial fibrillation (AF) has been detected by implanted cardiac devices (Glutzer & Ziegler, 2013) which is generally undetected in the clinical setting. Thus far, more than 10% of the general population older than 80 years have AF. Nonetheless, about 90 % of the elderly patients have none AF. Among other measures, long-term use of oral anticoagulants (OACs) for stroke prevention for people aged over 75 who have atrial fibrillation is required (Mant et al., 2007). The precise mechanisms involved in AF is not well understood. In particular, people who are living healthy lives and have no other medical problems develop sometimes atrial fibrillation to. In addition to cardiac causes such as congestive heart failure or valve disease other common risk factors (Wasmer et al., 2017) of atrial fibrillation like cigarette smoking, diabetes mellitus, hypertension (Kirchhof et al., 2012), prevalent coronary artery disease (Benjamin et al., 1994) chronic kidney disease (Psaty et al., 1997) have been identified and are well established. Less well-established risk factors are alcohol consumption, genetic factors (Fox et al., 2004), left atrial (LA) enlargement (Psaty et al, 1997) and other. The diagnosis of AF is secured by an cost-effective electrocardiogram (ECG) showing the typical pattern of absolutely irregular RR intervals and no discernible, distinct P waves while both symptomatic and asymptomatic episodes of AF (Kirchhof et al., 2009; Savelieva & Camm, 2000) must be distinguished. Further ECG monitoring is of further help for the management of AF. Several studies hypothesized even that inflammation (Aviles et al., 2003; Issac et al., 2007; Kirchhof et al., 2013) is involved in the pathogenesis and development of AF (Stroke Risk in Atrial Fibrillation Working Group, 2007; Cha et al., 2018).

## 2. Material and methods

### 2.1 Material

#### 2.1.1 Search Strategy

To answer the questions addressed in this paper, the electronic database PubMed was searched for appropriate studies conducted in any country which investigated the relationship between atrial fibrillation and inflammation. The search in PubMed was performed while using some medical key words. The articles found were saved as a \*.txt file while using the support of PubMed. The created \*.txt file was converted into a \*.pdf file. The abstracts

where studied within the \*.pdf file. Those articles were considered for a re-view which provided access to data without any data access barrier. Additionally the reference list of identified articles was used as a potential source of articles appropriate for this study.

### 2.1.2. The 2x2 Table

The meaning of the abbreviations  $a_t$ ,  $b_t$ ,  $c_t$ ,  $d_t$ ,  $N_t$  of the data table used are explained by a 2 by 2-table **Table 1**.

Table 1. The sample space of a contingency table.

		Conditioned $B_t$ (Outcome)		
		Yes = +1	Not = +0	Total
Condition $A_t$ (risk factor)	Yes =+1	$a_t$	$b_t$	$A_t$
	Not = +0	$c_t$	$d_t$	$\underline{A}_t$
	Total	$B_t$	$\underline{B}_t$	$N_t$

In general it is  $(a_t+b_t) = A_t$ ,  $(c_t+d_t) = \underline{A}_t$ ,  $(a_t+c_t) = B_t$ ,  $(b_t+d_t) = \underline{B}_t$  and  $a_t+b_t+c_t+d_t=N_t$ . Equally, it is  $B_t+\underline{B}_t = A_t + \underline{A}_t = N_t$ . In this context, it is  $p(a_t)=p(A_t \cap B_t)$ ,  $p(A_t) = p(a_t)+p(b_t)$  or  $p(A_t)= p(A_t \cap B_t)+ p(b_t) =p(A_t \cap B_t)+p(A_t \cap \underline{B}_t)$  while  $p(A_t)$  is not defined as  $p(a_t)$ . In the same context, it is  $p(B_t) = p(a_t)+p(c_t) = p(A_t \cap B_t) +p(c_t)$  and equally in the same respect  $p(\underline{B}_t) = 1- p(B_t) =p(b_t)+p(d_t)$ . Furthermore, the joint probability of  $A_t$  and  $B_t$  is denoted in general by  $p(A_t \cap B_t)$ . Thus far, it is  $p(A_t \cap B_t) = p(A_t) - p(b_t) = p(B_t) - p(c_t)$  or in other words it follows clearly that  $p(B_t) + p(b_t) - p(c_t) = p(A_t)$ . Thus far, define  $\Lambda = p(b_t) - p(c_t)$ , under conditions of probability theory and we obtain  $p(B_t) + \Lambda = p(A_t)$ . In general, it is  $p(a_t)+p(c_t)+p(b_t)+p(d_t) = 1$ .

### 2.1.3 The studies used to analyze the conventional risk factor

The table (**Table 2**) presents the data of some studies which analyzed the relationship between AF and some risk factors. The contradictions are signed by red colour.

Table 2. The relationship between some classical risk factors and AF.

Study Id	Year	Country	Risk Factor	Case_P	Case_T	Con_P	Con_T	k	p-val	X <sup>2</sup> (SINE)	X <sup>2</sup> (IMP)	X <sup>2</sup> (IMP^SINE)	X <sup>2</sup> (EXCL)
Wang et al.	2015	China	Family history	12	285	11	300	0.01398	0.15872	260.55	4.79	265.34	6.21
Wang et al.	2015	China	Alcohol	97	285	89	300	0.04688	0.03726	123.36	42.11	165.46	82.74
Wang et al.	2015	China	Smoke	147	285	159	300	-0.01422	0.06220	66.34	82.10	148.44	145.44
Chao et al.	2018	Taiwan	Diabetes mellitus	3770	15756	2884	14658	0.0513	3.5E-20	9117.30	1249.56	10366.86	3037.25
Wang et al.	2015	China	Diabetes mellitus	86	285	90	300	0.00191	0.0717	138.25	45.51	183.77	67.19
Wang et al.	2015	China	Dyslipidemia	117	285	112	300	0.03808	0.04422	98.44	54.29	152.73	106.89
Kim et al.	2014	USA	Hyperlipidemia	6439	20852	34561	104260	-0.01801	8.3E-12	9961.64	29132.39	39094.04	2999.10
Kim et al.	2014	USA	Hypertension	5876	20852	26360	104260	0.02468	3.4E-19	10755.11	21554.26	32309.38	2726.45
Chao et al.	2018	Taiwan	Hypertension	11679	15756	9005	14658	0.13594	4.5E-125	1054.70	3919.99	4974.69	15250.08
Wang et al.	2015	China	Hypertension	220	285	221	300	0.04091	0.04701	14.60	110.25	124.85	278.31
Wang et al.	2015	China	Cerebral infarction	91	285	82	300	0.05034	0.03448	131.38	38.39	169.77	76.08
Chao et al.	2018	Taiwan	Previous stroke/TIA	7772	15756	4724	14658	0.17366	2.1E-203	4045.21	1785.49	5830.69	8666.46
Wang et al.	2015	China	Arteriosclerosis	201	285	190	300	0.07636	0.01277	24.46	91.84	116.31	243.87
Chao et al.	2018	Taiwan	Prev. vascular disease	2004	15756	5230	14658	-0.2694	0	12002.02	3780.44	15782.45	809.64
Chao et al.	2018	Taiwan	Congestive heart failure	8680	15756	3042	14658	0.35252	0	3177.37	789.18	3966.55	11207.97
Kim et al.	2014	USA	Cardiovascular disease	1439	20852	5355	104260	0.02902	3.7E-24	18072.37	4220.00	22292.37	403.81

### 2.1.4 Methylprednisolone and permanent AF

Dernellis & Panaretou (Dernellis & Panaretou, 2004) investigated whether a low-dose glucocorticoid therapy (16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) compared to placebo is able to reduce AF.

Table 3. The relationship between a therapy with (methyl-) prednisolone and permanent AF

		The Study of Dernellis & Panaretou, 2004		
		Permanent AF <B>		
		Yes	No	Total
<A>	Methylprednisolone 16 mg	1	15	16
	No	51	37	88
Total		52	52	104

$$k = -0.3731$$

$$p \text{ value } (k) = 0.0001$$

$$95\% \text{ CI } (k) = (0.1538; 0.5924)$$

<A> EXCLUDES <B> and vice versa

$$p(\text{EXCL}) = 0.99038462$$

$$X^2(\text{EXCL}) = 0.0204$$

## 2.2 Methods

### 2.2.1 Statistical Analysis

All statistical analyses (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) were performed with Microsoft Excel version 14.0.7166.5000 (32-Bit) Software (Microsoft GmbH, Munich, Germany). The level of significance was set to 0.05. The probabilities of the contingency table are viewed by the following table (Table 4).

Table 4. The probabilities of a contingency table

		Conditioned		
		$B_t$		
		Yes = +1	No = +0	Total
Condition $A_t$	Yes = +1	$p(a_t) = p(A_t \cap B_t)$	$p(b_t)$	$p(A_t)$
	No = +0	$p(c_t)$	$p(d_t)$	$p(\underline{A}_t)$
Total		$p(B_t)$	$p(\underline{B}_t)$	1

### 2.2.2 Independence

In the case of independence of  $A_t$  and  $B_t$  (Kolmogoroff, 1933) it is generally valid that

$$p(A_t \cap B_t) \equiv p(A_t) \times p(B_t) \tag{1}$$

### 2.2.3 Exclusion ( $A_t$ Excludes $B_t$ and Vice Versa Relationship)

The mathematical formula of the *exclusion* relationship ( $A_t$  excludes  $B_t$  and vice versa) of a population was defined (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) as

$$p(A_t | B_t) \equiv \frac{b_t + c_t + d_t}{N_t} \equiv 1 - p(a_t) \equiv p(b_t) + p(c_t) + p(d_t) \equiv p(c_t) + (1 - p(B_t)) \equiv p(b_t) + (1 - p(A_t)) \equiv +1 \tag{2}$$

and used to proof the hypothesis:  $A_t$  excludes  $B_t$  and vice versa.

### 2.2.4 Necessary Condition (Conditio Sine Qua Non)

The mathematical formula of the *necessary* condition relationship (conditio sine qua non) of a population was defined (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) as

$$p(A_t \leftarrow B_t) \equiv \frac{a_t + b_t + d_t}{N_t} \equiv p(a_t) + p(b_t) + p(d_t) \equiv p(a_t) + (1 - p(B_t)) \equiv +1 \quad (3)$$

and used to proof the hypothesis: *without*  $A_t$  *no*  $B_t$ .

### 2.2.5. Sufficient Condition (Conditio per Quam)

The mathematical formula of the *sufficient* condition relationship (conditio per quam) of a population was defined (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) as

$$p(A_t \rightarrow B_t) \equiv \frac{a_t + c_t + d_t}{N_t} \equiv p(a_t) + p(c_t) + p(d_t) \equiv p(d_t) + p(B_t) \equiv +1 \quad (4)$$

and used to proof the hypothesis: *if*  $A_t$  *then*  $B_t$ .

### 2.2.6 The $X^2$ Goodness of Fit Test of a Necessary Condition

Under conditions where the chi-square goodness of fit test (Pearson, 1900) cannot be used it is possible to use an approximate and conservative (one sided) confidence interval known as *the rule of three* (Rumke, 1975; Hanley et al., 1983; Louis, 1981; Jovanovic et al., 1997). Using *the continuity correction* (Yates, 1934), the chi-square value of a conditio sine qua non distribution is derived as

$$\chi^2 (\text{SINE}) \equiv \frac{\left( c_t - \left( \frac{1}{2} \right) \right)^2}{(B_t)} + 0 = 0 \quad (5)$$

### 2.2.7 The $X^2$ Goodness of Fit Test of the Exclusion Relationship

The chi square value with degree of freedom  $2-1=1$  of the exclusion relationship (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) with a *continuity correction* (Yates, 1934) can be calculated as

$$\chi^2 (\text{EXCL}) = \frac{(-a_t - 0,5)^2}{A_t} + \frac{(-a_t - 0,5)^2}{B_t} \quad (6)$$

### 2.2.8 The Mathematical Formula of the Causal Relationship k

The mathematical formula of the causal relationship k (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006; Barukčić, 2016; Barukčić, 2017; Barukčić, 2018) is defined *at every single event, at every single Bernoulli trial t*, as

$$k(A_t, B_t) \equiv \frac{(p(A_t \cap B_t) - (p(A_t) \times p(B_t)))}{\sqrt{(p(A_t) \times p(\underline{A}_t)) \times (p(B_t) \times p(\underline{B}_t))}} \quad (7)$$

where  $A_t$  denotes the cause and  $B_t$  denotes the effect. The chi-square distribution can be applied to determine the significance of causal relationship k. Pearson's concept of correlation (Pearson, 1896) is not identical with causation, causation is not identical with correlation. In particular, the relationship between correlation and causation has already been discussed in many publications (Barukčić, 1989; Barukčić, 2005; Barukčić, 2006;

Barukčić, 2016; Barukčić, 2017; Barukčić, 2018). Thus far, repeating itself over and over again on this topic is only a waste of time and will not contribute anything new to further scientific progress.

### 2.2.9 The 95% Confidence Interval of the Causal Relationship k

A confidence interval (CI) of the causal relationship k calculated from the statistics of the observed data can help to estimate the true value of an unknown population parameter with a certain probability. Let the sample mean S be

$$S = \overline{k(A_t, B_t)} = \frac{k(A_1, B_1) + k(A_2, B_2) + \dots + k(A_n, B_n)}{n} = \frac{\sum_{t=1}^n k(A_t, B_t)}{n} \quad (8)$$

Since  $k(A_t, B_t)$  is Bernoulli(p) distributed with  $E(k(A_t, B_t)) = (1 \times p(k(A_t, B_t))) + (0 \times (1 - p(k(A_t, B_t)))) = p(k(A_t, B_t))$  where  $E(k(A_t, B_t))$  denotes the expected value of  $k(A_t, B_t)$  it is

$$E(S) = p(k(A_t, B_t)) \quad \text{and} \quad \sigma(S)^2 = \frac{p(k(A_t, B_t)) \times (1 - p(k(A_t, B_t)))}{n} \quad (9)$$

where  $\sigma(S)^2$  denotes the variance of the sampling distribution of  $p(k(A_t, B_t))$ . When the sample size is not too small, the central limit theorem based normal approximation can be used to estimate the confidence interval (CI) as

$$p(k(A_t, B_t)) \pm \left( Z \times \sqrt{\frac{p(k(A_t, B_t)) \times (1 - p(k(A_t, B_t)))}{n}} \right) = p(k(A_t, B_t)) \pm \left( \sqrt{\frac{Z^2}{n} \times p(k(A_t, B_t)) \times (1 - p(k(A_t, B_t)))} \right) \quad (10)$$

where  $p(k(A_t, B_t))$  denotes the proportion of successes in a Bernoulli trial process and Z is the  $(1 - (\alpha/2))$  quantile of a standard normal distribution. For a 95% confidence level  $Z \sim 1.96$ . For an unknown standard deviation the Student's t distribution t can be used as the critical value. Still, it is known that  $\sigma(S)^2$  has the maximum value  $(1/(4 \times n))$  when  $p=1/2$  and approximately we obtain

$$p(k(A_t, B_t)) \pm \left( \sqrt{\frac{Z^2}{4 \times n}} \right) \Leftrightarrow p(k(A_t, B_t)) \pm \left( \sqrt{\frac{1.96^2}{4 \times n}} \right) \approx p(k(A_t, B_t)) \pm \left( \sqrt{\frac{1}{n}} \right) \quad (11)$$

The proposed approximation is of use even under circumstances where  $p(\dots) = 0.9999 \dots 999 \sim p=1$ . In this context, we obtain the critical value  $p_{\text{critical}}$  approximately as  $p_{\text{critical}} = 1 - (1/n)^{1/2}$ . In particular, the concept of Chebyshev's inequality is profound because the same inequality is true for every distribution even if the distribution isn't normal. Thus far, Chebyshev's inequality allows calculating the 95% confidence of the causal relationship k and so by the Chebyshev inequality it is

$$p \left\{ p(k(A_t, B_t)) - c \times \sqrt{\sigma(S)^2} < S < p(k(A_t, B_t)) + c \times \sqrt{\sigma(S)^2} \right\} \geq 1 - \frac{1}{c^2} \quad (12)$$

were the right side has the value 0.95 when  $c = (20)^{1/2}$ . This is the case since  $(1 - (1/c^2)) = 0.95$  or  $0.05 = (1/c^2)$  or  $c^2 = (1/0.05)$  or  $c^2 = (100/5)$  or  $c^2 = 20$  or  $c = (20)^{1/2}$ . Thus far, if S does lie in the interval

$$\left\{ p(k(A_t, B_t)) - \sqrt{20 \times \sigma(S)^2}, p(k(A_t, B_t)) + \sqrt{20 \times \sigma(S)^2} \right\} \quad (13)$$

then  $p(k(A_t, B_t))$  itself must be in the interval

$$\left\{ S - \sqrt{20 \times \sigma(S)^2}, S + \sqrt{20 \times \sigma(S)^2} \right\} \quad (14)$$

which is equally the 95% confidence interval for an unknown parameter  $p(k(A_t, B_t))$ . Again,  $\sigma(S)^2$  has the maximum value  $(1/(4 \times n))$  when  $p=1/2$ , so we have

$$\left\{ S - \sqrt{\frac{20 \times 1}{4 \times n}}, S + \sqrt{\frac{20 \times 1}{4 \times n}} \right\} \quad (15)$$

or the 95% interval for the causal relationship  $k$  as

$$\left\{ k(A_t, B_t) - \sqrt{\frac{5}{n}}, k(A_t, B_t) + \sqrt{\frac{5}{n}} \right\} \quad (16)$$

### 2.2.10 Hypergeometric distribution

The probability of having exactly  $a_t$  (**Table 1**) successes or the significance of the causal relationship  $k$  can be tested under conditions of sampling without replacement by the hypergeometric distribution as

$$p(a_t) = \frac{\binom{A_t}{a_t} \times \binom{N_t - A_t}{n_t - a_t}}{\binom{N_t}{B_t}} \quad (17)$$

### 2.2.11 Odds Ratio

The odds ratio (OR) is given (Barukčić, 2018) by

$$OR(A_t, B_t) \equiv \frac{a_t / b_t}{c_t / d_t} = \frac{a_t \times d_t}{c_t \times b_t} \quad (18)$$

Under conditions were  $c_t=0$  we have a *conditio sine qua non* relationship while the odds ratio collapses. Under conditions were  $b_t=0$  we have a *conditio per quam* relationship but the odds ratio collapses again, since to date it is not generally accepted (Barukčić & Barukčić, 2016b; Barukčić, 2018d) to divide by zero. To avoid confusion on this issue, 0.5 is added to the cells  $a_t, b_t, c_t, d_t$  (Barukčić, 2018), if zero causes some problems with the calculation of the odds ratio or its standard error which is often very misleading. In point of fact, the odds ratio (OR) is nothing more but *Yule's coefficient of association Q* (Barukčić, 2018) re-written (Barukčić, 2018) in a non-normalized form and given by

$$Q(A_t, B_t) \equiv \frac{OR(A_t, B_t) - 1}{OR(A_t, B_t) + 1} = \frac{\frac{(a_t \times d_t)}{(b_t \times c_t)} - 1}{\frac{(a_t \times d_t)}{(b_t \times c_t)} + 1} = \frac{(a_t \times d_t) - (b_t \times c_t)}{(a_t \times d_t) + (b_t \times c_t)} = \frac{(a_t \times d_t) - (b_t \times c_t)}{(a_t \times d_t) - (b_t \times c_t)} \quad (19)$$

If  $Q = 0$  then there is no association. Yule's coefficient of association,  $Q$ , and thus far Odds ratio itself, has been severely criticized by Karl Pearson (1857–1925), the long-time and rarely challenged leader of statistical science and by Heron (Barukčić, 2018). The standard error and 95% confidence interval of the odds ratio (OR) can be calculated according to Altman (Altman, 1991). The standard error of the log odds ratio is given by

$$SE(\ln(OR(A_t, B_t))) \equiv \sqrt{\frac{1}{a_t} + \frac{1}{b_t} + \frac{1}{c_t} + \frac{1}{d_t}} \quad (20)$$

where  $\ln$  denotes the logarithmus naturalis. The 95% confidence interval of the odds ratio is given by

$$95\% \text{ CI} = \exp\left(\ln(OR(A_t, B_t)) - (1.96 \times SE(\ln(OR(A_t, B_t))))\right) \text{ to } \exp\left(\ln(OR(A_t, B_t)) + (1.96 \times SE(\ln(OR(A_t, B_t))))\right) \quad (21)$$

2.2.12 The unknown population proportion  $\pi_{upper}$

Tests of hypotheses concerning the sampling distribution of the sample proportion  $\mathbf{p}$  (i. e. *conditio sine qua non* p(SINE), *conditio per quam* p(IMP) et cetera) can be performed using the normal approximation. The calculation of the rejection region based on the sample proportion to construct a confidence interval for an unknown population proportion  $\pi_{upper}$  can be performed under conditions of *sampling without replacement* by the formula

$$\pi_{critical\ upper} = \left( \mathbf{p} - \frac{1}{2 \times \mathbf{n}} \right) - \left( Z \times \sqrt{\frac{\mathbf{p} \times (1 - \mathbf{p})}{\mathbf{n}} \times \left( \frac{\mathbf{N} - \mathbf{n}}{\mathbf{N} - 1} \right)} \right) \tag{22}$$

while the term  $((\mathbf{N} - \mathbf{n}) / (\mathbf{N} - 1))$  denotes *the finite population correction* (Isserlis, 1918).

2.2.13 The Chi Square Distribution

The following critical values of the chi square distribution as visualized by **Table 5** are used in this publication.

Table 5. The critical values of the chi square distribution (degrees of freedom: 1)

	<b>p-Value</b>	<b>One sided X<sup>2</sup></b>	<b>Two sided X<sup>2</sup></b>
	0.1000000000	1.642374415	2.705543454
	<b>0.0500000000</b>	<b>2.705543454</b>	<b>3.841458821</b>
	0.0400000000	3.06490172	4.217884588
	0.0300000000	3.537384596	4.709292247
	0.0200000000	4.217884588	5.411894431
	0.0100000000	5.411894431	6.634896601
The chi square distribution	0.0010000000	9.549535706	10.82756617
	0.0001000000	13.83108362	15.13670523
	0.0000100000	18.18929348	19.51142096
	0.0000010000	22.59504266	23.92812698
	0.0000001000	27.03311129	28.37398736
	0.0000000100	31.49455797	32.84125335
	0.0000000010	35.97368894	37.32489311
	0.0000000001	40.46665791	41.82145620

### 3. Results

#### 3.1 Classical risk factors (RF) and atrial fibrillation (AF)

Some of the studies analysed provided additional data about the relationship between classical risk factors like family history of AF, alcohol, smoking, diabetes mellitus, hyperlipidemia, hypertension, previous stroke/TIA, arteriosclerosis, congestive heart failure, cardiovascular disease and atrial fibrillation with conflicting results on its role as an etiological factor. The data are viewed by Table 2. The hypothesis *without* RF *no* AF was tested. The hypothesis if RF *then* AF was tested too. In fact, the null-hypothesis altogether had to be rejected.

#### Proof.

Without family history of AF no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If family history of AF then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without alcohol no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If alcohol then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without smoking no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If smoking then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without diabetes mellitus no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If diabetes mellitus then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without dyslipidemia no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If hyperlipidemia then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without hypertension no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If hypertension then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without perbral infarction no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If previous stroke/TIA then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without arteriosclerosis no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If previous vascular disease then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without congestive heart failure no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If congestive heart failure then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Without cardiovascular disease no atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

If cardiovascular disease then atrial fibrillation. (Null hypotheses rejected, no significant result (Table 2)).

Neither family history of AF nor alcohol nor smoking nor diabetes mellitus nor hyperlipidemia nor hypertension nor previous stroke/TIA nor arteriosclerosis nor congestive heart failure nor cardiovascular disease et cetera can be regarded as a cause or as the cause of AF. **Q. e. d.**

### 3.1 A therapy with (Methyl) prednisolone and atrial fibrillation(AF)

Little or none direct information is available on the effect of an anti-viral or anti-inflammatory therapy on the reduction of atrial fibrillation. According to Engelmann and Svendsen (Engelmann & Svendsen, 2005) a reatment with glucocorticoids seems to reduce AF. The systematic review and meta-analysis of observational studies of Liu et al. (Liu et al., 2007) suggests that increased CRP levels are related to greater risk of AF recurrence. Henningsen et al. (Henningsen et al., 2009) investigated the role of inflammatory processes in the development of atrial fibrillation and were able to provide evidence that patients with AF had elevated levels of inflammatory markers. In general, a low-dose glucocorticoid therapy (i. e. 16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) compared to placebo should have no effect on (recurrent or permanent) AF. In other words we do expect that a therapy with (methyl-) prednisolone and AF are independent of each other. Dernellis & Panaretou (Dernellis & Panaretou, 2004) investigated the effect of a low-dose glucocorticoid therapy (16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) compared to placebo on AF. According to Dernellis & Panaretou (Dernellis & Panaretou, 2004) a therapy with (methyl-) prednisolone was able to reduce recurrent AF from 50% in the placebo group to 9.6% in the glucocorticoid group. Furthermore, permanent AF was reduced by a therapy with (methyl-) prednisolone from 29% in the placebo group to 2% in the glucocorticoid group. The results were significant. The data of the study of Dernellis & Panaretou (Dernellis & Panaretou, 2004) were reanalysed (**Table 3**).

#### Null Hypothesis:

A therapy with (methyl-) prednisolone does not prevent from permanent AF. In other words,  $k \geq 0$ .

#### Alternative Hypothesis:

A therapy with (methyl-) prednisolone does prevent from permanent AF. In other words,  $k < 0$ .

The significance level (Alpha) below which the null hypothesis will be rejected is  $\alpha=0.05$ .

#### Proof.

The results of the re-analyses of the data of the study of Dernellis & Panaretou (Dernellis & Panaretou, 2004) which investigated the relationship between a low-dose glucocorticoid therapy (i. e. 16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) compared to placebo and the reduction of permanent AF are viewed by the table (**Table 3**). Altogether, the data of the study of Dernellis & Panaretou (Dernellis & Panaretou, 2004) provide highly significant evidence that a low-dose glucocorticoid therapy (i. e. 16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) compared to placebo **excludes** permanent AF ( $p(\text{Prednisolone excludes AF}) = 0.99038462$ ;  $X^2(\text{Prednisolone excludes AF}) = 0.0204$ ) while the causal relationship was  $k < 0$ . In other words, the data as published by Dernellis & Panaretou (Dernellis & Panaretou, 2004) are not self-contradictory (Barukčić, 2018) and can be used for causal analysis too. The causal relationship between a therapy with (methyl-) prednisolone and permanent AF is  $k = -0.3731$  while the exact p value ( $k$ ) calculated by the hypergeometric distribution is p value ( $k$ )  $< 0.0001$ . The 95% CI ( $k$ ) is  $-0.1538$  until  $-0.5924$ , a highly significant result. In other words, there is a highly causal relationship between a low-dose glucocorticoid therapy (i. e. 16 mg methylprednisolone for 4 weeks tapered to 4 mg for 4 months) and the absence of permanent AF. A low-dose glucocorticoid therapy is highly effective against permanent AF and an “anti-dot” against permanent AF. **Q. e. d.**

## 4. Discussion

Much attention has been devoted in the past years to inflammation (Aviles et al., 2003; Anderson et al., 2004; Conway et al., 2004; Asselbergs et al., 2005) as the pathophysiological mechanism underlying the genesis of AF which has been reinforced even by experimental studies (Kirchhof et al., 2013). Still, AF remains not even partially proofed. The pathophysiology of AF is complex, but inflammation can lead to a kind of “atrial myocarditis” with subsequent structural atrial changes resulting among other in AF (Korantzopoulos et al., 2003). Several case studies support this hypothesis. A 57-year-old patient with rheumatoid arthritis treated with a combination of etanercept and methotrexate and developed a new-onset of atrial fibrillation after treatment for five months on this therapy (Wooten et al., 2000). *Etanercept* is effective against cytomegalovirus (Hung et al., 2017) but not for sure against EBV. Thus far, according to this case, it is highly improbable, that cytomegalovirus is a cause or the cause of atrial fibrillation. The incidence of myocarditis as a complication of an EBV infection is not so high (Takano et al., 2008). However, a chronic active Epstein-Barr virus (CAEBV) infection characterized by high EBV DNA levels in the peripheral blood (Kawabe et al., 2018) can cause organ damage demonstrated by

endomyocardial biopsies with lymphocytic infiltration, necrosis of cardiomyocytes et cetera and end up in a chronic active myocarditis (Takano et al., 2008). Sometimes, an Epstein-Barr myocarditis (Antonakaki et al., 2016) is so severe that an urgent orthotopic heart transplantation is necessary. Finally, Aghenta et al. (Aghenta et al., 2008) published the case of a 19-year-old healthy male patient presented to hospital with clinical features of infectious mononucleosis (IM). Symptoms worsened despite therapy (ibuprofen, intravenous methylprednisolone, and intravenous fluids). On day 2 of admission Electrocardiogram (ECG) showed atrial fibrillation. An Epstein Barr virus infection is able to cause a symptomatic atrial fibrillation (Aghenta et al., 2008) as a result of viral invasion. One case is of course not enough to provide a proof of a cause effect relationship between EBV and AF.

To date, data are available which support the hypothesis that patients with rheumatoid arthritis have increased risk (Bacani et al., 2015) of atrial fibrillation. The incidence of atrial fibrillation (AF) among patients with rheumatoid arthritis compared to the entire Danish population was increased by a 40% (Lindhardsen et al., 2012). Bacani et al. (Bacani et al., 2015) found that the incidence of AF even after adjustment for AF risk factors is increased in patients with RA. Epstein–Barr virus (EBV) inflammation is the main cause of rheumatoid arthritis (RA).

The Epstein–Barr virus (EBV) was discovered by electron microscopy of cells cultured from Burkitt's lymphoma tissue by Epstein et al. (Epstein et al., 1964) Epstein-Barr virus (EBV), also known as human herpesvirus 4, is a gamma herpes virus which infects a large fraction of the human population. After the primary infection, EBV persists for life (Cohen, 2000) in memory B cells in the peripheral blood of human host while well controlled by host's immune system.

There is currently no treatment for removing EBV infections and very little success treating EBV infection with drugs. *Corticosteroids* are sometimes helpful. In the excellent paper, Dernellis and Panaretou (Dernellis & Panaretou, 2004) demonstrated that the use of low-dose Methylprednisolone successfully prevents recurrent and permanent AF.

Cell culture experiments support the thesis that *Vitamin D* may reduce the severity of viral infection by suppressing inflammation (Beard et al., 2011).

Acyclovir with limited gastrointestinal absorption had no significant effect on EBV-related symptoms (Bissell et al., 1980) while *Valacyclovir* treatment in Epstein-Barr virus infection was effective (Lerner et al., 2007). Valacyclovir inhibits EBV thymidine kinase and is absorbed more efficiently from the gastrointestinal tract than acyclovir. Valacyclovir was not able to reduce infant CMV acquisition or breastmilk shedding of cytomegalovirus (Roxby et al., 2014) while EBV reactivation was decreased with valganciclovir prophylaxis (Gill et al., 2014).

The direct anti-viral activity of *ascorbic acid* (vitamin C, ascorbate) against herpes simplex virus type 1, influenza virus type A, picornaviridae virus 1, herpes simplex viruses (types 1 and 2), cytomegalovirus, parainfluenza virus type 2 and other viruses are documented in vitro (Furuya et al., 2008). Vitamin C deficiency is associated with viral infections (Chen et al., 2009). The clinical study of (Mikirova & Hunninghake, 2014) demonstrated that an intravenous administration of high dose ascorbate (IVC) therapy according to Riordan IVC protocol (Riordan et al., 2003; Mikirova et al., 2013) is able to reduce EBV EA IgG and EBV VCA IgM antibody levels over time. High-dose IVC infusions at the 7.5, 15, 25, and 50 gram dosages adjusted by plasma ascorbic acid levels attained post infusion and patients' tolerance lead to hemolysis in patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency. Adding magnesium to high-dose IVC ascorbate infusions can help to reduce incidence of vein irritation and spasm.

*Zinc* (Zn) play a very important role in human immune system because of its pivotal role in the efficiency of modulating immune function (Mocchegiani & Muzzioli, 2000; Hirano et al., 2008). Zinc-deficient patients experience increased susceptibility to a variety of pathogens (Shankar & Prasad, 1998). Zinc supplementation may increase host resistance to infections. Arens and Travis (Arens and Travis, 2000) were able to inactivate herpes simplex virus (HSV) with zinc salts.

Despite good progress in the management of atrial fibrillation, AF remains one of the major causes of sudden death and morbidity in the world. For the patient, AF is associated with severe consequences such as stroke or death and other while the direct costs of AF for the society are already extraordinary high (Stewart et al., 2004; Kim et al., 2011). In the year 2011, the US-AF costs have been estimated to range from \$6.0 to \$26.0 billion (Kim et al., 2011). The complexity underlying AF and the relative inefficacy of the currently available pharmacotherapy requires development of new treatment strategies to try to *cure* AF (**Table 6**).

Patient:

**Table 6. Therapeutic schema to cure atrial fibrillation (experimental; not verified by trials)**

Drug	(Methyl-) prednisolon	Cimetidin	Valacyclovir	Zinc-histidine	Vitamin C	Vitamin D 20.000 I. U.
Dosage	40 mg 1-0-0	400 mg 1-0-1	500 mg 1-0-1	15 mg 0-1-0	1 g 0-1-0	1x1 per week
Week	a.c. (on 5 days per week)	a.c./c.c.	plus a lot of water	p.c.	c.c./p.c.	(i.e. Sundays)
1	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
2	-	x	x (7 days)	x	x	x
3	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
4	-	x	x (7 days)	x	x	x
5	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
6	-	x	x (7 days)	x	x	x
7	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
8	-	x	x (7 days)	x	x	x
9	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
12	-	x	x (7 days)	x	x	x
11	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
12	-	x	x (7 days)	x	x	x
13	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
14	-	x	x (7 days)	x	x	x
15	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
16	-	x	x (7 days)	x	x	x
17	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
18	-	x	x (7 days)	x	x	x
19	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
20	-	x	x (7 days)	x	x	x
21	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
22	-	x	x (7 days)	x	x	x
23	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
24	-	x	x (7 days)	x	x	x
26	x (Mo, Tue, Wen, Thur, Fri)	x	-	x	x	x
26	-	x	x (7 days)	x	x	x

Start of therapy on:

The drugs taken are cancelled. "x" denotes the necessity of taking the drug, "-" denotes the necessity not to take the drug.

## 5. Conclusion

Given the complexity of AF pathophysiology the findings of this study provide evidence of an impact of EBV infection on atrial fibrillation. Undoubtedly, the hypothesis: *without* EBV infection *no* AF remains to be elucidated and more systematic, regional studies of AF are required to better understand the relationship between an EBV infection and AF.

## Conflict of interest

Author declare no conflict of interests for this article.

## Acknowledgement

None.

## References

- Aghenta, A., Osowo, A., & Thomas, J. (2008). Symptomatic atrial fibrillation with infectious mononucleosis *Canadian Family Physician Medecin De Famille Canadien*, 54(5), 695–696.
- Anderson, J. L., Allen Maycock, C. A., Lappé, D. L., Crandall, B. G., Horne, B. D., Bair, T. L., . . . Muhlestein, J. B. (2004). Frequency of elevation of C-reactive protein in atrial fibrillation *The American Journal of Cardiology*, 94, 1255–1259. <https://doi.org/10.1016/j.amjcard.2004.07.108>
- Antonakaki, D., Mirza, S. A., Sundar, S., & Grapsa, J. (2016). Epstein-Barr futile myocarditis requiring urgent orthotopic heart transplantation *Perfusion*, 31, 431–432. <https://doi.org/10.1177/0267659115618460>
- Arens, M., & Travis, S. (2000). Zinc salts inactivate clinical isolates of herpes simplex virus in vitro *Journal of Clinical Microbiology*, 38(5), 1758–1762.
- Armitage, P., & Colton, T. (Eds.). (2005). *Encyclopedia of biostatistics* (2nd ed.). Chichester, West Sussex, England, Hoboken, NJ: John Wiley.
- Asselbergs, F. W., van den Berg, M. P., Diercks, G. F. H., van Gilst, W. H., & van Veldhuisen, D. J. (2005). C-reactive protein and microalbuminuria are associated with atrial fibrillation *International Journal of Cardiology*, 98, 73–77. <https://doi.org/10.1016/j.ijcard.2003.12.028>
- Aviles, R. J., Martin, D. O., Apperson-Hansen, C., Houghtaling, P. L., Rautaharju, P., Kronmal, R. A., . . . Chung, M. K. (2003). Inflammation as a risk factor for atrial fibrillation *Circulation*, 108, 3006–3010. <https://doi.org/10.1161/01.CIR.0000103131.70301.4F>
- Bacani, A. K., Crowson, C. S., Roger, V. L., Gabriel, S. E., & Matteson, E. L. (2015). Increased incidence of atrial fibrillation in patients with rheumatoid arthritis *BioMed Research International*, 2015, 809514. <https://doi.org/10.1155/2015/809514>
- Barukčić, I. (1989). *Die Kausalität* (1. Aufl.). Hamburg: Wiss.-Verl.
- Barukčić, I. (1997). *Die Kausalität* (2., völlig überarb. Aufl.). Wilhelmshaven: Scientia.
- Barukčić, I. (2005). *Causality: New statistical methods*. Norderstedt, Germany: Books on Demand GmbH.
- Barukčić, I. (2005). *Causality: New statistical methods*. Norderstedt, Germany: Books on Demand GmbH.
- Barukčić, I. (2006). *Causality: New statistical methods* (2. Aufl.). Norderstedt: Books on Demand.
- Barukčić, I. (2006). New Method for Calculating Causal Relationships *XXIIIrd International Biometric Conference, Montréal, Québec, Canada, July 16 — 21, 2006*, 23, 49. Retrieved from <http://www.medicine.mcgill.ca/epidemiology/hanley/ibc2006program.pdf>
- Barukčić, I. (2016). The Mathematical Formula of the Causal Relationship *International Journal of Applied*

- Physics and Mathematics*, 6, 45–65. <https://doi.org/10.17706/ijapm.2016.6.2.45-65>
- Barukčić, I. (2017). *Die Kausalität*. Norderstedt: Books on Demand.
- Barukčić, I. (2017). Helicobacter pylori—The Cause of Human Gastric Cancer *Journal of Biosciences and Medicines*, 05, 1–9. <https://doi.org/10.4236/jbm.2017.52001>
- Barukčić, I. (2017). *Theoriae causalitatis principia mathematica*. Norderstedt: Books on Demand.
- Barukčić, I. (2018). Epstein Bar Virus—The Cause of Hodgkin's Lymphoma *Journal of Biosciences and Medicines*, 06, 75–100. <https://doi.org/10.4236/jbm.2018.61008>
- Barukčić, I. (2018). Fusobacterium nucleatum —The Cause of Human Colorectal Cancer *Journal of Biosciences and Medicines*, 06, 31–69. <https://doi.org/10.4236/jbm.2018.63004>
- Barukčić, I. (2018). Gastric Cancer and Epstein-Barr Virus Infection *Modern Health Science*, 1, 1-18. <https://doi.org/10.30560/mhs.v1n2p1>
- Barukčić, I. (2018). Helicobacter Pylori is the Cause of Gastric Cancer *Modern Health Science*, 1, 43-50. <https://doi.org/10.30560/mhs.v1n1p43>
- Barukčić, I. (2018). Human Cytomegalovirus is the Cause of Glioblastoma Multiforme *Modern Health Science*, 1, p19. <https://doi.org/10.30560/mhs.v1n2p19>
- Barukčić, I. (2018). Human Papillomavirus—The Cause of Human Cervical Cancer *Journal of Biosciences and Medicines*, 06, 106–125. <https://doi.org/10.4236/jbm.2018.64009>
- Barukčić, I. (2018). Mycobacterium Avium Subspecies Paratuberculosis: The Cause Of Croh's Disease. *Modern Health Science*, 1, 19-34. <https://doi.org/10.30560/mhs.v1n1p19>
- Barukčić, K., & Barukčić, I. (2016). Epstein Barr Virus—The Cause of Multiple Sclerosis *Journal of Applied Mathematics and Physics*, 04, 1042–1053. <https://doi.org/10.4236/jamp.2016.46109>
- Beard, J. A., Bearden, A., & Striker, R. (2011). Vitamin D and the anti-viral state *Journal of Clinical Virology : the Official Publication of the Pan American Society for Clinical Virology*, 50, 194–200. <https://doi.org/10.1016/j.jcv.2010.12.006>
- Benjamin, E. J., Levy, D., Vaziri, S. M., D'Agostino, R. B., Belanger, A. J., & Wolf, P. A. (1994). Independent risk factors for atrial fibrillation in a population-based cohort. The Framingham Heart Study *JAMA*, 271(11), 840–844.
- Bissell, M. J., Hatie, C., Farson, D. A., Schwarz, R. I., & Soo, W. J. (1980). Ascorbic acid inhibits replication and infectivity of avian RNA tumor virus *Proceedings of the National Academy of Sciences of the United States of America*, 77(5), 2711–2715.
- Bolboacă, S. D., Jäntschi, L., Sestraș, A. F., Sestraș, R. E., & Pamfil, D. C. (2011). Pearson-Fisher Chi-Square Statistic Revisited *Information*, 2, 528–545. <https://doi.org/10.3390/info2030528>
- Bombardier, C., Laine, L., Reicin, A., Shapiro, D., Burgos-Vargas, R., Davis, B., . . . Schnitzer, T. J. (2000). Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. VIGOR Study Group *The New England Journal of Medicine*, 343, 1520-8, 2 p following 1528. <https://doi.org/10.1056/NEJM200011233432103>
- Centers for Disease Control and Prevention. (2018). Epstein-Barr Virus and Infectious Mononucleosis. Laboratory Testing. *National Center for Immunization and Respiratory Diseases*. Retrieved from <https://www.cdc.gov/epstein-barr/index.html>
- Cha, M.-J., Seo, H.-M., Choi, E.-K., Lee, J. H., Han, K., Lee, S.-R., . . . Oh, S. (2018). Increased Risk of Atrial Fibrillation in the Early Period after Herpes Zoster Infection: a Nationwide Population-based Case-control Study *Journal of Korean Medical Science*, 33, e160. <https://doi.org/10.3346/jkms.2018.33.e160>
- Chao, T.-F., Liu, C.-J., Lin, Y.-J., Chang, S.-L., Lo, L.-W., Hu, Y.-F., . . . Chen, S.-A. (2018). Oral Anticoagulation in Very Elderly Patients With Atrial Fibrillation: A Nationwide Cohort Study *Circulation*, 138, 37–47. <https://doi.org/10.1161/CIRCULATIONAHA.117.031658>
- Chen, J.-Y., Chang, C.-Y., Feng, P.-H., Chu, C.-C., So, E. C., & Hu, M.-L. (2009). Plasma vitamin C is lower in postherpetic neuralgia patients and administration of vitamin C reduces spontaneous pain but not brush-evoked pain *The Clinical Journal of Pain*, 25, 562–569. <https://doi.org/10.1097/AJP.0b013e318193cf32>

- Chugh, S. S., Havmoeller, R., Narayanan, K., Singh, D., Rienstra, M., Benjamin, E. J., . . . Murray, C. J. L. (2014). Worldwide epidemiology of atrial fibrillation: a Global Burden of Disease 2010 Study *Circulation*, *129*, 837–847. <https://doi.org/10.1161/CIRCULATIONAHA.113.005119>
- Cohen, J. I. (2000). Epstein-Barr virus infection *The New England Journal of Medicine*, *343*, 481–492. <https://doi.org/10.1056/NEJM200008173430707>
- Conway, D. S. g., Buggins, P., Hughes, E., & Lip, G. Y. H. (2004). Prognostic significance of raised plasma levels of interleukin-6 and C-reactive protein in atrial fibrillation *American Heart Journal*, *148*, 462–466. <https://doi.org/10.1016/j.ahj.2004.01.026>
- Dernellis, J., & Panaretou, M. (2004). Relationship between C-reactive protein concentrations during glucocorticoid therapy and recurrent atrial fibrillation *European Heart Journal*, *25*, 1100–1107. <https://doi.org/10.1016/j.ehj.2004.04.025>
- Engelmann, M. D. M., & Svendsen, J. H. (2005). Inflammation in the genesis and perpetuation of atrial fibrillation *European Heart Journal*, *26*, 2083–2092. <https://doi.org/10.1093/eurheartj/ehi350>
- EPSTEIN, M. A., ACHONG, B. G., & BARR, Y. M. (1964). Virus particles in cultured lymphoblasts from Burkitt's lymphoma *Lancet (London, England)*, *1*(7335), 702–703.
- Fox, C. S., Parise, H., D'Agostino, R. B., Lloyd-Jones, D. M., Vasan, R. S., Wang, T. J., . . . Benjamin, E. J. (2004). Parental atrial fibrillation as a risk factor for atrial fibrillation in offspring *JAMA*, *291*, 2851–2855. <https://doi.org/10.1001/jama.291.23.2851>
- Furuya, A., Uozaki, M., Yamasaki, H., Arakawa, T., Arita, M., & Koyama, A. H. (2008). Antiviral effects of ascorbic and dehydroascorbic acids in vitro *International Journal of Molecular Medicine*, *22*(4), 541–545.
- Gill, H., Hwang, Y.-Y., Chan, T. S. Y., Pang, A. W. K., Leung, A. Y. H., Tse, E., & Kwong, Y.-L. (2014). Valganciclovir suppressed Epstein Barr virus reactivation during immunosuppression with alemtuzumab *Journal of Clinical Virology : the Official Publication of the Pan American Society for Clinical Virology*, *59*, 255–258. <https://doi.org/10.1016/j.jcv.2014.01.005>
- Glotzer, T. V., & Ziegler, P. D. (2013). Silent atrial fibrillation as a stroke risk factor and anticoagulation indication *The Canadian Journal of Cardiology*, *29*, S14-23. <https://doi.org/10.1016/j.cjca.2013.03.023>
- Hanley, J. A. (1983). If Nothing Goes Wrong, Is Everything All Right? *JAMA*, *249*, 1743. <https://doi.org/10.1001/jama.1983.03330370053031>
- Hanley, J. A. (1983). If Nothing Goes Wrong, Is Everything All Right? *JAMA*, *249*, 1743. <https://doi.org/10.1001/jama.1983.03330370053031>
- Henningsen, K. M. A., Therkelsen, S. K., Bruunsgaard, H., Krabbe, K. S., Pedersen, B. K., & Svendsen, J. H. (2009). Prognostic impact of hs-CRP and IL-6 in patients with persistent atrial fibrillation treated with electrical cardioversion *Scandinavian Journal of Clinical and Laboratory Investigation*, *69*, 425–432. <https://doi.org/10.1080/00365510802676848>
- Hirano, T., Murakami, M., Fukada, T., Nishida, K., Yamasaki, S., & Suzuki, T. (2008). Roles of zinc and zinc signaling in immunity: zinc as an intracellular signaling molecule *Advances in Immunology*, *97*, 149–176. [https://doi.org/10.1016/S0065-2776\(08\)00003-5](https://doi.org/10.1016/S0065-2776(08)00003-5)
- Hung, Y.-M., Lin, L., Chen, C.-M., Chiou, J.-Y., Wang, Y.-H., Wang, P. Y.-P., & Wei, J. C.-C. (2017). The effect of anti-rheumatic medications for coronary artery diseases risk in patients with rheumatoid arthritis might be changed over time: A nationwide population-based cohort study *PloS One*, *12*, e0179081. <https://doi.org/10.1371/journal.pone.0179081>
- Issac, T. T., Dokainish, H., & Lakkis, N. M. (2007). Role of inflammation in initiation and perpetuation of atrial fibrillation: a systematic review of the published data *Journal of the American College of Cardiology*, *50*, 2021–2028. <https://doi.org/10.1016/j.jacc.2007.06.054>
- Isserlis, L. (1918). On the Value of a Mean as Calculated from a Sample *Journal of the Royal Statistical Society*, *81*, 75. <https://doi.org/10.2307/2340569>
- Jovanovic, B. D., & Levy, P. S. (1997). A Look at the Rule of Three *The American Statistician*, *51*, 137–139. <https://doi.org/10.1080/00031305.1997.10473947>
- Kawabe, A., Nakano, K., Miyata, H., Shibuya, R., Matsuyama, A., Ogoshi, T., & Tanaka, Y. (2018). Fatal Chronic Active Epstein-Barr Virus Infection in a Rheumatoid Arthritis Patient Treated with Abatacept

- Internal Medicine (Tokyo, Japan)*. Advance online publication. <https://doi.org/10.2169/internalmedicine.1280-18>
- Kim, M. H., Johnston, S. S., Chu, B.-C., Dalal, M. R., & Schulman, K. L. (2011). Estimation of total incremental health care costs in patients with atrial fibrillation in the United States *Circulation. Cardiovascular Quality and Outcomes*, 4, 313–320. <https://doi.org/10.1161/CIRCOUTCOMES.110.958165>
- Kim, S. C., Liu, J., & Solomon, D. H. (2014). The risk of atrial fibrillation in patients with rheumatoid arthritis *Annals of the Rheumatic Diseases*, 73, 1091–1095. <https://doi.org/10.1136/annrhumdis-2013-203343>
- Kirchhof, P., Breithardt, G., Aliot, E., Al Khatib, S., Apostolakis, S., Auricchio, A., . . . Camm, A. J. (2013). Personalized management of atrial fibrillation: Proceedings from the fourth Atrial Fibrillation competence NETwork/European Heart Rhythm Association consensus conference *Europace : European Pacing, Arrhythmias, and Cardiac Electrophysiology : Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*, 15, 1540–1556. <https://doi.org/10.1093/europace/eut232>
- Kirchhof, P., Lip, G. Y. H., van Gelder, I. C., Bax, J., Hylek, E., Kaab, S., . . . Camm, A. J. (2012). Comprehensive risk reduction in patients with atrial fibrillation: emerging diagnostic and therapeutic options--a report from the 3rd Atrial Fibrillation Competence NETwork/European Heart Rhythm Association consensus conference *Europace : European Pacing, Arrhythmias, and Cardiac Electrophysiology : Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*, 14, 8–27. <https://doi.org/10.1093/europace/eur241>
- Klutts, J. S., Ford, B. A., Perez, N. R., & Gronowski, A. M. (2009). Evidence-based approach for interpretation of Epstein-Barr virus serological patterns *Journal of Clinical Microbiology*, 47, 3204–3210. <https://doi.org/10.1128/JCM.00164-09>
- Kocher, M. S., & Zurakowski, D. (2004). Clinical epidemiology and biostatistics: a primer for orthopaedic surgeons *The Journal of Bone and Joint Surgery. American Volume*, 86-A(3), 607–620.
- Kolmogoroff, A. (1933). *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Berlin, Heidelberg: Springer Berlin Heidelberg.
- Korantzopoulos, P., Kolettis, T., Siogas, K., & Goudevenos, J. (2003). Atrial fibrillation and electrical remodeling: the potential role of inflammation and oxidative stress *Medical Science Monitor : International Medical Journal of Experimental and Clinical Research*, 9(9), RA225-9.
- Krumholz, H. M., Ross, J. S., Presler, A. H., & Egilman, D. S. (2007). What have we learnt from Vioxx? *BMJ (Clinical Research Ed.)*, 334, 120–123. <https://doi.org/10.1136/bmj.39024.487720.68>
- LANCASTER, H. O., & Seneta, E. (2005). Chi-Square Distribution. In P. Armitage & T. Colton (Eds.), *Encyclopedia of biostatistics* (2nd ed.). Chichester, West Sussex, England, Hoboken, NJ: John Wiley. <https://doi.org/10.1002/0470011815.b2a15018>
- Leemis, L. M., & Trivedi, K. S. (1996). A Comparison of Approximate Interval Estimators for the Bernoulli Parameter *The American Statistician*, 50, 63. <https://doi.org/10.2307/2685046>
- Lerner, A. M., Beqaj, S. H., Deeter, R. G., & Fitzgerald, J. T. (2007). Valacyclovir treatment in Epstein-Barr virus subset chronic fatigue syndrome: thirty-six months follow-up *In Vivo (Athens, Greece)*, 21(5), 707–713.
- Lindhardsen, J., Ahlehoff, O., Gislason, G. H., Madsen, O. R., Olesen, J. B., Svendsen, J. H., . . . Hansen, P. R. (2012). Risk of atrial fibrillation and stroke in rheumatoid arthritis: Danish nationwide cohort study *BMJ (Clinical Research Ed.)*, 344, e1257.
- Liu, T., Li, G., Li, L., & Korantzopoulos, P. (2007). Association between C-reactive protein and recurrence of atrial fibrillation after successful electrical cardioversion: a meta-analysis *Journal of the American College of Cardiology*, 49, 1642–1648. <https://doi.org/10.1016/j.jacc.2006.12.042>
- Louis, T. A. (1981). Confidence Intervals for a Binomial Parameter after Observing No Successes *The American Statistician*, 35, 154. <https://doi.org/10.1080/00031305.1981.10479337>
- Mant, J., Hobbs, F. R., Fletcher, K., Roalfe, A., Fitzmaurice, D., Lip, G. Y. H., & Murray, E. (2007). Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial *The*

- Lancet*, 370, 493–503. [https://doi.org/10.1016/S0140-6736\(07\)61233-1](https://doi.org/10.1016/S0140-6736(07)61233-1)
- Mikirova, N., Casciari, J., Riordan, N., & Hunninghake, R. (2013). Clinical experience with intravenous administration of ascorbic acid: achievable levels in blood for different states of inflammation and disease in cancer patients *Journal of Translational Medicine*, 11, 191. <https://doi.org/10.1186/1479-5876-11-191>
- Mikirova, N., & Hunninghake, R. (2014). Effect of high dose vitamin C on Epstein-Barr viral infection *Medical Science Monitor : International Medical Journal of Experimental and Clinical Research*, 20, 725–732. <https://doi.org/10.12659/MSM.890423>
- Mocchegiani, E., & Muzzioli, M. (2000). Therapeutic application of zinc in human immunodeficiency virus against opportunistic infections *The Journal of Nutrition*, 130, 1424S–31S. <https://doi.org/10.1093/jn/130.5.1424S>
- Nystad, T. W., & Myrnel, H. (2007). Prevalence of primary versus reactivated Epstein-Barr virus infection in patients with VCA IgG-, VCA IgM- and EBNA-1-antibodies and suspected infectious mononucleosis *Journal of Clinical Virology : the Official Publication of the Pan American Society for Clinical Virology*, 38, 292–297. <https://doi.org/10.1016/j.jcv.2007.01.006>
- Odumade, O. A., Hogquist, K. A., & Balfour, H. H. (2011). Progress and problems in understanding and managing primary Epstein-Barr virus infections *Clinical Microbiology Reviews*, 24, 193–209. <https://doi.org/10.1128/CMR.00044-10>
- Pearson, K. (1900). X. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 50, 157–175. <https://doi.org/10.1080/14786440009463897>
- Psaty, B. M., Manolio, T. A., Kuller, L. H., Kronmal, R. A., Cushman, M., Fried, L. P., . . . Rautaharju, P. M. (1997). Incidence of and risk factors for atrial fibrillation in older adults *Circulation*, 96(7), 2455–2461.
- Riordan, H. D., Hunninghake, R. B., Riordan, N. H., Jackson, J. J., Meng, X., Taylor, P., . . . Rivera, A. (2003). Intravenous ascorbic acid: protocol for its application and use *Puerto Rico Health Sciences Journal*, 22(3), 287–290.
- Roxby, A. C., Atkinson, C., Asbjörnsdóttir, K., Farquhar, C., Kiarie, J. N., Drake, A. L., . . . Slyker, J. A. (2014). Maternal valacyclovir and infant cytomegalovirus acquisition: a randomized controlled trial among HIV-infected women *PloS One*, 9, e87855. <https://doi.org/10.1371/journal.pone.0087855>
- Ruiz Espejo, M. (2007). Review of Causality: New Statistical Methods, 2nd edn (by Ilija Barukčić; Books on Demand, Norderstedt DE, 2006): 34:1013-1014 *Journal of Applied Statistics*, 34, 1013–1014. <https://doi.org/10.1080/02664760701590707>
- Rumke, C. L. (1975). Implications of the Statement: No Side Effects Were Observed *The New England Journal of Medicine*, 292, 372–373. <https://doi.org/10.1056/NEJM197502132920723>
- Rumke, C. L. (1975). Implications of the Statement: No Side Effects Were Observed *The New England Journal of Medicine*, 292, 372–373. <https://doi.org/10.1056/NEJM197502132920723>
- Savelieva, I., & Camm, A. J. (2000). Clinical relevance of silent atrial fibrillation: prevalence, prognosis, quality of life, and management *Journal of Interventional Cardiac Electrophysiology : an International Journal of Arrhythmias and Pacing*, 4(2), 369–382.
- Shankar, A. H., & Prasad, A. S. (1998). Zinc and immune function: the biological basis of altered resistance to infection *The American Journal of Clinical Nutrition*, 68, 447S–463S. <https://doi.org/10.1093/ajcn/68.2.447S>
- Sibbald, B. (2004). Rofecoxib (Vioxx) voluntarily withdrawn from market *CMAJ : Canadian Medical Association Journal = Journal De L'Association Medicale Canadienne*, 171, 1027–1028. <https://doi.org/10.1503/cmaj.1041606>
- Şimşek, B., Altay, S., Özbilgin, N., & Onat, A. (2018). Autoimmune activation as a determinant of atrial fibrillation among Turks: A prospective evaluation *Medicine*, 97, e11779. <https://doi.org/10.1097/MD.00000000000011779>
- Slomiany, B. L., & Slomiany, A. (2017). Role of LPS-elicited signaling in triggering gastric mucosal inflammatory responses to *H. pylori*: modulatory effect of ghrelin *Inflammopharmacology*, 25, 415–429. <https://doi.org/10.1007/s10787-017-0360-1>

- Slomiany, B. L., & Slomiany, A. (2018). Helicobacter pylori LPS-induced gastric mucosal spleen tyrosine kinase (Syk) recruitment to TLR4 and activation occurs with the involvement of protein kinase C $\delta$  *Inflammopharmacology*, 26, 805–815. <https://doi.org/10.1007/s10787-017-0430-4>
- Stroke Risk in Atrial Fibrillation Working Group. (2007). Independent predictors of stroke in patients with atrial fibrillation: a systematic review *Neurology*, 69, 546–554. <https://doi.org/10.1212/01.wnl.0000267275.68538.8d>
- Takano, H., Nakagawa, K., Ishio, N., Daimon, M., Daimon, M., Kobayashi, Y., . . . Komuro, I. (2008). Active myocarditis in a patient with chronic active Epstein-Barr virus infection *International Journal of Cardiology*, 130, e11-3. <https://doi.org/10.1016/j.ijcard.2007.07.040>
- Thompson, M. E. (2006). Ilija Barukčić. Causality. New Statistical Methods. A Book Review. *International Statistical Institute - Short Book Review*, 26(01), 6. Retrieved from [http://isi.cbs.nl/sbr/images/V26-1\\_Apr06.pdf](http://isi.cbs.nl/sbr/images/V26-1_Apr06.pdf)
- Wang, D.-Z., Chen, W., Yang, S., Wang, J., Li, Q., Fu, Q., . . . Chen, B.-X. (2015). Helicobacter pylori infection in Chinese patients with atrial fibrillation *Clinical Interventions in Aging*, 10, 813–819. <https://doi.org/10.2147/CIA.S72724>
- Wasmer, K., Eckardt, L., & Breithardt, G. (2017). Predisposing factors for atrial fibrillation in the elderly *Journal of Geriatric Cardiology : JGC*, 14, 179–184. <https://doi.org/10.11909/j.issn.1671-5411.2017.03.010>
- Wooten, M. D., Reddy, G. V., & Johnson, R. D. (2000). Atrial fibrillation occurring in a patient taking etanercept plus methotrexate for rheumatoid arthritis *Delaware Medical Journal*, 72(12), 517–519.
- Yamane, T. (Ed.). (1964). *Statistics. An introductory analysis*: Harper International Edition.
- Yates, F. (1934). Contingency Tables Involving Small Numbers and the  $\chi^2$  Test *Supplement to the Journal of the Royal Statistical Society*, 1, 217. <https://doi.org/10.2307/2983604>