

Theorem for w^n and Fermat's last theorem

Nguyen Van Quang
Hue - Vietnam, 2018

Abstract

We give expression of w^n and the possible to apply for solving Fermat's Last theorem

Theorem 1. $w^n = (u \pm v)^n$ can be always expressed as $(u \pm v)^n = u.F^2 \pm v.G^2$ when n is odd natural number, and can be always expressed as $(u \pm v)^n = F^2 \pm u.v.G^2$ when n is even natural number.

Proof. n is odd, $n = 2m + 1$

Write : $u - v = (\sqrt{u} + \sqrt{v})(\sqrt{u} - \sqrt{v})$, then :

$$\begin{aligned} (u - v)^{2m+1} &= (\sqrt{u} + \sqrt{v})^{2m+1}(\sqrt{u} - \sqrt{v})^{2m+1} \\ &= \left[\frac{(\sqrt{u} + \sqrt{v})^{2m+1} + (\sqrt{u} - \sqrt{v})^{2m+1}}{2} \right]^2 - \left[\frac{(\sqrt{u} + \sqrt{v})^{2m+1} - (\sqrt{u} - \sqrt{v})^{2m+1}}{2} \right]^2 \\ &= u.F^2 - v.G^2 \end{aligned}$$

Write: $u + v = (\sqrt{u} + i\sqrt{v})(\sqrt{u} - i\sqrt{v})$, then :

$$\begin{aligned} (u + v)^{2m+1} &= (\sqrt{u} + i\sqrt{v})^{2m+1}(\sqrt{u} - i\sqrt{v})^{2m+1} \\ &= \left[\frac{(\sqrt{u} + i\sqrt{v})^{2m+1} + (\sqrt{u} - i\sqrt{v})^{2m+1}}{2} \right]^2 - \left[\frac{(\sqrt{u} + i\sqrt{v})^{2m+1} - (\sqrt{u} - i\sqrt{v})^{2m+1}}{2} \right]^2 \\ &= u.F^2 + v.G^2 \end{aligned}$$

n is even, $n = 2m$ then:

$$\begin{aligned} (u - v)^{2m} &= (\sqrt{u} + \sqrt{v})^{2m}(\sqrt{u} - \sqrt{v})^{2m} \\ &= \left[\frac{(\sqrt{u} + \sqrt{v})^{2m} + (\sqrt{u} - \sqrt{v})^{2m}}{2} \right]^2 - \left[\frac{(\sqrt{u} + \sqrt{v})^{2m} - (\sqrt{u} - \sqrt{v})^{2m}}{2} \right]^2 \\ &= F^2 - u.v.G^2 \end{aligned}$$

And:

$$\begin{aligned} (u + v)^{2m} &= (\sqrt{u} + i\sqrt{v})^{2m}(\sqrt{u} - i\sqrt{v})^{2m} \\ &= \left[\frac{(\sqrt{u} + i\sqrt{v})^{2m} + (\sqrt{u} - i\sqrt{v})^{2m}}{2} \right]^2 - \left[\frac{(\sqrt{u} + i\sqrt{v})^{2m} - (\sqrt{u} - i\sqrt{v})^{2m}}{2} \right]^2 \\ &= F^2 + u.v.G^2 \end{aligned}$$

Here:

i : imaginary unit $i^2 = -1$; $i^{4k} = 1$; $i^{4k+2} = -1$, $F = f(u, v)$, $G = g(u, v)$ will not contain i (Since $i^{4k+1} = i$; $i^{4k+3} = -i$ is lost).

Special cases:

$$u = u_0^2, v = v_0^2:$$

$$(u \pm v)^n = (u_0^2 \pm v_0^2)^n = u_0^2.F^2 \pm v_0^2.G^2 = (u_0F)^2 \pm (v_0G)^2 \text{ for } n \text{ is odd}$$

$$(u \pm v)^n = (u_0^2 \pm v_0^2)^n = F^2 \pm u_0^2v_0^2.G^2 = F^2 \pm (u_0v_0G)^2 \text{ for } n \text{ is even}$$

Consequently,

Theorem 2. The equation $x^2 \pm y^2 = z^n$ always has infinitive solutions in integer for any positive integer n

Note:

Above expression is the only way or not, it depends on w (even or odd), u and v (square e^2 or not square e, fe^2).

So that, be carefully when apply for specific case.

For the case w is odd, u and v are squares, $u = a^2, v = b^2$, a and b different parity, $w = a^2 - b^2$, Above expression is the only way.

However, the case below:

$$3 = 5 - 2, \text{ then } 3^3 = (5 - 2)^3 = \left[\frac{(\sqrt{5} + \sqrt{2})^3 + (\sqrt{5} - \sqrt{2})^5}{2} \right]^2 - \left[\frac{(\sqrt{5} + \sqrt{2})^3 - (\sqrt{5} - \sqrt{2})^3}{2} \right]^2$$

$$= 5.11^2 - 2.17^2$$

But, there is other way such that:

$$3^3 = (5 - 2)^3 = 5.5^2 - 2.7^2.$$

1 Applying for FLt

$$x^n + y^n = z^n \quad (1)$$

n is odd, $n = 2m + 1$

The left hand side:

$$x^{2m+1} + y^{2m+1} = (x + y)(x^{2m} - x^{2m-1}y + x^{2m-2}y^2 - \dots + y^{2m}) \quad (2)$$

we can write $x^{2m+1} + y^{2m+1} = (x + y)Q$, here $Q = x^{2m} - x^{2m-1}y + x^{2m-2}y^2 - \dots + y^{2m}$
to consider FLt, it is enough to consider n prime, $n = p$.

Assume x and y are odd, we express Q as one of two formulas below:

$$Q_p = M^2 + pN^2 \quad (3)$$

or ;

$$Q_p = M^2 - pN^2 \quad (4)$$

Here: $M = f(a, b), N = g(a, b), a + b = x, a - b = y$. M and N are coprime.

For p = 3:

$$Q_3 = a^2 + 3b^2$$

For p = 5:

$$Q_5 = (a^2 + 5b^2)^2 - 5(2b^2)^2$$

For p = 7:

$$Q_7 = a^2(a^2 + 7b^2)^2 + 7b^2(b^2 - a^2)^2$$

For p = 11:

$$Q_{11} = a^2(a^4 - 22a^2b^2 - 11b^4)^2 + 11b^2(b^4 + 2a^2b^2 - 3a^4)^2$$

...

Since $x^p + y^p = z^p$, then $Q_p = w^p$ or $Q_p = pw^p$

w is not divisible by p.

Two equations must be considered :

$$M^2 + pN^2 = w^p \text{ (or } M^2 - pN^2 = w^p) \quad (5)$$

$$M^2 + pN^2 = pw^p \text{ (or } M^2 - pN^2 = pw^p) \quad (6)$$

For (6), $M = pM_0$, it yields: $pM_0^2 + N^2 = w^p$ (or $pM_0^2 - N^2 = w^p$)

2 The algorithm

Express w^p as:

$$w^p = M'^2 + pN'^2 \quad (7)$$

or

$$w^p = M'^2 - pN'^2 \quad (8)$$

Apply theorem above, let $w = c^2 + pd^2$ or $w = c^2 - pd^2$

For $p = 3$:

$$\begin{aligned} w^3 &= (c^2 + 3d^2)^3 = (c + i\sqrt{3}d)^3(c - i\sqrt{3}d)^3 \\ &= \left[\frac{(c + i\sqrt{3}d)^3 + (c - i\sqrt{3}d)^3}{2} \right]^2 - \left[\frac{(c + i\sqrt{3}d)^3 - (c - i\sqrt{3}d)^3}{2} \right]^2 \\ &= c^2(c^2 - 9d^2)^2 + 3 \cdot 3^2 d^2(c^2 - d^2)^2 \\ &(a = c(c^2 - 9d^2) \text{ and } b = 3d(c^2 - d^2); \text{ Euler' proof-1770 year}) \end{aligned}$$

For $p = 5$:

$$\begin{aligned} w^5 &= (c^2 - 5d^2)^3 = (c + \sqrt{5}d)^5(c - \sqrt{5}d)^5 \\ &= \left[\frac{(c + \sqrt{5}d)^5 + (c - \sqrt{5}d)^5}{2} \right]^2 - \left[\frac{(c + \sqrt{5}d)^5 - (c - \sqrt{5}d)^5}{2} \right]^2 \\ &= c^2(c^4 + 50c^2d^2 + 125d^4)^2 - 5 \cdot 5^2 d^2(c^4 + 10c^2d^2 + 5d^4)^2 \\ &(a^2 + 5b^2 = c(c^4 + 50c^2d^2 + 125d^4) \text{ and } 2b^2 = 5d(c^4 + 10c^2d^2 + 5d^4); \text{ Dirichlet's proof-1825 year}) \\ &\text{and} \end{aligned}$$

For $p = 7$:

$$\begin{aligned} w^7 &= (c^2 + 7d^2)^7 = (c + i\sqrt{7}d)^7(c - i\sqrt{7}d)^7 \\ &= \left[\frac{(c + i\sqrt{7}d)^7 + (c - i\sqrt{7}d)^7}{2} \right]^2 - \left[\frac{(c + i\sqrt{7}d)^7 - (c - i\sqrt{7}d)^7}{2} \right]^2 \\ &= c^2(c^6 - 3 \cdot 7^2 c^4 d^2 + 5 \cdot 7^3 c^2 d^4 - 7^4 d^6)^2 + 7 \cdot 7^2 d^2(c^6 - 5 \cdot 7 c^4 d^2 + 3 \cdot 7^2 c^2 d^4 - 7^2 d^6)^2 \\ &\dots \end{aligned}$$

If it is the only way for the specific case, then there is only one choice, and not more. We obtain the two equations below:

$$M = M'$$

$$N = N'$$

If they have no solution in integer, FLt is true for that case, if they have a solution in integer, then continue consider if it satisfy to condition $2a = w^p$ ($w'w = z$) when $p \nmid z$ (or $2a = p^{p-1}w^p$, when $p \mid z$) or not.

If the only way is not shown, then the proof of FLt by algorithm above is not completed (flawed)!!

3 About Fermat's margin-notes

Around 1637, Fermat wrote his Last Theorem in the margin of his copy of the Arithmetica next to Diophantu's sum - of- squares problem:

It is impossible to separate a cube in two cubes, or a fourth power into two fourth powers, or in general, any power higher than second,into two like powers. I have discovered a truly marvelous proof of this, which this margin is too narrow to contain.

It is not known whether Fermat had actually a valid proof for all exponents n.

I am Quang, Math independent researcher. In the letter was sent to The Annal of Math in 2015 year, I supposed that the short proof of Flt will appear , and Fermat could have a proof of FLt as he wrote (margin - notes). Indeed the short proof of FLt was found.

In my opinion, Fermat is famous enough , if he had a proof of Flt, publishing a proof of Flt or not, no problem for him, but for us . The short proof could be kept in mind without writing for memory.

4 Acknowledgement

I published a proof of the four color theorem in 2016 year, I think that professional and none -professional mathematicians could understand and verify it . I am very happy if my proof of the four color theorem by induction is correct, was verified and recognized before I publish the short proof of Flt.Thank you!

□

References

Quang N V, A proof of the four color theorem by induction Vixra: 1601.0247 (CO)

APPENDIS

About proof of the FLt for n = 5

Dirichlet have proved FLt for n = 5 by infinitive descent, his proof is correct if $w = (c^2 - 5d^2)^5 = c^2(c^4 + 50c^2d^2 + 125d^4)^2 - 5 \cdot 5^2d^2(c^4 + 10c^2d^2 + 5d^4)^2$ is the only way for expression $w = M^2 - 5N^2$.* If the condition* is true was shown! I give a very simple poof of FLt for n =5 without using infinitive descent below:

Since $a^2 + 5b^2 = c(c^4 + 50c^2d^2 + 125d^4)$, and $2b^2 = 5d(c^4 + 10c^2d^2 + 5d^4)$, then $5 \mid b$, that means $x = a + b$ and $y = a - b$ is not divisible by 5. In other hand, if $x^5 + y^5 = z^5$, then one of x,y and z must divisible by y, it yields $5 \mid z$

It gives:

$5 \mid a^2 + 5b^2$, hence $5 \mid a$, it yields $5 \mid x$; $5 \mid y$ and $5 \mid z$,that means x,y and z have a common factor, a contradiction!

Email:

nguyenvquang67@gmail.com

quangnhu67@yahoo.com.vn