

Variable Polytropic gas cosmology

M. Salti*,¹ E. E. Kangal†,² and O. Aydogdu‡¹

¹*Department of Physics, Faculty of Arts and Science,*

Mersin University, Mersin, TR 33343, Turkey

²*Computer Technology and Information Systems,*

School of Applied Technology and Management of Erdemli,

Mersin University, Mersin, TR 33740, Turkey

Abstract

We mainly study a cosmological scenario defining by the variable Polytropic gas (VPG) unified energy density proposal. To reach this aim, we start with reconstructing a generalized form of the original Polyrtpropic gas (OPG) definition. Later, we fit the auxiliary parameters given in the model and discuss essential cosmological features of the VPG proposal. Besides, we compare the VPG with the OPG by focusing on recent observational dataset given in literature including Planck 2018 results. We see that the VPG model yields better results than the OPG description and it fits very well with the recent experimental data. Moreover, we discuss some thermodynamical features of the VPG and conclude that the model describes a thermodynamically stable system.

PACS numbers: 04.50.+h, 95.35.+d, 98.80.-k

Keywords: Cosmology, dark energy, variable, Polytropic gas, Chaplygin gas.

* msalti@mersin.edu.tr (Corresponding author)

† evrimersin@gmail.com

‡ oktaydogdu@mersin.edu.tr

I. INTRODUCTION

There are plenty of astrophysical data that signal us to a universe entered in a speedy expansion phase[1–10]. In order to understand the accelerated expansion phase of our universe, a mysterious type of energy called dark energy is to be needed. The dark energy dominates approximately 68.3% of the space-time tissue. Assumption of a dominant component which resembles familiar forms of matter or energy has not been justified yet, because the exotic dark content cannot be detected directly. Diverse proposals have been introduced in literature to express the speedy expansion behavior of our universe. The earliest and simplest description for the mysterious type of energy mentioned above is the cosmological constant[11], but it includes both the fine-tuning and the cosmic coincidence problems. To remove these issues, different ideas to identify the dark energy were also introduced such as braneworld models[12, 13], scalar fields[14–18], modified gravity theories[19, 20], assuming extra dimensions[21–23] and so on. For a convenient brief about theoretical dark energy models, one can read Ref.[24] and references therein.

An energy-momentum tensor must be defined on the right-hand side of a gravitational field equation when the compactification is due to matter fields. In order to add dark content's contributions to the field equation, various new descriptions, like ghost dark energy[25–28], holographic energy density[29, 30], Hobbit model[31], new agegraphic dark energy[32], Chaplygin gas[33, 34], Polytropic gas[35, 36], etc., have been introduced in literature. Among these formulations, Hobbit[31], Chaplygin gas[33, 34] and Polytropic gas[35, 36] models have a significant future which leads interesting cosmological conclusions. These three proposals unify different fluids of the standard cosmological model and can express both dark matter and dark energy with a single fluid (consequently automatically removing the coincidence issue)

Constraining the auxiliary parameters given a theoretical model is one of the significant tasks in contemporary theoretical cosmology. The most often considered technique is analyzing the luminosity distance measurements for a specific family of objects[37–40]. In some recent research papers, a new way including observational values of the Hubble parameter (OHV) has been considered to check some cosmological tests[41–46].

The layout of the paper is as follows. In the next section, we give some preliminary relations and definitions. Next, in the third section, we introduce the VPG model. In

the fourth section of the paper, we fit the free parameters of the model by considering the recent astrophysical observations. In the fifth section, we test the VPG model and compare it with the OPG proposal. In the sixth section, we investigate thermodynamical features and discuss the stability of the VPG model. The final section is devoted to the closing remarks. All numerical calculations and analyzes are performed by using MATHEMATICA software[47].

II. PRELIMINARIES: COSMOLOGICAL SCENARIO

Here, as a first step, we assume the Friedmann-Robertson-Walker (FRW) universe is filled with the VPG and the baryonic matter and its line-element is written as

$$ds^2 = dt^2 - a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right], \quad (1)$$

where the time-dependent function $a(t)$ indicates the cosmic scale factor while k implies the curvature parameter for the flat ($k = 0$), closed ($k = -1$) and open ($k = +1$) universe types. We also suppose that the FRW universe is filled with perfect fluid which is defined by

$$T_{\mu\nu} = (\rho + p)u_\mu u_\nu - g_{\mu\nu}p. \quad (2)$$

Here, $\rho = \rho_{vpg} + \rho_{bm}$ and $p = p_{vpg} + p_{bm}$ are total energy density and pressure, respectively. The subscripts vpg and bm denote the VPG and the baryonic matter, respectively. It is important to mention here that the VPG is a unification of the dark matter and dark energy. Thus, we can write $\rho_{vpg} = \rho_{dm} + \rho_{de}$ and $p_{vpg} = p_{de}$. Note that, now, the subscripts dm and de mean the dark matter and the dark energy, respectively. Moreover, u_μ is the four-velocity vector and we have $u^\mu u_\mu = 1$.

The Einstein field equations can be written in the following form

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G [(\rho + p)u_\mu u_\nu - g_{\mu\nu}p] \quad (3)$$

where $R_{\mu\nu}$, $g_{\mu\nu}$, R and G show the Ricci tensor, metric tensor, Ricci scalar and the gravitational constant, respectively.

The recent observational data obtained by SNe-Ia[1], WMAP[4–6], SDSS[7], X-ray[48] and Planck-results[8–10] have strongly suggested that the geometry of the universe is spatially flat. From this point of view, we take $k = 0$ in the further calculations. Hence, making

use of equations (1) and (3), it follows that

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho, \quad (4)$$

where H is the Hubble expansion parameter.

Next, the continuity equation, i.e. $T_{;\nu}^{\mu\nu} = 0$, yields

$$\dot{\rho}_{bm} + 3H\rho_{bm} = 0, \quad (5)$$

$$\dot{\rho}_{vpg} + 3H(\rho_{vpg} + p_{vpg}) = 0. \quad (6)$$

Additionally, we can also write

$$\dot{\rho}_{dm} + 3H\rho_{dm} = 0, \quad (7)$$

$$\dot{\rho}_{de} + 3H(\rho_{de} + p_{de}) = 0, \quad (8)$$

with the fact that equation-of-state (EoS) parameters of the baryonic matter and the dark matter are taken as $\omega_{bm} = \omega_{dm} = 0$. From equations (5) and (7), one can easily obtain that

$$\rho_{bm} = \rho_0^b a^{-3}, \quad (9)$$

$$\rho_{dm} = \rho_0^m a^{-3}, \quad (10)$$

where ρ_0^b and ρ_0^m show present values of the baryonic matter and the dark matter, respectively.

In addition to the above calculations, introducing dimensionless density parameters helps us to rewrite the Friedmann equation (4) in a very useful and elegant form. On this purpose, first, we define

$$\Omega_{bm} = \frac{8\pi G}{3H^2}\rho_{bm}, \quad \Omega_{vpg} = \frac{8\pi G}{3H^2}\rho_{vpg}, \quad (11)$$

$$\Omega_{dm} = \frac{8\pi G}{3H^2}\rho_{dm}, \quad \Omega_{de} = \frac{8\pi G}{3H^2}\rho_{de}. \quad (12)$$

Subsequently, in the second step, it can be concluded that

$$\sum_{i=bm, dm, de} \Omega_i \equiv 1, \quad (13)$$

where $\Omega_i \equiv (\Omega_{bm}, \Omega_{dm}, \Omega_{de})$ and $\Omega_{vpg} = \Omega_{dm} + \Omega_{de}$.

It is known that we can fix the free parameters given in the model according to the current cosmological measurements by using the red shift parameter in corresponding calculations. The red shift parameter z is related to the cosmic scale factor $a(t)$ by $z = \frac{1}{a} - 1$. So, we get

$$\rho_{bm} = \rho_0^b (1+z)^3, \quad (14)$$

$$\rho_{dm} = \rho_0^m (1+z)^3. \quad (15)$$

III. THE VPG PROPOSAL

The OPG model is described by the following EoS[35, 36]

$$p_{pg} = \beta \rho_{pg}^{1+\frac{1}{\xi}}, \quad (16)$$

where both β and the polytropic index ξ denote real constants. Here, we define another form of the PG model in order to check whether we can get better conclusions. So, we assume

$$p_{vpg} = \beta a^{-n} \rho_{vpg}^{1+\frac{1}{\xi}}, \quad (17)$$

where n is a constant, so, it can be seen that we have three free parameters. It is significant to mention here that this new proposal can be reduced some of other unified dark matter-energy models. In the case $n = 0$, the above relation recovers the OPG model[35, 36]. Assuming $n = 0$, $\beta = -\kappa$ and $\xi = -\frac{1}{2}$ reduces the above expression into the form of the original Chaplygin (OCG)[33, 34] which is given by $p_{ocg} = -\frac{\kappa}{\rho_{ocg}}$. Next, taking $n = 0$, $\beta = -\kappa$ and $\xi = -\frac{1}{1+\alpha}$ transforms the expression of VPG into the form of the generalized Chaplygin (GCG)[50–52] model, i.e. $p_{gcg} = -\frac{\kappa}{\rho_{gcg}^\alpha}$. On the other hand, the case including $n \neq 0$, $\xi = -\frac{1}{1+\alpha}$ and $\beta = -\kappa$ yields the variable generalized Chaplygin gas (VGCG)[53, 54] model, i.e. $p_{vgcg} = -\frac{\kappa a^{-n}}{\rho_{vgcg}^\alpha}$.

Now, we can focus on the equation (6) to find an exact expression for the corresponding energy density. We can rewrite the equation (6) in a more convenient form as given below

$$d(\rho_{vpg} a^3) + p_{vpg} d(a^3) = 0. \quad (18)$$

Consequently, using the above equation, we can derive the energy density of the VPG model:

$$\rho_{vpg} = \rho_0^p \left[\Delta(1+z)^n + (1-\Delta)(1+z)^{-\frac{3}{\xi}} \right]^{-\xi}, \quad (19)$$

where ρ_0^p indicates the present value of VPG energy density and Δ is defined as

$$\Delta = -\frac{\beta}{1 + \frac{n\xi}{3}} \sqrt[\xi]{\rho_0^p}. \quad (20)$$

For an expanding spacetime model, it must be $n > 0$ and $\xi < 0$. Otherwise, $a \rightarrow \infty$ yields $\rho_{vpg} \rightarrow \infty$ which cannot define an expanding spacetime. Now, in order to investigate the evolution propensity of the dark matter and the dark energy and study cosmological features

of the dark energy, one can consider the decomposition of the VPG fluid: $\rho_{vpg} = \rho_{dm} + \rho_{de}$ and $p_{vpg} = p_{de}$. Then, using equations (14), (15) and (19), we get

$$\begin{aligned}\rho_{de} &= \rho_{vpg} - \rho_{dm} \\ &= \rho_0^p \left[\Delta(1+z)^n + (1-\Delta)(1+z)^{-\frac{3}{\xi}} \right]^{-\xi} - \rho_0^m (1+z)^3.\end{aligned}\quad (21)$$

On the other hand, making use of equations (4), (11), (12), (14), (15) and (19), we obtain the following result

$$H^2 = H_0^2 \left\{ \Omega_{vpg}^0 \left[\Delta(1+z)^n + (1-\Delta)(1+z)^{-\frac{3}{\xi}} \right]^{-\xi} + \Omega_{bm}^0 (1+z)^3 \right\}, \quad (22)$$

where H_0 denotes the present value of the Hubble parameter and

$$\Omega_{vpg}^0 + \Omega_{bm}^0 = 1, \quad (23)$$

with

$$\Omega_{vpg}^0 = \frac{8\pi G}{3H_0^2} \rho_{vpg}, \quad \Omega_{bm}^0 = \frac{8\pi G}{3H_0^2} \rho_{bm}. \quad (24)$$

IV. FITTING THE MODEL PARAMETERS

In this section, on the basis of the result (19), we consider the recent observational datasets such as the Supernova type Ia (SN Ia) sample, the OHV and the baryon acoustic oscillations (BAO) data to constrain the VPG proposal and investigate the evolutionary behavior of our universe.

A. Analysis methods

1. SN Ia data

Here, we consider the observational SN Ia dataset which consists information about the luminosity distance. The Hubble-free definition of the luminosity parameter is written as

$$d_L = (1+z) \int_0^z \frac{H_0 dz'}{H(z')}. \quad (25)$$

Additionally, for the SN Ia dataset, χ^2 function is given by[55]

$$\chi_{SN}^2 = \sum_i^{580} \frac{[\mu_{obs}(z_i) - \mu_{theo}(z_i)]^2}{\sigma_i^2}, \quad (26)$$

where the theoretical distance modulus is written as

$$\mu_{theo} = 5 \log_{10} d_L(z_i) + \mu_0, \quad (27)$$

with $\mu_0 = 42.38 - 5 \log_{10} h$. Note that $\mu_{obs}(z_i)$, σ_i and $h = H_0/100/[km\,sec^{-1}Mpc^{-1}]$ show the observed distance modulus, the uncertainty in the distance modulus and the then-favored dimensionless Hubble parameter, respectively. For the minimization of χ_{SN}^2 with respect to μ_0 for 580 data points of the SN Ia measurements[56], we get[57]

$$\tilde{\chi}_{SN}^2 = P - \frac{Q^2}{R}, \quad (28)$$

where

$$P = \sum_i^{580} \frac{[\mu_{obs}(z_i) - \mu_{theo}(z_i; \mu_0 = 0)]^2}{\sigma_i^2}, \quad (29)$$

$$Q = \sum_i^{580} \frac{[\mu_{obs}(z_i) - \mu_{theo}(z_i; \mu_0 = 0)]}{\sigma_i^2}, \quad (30)$$

$$R = \sum_i^{580} \frac{1}{\sigma_i^2}. \quad (31)$$

2. OHV

Considering some observed[58–71] $H(z)$ values given in TABLE I, we can investigate the validity of the constraints on the auxiliary parameters given in the VPG definition. One can minimize χ_{OHD}^2 which can be defined as given below

$$\chi_{OHV}^2 = \sum_i^{27} \frac{[H_{obs}(z_i) - H_{theo}(z_i)]^2}{\sigma_i^2} \quad (32)$$

where H_{obs} and H_{theo} describe observational and theoretical values of the cosmic Hubble parameter, respectively.

3. BAO data

It is known that our universe includes a fraction of baryon, so the acoustic oscillations in the relativistic plasma may be stamped on the late-time power spectrum of the non-relativistic matter[54, 72]. We can minimize the χ_{BAO}^2 which is written as

$$\chi_{BAO}^2 = \frac{[\Gamma(\theta) - \Gamma_{obs}]^2}{\sigma_A^2}, \quad (33)$$

TABLE I: The recent observable $H(z)$ dataset.

z	H_{obs}	σ	Ref.	z	H_{obs}	σ	Ref.
0.0708	69.00	∓ 19.68	[59]	0.5700	92.40	∓ 4.500	[67]
0.1200	68.60	∓ 26.20	[59]	0.5930	104.0	∓ 13.00	[61]
0.1700	83.00	∓ 8.000	[60]	0.6800	92.00	∓ 8.000	[61]
0.1990	75.00	∓ 5.000	[61]	0.7300	97.30	∓ 7.000	[68]
0.2400	79.69	∓ 2.650	[62]	0.7810	105.0	∓ 12.00	[61]
0.2800	88.80	∓ 36.60	[59]	0.8750	125.0	∓ 17.50	[61]
0.3500	84.40	∓ 7.000	[63]	0.9000	117.0	∓ 23.00	[60]
0.3802	83.00	∓ 13.50	[61]	1.3000	168.0	∓ 17.00	[69]
0.4000	95.00	∓ 17.00	[60]	1.4300	177.0	∓ 18.00	[69]
0.4247	87.10	∓ 11.20	[64]	1.5300	140.0	∓ 14.00	[60]
0.4300	86.45	∓ 3.680	[62]	1.7500	202.0	∓ 40.00	[70]
0.4497	92.80	∓ 12.90	[64]	1.9650	186.5	∓ 50.40	[61]
0.4783	80.90	∓ 9.000	[64]	2.3400	222.0	∓ 7.000	[71]
0.4800	97.00	∓ 62.00	[65, 66]				

where

$$\Gamma(\theta) = \frac{\sqrt{\Omega_{dm}^0}}{\sqrt[3]{E(z_{BAO})}} \left\{ \frac{1}{z_{BAO}} \int_0^z \frac{dz'}{E(z'; \theta)} \right\}^{\frac{2}{3}}, \quad (34)$$

where $E(z) = \frac{H(z)}{H_0}$. Remember that Ω_{dm}^0 is not explicitly included in the VPG proposal. According to the Planck results[9, 10], here we take $H_0 = 67.4 \pm 0.5 kmsec^{-1} Mpc^{-1}$ and $\Omega_{dm}^0 = 0.278$. Next, in the 1σ confidence region, it was measured that $\Gamma_{obs} = 0.469(\frac{0.98}{n_s})^{0.35} \pm 0.017$ from the SDSS observations at $z_{BAO} = 0.35$. Here, $n_s = 0.96$ indicates the scalar spectral index[73].

B. Constraints on the free parameters

We focus on a combined constraint on the VPG model. So, we have $\chi^2 = \tilde{\chi}_{SN}^2 + \chi_{OHV}^2 + \chi_{BAO}^2$ where $\tilde{\chi}_{SN}^2$, χ_{OHV}^2 and χ_{BAO}^2 are defined by equations (28), (32) and (33). From this point of view, making us of the observational datasets, one can calculate the best fit values

of auxiliary parameters:

$$(\beta, n, \xi) = (-0.031, -1.869, 2.990) \quad (35)$$

with $\chi_{min}^2 = 324.691$. In FIGs. 1 and 2, we depict confidence contours on $\beta - n$, $\xi - n$ and $\beta - \xi$ parameter spaces for SN Ia+OHV+BAO datasets.




FIG. 1: Plots of 1σ (pink), 2σ (light pink) and 3σ (white) confidence contours on $\beta - n$ and $\xi - n$ parameter spaces for SN Ia+OHV+BAO datasets in the VPG dark energy description.




FIG. 2: Plots of 1σ (pink), 2σ (light pink) and 3σ (white) confidence contours on $\beta - \xi$ parameter space for SN Ia+OHV+BAO datasets in the VPG dark energy description.

V. TESTING THE MODEL

FIGs. 3 and 4 illustrate the evolutionary nature of the cosmic Hubble parameter according to the OPG and the VPG models in the 1σ confidence region, respectively. Note that, in FIGs. 3 and 4, the circles show the recent observable values.




FIG. 3: $H \sim z$ relation for SN Ia+OHV+BAO datasets in the OPG dark energy model.




FIG. 4: $H \sim z$ relation for SN Ia+OHV+BAO datasets in the VPG proposal.

In order to check correlation between H_{obs} and H_{theo} datasets, we can calculate the corresponding correlation coefficient r which is used in statistics to measure how strong a relationship is between two different variables. Thus, we focus on the following relations:

$$S_x = \sqrt{\frac{\sum_{n=1}^{27} (H_{obs}(z_i) - \bar{H}_{obs})^2}{n-1}}, \quad (36)$$

$$S_y = \sqrt{\frac{\sum_{n=1}^{27} (H_{theo}(z_i) - \bar{H}_{theo})^2}{n-1}}, \quad (37)$$

where \bar{H}_{obs} and \bar{H}_{theo} indicate mean values of observational and theoretical $H(z)$ datasets, respectively. Making use of the above relations, one can define the correlation coefficient as given below

$$r = \frac{\sum_{n=1}^{27} (H_{obs}(z_i) - \bar{H}_{obs}) (H_{theo}(z_i) - \bar{H}_{theo})}{(n-1)S_x S_y}. \quad (38)$$

Note that the correlation coefficient should always take values between -1 and $+1$. The case $r = +1$ ($r = -1$) means the points are on a perfect straight line with positive (negative) slope. The case of zero implies no relationship at all. Absolute value of the correlation coefficient, i.e. $|r|$, indicates the relationship strength. It is known that the larger the absolute value of correlation coefficient, the stronger the linear relationship. Considering numerical values given in TABLE I, we find that $r = 0.968769$ which means there is a strong positive relationship between H_{obs} and H_{theo} .

VI. THERMODYNAMICS OF THE VPG

Here, we consider some thermodynamical relations in order to check thermodynamical features of the model. We start with the energy density relation $\rho = \frac{U}{V}$ where U indicates the internal energy and $V = \frac{4}{3}\pi r_h^3$ with the dynamical apparent horizon $r_h = \frac{1}{\sqrt{H^2+k(1+z)^2}}$ denotes the volume of the system. Remember that it should be taken $k = 0$ due to the recent astrophysical observations[1, 4–10, 48]. For the flat FRW universe, we can write $r_h = \frac{1}{H}$ yclept the dynamical Hubble horizon. In this section, we also focus on another important relation defining the entropy, i.e. $S = \frac{A}{4G}$ where $A = 4\pi r_h^2$ is the surface area. Consequently, making use of the equation (22) defining the theoretical Hubble parameter, we get

$$\begin{aligned} S &= \frac{\pi}{GH_0} \left[\Omega_{vpg}^0 \left[\Delta(1+z)^n + (1-\Delta)(1+z)^{-\frac{3}{\xi}} \right]^{-\xi} + \Omega_{bm}^0 (1+z)^3 \right]^{-2} \\ &= \frac{\pi^{\frac{1}{3}}}{G} \left(\frac{3V}{4} \right)^{\frac{2}{3}}. \end{aligned} \quad (39)$$

It is clearly seen from the above result that the entropy is an incensing function of volume which means it is also an increasing function of time. Therefore, the second thermodynamical law is valid as it is expected.

Next, we can also check the thermodynamical stability of the VPG model. In order to reach this aim, one can check the validity of $T \frac{\partial S}{\partial T} > 0$. As a matter of fact, this case leads to the relation of heat capacity which can be rewritten in terms of volume[74]:

$$C = V \frac{\partial \rho}{\partial V} \left[\frac{\partial T}{\partial V} \right]^{-1}. \quad (40)$$

Hence, for the VPG proposal, one can calculate that

$$C = \frac{6\pi V}{3^{\frac{1}{3}}(8\pi)^{\frac{2}{3}}G} \left[\frac{\sqrt{V}}{2} + \beta a^{-n} \left(\frac{1}{\xi} + \frac{1}{2} \right) \left(\frac{3}{8\pi G} \right)^{-\frac{1}{3\xi}} V^{\frac{1}{3} - \frac{2}{3\xi}} \right]^{-1}. \quad (41)$$

In FIG. 5, we plot $S \sim V$ (blue dashed line) and $C \sim V$ (red solid line) relations and conclude that the VPG model is thermodynamically stable.




FIG. 5: $S \sim V$ (blue dashed line) and $C \sim V$ (red solid line) relations for SN Ia+OHD+BAO datasets in the VPG proposal. Here, we take $z = 0.35$ and $8\pi G = 1$.

On the other hand, according to the first thermodynamical law, the temperature is described as

$$T = (\rho + p) \frac{V}{S}. \quad (42)$$

It is significant to write relations among ρ , S and T , because it leads very interesting cosmological implications. Making use of our previous computations, one can find that

$$T(\rho) = \frac{(8\pi G)^{\frac{5}{6}}}{6\pi 3^{-\frac{1}{6}}} \left(1 + \beta a^{-n} \rho^{\frac{1}{\xi}} \right) \sqrt{\rho}. \quad (43)$$

In FIG. 6, we depict the evolution temperature as a function of energy density according to the best fit values of auxiliary parameters β , n and ξ . It is seen that temperature of

the FRW universe dominated by the VPG increases by increasing energy density just as expected. Thence, we may interpret this conclusion as the third law of thermodynamics is satisfied for the VPG.




FIG. 6: Graphical analyze of $T \sim \rho$ relation with the best fit values of the free parameters given in the VPG model. Here, we assume that $z = 0.35$ and $8\pi G = 1$.

VII. CLOSING REMARKS

We introduce the VPG model as a unification of the dark matter and the dark energy and investigate its observational constraint by making use of a union case of the SN Ia sample, the OHV and the BAO data. Based on the best fit values of auxiliary parameters for the VPG proposal, it is concluded that our universe will not end up with big rip in the future. It is shown that the VPG model indicates a thermodynamically stable system and is consistent with the current astrophysical observations. We also discriminate the VPG model with the other unified energy density descriptions by making use of numerical and statistical analyzes. It is important to emphasize here that there is an obvious difference between the VPG and the OPG models. Compatibility of the VPG model with the current observations is better than the OPG proposal.

[1] S. Perlmutter, et al., Nature 391 (1998) 51; Astrophys. J. 517 (1999) 565.

- [2] A.D. Miller, et al., *Astrophys. J. Lett.* 524 (1999) L1.
- [3] N.A. Bahcall, J.P. Ostriker, S. Perlmutter and P.J. Steinhardt, *Science* 284 (1999) 1481.
- [4] C.L. Bennett, et al., *Astrophys. J. Suppl.* 148 (2003) 1.
- [5] S.L. Brile, O. Lahav, J.P. Ostriker and P.J. Steinhardt, *Science* 299 (2003) 1532.
- [6] D.N. Spergel, et al., *The Astrophysical Journal Supplement Series* 148 (2003) 175.
- [7] M. Tegmark, et al., *Phys. Rev. D* 69 (2004) 103501.
- [8] Planck Collaboration (P.A.R. Ade, et al.), *A&A* 571 (2014) A16.
- [9] Planck Collaboration (P.A.R. Ade, et al.), *A&A* 594 (2016) A13.
- [10] Planck Collaboration (N. Aghanim, et al.), e-Print: 1807.06209.
- [11] S. Weinberg, *Mod. Phys. Rev.* 61 (1989) 527.
- [12] L. Randall, R. Sundrum, *Phys. Rev. Lett.* 83 (1999) 4690.
- [13] Z.H. Zhu, J.S. Alcaniz, *Astrophys. J.* 620 (2005) 7.
- [14] T. Chiba, T. Okabe and M. Yamaguchi, *Phys. Rev. D* 62 (2000) 023511.
- [15] A. Sen, *Lect. Notes Phys.* 653 (2004) 141
- [16] B. Feng, X.L. Wang and X.M. Zhang, *Phys. Lett. B* 607 (2005) 35.
- [17] R.R. Caldwell, *Phys. Lett. B* 545 (2002) 204.
- [18] E.J. Copeland, M. Sami and S. Tsujikawa, *Int. J. Mod. Phys. D* 15 (2006) 1753.
- [19] B. Boisseau, G. Esposito-Farese, D. Polarski and A.A. Starobinsky, *Phys. Rev. Lett.* 85 (2000) 2236.
- [20] S. Capozziello, *Int. J. Mod. Phys. D* 11 (2002) 483.
- [21] T. Kaluza, *Sits. Press. Akad. Wiss. Math. Phys. K* 1 (1921) 895.
- [22] O. Klein, *Zeits. Phys.* 37 (1926) 895.
- [23] G. Calcagni, *Phys. Rev. Lett.* 104 (2010) 251301.
- [24] Y.-F. Cai, S. Capozziello, M. De Laurentis and E.N. Saridakis, *Rep. Progr. Phys.* 79 (2016) 106901.
- [25] F.R. Urban and A.R. Zhitnitsky, *JCAP* 0909 (2009) 018.
- [26] N. Ohta, *Phys. Lett. B* 695 (2011) 41.
- [27] C.-J. Feng, X.-Z. Li and P. Xi, *JHEP* 1205 (2012) 046.
- [28] G. Veneziano, *Nucl. Phys. B* 159 (1979) 213.
- [29] M. Li, *Phys. Lett. B* 603 (2004) 1.
- [30] M. Li et al., *Commun. Theor. Phys.* 56 (2011) 525.

- [31] V.F. Cardone, A. Troisi and S. Capozziello, *Phys. Rev. D* 69 (2004) 083517.
- [32] H. Wei and R.-G. Cai, *Phys. Lett. B* 660 (2008) 113.
- [33] A. Yu. Kamenshchik, U. Moschella and V. Pasquier, *Phys. Lett. B* 511 (2001) 265.
- [34] M. C. Bento, O. Bertolami and A. A. Sen, *Phys. Rev. D* 66 (2002) 043507.
- [35] S. Chandrasekhar, "An Introduction to the Study of Stellar Structure", (1939, Dover editions: 1958, 1967)
- [36] J. Christensen-Dalsgaard, "Lecture Notes on Stellar Structure and Evolution", (Aarhus Univ. Press, Aarhus, (2004) p.13.
- [37] S. Nesseris and L. Perivolaropoulos, *Phys. Rev. D* 70 (2004) 043531.
- [38] Z. G. Dai, E. W. Liang and D. Xu, *Ap. J. Lett.* 612 (2004) L101.
- [39] Z. H. Zhu, M. K. Fujimoto and X. T. He, *Astron. Astrophys.* 417 (2004) 833.
- [40] D.N. Spergel et al., *Astrophys. J. Suppl.* 170 (2007) 377.
- [41] Z.-L. Yi and T.-J. Zhang, *Mod. Phys. Lett. A* 22 (2007) 41.
- [42] L. Samushia and B. Ratra, *Ap. J. Lett.* 650 (2006) L5.
- [43] J.F. Zhang, X. Zhang and H.Y. Liu, *Eur. Phys. J. C* 52 (2007) 693.
- [44] M.A. Dantas, J.S. Alcaniz, D. Jain and A. Dev, *Astron. Astrophys.* 467 (2007) 421.
- [45] P.X. Wu and H.W. Yu, *Phys. Lett. B* 644 (2007) 16.
- [46] H. Lin, et al., *Mod. Phys. Lett. A* 24 (2009) 1699.
- [47] Wolfram Research Inc., *Mathematica 9.0*, 2012.
- [48] S.W. Allen, et al., *Mon. Not. Roy. Astron. Soc.* 353 (2004) 457.
- [49] M. Salti, O. Aydogdu, E.E. Kangal, A. Tas and K. Sogut, "Does the Polytopic proposal give better results in the 5D cosmology?", 2018, Submitted to a journal.
- [50] V. Gorini, A. Kamenshchik and U. Moschella, *Phys. Rev. D* 67 (2003) 063509.
- [51] U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, *Mon. Not. Roy. Astron. Soc.* 344 (2003) 1057.
- [52] V. Sahni, T. D. Saini, A. A. Starobinsky and U. Alam, *JETP Lett.* 77 (2003) 201.
- [53] X.Y. Yang, Y.B. Wu, J.B. Lu and S. Li, *Chin. Phys. Lett.* 24 (2007) 302.
- [54] J. Lu, *Phys. Lett. B* 680 (2009) 404.
- [55] L. Perivolaropoulos, *Phys. Rev. D* 71 (2005) 063503.
- [56] N. Suzuki, et al., *Astrophys. J.* 746 (2012) 85.
- [57] A. Banijamali, S. Bellucci, B. Fazlpour and M. Solbi, *Astrophys. Space Sci.* 363 (2018) 59.

- [58] S.-L. Cao, S. Li, H.-R. Yu and T.-J. Zhang, RAA (Research in Astronomy and Astrophysics) 18 (2018) 26.
- [59] C. Zhang, et al., RAA (Research in Astronomy and Astrophysics) 14 (2014) 1221.
- [60] J. Simon, L. Verde and R. Jimenez, Phys. Rev. D 71 (2005) 123001.
- [61] M. Moresco, L. Verde, L. Pozzetti, R. Jimenez, and A. Cimatti, J. Cosmol. Astropart. Phys. 7 (2012) 053.
- [62] E. Gaztañaga, A. Cabré and L. Hui, MNRAS 399 (2009) 1663.
- [63] X. Xu, A. J. Cuesta, N. Padmanabhan, D. J. Eisenstein and C. K. McBride, MNRAS 431 (2013) 2834.
- [64] M. Moresco, et al., J. Cosmol. Astropart. Phys. 5 (2016) 014.
- [65] D. Stern, R. Jimenez, L. Verde, M. Kamionkowski and S.A. Stanford, JCAP 02 (2010) 008.
- [66] R. Jimenez, L. Verde, T. Treu and D. Stern, Astrophys. J. 593 (2003) 622.
- [67] L. Samushia, et al., MNRAS 429 (2013) 1514.
- [68] C. Blake, et al., MNRAS 425 (2012) 405.
- [69] O. Farooq and B. Ratra, Astrophys. J. 766 (2013) L7.
- [70] P. Thakur, S. Ghose and B.C. Paul, Mon. Not. R. Astron. Soc. 397 (2009) 1935.
- [71] T. Delubac, et al., A&A 574 (2015) A59.
- [72] D.J. Eisenstein and W. Hu, Astrophys. J. 496 (1998) 605.
- [73] E. Komatsu, et al., Astrophys. J. Suppl. 180 (2009) 330.
- [74] F.C. Santos, M.L. Bedran, V. Soares, Phys. Lett. B 646 (2007) 215.