

Mass deficit and topology of nucleons

Michael Tzoumpas

Mechanical and Electrical Engineer
National Technical University of Athens
Irinis 2, 15234 Athens, Greece
E-mail: m.tzoumpas@gmail.com

April 2019

Abstract. Nucleus is identical with the lower inverse electric-nuclear field where a rapid increase of its potential occurs with a corresponding reduction of the space cohesive pressure, resulting to the mass deficit of the neutron entering the nucleus and to the finding of its location potential. Therefore, the so called topology of the nucleons can be now found. So, the neutrons are stable into the lower inverse nuclear field where a reduced cohesive pressure prevails. Moreover, there would be no nuclei without the presence of neutrons that reduce the negativity of the protons field, while neutrons are those that move into the nuclei (with the remaining half of their kinetic energy) on circular orbits around immobilized protons which have spin only.

Keywords: Inverse electric field; cohesive pressure; mass deficit; topology of nucleons.

PACS numbers: 03.50.Kk, 12.10.-g

1. Mass deficit Δm of neutrons

By the unified theory of dynamic space^{1,2} the inverse electric field³ of nucleus is described (Fig. 1), whose the change of its relative electric density³ ρ affects directly the cohesive pressure^{1,2} of proximal space, since it depends on the number of pairs of electrically opposite elementary units,^{1,2} which have remained in the electric field and caused from tensions

$$F = kL_0 \tag{1}$$

between these electric units where

$$L_0 = 0,558 \cdot 10^{-54} m \tag{2}$$

is the quantum dipole length.⁴ Specifically, the remaining cohesive pressure P is proportional to these pairs of electrically opposite units, which have remained in the electric field and especially in the lower one (section 2), where this change happens rapidly. So, if ρ is the relative electric density at a position of the inverse electric field, then the absolute electric density $\rho_0 - \rho$ is proportional to the number of the above pairs

of units, whose the attractive forces create the remaining cohesive pressure P at this position. Consequently, the cohesive pressures P_0 and P are respectively proportional to the background electric density ρ_0 and the absolute one $\rho_0 - \rho$, that is

$$\frac{P}{P_0} = \frac{\rho_0 - \rho}{\rho_0} \Rightarrow P = P_0 \frac{\rho_0 - \rho}{\rho_0}. \quad (3)$$




Figure 1. Dynamics of the upper and lower inverse electric field of proton by entering another proton into these fields where ρ_1, ρ_2, ρ_3 and ρ_4 the relative electric densities, F_1, F_2, F_3 and F_4 the Coulomb electric forces, P_1, P_2, P_3 and P_4 the cohesive pressures, F'_a and F_a the antigravity forces,⁵ r_{el} the electric radius of proton and B the potential barrier

It has been found that the cohesive pressure P_0 causes at the core vacuum^{1,2} of neutron (of a radius r) a total gravity force⁴

$$F_0 = 4\pi r^2 P_0, \quad (4)$$

which, due to $E_0 = F_0 L_0$ (Eq. 6),⁴ is identical with its gravity⁶ mass[‡]

$$m_0 = \frac{E_0}{C_0^2} = \frac{F_0 L_0}{C_0^2} \Rightarrow m_0 = \frac{F_0 L_0}{C_0^2}. \quad (5)$$

‡ $F_f^2 = F_0^2 + F_s^2$,⁶ where for the E/M wave⁷ applies $F_0 = 0$, therefore $F_f = F_s$, namely the final force F_f of the formation is equal to the accumulated force⁶ F_s , where $F_f = E/L_0$ represents the energy of the E/M wave and $F_s = pC_0/L_0$ represents its momentum.⁶ Substituting in the above $F_f = F_s$ we have $E/L_0 = pC_0/L_0$, where $p = mC_0$ is the momentum of the formation, so $m = E/C_0^2$.

Between M and O the potential of nuclear field is negative and takes very large values. Consequently, from M to E there exists the upper inverse electric-nuclear field and from M to O the lower one, where the increase of potential is rapid.

The potential V of the electric-nuclear field is⁹

$$V = 2K \frac{Ze}{r} - K \frac{Ze}{x}, \quad (7)$$

where r is the electric radius³ of nucleus, Ze its electric charge and K the Coulomb constant.




Figure 3. Limited inverse electric field of the calculated¹⁰ negative electric charge $q_n = -0,685e$ of neutron with its cloud of positive electrical units

Nucleus is identical with the lower inverse electric-nuclear field where a rapid increase of potential occurs with a corresponding reduction of cohesive pressure. By the mass deficit of the neutron (section 1) entering the nucleus, the location potential and the topology of the nucleons can be now found. The proton entering the nucleus can be cleaved (beta decay β^+) at the strong negative potential of field, which though reduced by entering of the neutron in the nucleus. This happens at the scale of nucleus where, due to the neutron's magnetic dipole moment¹⁰

$$\mu = -1,913\mu_n, \quad (8)$$

the calculated¹⁰ negative electric charge of neutron (Fig. 3)

$$q_n = -0,685e \quad (9)$$

behaves as a positively charged particle with the positive potential of its inverse electric field as a cloud of positive electrical units. So, the result is to affect the nuclear field and the cohesive pressure of the proximal space. This reduction in the negative potential of the field allows protons to be present in the nuclei.

It is noted that two protons can not exist in the nucleus without the presence of a neutron, because the increased negative potential of field causes a cleaving (beta decay β^+) of one proton. There would be no nuclei without the presence of neutrons that reduce the negativity of the protons field.

Oppositely, a greater reduction in the negativity of nuclear field may cause a neutron cleaving (beta decay β^-) and a reduction in mass deficit due to increased cohesive pressure.

Moreover, the neutron as a positively charged particle at the nucleus scale repels the closest proton, which is now moving on a helical orbit emitting gamma radiation and is finally immobilized. This radiant energy of the proton transmitted by the neutron is measured as mass deficit Δm and is equal to half of the kinetic energy of the neutron. Therefore, neutrons are those that move into the nuclei¹¹ (with the remaining half of their kinetic energy) on circular orbits around immobilized protons which have spin¹² only.

3. References

- [1] N.I.Gosdas, *The Unified Theory of Dynamic Space*, Greek Edition (Trohalia, Athens, 1999).
- [2] M.Tzoumpas, *Hubble's Law and antigravity - Higgs boson and gravity*, <http://viXra.org/abs/1710.0082> [Quantum Gravity and String Theory].
- [3] M.Tzoumpas, *Inverse electric-nuclear field*, <http://viXra.org/abs/1902.0266> (sections 1 and 2) [High Energy Particle Physics].
- [4] M.Tzoumpas, *Hubble's Law and antigravity - Higgs boson and gravity*, <http://viXra.org/abs/1710.0082> (subsection 3.2) [Quantum Gravity and String Theory].
- [5] M.Tzoumpas, *Universe expansion - Black holes - Nuclear forces*, <http://viXra.org/abs/1903.0248> (sections 4) [Relativity and Cosmology].
- [6] M.Tzoumpas, *Time as motion phenomenon - Physics Laws do not apply to inertial systems*, <http://viXra.org/abs/1802.0372> (sections 4 and 5) [Relativity and Cosmology].
- [7] M. Tzoumpas, *Unified E/M waves theory*, <http://viXra.org/abs/1905.0079> [Quantum Gravity and String Theory].
- [8] M.Tzoumpas, *Time as motion phenomenon - Physics Laws do not apply to inertial systems*, <http://viXra.org/abs/1802.0372> (sections 2) [Relativity and Cosmology].
- [9] M.Tzoumpas, *Inverse electric-nuclear field*, <http://viXra.org/abs/1902.0266> (sections 4) [High Energy Particle Physics].
- [10] M.Tzoumpas, *What is the spin of the particles?*, <http://viXra.org/abs/1810.0281> (section 1) [High Energy Particle Physics].
- [11] N.I.Gosdas, *The Structure of Nuclei* (sections 4.7-4.12, pages 106-201), Greek Edition (SALTO, Thessaloniki, 2001).
- [12] M.Tzoumpas, *What is the spin of the particles?*, <http://viXra.org/abs/1810.0281> (sections 2 and 3) [High Energy Particle Physics].