

Proof of Twin Prime Conjecture

K.H.K. Geerasee Wijesuriya

Research Scientist in Physics and Astronomy

PhD student in Astrophysics, Belarusian National Technical University

BSc (Hons) in Physics and Mathematics University of Colombo, Sri Lanka

Doctorate Degree in Physics

geeraseew@gmail.com , geerasee1@gmail.com

October 2019

Author's Biography

The author of this research paper is K.H.K. Geerasee Wijesuriya . And this proof of twin prime conjecture is completely K.H.K. Geerasee Wijesuriya's proof.

Geerasee she studied before at Faculty of Science, University of Colombo Sri Lanka. And she graduated with BSc (Hons) in Physics and Mathematics from the University of Colombo, Sri Lanka in 2014. And in March 2018, she completed her first Doctorate Degree in Physics with first class recognition. Now she is following her second PhD in Astrophysics with Belarusian National Technical University.

Geerasee has been invited by several Astronomy/Physics institutions and organizations worldwide, asking to get involve with them. Also, She has received several invitations from some private researchers around the world asking to contribute to their researches. She worked as Mathematics tutor/Instructor at Mathematics department, Faculty of Engineering, University of Moratuwa, Sri Lanka. Furthermore she has achieved several other scientific achievements already.

List of Affiliations

Faculty of Science, University of Colombo, Sri Lanka , Belarusian National Technical University
Belarus

Acknowledgement

I would be thankful to my parents who gave me the strength to go forward with mathematics and Physics knowledge and achieve my scientific goals.

Keywords: prime; contradiction; greater than ; natural number

Abstract

Twin prime numbers are two prime numbers which have the difference of 2 exactly. In other words, twin primes is a pair of prime that has a prime gap of two. Sometimes the term twin prime is used for a pair of twin primes; an alternative name for this is prime twin or prime pair. Up to date there is no any valid proof/disproof for twin prime conjecture. Through this research paper, my attempt is to provide a valid proof for twin prime conjecture.

Literature Review

The question of whether there exist infinitely many twin primes has been one of the great open questions in number theory for many years. This is the content of the twin prime conjecture, which states that there are infinitely many primes p such that $p + 2$ is also prime. In 1849, de Polignac made the more general conjecture that for every natural number k , there are infinitely many primes p such that $p + 2k$ is also prime. The case $k = 1$ of de Polignac's conjecture is the twin prime conjecture.

A stronger form of the twin prime conjecture, the Hardy–Littlewood conjecture, postulates a distribution law for twin primes akin to the prime number theorem. On April 17, 2013, Yitang Zhang announced a proof that for some integer N that is less than 70 million, there are infinitely many pairs of primes that differ by N . Zhang's paper was accepted by *Annals of Mathematics* in early May 2013. Terence Tao subsequently proposed a Polymath Project collaborative effort to optimize Zhang's bound. As of April 14, 2014, one year after Zhang's announcement, the bound has been reduced to 246. Further, assuming the Elliott–Halberstam conjecture and its generalized form, the Polymath project wiki states that the bound has been reduced to 12 and 6, respectively. These improved bounds were discovered using a different approach that was simpler than Zhang's and was discovered independently by James Maynard and Terence Tao.

Assumption

Let's assume that there are finitely many twin prime numbers.

Therefore we proceed by considering that there are finitely many twin prime numbers. Then let the highest twin prime numbers are P_{n-1} and $(P_{n-1} + 2)$. Then for all prime numbers P_n greater than P_{n-1} , $(P_n - 2)$ is not a prime number.

Methodology

With this mathematical proof, I use the contradiction method to prove the twin prime conjecture.

Let P_n is an arbitrary prime number greater than P_{n-1} (because there are infinite number of prime numbers). Then according to our consideration, $(P_n - 2)$ is not a prime number (And P_n is a prime such that $(P_n - 2)$ is divisible by x_3 . To see the meaning of x_3 and P_3 , please refer the below content). To see the existence of prime P_n (greater than P_{n-1}) such that $(P_n - 2)$ is divisible by x_3 , please refer the 'Proof' below. Since $P_n > 2$ and since P_n is a prime number and since P_n is an odd number, for all prime numbers P_i :

$$P_i (< P_n / 2): P_n / P_i = r_1$$

$$\text{Thus } P_n = P_i * r_1 \dots \dots \dots (01.0)$$

Where r_1 is a rational number (which is not a natural number)

But according to our consideration, $(P_n - 2)$ is not a prime number. Also since P_n is a prime number greater than 2, $(P_n - 2)$ is an odd number.

Thus for some prime number $P_1 (< [(P_n - 2) / 2])$; $(P_n - 2) / P_1 = x_1$. Where we choose P_1 such that x_1 is a natural number. But since previously chose P_i is any arbitrary prime number less than $(P_n / 2)$; now we consider $P_1 = P_i$

$$\text{Then } (P_n - 2) = P_1 * x_1 \dots \dots \dots (02) \text{ and } P_n = P_1 * r_1 \dots \dots \dots (01)$$

Let P_N is a prime number greater than P_{n-1} (which is not P_n). Then according to our assumption, $(P_N + 2)$ is not a prime number. Here P_N is a prime number such that $(P_N + 2)$ is dividing by prime number P_2 . $\dots \dots \dots (1.1)$

*** Here we should consider a prime number P_N such that $P_3 \neq x_3$ and x_3 is not divisible by P_3 ; whenever $(P_N - 2) = P_3 \cdot x_3$. See the below content in the 'Proof' to see the verification of the existence of prime number P_N such that $P_3 \neq x_3$ and x_3 is not divisible by P_3 ; whenever $(P_N - 2) = P_3 \cdot x_3$.

Thus $(P_N + 2) = P_2 \cdot x_2$ for some x_2 natural number. Because there are infinitely many prime numbers. Since P_N is a prime number, for some r_2 (rational number which is not a natural number): $P_N / r_2 = P_2$.

Thus $(P_N + 2) = P_2 \cdot x_2$ (03) and $P_N = r_2 \cdot P_2$ (04)

x_1 and x_2 are natural numbers and P_1 and P_2 are prime numbers.

Since P_N is a prime number , $(P_N - 2)$ is also not a prime number (Since $P_N - 2 > P_{n-1}$)

Then for some prime P_3 , $(P_N - 2) / P_3 = x_3$, Here we should considered prime number P_N such that $P_3 \neq x_3$ and x_3 is not divisible by P_3 ; whenever $(P_N - 2) = P_3 \cdot x_3$. See the below content in the 'Proof' to see the verification of the existence of prime number P_N such that $P_3 \neq x_3$ and x_3 is not divisible by P_3 ; whenever $(P_N - 2) = P_3 \cdot x_3$

$(P_N - 2) = P_3 \cdot x_3$ (05)

By (04) and (05): $P_3 \cdot x_3 = P_2 \cdot r_2 - 2$ (06)

But according to the below induction method proof which is in the "Proof" below, there exists the prime P_n such that $(P_n - 2)$ divides by x_3 .

But $(P_N + 2)$, $(P_n - 2)$ both are odd numbers. Thus $(P_N + 2) = (P_n - 2) + 2.l$ for some l integer number.....(06)'

Then $(P_N - 2) = (P_n - 2) + 2.l - 4 = P_n + 2.l - 6 = P_n + 2 \cdot (l - 3)$ (6.1)'

Since $(P_N - 2)$ is divisible by P_3 , $[P_n + 2 \cdot (l - 3)]$ is divisible by P_3 (6.1)

And we know that $(P_N + 2) = (P_n - 2) + 2.l \rightarrow P_N = P_n + 2.l - 4$ (*)

Thus by (*): $P_1 \cdot r_1 + 2.l - 4 = r_2 \cdot P_2$. Thus by (06): $P_3 \cdot x_3 + 2 = P_1 \cdot r_1 + 2.l - 4$

Thus $P_3 \cdot x_3 - 2.l + 6 = P_1 \cdot r_1 = P_n$ (6.1.0)

Thus $P_3 * x_3 + 2. (l - 3) = P_n + 4. (l - 3) = P_n + 2.P_N - 4 - 2.P_n = 2.P_N - 4 - P_n$ (by (6.1)')

Thus $P_3 * x_3 + 2. (l - 3) = 2.P_N - 4 - P_n = P_n$,

Thus $P_3 * x_3 + 2. (l - 3) = P_n$,(7)

Thus $P_3 * x_3 + 2.l = 6 + 2. P_3 * x_3 - P_n$

$P_3*(x_3 + 1)+(2.l - P_3) = (6 - P_n) + 2. P_3 * x_3$

$P_3*(x_3 + 1)+(2.l - P_3 + M) = (6 + M - P_n) + 2. P_3 * x_3$; Where M is an integer.....(8)

But we chose M such that $(P_n - 6 - M) = x_3 .m_1$ for some integer m_1 (m_1 is divisible by x_3). But we choose an integer F (not divisible by x_3) such that $(P_3 + F)$ is not divisible by x_3 and we chose integer F and m_1 such that $(F + x_3 .m_1 + P_3)$ divisible by x_3 . **Because $(x_3 .m_1)$ has the order of x_3 as at least 2 , since m_1 is divisible by x_3 .**

And we chose P_n such that $(P_n - 2)$ divides by x_3(9) But here $P_3 \neq x_3$.

Thus $(P_n - 6 - M) = x_3 .m_1$ and $(P_n - 2) = x_3.m_0$ for some integer m_1 and m_0 .

But $(P_N - 2)$ divides by x_3 . Thus according to our choice, $[(P_N - 2) - (P_n - 2)]$ divides by x_3 . i.e. $(P_N - P_n)$ divides by x_3 . Thus $P_N - P_n = x_3. m_2$; for some integer m_2 .

Thus according to our choice: $[(P_n - 6 - M) = x_3 .m_1]$ and $[(P_N - P_n) = x_3. m_2]$ for some integer m_1 and integer m_2 .

But $(P_n - 6 - M) = x_3 .m_1$ and $(P_n - 2) = x_3.m_0$ for some integer m_1 and integer m_0 .

Thus $(4 + M) = x_3.m_3$; $m_3 = (m_0 - m_1)$ for an integer m_3 (10)

By (*): $P_N - P_n + 4 = 2.l$

Thus $2.l - P_3 + M = (P_N - P_n) + 4 - P_3 + M = x_3. m_2 + (M + 4 - P_3) = x_3 . r'$ (since $[M + 4]$ is divisible by x_3 and since P_3 is a prime number which is not equal to x_3). And here r' is not an integer. But r' is a rational number.

Then $2.l - P_3 + M - F = (P_N - P_n) + 4 - P_3 + M - F = x_3. m_2 + (M + 4 - P_3 - F) = x_3 . r''$; r'' is not an integer. Because $(P_3 + F)$ is not divisible by x_3 .

Thus by (8): $(x_3+1)* P_3 + x_3.r'' = - x_3 .m_1 - F + 2. P_3 * x_3$ (11)

But we chose F and m_1 such that $(x_3 \cdot m_1 + P_3 + F)$ is divisible by x_3 . Because $(x_3 \cdot m_1)$ has order of x_3 as 2, since m_1 is divisible by x_3 . Let $(x_3 \cdot m_1 + P_3 + F) = x_3 \cdot m_4$; where m_4 is an integer.

Thus by (11): $P_3 \cdot x_3 + [x_3 \cdot r''] = 2 \cdot P_3 \cdot x_3 - x_3 \cdot m_4$ where m_4 is an integer.

Then $r'' = P_3 - m_4 \dots\dots\dots(12)$

But m_4 and P_3 both are integers. But r'' is not an integer. Thus by (12), we have a contradiction. Therefore the only possibility is: our assumption is false. Therefore there are infinitely many Twin Prime Numbers.

Proof

Now let's prove that there exists infinite number of Prime numbers P_n such that $(P_n - 2) = x_3 \cdot m_0$ for an integer m_0 .

Let's consider the statement $Q(n) : [P(n) - 2] / x_3 = x(n)$; where $P(n)$ is the n th prime number which obeys $[P(n) - 2] = x_3 \cdot x(n)$. And $x(n)$ is the n th integer which is in the form of $[P(n) - 2] / x_3$.

For $n = 1$, L.H.S. of $Q(1) = [2 - 2] / x_3 = 0$. But for $x(n) = 0$ (which is an integer), R.H.S. of $Q(1) = 0$. Thus for $n = 1$, R.H.S. of $Q(1) =$ L.H.S. of $Q(1)$. Thus for $n = 1$, the result holds.

Now assume for $n = s$, the result $Q(s)$ holds. Then $[P_s - 2] / x_3 = x(s) =$ natural number, where $x(s) =$ integer number.

Here we must consider $n = s$ part as below.

Let ϵ_s is a positive real number $\epsilon_s = [- B + P_s + C_s - 2 + x_3 \cdot k'] / P_s > 0$ for all $s > (L - 2)$, $h_s < P_s \cdot \epsilon_s$ (since the only existing $s > (L - 2)$ is $(L - 1)$; " for all $s > (L - 2)$ means $s = (L - 1)$)".

Where k' is an integer number. Here the chosen k' integer number is responsible for $h_s < P_s \cdot \epsilon_s$ for all $s > (L - 2)$ and k' is responsible for $\epsilon_{L-1} > 0$. That means here the value of k' is responsible to say: " ϵ_s is existing such that $h_s < P_s \cdot \epsilon_s$, for $s = (L - 1)$ ". Here $h_j = b_j$ for all $j < (L - 1) = s$. And where $\sum b_j = B$ for $j < (L - 1) = s$. Then for ϵ_s , $h_s = P_s \cdot \epsilon_s - C_s$; here $s \equiv L - 1$. *** the meaning of 'j' is the order number and h_j is the prime gap between P_{j+1} and P_j , please refer the below content and the 2nd reference.

But $s \equiv (L - 1)$. But here we chose C_{L-1} such that $h_{L-1} = P_{L-1} * C_{L-1} - C_{L-1}$

But $h_{L-1} = P_{L-1} * C_{L-1} - C_{L-1} = (P_s - B - 2 + x_3 .k')$. Where k' is an integer number.

Then let's show for $n = s + 1$, $Q(s+1)$ holds. We denote $P(s+1) = P_L$

But we know $[P_s - 2] / x_3 = x(s)$ (13)

Now let's use the 2nd reference to proceed further.

By 2nd reference, $P_L = 2 + \sum_{j=1}^{L-1} h_j$ (i)

But we know already that for $C_{L-1} > 0$, $h_{L-1} < P_{L-1} * C_{L-1}$ for $L - 1 = s$.

Here $s \equiv (L - 1)$

(*** refer the 2nd reference below)

Then we already know that for some C_{L-1} positive number, $h_{L-1} = P_{L-1} * C_{L-1} - C_{L-1}$.

But $h_{L-1} = P_{L-1} * C_{L-1} - C_{L-1}$ for $(L - 1) \equiv s$

We know already that $C_{L-1} = [P_s - B + C_{L-1} - 2 + x_3 .k'] / P_{L-1} > 0$.

And $h_{L-1} = P_{L-1} * C_{L-1} - C_{L-1} = (- B + P_s - 2 + x_3 .k')$. Where k' is an integer number. We know already that the chosen k' integer number is responsible for $C_{L-1} > 0$.

We know that $h_j = b_j$ for all $j < (L - 1)$. Where b_j is a natural number. Also we know that $\sum b_j = B$ for $j < L - 1$.

Thus by (i): $P_L = 2 + P_s + x_3.k' - B - 2 + B = x_3 .k' + P_s$

Thus $(P_L - 2) = (P_s - 2) + x_3.k'$ (14)

But $(P_s - 2)$ is divisible by x_3 and $(P_s - 2) / x_3 = x(s)$ according to (13). Thus $(P_L - 2)$ is divisible by x_3 according to (14), since $x_3.k'$ is divisible by x_3 .

Thus $(P_L - 2)$ is divisible by x_3 . i.e. $[P(s+1) - 2]$ is divisible by x_3 .

Thus for $n = s + 1 (= L)$, the result $Q(n + 1)$ holds. Thus by mathematical induction method:

There exists infinite number of prime numbers P_L such that $(P_L - 2) = x_3 \cdot m'_0$ for some integer number m'_0 .

Thus there exists P_n prime (where we consider that as prime numbers greater than P_{n-1}) such that $(P_n - 2)$ is divisible by x_3 and $(P_n - 2) = x_3 \cdot m_0$ for some integer number m_0 , whenever $P_3 \neq x_3$.

Verification of existence of prime number P_N (greater than P_{n-1}) such that $(P_N - 2) = P_3 \cdot x_3$; for the integer number x_3 which is not divisible by P_3

Let C_s is a positive real number $C_s = [-A + C_s + (P_3) \cdot t_s] / P_s > 0$ for all $s > (R - 2)$, $g_s < P_s * C_s$ (since the only existing $s > (R - 2)$ is $(R - 1)$; "for all $s > (R - 2)$ means $s = (R - 1)$ "). Where t_s is an integer number such that t_s is not divisible by P_3 . Here the chosen t_s integer number is responsible for $g_s < P_s * C_s$ for all $s > (R - 2)$ and t_s is responsible for $C_{R-1} > 0$. That means here the value of t_s is responsible to say: " C_s is existing such that $g_s < P_s * C_s$, for $s = (R-1)$ ". Here $g_j = a_j$ for all $j < (R - 1) = s$. And where $\sum a_j = A$ for $j < (R - 1) = s$. Then for C_s , $g_s = P_s * C_s - C_s$; here $s \equiv R - 1$. *** the meaning of 'j' is the order number and g_j is the prime gap between P_{j+1} and P_j , please refer the below content and the 2nd reference.

But $s \equiv (R - 1)$. But here we chose C_{R-1} such that $g_{R-1} = P_{R-1} * C_{R-1} - C_{R-1}$

But $g_{R-1} = P_{R-1} * C_{R-1} - C_{R-1} = (-A + P_3 \cdot t_s)$. Where t_s is an integer number which is not divisible by P_3 .

Now let's use the 2nd reference to proceed further.

By 2nd reference, $P_R = 2 + \sum_{j=1}^{R-1} g_j$ (ii)

But we know already that for $C_{R-1} > 0$, $g_{R-1} < P_{R-1} * C_{R-1}$ for $R - 1 = s$.

Here $s \equiv (R - 1)$

(*** refer the 2nd reference below)

Then we already know that for some C_{R-1} positive number, $g_{R-1} = P_{R-1} * C_{R-1} - C_{R-1}$.

But $g_{R-1} = P_{R-1} * C_{R-1} - C_{R-1}$ for $(R - 1) \equiv s$

We know already that $C_{R-1} = [- A + C_{R-1} + P_3.t_s] / P_{R-1} > 0$.

And $g_{R-1} = P_{R-1} * C_{R-1} - C_{R-1} = (- A + P_3.t_s)$. Where t_s is an integer number that is not divisible by P_3 . We know already that the chosen t_s integer number is responsible for $C_{R-1} > 0$.

We know that $g_j = a_j$ for all $j < (R - 1)$. Where a_j is a natural number. Also we know that $\sum a_j = A$ for $j < R - 1$.

Thus by (ii): $P_R = 2 + P_3.t_s - A + A = P_3.t_s + 2$

Thus there exists prime number P_R such that $(P_R - 2) = P_3.t_s \dots\dots\dots(15)$ where t_s is not divisible by P_3 .

Now put $N \equiv R$. Then we can state that $(P_N - 2) = P_3.x_3$; for some integer x_3 which is not divisible by P_3 .

Discussion

We assumed initially that there are finitely many twin primes. After proceeding with that, I ended up with a contradiction. But to get the contradiction, I used that P_n and P_N as primes numbers greater than P_{n-1} . Also to get the contradiction, I used the facts that $(P_n - 2)$ and $(P_N + 2)$ and $(P_N - 2)$ as non-primes. And also I have used that x_1 , x_2 and x_3 as natural numbers (since $(P_n - 2)$, $(P_N + 2)$ and $(P_N - 2)$ are not prime numbers). Therefore to get the contradiction, I have used the facts got from our assumption. Then the only possibility is our assumption is false.

Results

Therefore I have used our assumption to get a contradiction finally as showed in (A.6). Therefore it is possible to conclude that our assumption is false.

Thus there are infinitely many twin prime numbers.

Appendix

Prime number: A natural number which divides by 1 and itself only.

Twin Prime Numbers: Two prime numbers which have the difference exactly 2.

We denote 'i' th prime gap $g_i = P_{i+1} - P_i$

Then according to the 2nd reference; Prime number $P_N = 2 + \sum_{j=1}^{N-1} g_j$

Also by 2nd reference: for all $\epsilon > 0$, there is a natural number 'n' such that for all $N - 1 > n$;

$$g_{N-1} < P_{N-1} \cdot \epsilon$$

References

1. Zhang, Yitang (2014). "Bounded gaps between primes". *Annals of Mathematics*. 179 (3): 1121–1174.
2. https://ipfs.io/ipfs/QmXoypizjW3WknFiJnKLwHCnL72vedxjQkDDP1mXWo6uco/wiki/Prime_gap.html
3. Terry Tao, Small and Large Gaps between the Primes
4. Maynard, James (2015), "Small gaps between primes", *Annals of Mathematics, Second Series*, **181** (1): 383–413
5. Tchudakoff, N. G. (1936). "On the difference between two neighboring prime numbers". *Math. Sb.* **1**: 799–814.
6. Ingham, A. E. (1937). "On the difference between consecutive primes" *Quarterly Journal of Mathematics. Oxford Series.* **8** (1): 255–266.