

Proof of

$$\sum_{n=1}^{\infty} (-1)^n = -\frac{1}{2}$$

July 29, 2019
Yuji Masuda
[\(y_masuda0208@yahoo.co.jp\)](mailto:y_masuda0208@yahoo.co.jp)

First, $\pm\infty$ is constant at any observation point (position).

If a set of real numbers is R , then,

$$R \times (\pm\infty) = \pm\infty$$

$$R + (\pm\infty) = \pm\infty$$

$$(-1) \times (\pm\infty) \neq \mp\infty$$

On the other hand, when x ($\in R$) is taken on a number line, the absolute value X becomes larger toward $\pm\infty$ as the absolute value X is expanded.

Similarly, as the size decreases, the absolute value X decreases toward 0. Furthermore, $\times (-1)$ represents the reversal of the direction of the axis.

$$\boxed{\begin{aligned}\frac{1}{\pm\infty} &= (-1) \cdot (\pm\infty) = i \\ (\pm\infty) \cdot i - 1 &= 0\end{aligned}}$$

$$(-1) \cdot (\pm\infty) = \frac{1}{\pm\infty}$$

$$i^2 = (\pm\infty)^2 \rightarrow i = \pm(\pm\infty)$$

$$\therefore i = -(\pm\infty) = (-1)(\pm\infty) = \frac{1}{\pm\infty}, (\because i \neq \pm(\pm\infty))$$

Next,


$$\boxed{\pi = \frac{2}{\pi} + 2 \arctan \left(\frac{1}{\tan \left(\frac{1}{x} \right)} \right), (\because x \geq \frac{1}{\pi})}$$

$$x = \frac{2}{\pi} \left(\geq \frac{1}{x} \right)$$

$$\pi = \left(\frac{2}{\pi} \right) + 2 \arctan \left(\frac{1}{\tan \left(\frac{\pi}{2} \right)} \right) = \pi + 2 \arctan \left(\frac{1}{\pm\infty} \right)$$

$$\arctan \left(\frac{1}{\pm\infty} \right) = \arctan(i) = 0$$

$$\therefore \tan 0 = \frac{1}{\pm\infty} = (-1)(\pm\infty) = i$$



Second, we consider the figure below.




Fig.2

From the figure above, I got the following equation.

$$\theta = \arcsin \left(\frac{1}{\frac{2 \cdot (x+1)^2}{2x+1} - 1} \right)$$

Here, when take $\pm\infty$ to the consideration,

$$\boxed{\begin{aligned}\tan \theta &= \frac{2x+1}{2x(x+1)} = i \left(= (-1) \cdot (\pm\infty) = \frac{1}{\pm\infty} \right) \\ \therefore x &= -\frac{1}{2}(1+i)\end{aligned}}$$

Here, when we consider the figure above,

$$\frac{x^2}{(x+1)^2} = -1$$

So, when we put $x=2/i=-2i$,

Here, from Fig.1, $i+(\pm\infty)=0$.

$$\boxed{\begin{aligned}-2i + 2 \cdot (-1) \cdot (-2i) + 2 \cdot (-2i) + 2 \cdot (-1) \cdot (-2i) + 2 \cdot (-2i) + \dots &= i + (\pm\infty) = 0 \\ -2i \cdot (1 - 2 + 2 - 2 + 2 - 2 + \dots) &= i + (\pm\infty) = 0 \\ 1 - 2 + 2 - 2 + 2 - 2 + \dots &= \frac{1}{2} \left(-1 - \frac{\pm\infty}{i} \right) = \frac{1}{2} (-1 - (-1)) = 0 \\ 1 + 2((-1) + 1 + (-1) + 1 + (-1) + \dots) &= 0 \\ \frac{1}{2} + \sum_{n=1}^{\infty} (-1)^n &= 0\end{aligned}}$$

$$\boxed{\therefore \sum_{n=1}^{\infty} (-1)^n = -\frac{1}{2}}$$