

Merging Galaxies

In a new paper based on Illustris simulated images of merger galaxies, the astronomers present a way to help identify when imaged systems are mergers. [11]

Scientists, including from The Australian National University (ANU), say they have detected a black hole swallowing a neutron star for the first time. [10]

Gravitational waves from the merger of a black hole and a neutron star may have been spotted for the first time by the LIGO and Virgo detectors. [9]

Astrophysicists at Frankfurt, the Max Planck Institute for Radio Astronomy in Bonn, and Nijmegen, collaborating in the project BlackHoleCam, answer this question by computing the first images of feeding non-Einsteinian black holes: it is presently hard to tell them apart from standard black holes. [8]

Using data from the first-ever gravitational waves detected last year, along with a theoretical analysis, physicists have shown that gravitational waves may oscillate between two different forms called "g" and "f"-type gravitational waves. [7]

Astronomy experiments could soon test an idea developed by Albert Einstein almost exactly a century ago, scientists say. [6]

It's estimated that 27% of all the matter in the universe is invisible, while everything from PB&J sandwiches to quasars accounts for just 4.9%. But a new theory of gravity proposed by theoretical physicist Erik Verlinde of the University of Amsterdam found out a way to dispense with the pesky stuff. [5]

The proposal by the trio though phrased in a way as to suggest it's a solution to the arrow of time problem, is not likely to be addressed as such by the physics community—it's more likely to be considered as yet another theory that works mathematically, yet still can't answer the basic question of what is time. [4]

The Weak Interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and Time reversal symmetry.

The Neutrino Oscillation of the Weak Interaction shows that it is a General electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

Spotting merging galaxies.....	3
Scientists detect a black hole swallowing a neutron star	4
Were gravitational waves from a black-hole–neutron-star merger detected on 14 August?	5
The current ability to test theories of gravity with black hole shadows.....	6
Gravitational waves may oscillate, just like neutrinos.....	8
Two gravitons instead of one.....	8
Oscillations.....	9
Quest to settle riddle over Einstein's theory may soon be over.....	10
NEW THEORY OF GRAVITY DOES AWAY WITH NEED FOR DARK MATTER	11
Identification of a Gravitational Arrow of Time	12
Asymmetry in the interference occurrences of oscillators	13
Spontaneously broken symmetry in the Planck distribution law.....	15
The structure of the proton.....	16
The Weak Interaction	17
The General Weak Interaction	18
Fermions and Bosons	18
The fermions' spin	19
The source of the Maxwell equations	19
The Special Relativity.....	20
The Heisenberg Uncertainty Principle	20
The Gravitational force.....	20
The Graviton.....	21
What is the Spin?	22
The Casimir effect	22
The Fine structure constant	22
Conclusions.....	23
References	24

Author: George Rajna

Spotting merging galaxies

Over 30 years ago, the Infrared Astronomy Satellite discovered that the universe contained many extremely luminous galaxies, some more than a thousand times brighter than our own Milky Way, but which are practically invisible at optical wavelengths. These galaxies are powered by bursts of star formation buried deep within clouds of dust and gas. The dust absorbs the ultraviolet light while radiating at infrared wavelengths. In many cases the hyperactivity was triggered by a collisional encounter between galaxies that facilitated the collapse of interstellar gas into new stars.

Collisions between galaxies are common. Indeed, most galaxies have probably been involved in one or more encounters during their lifetimes, making these interactions an important phase in galaxy evolution and the formation of stars in the universe. The Milky Way, for example, is bound by gravity to the Andromeda galaxy and is approaching it at a speed of about 50 kilometers per second; we are expected to meet in another billion years or so. In the local universe about five percent of galaxies are currently in a merger, and mergers usually can be easily identified by the visible morphological distortions they produce such as tidal tails sweeping out from the galactic discs.

Not all infrared [luminous galaxies](#) show such distortions, however, and the issue of identifying (and classifying) mergers becomes especially problematic for studies of earlier cosmic epochs when the star formation rates were much higher than today, and when the merger rate of galaxies was also higher. (Moreover, such systems are preferentially discovered in deep galaxy surveys precisely because they are so luminous.) But galaxies in the distant cosmos are too remote to detect spatial signatures like tidal arms (at least with current telescopes). It is possible that other processes besides merger-induced star formation are lighting up some of these bright galaxies, for example accreting [supermassive black holes](#) can emit copious amounts of ultraviolet radiation. Because of such cases, estimates of star formation in the early universe based on luminosity measurements alone could be incorrect.

CfA astronomer Lars Hernquist is a pioneer in the development of computer simulations of merging galaxies. Several years ago he and a team of colleagues produced a massive new simulation of the formation and evolution of galaxies in the universe, called Illustris. In a new paper based on Illustris simulated images of [merger](#) galaxies, the astronomers present a way to help identify when imaged systems are mergers. They created about one million synthetic Hubble and James Webb Space Telescope images from their simulated mergers, and then looked for common morphological indicators of merging. They developed an algorithm that successfully identified mergers at roughly a seventy percent level of completeness out to distances of as much as eighty-five billion light-years (the current distance value), corresponding to light dating from the epoch about 2 billion years after the big bang. Results from the algorithm indicated that spatial features associated with strong central concentrations (or bulges) were most important for selecting past mergers, while double nuclei and asymmetries were most important for selecting future mergers (that is, sometime in the next 250 million years). The new algorithm will be particularly valuable when applied to future Webb images of very distant mergers. [11]

Scientists detect a black hole swallowing a neutron star

Scientists, including from The Australian National University (ANU), say they have detected a black hole swallowing a neutron star for the first time.

Neutron stars and black holes are the super-dense remains of dead stars.

On Wednesday 14 August 2019, gravitational-wave discovery machines in the United States and Italy detected ripples in space and time from a cataclysmic event that happened about 8,550 million trillion kilometres away from Earth.

Professor Susan Scott, from the ANU Research School of Physics, said the achievement completed the team's trifecta of observations on their original wish list, which included the merger of two black holes and the collision of two neutron stars.

"About 900 million years ago, this black hole ate a very dense star, known as a neutron star, like Pac-man—possibly snuffing out the star instantly," said Professor Scott, Leader of the General Relativity Theory and Data Analysis Group at ANU and a Chief Investigator with the ARC Centre of Excellence for Gravitational Wave Discovery (OzGrav).

"The ANU SkyMapper Telescope responded to the detection alert and scanned the entire likely region of space where the event occurred, but we've not found any visual confirmation."

Scientists are still analysing the data to confirm the exact size of the two objects, but initial findings indicate the very strong likelihood of a black hole enveloping a neutron star. The final results are expected to be published in scientific journals.

"Scientists have never detected a black hole smaller than five solar masses or a neutron star larger than about 2.5 times the mass of our Sun," Professor Scott said.

"Based on this experience, we're very confident that we've just detected a black hole gobbling up a neutron star.

"However, there is the slight but intriguing possibility that the swallowed object was a very light black hole—much lighter than any other black hole we know about in the Universe. That would be a truly awesome consolation prize."

ANU plays a lead role in Australia's partnership with the Advanced Laser Interferometer Gravitational-wave Observatory (LIGO), which is the most sensitive scientific instrument ever built and comprises twin detectors in the US.

The European Gravitational Observatory has a gravitational-wave detector in Italy called Virgo. [10]

Were gravitational waves from a black-hole–neutron-star merger detected on 14 August?

Gravitational waves from the merger of a black hole and a neutron star may have been spotted for the first time by the LIGO and Virgo detectors. The signal was detected on 14 August and appears to come from an event that occurred about 900 million light-years away.



PHYSICAL
REVIEW
RESEARCH

APS
physics

A new journal welcoming the
full spectrum of research topics

Open access

Submit your research today

Advertisement

The detection has been logged on the [Gravitational-Wave Candidate Event Database](#), where the LIGO–Virgo team say that there is a greater than 99% probability that the source of the gravitational waves are from a black-hole–neutron-star merger. It is likely that the larger object was heavier than five solar masses and the smaller object lighter than three solar masses.

There is no indication so far that any electromagnetic observations of the object have been made by any other telescopes. This will come as a disappointment to the astronomy community because “multimessenger” observations across the electromagnetic spectrum could provide important insights into black holes, neutron stars and how they merge. So far, the only multimessenger observations to involve gravitational waves were made in 2017 when LIGO–Virgo spotted a signal from the [merger of two neutron stars](#).



A new cosmic messenger

This is not the first time that LIGO–Virgo has reported evidence of a black-hole–neutron-star merger. Earlier this year on 26 April the detectors recorded a [possible signal](#) from such an event, which may have occurred 1.2 billion light-years away. The signal from this object, however was too weak to confirm.

After upgrades that started in mid-2017, the LIGO and Virgo detectors were back in operation on 1 April 2019. By the end of July, they had spotted 18 binary black hole merger candidates and four binary neutron star merger candidates. However, unlike the neutron star merger of [2017](#), no accompanying electromagnetic radiation from the four neutron-star mergers have been detected. Work is ongoing to confirm whether any of these candidates can be considered full-fledged discoveries. [9]

The current ability to test theories of gravity with black hole shadows

Astrophysicists at Frankfurt, the Max Planck Institute for Radio Astronomy in Bonn, and Nijmegen, collaborating in the project BlackHoleCam, answer this question by computing the first images of feeding non-Einsteinian black holes: it is presently hard to tell them apart from standard black holes.

Their findings are published as Advance Online Publication (AOP) on the *Nature Astronomy* website on 16 April 2018.

One of the most fundamental predictions of Einstein's theory of relativity is the existence of black holes. In spite of the recent detection of gravitational waves from [binary black holes](#) by LIGO, direct evidence using

electromagnetic waves remains elusive and astronomers are looking for it with radio telescopes. For the first time, collaborators in the ERC funded project BlackHoleCam, including astrophysicists at Goethe University Frankfurt, Max Planck Institute for Radio Astronomy (MPIfR) Bonn, and Radboud University Nijmegen, have compared self-consistent and realistic images of the shadow of an accreting [supermassive black hole](#) – such as the black-hole candidate Sagittarius A* (Sgr A*) in the heart of our Galaxy – both in general relativity and in a different theory of gravity. The goal was to test if Einsteinian black holes can be distinguished from those in alternative theories of gravity.

Not all of the light rays (or photons) produced by matter falling onto a black hole are trapped by the event horizon, a region of spacetime from which nothing can escape. Some of these photons will reach distant observers, so that when a black hole is observed directly, a "shadow" is expected against the background sky. The size and shape of this shadow will depend on the black hole's properties and on the theory of gravity.

Because the largest deviations from Einstein's theory of relativity are expected very close to the [event horizon](#), and since alternative theories of gravity make different predictions on the properties of shadow, direct observations of Sgr A* represent a very promising approach for testing gravity in the strongest regime. Making such images of the black-hole shadow is the prime goal of the international Event Horizon Telescope Collaboration (EHTC), which combines radio data from telescopes around the world.

Scientists from the BlackHoleCam Team in Europe, who are part of the EHTC, have now have gone a step further and investigated whether it is possible to distinguish between a "Kerr" black hole from Einstein's gravity and a "dilaton" black hole, which is a possible solution of an alternative [theory](#) of gravity.

The researchers studied the evolution of matter falling into the two very different types of black holes and calculated the radiation emitted to construct the images. Furthermore, real-life physical conditions in the telescopes and interstellar medium were used to create physically realistic images. "To capture the effects of different black holes we used realistic simulations of accretion disks with near-identical initial setups. These expensive numerical simulations used state-of-the-art codes and several months on the Institute's supercomputer LOEWE," says Dr. Mizuno, lead author of the study.

Moreover, expected radio images obviously have a limited resolution and image fidelity. When using realistic image resolutions, the scientists found, to their surprise, that even highly non-Einsteinian black holes could disguise themselves as normal black holes.

"Our results show that there are theories of [gravity](#) in which black holes can masquerade as Einsteinian, so new techniques of analyzing EHT data may be needed to tell them apart," remarks Luciano Rezzolla, professor at Goethe University and leader of the Frankfurt team. "While we believe general relativity is correct, as scientists we need to be open-minded. Luckily, future observations and more advanced techniques will eventually settle these doubts," concludes Rezzolla.

"Indeed, independent information from an orbiting pulsar, which we are actively searching for, will help eliminate these ambiguities," says Michael Kramer, director at the MPI for Radio Astronomy in Bonn. Heino Falcke (professor at Radboud University), who 20 years ago proposed using radio telescopes to image the shadow of [black holes](#), is optimistic. "There is little doubt that the EHT will eventually obtain strong evidence of a black hole shadow. These results encourage us to refine our techniques beyond the current state of the art and thus make even sharper images in the future. [8]

Gravitational waves may oscillate, just like neutrinos

Using data from the first-ever gravitational waves detected last year, along with a theoretical analysis, physicists have shown that gravitational waves may oscillate between two different forms called "g" and "f"-type gravitational waves. The physicists explain that this phenomenon is analogous to the way that neutrinos oscillate between three distinct flavors—electron, muon, and tau. The oscillating gravitational waves arise in a modified theory of gravity called bimetric gravity, or "bigravity," and the physicists show that the oscillations may be detectable in future experiments.

The researchers, Kevin Max, a PhD student at Scuola Normale Superiore di Pisa and INFN Pisa, Italy; Moritz Platscher, a PhD student at the Max Planck Institute for Nuclear Physics, Germany; and Juri Smirnov, a postdoc at the University of Florence, Italy, have published a paper on their analysis of gravitational wave oscillations in a recent issue of *Physical Review Letters*.

As the physicists explain, the work may help answer the question of what "the other 95%" of the universe is made of, by suggesting that the answer may lie in modifications to gravity rather than new particles.

"Only 5% of matter is of a type we think to understand properly," Smirnov told Phys.org. "To address the question of what our universe is made of ('dark matter' and 'dark energy'), most authors discuss alternative particle physics models with new particles. However, experiments such as the ones at the LHC [Large Hadron Collider] haven't detected any exotic particles, yet. This raises the question if maybe the gravitational side needs to be modified.

"In our work, we ask what signals we could expect from a modification of gravity, and it turns out that bigravity features a unique such signal and can therefore be discriminated from other theories. The recent detection of gravitational waves by LIGO [Laser Interferometer Gravitational-Wave Observatory] has opened a new window on the dark sectors of the universe for us. Whether Nature has chosen general relativity, bigravity, or any other theory is a different question in the end. We can only study possible signals for experimentalists to look for."

Two gravitons instead of one

Currently, the best theory of gravity is Einstein's theory of general relativity, which uses a single metric to describe spacetime. As a result, gravitational interactions are mediated by a single hypothetical particle called a graviton, which is massless and so travels at the speed of light.

The main difference between general relativity and bigravity is that bigravity uses two metrics, g and f. Whereas g is a physical metric and couples to matter, f is a sterile metric and does not couple to matter. In bigravity, gravitational interactions are mediated by two gravitons, one of which has mass and the other of which is massless. The two gravitons are composed of different combinations (or superpositions) of the g and f metrics, and so they couple to the surrounding

matter in different ways. The existence of two metrics (and two gravitons) in the bigravity framework eventually leads to the oscillation phenomenon.

As the physicists explain, the idea that there might exist a graviton with mass has been around since almost as long general relativity itself.

"Einstein's theory of general relativity predicts one mediator (the 'graviton') of the gravitational interactions, which travels at the speed of light, i.e., which is massless," Max said. "Back in the late 1930s, people were already trying to find a theory containing a mediator that has a mass, and thus travels at a speed less than the speed of light. This turned out to be a very difficult task and was only recently accomplished in 2010. Bigravity is a variation of this 2010 framework, which features not one, but two dynamical metrics. Only one of them couples to matter while the other doesn't; and a linear combination of them becomes massive (slower than the speed of light) while the other is massless (speed of light)."

Oscillations

The physicists show that, in the framework of bigravity, as gravitational waves are produced and propagate through space, they oscillate between the g- and f-types—though only the g-type can be detected. Although previous research has suggested that these oscillations might exist, it appeared to lead to unphysical results, such as a violation of energy conservation. The new study shows that the oscillations can theoretically emerge in a realistic physical scenario when considering graviton masses that are large enough to be detected by current astrophysical tests.

In order to understand these oscillations, the scientists explain that in many ways they resemble neutrino oscillations. Although neutrinos come in three flavors (electron, muon, and tau), typically the neutrinos produced in nuclear reactions are electron neutrinos (or electron anti-neutrinos) because the others are too heavy to form stable matter. In a similar way, in bigravity only the g metric couples to matter, so the gravitational waves produced by astrophysical events, such as black hole mergers, are g-type since f-type gravitational waves do not couple to matter.

"The key to understanding the oscillation phenomenon is that electron neutrinos do not have a definite mass: they are a superposition of the three neutrino mass eigenstates," Platscher explained. "More mathematically speaking, the mass matrix is not diagonal in the flavor (electron-muon-tau) basis. Therefore, the wave equation that describes how they move through space will mix them up and therefore they 'oscillate.'"

"The same is true in bigravity: g is a mixture of the massive and the massless graviton, and therefore as the gravitational wave travels through the Universe, it will oscillate between g- and f-type gravitational waves. However, we can only measure the former with our detectors (which are made of matter), while the latter would pass through us unseen! This would, if bigravity is a correct description of Nature, leave an important imprint in the gravitational wave signal, as we have shown."

As the physicists note, the similarity between neutrinos and gravitational waves holds even though neutrino oscillation is a quantum mechanical phenomenon that is described by the Schrödinger wave equation, whereas gravitational wave oscillation is not a quantum effect and instead is described by a classical wave equation.

One particular effect that the physicists predict is that gravitational wave oscillations lead to larger strain modulations compared to those predicted by general relativity. These results suggest a path toward experimentally detecting gravitational wave oscillations and finding support for bigravity.

"Since bigravity is a very young theory, there is still a lot to be done, and its potential to address our theories' shortcomings needs to be explored," Smirnov said. "There has been some work along these lines, but certainly a lot is yet to be done and we hope to contribute in the future as well!"
[7]

Quest to settle riddle over Einstein's theory may soon be over

Astronomy experiments could soon test an idea developed by Albert Einstein almost exactly a century ago, scientists say.

Tests using advanced technology could resolve a longstanding puzzle over what is driving the accelerated expansion of the Universe.

Researchers have long sought to determine how the Universe's accelerated expansion is being driven. Calculations in a new study could help to explain whether dark energy- as required by Einstein's theory of general relativity - or a revised theory of gravity are responsible.

Einstein's theory, which describes gravity as distortions of space and time, included a mathematical element known as a Cosmological Constant. Einstein originally introduced it to explain a static universe, but discarded his mathematical factor as a blunder after it was discovered that our Universe is expanding.

Research carried out two decades ago, however, showed that this expansion is accelerating, which suggests that Einstein's Constant may still have a part to play in accounting for dark energy. Without dark energy, the acceleration implies a failure of Einstein's theory of gravity across the largest distances in our Universe.

Scientists from the University of Edinburgh have discovered that the puzzle could be resolved by determining the speed of gravity in the cosmos from a study of gravitational waves -space-time ripples propagating through the universe.

The researchers' calculations show that if gravitational waves are found to travel at the speed of light, this would rule out alternative gravity theories, with no dark energy, in support of Einstein's Cosmological Constant. If however, their speed differs from that of light, then Einstein's theory must be revised.

Such an experiment could be carried out by the Laser Interferometer Gravitational-Wave Observatory (LIGO) in the US, whose twin detectors, 2000 miles apart, directly detected gravitational waves for the first time in 2015.

Experiments at the facilities planned for this year could resolve the question in time for the 100th anniversary of Einstein's Constant.

The study, published in *Physics Letters B*, was supported by the UK Science Technology Facilities Council, the Swiss National Science Foundation, and the Portuguese Foundation of Science and Technology.

Dr Lucas Lombriser, of the University of Edinburgh's School of Physics and Astronomy, said: "Recent direct gravitational wave detection has opened up a new observational window to our Universe. Our results give an impression of how this will guide us in solving one of the most fundamental problems in physics." [6]

NEW THEORY OF GRAVITY DOES AWAY WITH NEED FOR DARK MATTER

Let's be honest. Dark matter's a pain in the butt. Astronomers have gone to great lengths to explain why it must exist and exist in huge quantities, yet it remains hidden. Unknown. Emitting no visible energy yet apparently strong enough to keep galaxies in clusters from busting free like wild horses, it's everywhere in vast quantities. What is the stuff – axions, WIMPS, gravitinos, Kaluza Klein particles?

It's estimated that 27% of all the matter in the universe is invisible, while everything from PB&J sandwiches to quasars accounts for just 4.9%. But a new theory of gravity proposed by theoretical physicist Erik Verlinde of the University of Amsterdam found out a way to dispense with the pesky stuff.

Unlike the traditional view of gravity as a fundamental force of nature, Verlinde sees it as an emergent property of space. Emergence is a process where nature builds something large using small, simple pieces such that the final creation exhibits properties that the smaller bits don't. Take a snowflake. The complex symmetry of a snowflake begins when a water droplet freezes onto a tiny dust particle. As the growing flake falls, water vapor freezes onto this original crystal, naturally arranging itself into a hexagonal (six-sided) structure of great beauty. The sensation of temperature is another emergent phenomenon, arising from the motion of molecules and atoms.

So too with gravity, which according to Verlinde, emerges from entropy. We all know about entropy and messy bedrooms, but it's a bit more subtle than that. Entropy is a measure of disorder in a system or put another way, the number of different microscopic states a system can be in. One of the coolest descriptions of entropy I've heard has to do with the heat our bodies radiate. As that energy dissipates in the air, it creates a more disordered state around us while at the same time decreasing our own personal entropy to ensure our survival. If we didn't get rid of body heat, we would eventually become disorganized (overheat!) and die.

The more massive the object, the more it distorts space-time, shown here as the green mesh. Earth orbits the Sun by rolling around the dip created by the Sun's mass in the fabric of space-time. It doesn't fall into the Sun because it also possesses forward momentum. Credit: LIGO/T. Pyle

Emergent or entropic gravity, as the new theory is called, predicts the exact same deviation in the rotation rates of stars in galaxies currently attributed to dark matter. Gravity emerges in Verlinde's view from changes in fundamental bits of information stored in the structure of space-time, that four-dimensional continuum revealed by Einstein's general theory of relativity. In a word, gravity is a consequence of entropy and not a fundamental force.

Space-time, comprised of the three familiar dimensions in addition to time, is flexible. Mass warps the 4-D fabric into hills and valleys that direct the motion of smaller objects nearby. The Sun doesn't so much "pull" on the Earth as envisaged by Isaac Newton but creates a great pucker in space-time that Earth rolls around in.

In a 2010 article, Verlinde showed how Newton's law of gravity, which describes everything from how apples fall from trees to little galaxies orbiting big galaxies, derives from these underlying microscopic building blocks.

His latest paper, titled *Emergent Gravity and the Dark Universe*, delves into dark energy's contribution to the mix. The entropy associated with dark energy, a still-unknown form of energy responsible for the accelerating expansion of the universe, turns the geometry of spacetime into an elastic medium.

"We find that the elastic response of this 'dark energy' medium takes the form of an extra 'dark' gravitational force that appears to be due to 'dark matter,'" writes Verlinde. "So the observed dark matter phenomena is a remnant, a memory effect, of the emergence of spacetime together with the ordinary matter in it."

Rotation curve of the typical spiral galaxy M 33 (yellow and blue points with errorbars) and the predicted one from distribution of the visible matter (white line). The discrepancy between the two curves is accounted for by adding a dark matter halo surrounding the galaxy. Credit: Public domain / Wikipedia

This diagram shows rotation curves of stars in M33, a typical spiral galaxy. The vertical scale is speed and the horizontal is distance from the galaxy's nucleus. Normally, we expect stars to slow down the farther they are from galactic center (bottom curve), but in fact they revolve much faster (top curve). The discrepancy between the two curves is accounted for by adding a dark matter halo surrounding the galaxy.

I'll be the first one to say how complex Verlinde's concept is, wrapped in arcane entanglement entropy, tensor fields and the holographic principle, but the basic idea, that gravity is not a fundamental force, makes for a fascinating new way to look at an old face.

Physicists have tried for decades to reconcile gravity with quantum physics with little success. And while Verlinde's theory should be rightly be taken with a grain of salt, he may offer a way to combine the two disciplines into a single narrative that describes how everything from falling apples to black holes are connected in one coherent theory. [5]

Identification of a Gravitational Arrow of Time

The proposal by the trio though phrased in a way as to suggest it's a solution to the arrow of time problem, is not likely to be addressed as such by the physics community—it's more likely to be considered as yet another theory that works mathematically, yet still can't answer the basic question of what is time.

For all the advances made in understanding the world around us, there are still two very basic fundamental concepts that have defied explanation: time and gravity. Though we have progressed

greatly in measuring both and using both to understand other concepts, we still today are no closer to understanding either than we were when we first conceptualized them. Such an acknowledgment suggests that we likely have a major flaw in our understanding of the universe. In considering such a possibility, the three physicists with this new effort suggest we might look at time in a completely new way—by dividing a dynamically closed universe (ala the Newtonian N-body problem) into two halves with shape complexity growing from a single point—each solution to the problem can then be considered as having one past but two distinctly futures. In such a scenario, an observer would of necessity have to exist on one side or the other, and thus would only ever have that perspective. Critical to this idea is that the all of the energy and angular momentum in such a system would have to be zero.

In essence, the team has removed time from mathematical functions that describe the energy of the universe—that's what allows for splitting the equations that have been created to describe the evolution of the universe into two parts, with both having initial low complexity moving to higher complexity (similar in some respects to theories of time based on entropy).

The proposal by the trio though phrased in a way as to suggest it's a solution to the arrow of time problem, is not likely to be addressed as such by the physics community—it's more likely to be considered as yet another theory that works mathematically, yet still can't answer the basic question of what is time. [4]

Asymmetry in the interference occurrences of oscillators

The asymmetrical configurations are stable objects of the real physical world, because they cannot annihilate. One of the most obvious asymmetry is the proton – electron mass rate $M_p = 1840 M_e$ while they have equal charge. We explain this fact by the strong interaction of the proton, but how remember it his strong interaction ability for example in the H – atom where are only electromagnetic interactions among proton and electron.

This gives us the idea to origin the mass of proton from the electromagnetic interactions by the way interference occurrences of oscillators. The uncertainty relation of Heisenberg makes sure that the particles are oscillating.

The resultant intensity due to n equally spaced oscillators, all of equal amplitude but different from one another in phase, either because they are driven differently in phase or because we are looking at them an angle such that there is a difference in time delay:

$$(1) \quad I = I_0 \sin^2 n \varphi/2 / \sin^2 \varphi/2$$

If φ is infinitesimal so that $\sin\varphi = \varphi$ than

$$(2) \quad I = n^2 I_0$$

This gives us the idea of

$$(3) \quad M_p = n^2 M_e$$




Fig. 30-3. A linear array of n equal oscillators, driven with phases $\alpha_s = s\alpha$.

Figure 1.) A linear array of n equal oscillators

There is an important feature about formula (1) which is that if the angle φ is increased by the multiple of 2π it makes no difference to the formula.

So

$$(4) \quad d \sin \theta = m \lambda \text{ and we get } m\text{-order beam if } \lambda \text{ less than } d. [6]$$

If d less than λ we get only zero-order one centered at $\theta = 0$. Of course, there is also a beam in the opposite direction. The right choices of d and λ we can ensure the conservation of charge.

For example

$$(5) \quad 2(m+1) = n$$

Where $2(m+1) = N_p$ number of protons and $n = N_e$ number of electrons.

In this way we can see the H_2 molecules so that $2n$ electrons of n radiate to $4(m+1)$ protons, because $d_e > \lambda_e$ for electrons, while the two protons of one H_2 molecule radiate to two electrons of them, because of $d_e < \lambda_e$ for this two protons.

To support this idea we can turn to the Planck distribution law, that is equal with the Bose – Einstein statistics.

Spontaneously broken symmetry in the Planck distribution law

The Planck distribution law is temperature dependent and it should be true locally and globally. I think that Einstein's energy-matter equivalence means some kind of existence of electromagnetic oscillations enabled by the temperature, creating the different matter formulas, atoms molecules, crystals, dark matter and energy.

Max Planck found for the black body radiation

As a function of wavelength (λ), Planck's law is written as:

$$B_{\lambda}(T) = \frac{2hc^2}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda k_B T}} - 1}$$




Figure 2. The distribution law for different T temperatures

We see there are two different λ_1 and λ_2 for each T and intensity, so we can find between them a d so that $\lambda_1 < d < \lambda_2$.

We have many possibilities for such asymmetrical reflections, so we have many stable oscillator configurations for any T temperature with equal exchange of intensity by radiation. All of these configurations can exist together. At the λ_{\max} is the annihilation point where the configurations are symmetrical. The λ_{\max} is changing by the Wien's displacement law in many textbooks.

$$(7) \quad \lambda_{\max} = \frac{b}{T}$$

where λ_{\max} is the peak wavelength, T is the absolute temperature of the black body, and b is a constant of proportionality called *Wien's displacement constant*, equal to $2.8977685(51) \times 10^{-3} \text{ m} \cdot \text{K}$ (2002 CODATA recommended value).

By the changing of T the asymmetrical configurations are changing too.

The structure of the proton

We must move to the higher T temperature if we want look into the nucleus or nucleon arrive to $d < 10^{-13} \text{ cm}$. If an electron with $\lambda_e < d$ move across the proton then by (5) $2(m+1) = n$ with $m = 0$ we get $n = 2$ so we need two particles with negative and two particles with positive charges. If the proton can fraction to three parts, two with positive and one with negative charges, then the reflection of oscillators are right. Because this very strange reflection where one part of the proton with the electron together on the same side of the reflection, the all parts of the proton must be quasi lepton so $d > \lambda_q$. One way dividing the proton to three parts is, dividing his oscillation by the three direction of the space. We can order $1/3$ e charge to each coordinates and $2/3$ e charge to one plane oscillation, because the charge is scalar. In this way the proton has two $+2/3$ e plane oscillation and one linear oscillation with $-1/3$ e charge. The colors of quarks are coming from the three directions of coordinates and the proton is colorless. The flavors of quarks are the possible oscillations differently by energy and if they are plane or linear oscillations. We know there is no possible reflecting two oscillations to each other which are completely orthogonal, so the quarks never can be free, however there is an asymptotic freedom while their energy are increasing to turn them to the orthogonally. If they will be completely orthogonal then they lose this reflection and take new partners from the vacuum. Keeping the symmetry of the vacuum the new oscillations are keeping all the conservation laws, like charge, number of baryons and leptons. The all features of gluons are coming from this model. The mathematics of reflecting oscillators show Fermi statistics.

Important to mention that in the Deuteron there are 3 quarks of $+2/3$ and $-1/3$ charge, that is three u and d quarks making the complete symmetry and because this its high stability.

The Pauli Exclusion Principle says that the diffraction points are exclusive!

The Weak Interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse order, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a $1/2$ spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $1/2$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T-symmetry breaking. This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's

interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating; it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change. There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

Fermions and Bosons

The fermions are the diffraction patterns of the bosons such a way that they are both sides of the same thing.

The Higgs boson or Higgs particle is a proposed elementary particle in the Standard Model of particle physics. The Higgs boson's existence would have profound importance in particle physics because it would prove the existence of the hypothetical Higgs field - the simplest of several proposed explanations for the origin of the symmetry-breaking mechanism by which elementary particles gain mass. [3]

The fermions' spin

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light.

The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $1/2 \hbar = \Delta x \Delta p$ or $1/2 \hbar = \Delta t \Delta E$, that is the value of the basic energy status.

What are the consequences of this in the weak interaction and how possible that the neutrinos' velocity greater than the speed of light?

The neutrino is the one and only particle doesn't participate in the electromagnetic interactions so we cannot expect that the velocity of the electromagnetic wave will give it any kind of limit.

The neutrino is a $1/2$ spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $1/2$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell-Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The source of the Maxwell equations

The electrons are accelerating also in a static electric current because of the electric force, caused by the potential difference. The magnetic field is the result of this acceleration, as you can see in [2].

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed.

Also an interesting question, how the changing magnetic field creates a negative electric field? The answer also the accelerating electrons will give. When the magnetic field is increasing in time by increasing the electric current, then the acceleration of the electrons will increase, decreasing the charge density and creating a negative electric force. Decreasing the magnetic field by decreasing the electric current will decrease the acceleration of the electrons in the electric current and increases the charge density, creating an electric force also working against the change. In this way we have explanation to all interactions between the electric and magnetic forces described in the Maxwell equations.

The second mystery of the matter is the mass. We have seen that the acceleration change of the electrons in the flowing current causing a negative electrostatic force. This is the cause of the relativistic effect - built-in in the Maxwell equations - that is the mass of the electron growing with its acceleration and its velocity never can reach the velocity of light, because of this growing

negative electrostatic force. The velocity of light is depending only on 2 parameters: the magnetic permeability and the electric permittivity.

There is a possibility of the polarization effect created by electromagnetic forces creates the negative and positive charges. In case of equal mass as in the electron-positron pair it is simply, but on higher energies can be asymmetric as the electron-proton pair of neutron decay by weak interaction and can be understood by the Feynman graphs.

Anyway the mass can be electromagnetic energy exceptionally and since the inertial and gravitational mass are equals, the gravitational force is electromagnetic force and since only the magnetic force is attractive between the same charges, is very important for understanding the gravitational force.

The Uncertainty Relations of Heisenberg gives the answer, since only this way can be sure that the particles are oscillating in some way by the electromagnetic field with constant energies in the atom indefinitely. Also not by chance that the uncertainty measure is equal to the fermions spin, which is one of the most important feature of the particles. There are no singularities, because the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that they product is about the half Planck reduced constant. For the proton this Δx much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greatest proton mass.

The Special Relativity

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed.

The Heisenberg Uncertainty Principle

Moving faster needs stronger acceleration reducing the Δx and raising the Δp . It means also mass increasing since the negative effect of the magnetic induction, also a relativistic effect!

The Uncertainty Principle also explains the proton – electron mass rate since the Δx is much less requiring bigger Δp in the case of the proton, which is partly the result of a bigger mass m_p because of the higher electromagnetic induction of the bigger frequency (impulse).

The Gravitational force

The changing magnetic field of the changing current causes electromagnetic mass change by the negative electric field caused by the changing acceleration of the electric charge.

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass ratio $M_p = 1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass. [1]

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only

experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Casimir effect

The Casimir effect is related to the Zero-point energy, which is fundamentally related to the Heisenberg uncertainty relation. The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $1/2 h = \Delta x \Delta p$ or $1/2 h = \Delta t \Delta E$, that is the value of the basic energy status.

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light. In the atomic scale the Heisenberg uncertainty relation gives the same result, since the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that they product is about the half Planck reduced constant. For the proton this Δx much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greater proton mass. This means that the electron is not a point like particle, but has a real charge distribution.

Electric charge and electromagnetic waves are two sides of the same thing; the electric charge is the diffraction center of the electromagnetic waves, quantified by the Planck constant h .

The Fine structure constant

The Planck constant was first described as the proportionality constant between the energy (E) of a photon and the frequency (ν) of its associated electromagnetic wave. This relation between the energy and frequency is called the **Planck relation** or the **Planck–Einstein equation**:

$$E = h\nu .$$

Since the frequency ν , wavelength λ , and speed of light c are related by $\lambda\nu = c$, the Planck relation can also be expressed as

$$E = \frac{hc}{\lambda} .$$

Since this is the source of Planck constant, the electric charge countable from the Fine structure constant. This also related to the Heisenberg uncertainty relation, saying that the mass of the proton should be bigger than the electron mass because of the difference between their wavelengths.

The expression of the fine-structure constant becomes the abbreviated

$$\alpha = \frac{e^2}{\hbar c}$$

This is a dimensionless constant expression, $1/137$ commonly appearing in physics literature.

This means that the electric charge is a result of the electromagnetic waves diffractions, consequently the proton – electron mass rate is the result of the equal intensity of the corresponding electromagnetic frequencies in the Planck distribution law, described in my diffraction theory.

Conclusions

In essence, the team has removed time from mathematical functions that describe the energy of the universe—that's what allows for splitting the equations that have been created to describe the evolution of the universe into two parts, with both having initial low complexity moving to higher complexity (similar in some respects to theories of time based on entropy). [4]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

References

- [1] http://www.academia.edu/3834454/3_Dimensional_String_Theory
- [2] http://www.academia.edu/3833335/The_Magnetic_field_of_the_Electric_current
- [3] http://www.academia.edu/4158863/Higgs_Field_and_Quantum_Gravity
- [4] <http://phys.org/news/2014-11-physicists-identification-gravitational-arrow.html> [5] NEW THEORY OF GRAVITY DOES AWAY WITH NEED FOR DARK MATTER
<http://www.universetoday.com/131901/new-theory-of-gravity-does-away-with-need-for-darkmatter/>
- [6] Quest to settle riddle over Einstein's theory may soon be over
<https://phys.org/news/2017-02-quest-riddle-einstein-theory.html>
- [7] Gravitational waves may oscillate, just like neutrinos
<https://phys.org/news/2017-09-gravitational-oscillate-neutrinos.html>
- [8] The current ability to test theories of gravity with black hole shadows
<https://phys.org/news/2018-04-current-ability-theories-gravity-black.html>
- [9] Were gravitational waves from a black-hole–neutron-star merger detected on 14 August?
<https://physicsworld.com/a/were-gravitational-waves-from-a-black-hole-neutron-star-merger-detected-on-14-august/>
- [10] Scientists detect a black hole swallowing a neutron star
<https://phys.org/news/2019-08-scientists-black-hole-swallowing-neutron.html>
- [11] Spotting merging galaxies
<https://phys.org/news/2019-08-merging-galaxies.html>