

A New numerical method for multi-roots finding with the R software

Kouider Mohammed Ridha

Department of Mathematics, Applied Mathematics Laboratory , University of Mohamed Khider, Biskra,
Algeria, mohakouider@gmail.com

Abstract

The most basic problem in numerical analysis (methods) is the root finding problem. In this paper we are interesting in new numerical method which based on the Modified Bisection Method (MBM) referred to by Tanakan superficially and he didn't know her as a numerical method for finding the roots of a function. Hence in this study we define a new numerical method base on MBM with error bound and the number iterations necessary. Finally we present our new MBM for multi-roots with the R software.

Keywords: Nonlinear equation , Linear interpolation , Bisection Method , Modified Bisection Algorithm.

1. Introduction

The process of finding the roots of function is involves finding the value of x for which $f(x)=0$. If the function equals zero, x is the root of the function. The simplest root-finding algorithm is the bisection method is also called the interval halving method. It works when $f(x)$ is a continuous function and it requires previous knowledge of two initial guesses a and b such that $f(a)$ and $f(b)$ have opposite signs, then find the midpoint of $[a, b]$, and then decide whether the root lies on $[a, (a+b)/2]$ or $[(a+b)/2, b]$ repeat until the interval is sufficiently small. However, the bisection method is reliable, but it converges slowly, gaining one bit of accuracy with each iteration. In this paper we interesting in new numerical method based on MBM and like a linear interpolation method is also known as the method of *regula falsi* (false position) see [1], but with other way.

Tanakan, (2013, [3]) in his paper proposed algorithm base in the bisection method and the value of x_s as we defined in (1) and (2) for solve nonlinear equation, but his study did not define the value of x_{s+1} in term of x_s he defined in term of the midpoint of $[a, b]$, and bending the bisection method and the value of x_s , he named modified bisection algorithm [for more detail see Tanakan, (2013, [3])]. Firstly we present the value of x_s and define a new MBM with her algorithm and the error bound with the

iterations necessary for finding the root of function. In section (2) we programming the new MBM for multi-roots finding with the R software

By the intermediate value theorem implies that a number x^* exists in $[a, b]$ with $f(x^*)=0$ if $f(a)f(b)<0$, mean that $f(a)f(b)<0$, which means that one of them is above the x -axis and the other one below the x -axis.

Let $f(x)$ be a continuous function and defined on interval $[a, b]$ with $f(a)f(b)<0$

Firstly, we set $a=\alpha$ and $b=\beta$. For an integer $s \geq 1$. Then, we can find the equation of straight line from the points $(\alpha_s^*, f(\alpha_s^*))$ and $(\beta_s^*, f(\beta_s^*))$ as $y = ax + b$ where

$$a = \frac{f(\beta_s^*) - f(\alpha_s^*)}{\beta_s^* - \alpha_s^*} \text{ or } a = \frac{f(\beta_s^*) - f(\alpha_s^*)}{\beta_s^* - \alpha_s^*}$$

and

$$b = f(\beta_s^*) - a\beta_s^* \text{ or } b = f(\alpha_s^*) - a\alpha_s^*$$

Hence, the x -intercept of the straight line is at a point $x_s = \frac{-b}{a}$ That is

$$x_s = \beta_s^* - f(\beta_s^*) \frac{\beta_s^* - \alpha_s^*}{f(\beta_s^*) - f(\alpha_s^*)} \quad (1)$$

or

$$x_s = \alpha_s^* - f(\alpha_s^*) \frac{\beta_s^* - \alpha_s^*}{f(\beta_s^*) - f(\alpha_s^*)} \quad (2)$$

It's clearly that

$$x_1 = \alpha - f(\alpha) \frac{\beta - \alpha}{f(\beta) - f(\alpha)} \quad (3)$$

or

$$x_1 = \beta - f(\beta) \frac{\beta - \alpha}{f(\beta) - f(\alpha)} \quad (4)$$

By the intermediate value theorem when f is a continuous function and monotone on $[\alpha, \beta]$, we have for $s \geq 1$, $f(\alpha) < f(x_s) < f(\beta)$ (or $f(\alpha) > f(x_s) > f(\beta)$), then we can defined two cases.

For the case1, if $f(\alpha)$ and $f(x_1)$ have opposite signs implies that $f(x_1)$ and $f(\beta)$ have same signs so the work remains on the interval $[\alpha; x_1]$ and for all $x_s \in [\alpha, x_1]$ as defined in the relationship (2) we get $f(\alpha)f(x_s) < 0$ then x_s in (2) becomes defined for $x_0 = \beta$ by

$$x_{s+1} = \alpha - f(\alpha) \frac{x_s - \alpha}{f(x_s) - f(\alpha)} \quad (5)$$

Finally, we choose the new subinterval for the next iteration $[\alpha; x_{s+1}^*] = [\alpha; x_{s+1}]$

We stay in the same process until we find a solution or a approximate value for it as we define in theorem1 .

And the same thing we find in the other case $x_s \in [x_1; \beta]$ for $x_0 = \alpha$ with

$$x_{s+1} = \beta - f(\beta) \frac{\beta - x_s}{f(\beta) - f(x_s)} \quad (6)$$

and we choose the new subinterval for the next iteration $[x_{s+1}^*; \beta] = [x_{s+1}; \beta]$

1.1 Algorithm for a new Modified Bisection Method

The steps to apply the bisection method to find the root of the equation $f(x)=0$ are

- Step1: Choose α, β for $\alpha < \beta$ where $f(\alpha)f(\beta) < 0$. and tolerance 1×10^{-7}
- Step2: Compute

$$x_1 = \alpha - f(\alpha) \frac{\beta - \alpha}{f(\beta) - f(\alpha)}$$

- Step3: If $f(\alpha)f(x_1) < 0$ for $n \geq 1$, compute

$$x_{n+1} = \alpha - f(\alpha) \frac{x_n - \alpha}{f(x_n) - f(\alpha)}$$

ELSE

$$x_{n+1} = \beta - f(\beta) \frac{\beta - x_n}{f(\beta) - f(x_n)}$$

- Step 4: If $|x_{n+1} - x^*| < 1 \times 10^{-7}$, then stop program (i.e $x_{n+1} = x^*$) the zero is x_{n+1} .
ELSE if $f(\alpha)f(x_1) < 0$, $\alpha = \alpha; x_{n+1}^* = x_{n+1}$

else

$$\beta = \beta; x_{n+1}^* = x_{n+1}$$

and set $n = n + 1$, GOTO Step3.

Next we present theorems for the error bound of a new MBM, but before we going to this

theorems let us recall same definitions

We say that a function f defined on interval I to \mathbb{R} admits a maximum in $\gamma \in I$ if for all $x \in I$ we have $f(x) \leq f(\gamma)$ and we say that $f(\gamma)$ is the global maximum of f .

If I has a distance, or a norm , we say that f admits a local maximum, or a relative maximum in $\gamma \in I$ if there is a neighborhood V of γ such that for all $x \in V$, we have $f(x) \leq f(\gamma)$.

Of course, by changing the inequalities of meaning we can define a global minimum and local minimum.

An extremum is maximum or a minimum. The search for extremum is related to the differential calculus. Thus it if f is defined on interval I of \mathbb{R} and if f is differentiable in γ , then

$f'(\gamma) = 0$. The study for the sign of f' on the neighborhood of γ often makes it possible to conclude as to existence of a maximum or minimum.

Theorem 1: Let f be a continuous function and defined on $[\alpha; \beta]$ which $f(\alpha)f(\beta) < 0$. A new Modified Bisection generate s a sequence $\{x_n\}_{n=1}^{\infty}$ approximating a zero x^* of f for

$$\lambda = \frac{f(\gamma_2)}{f(\gamma_1)} < 0 \quad (7)$$

where $f(\gamma_2)$ is a maximum of f in $[\alpha; \beta]$ and $f(\gamma_1)$ is a minimum of f in $[\alpha; \beta]$ respectively

Case1, for $f(\alpha)f(x_1) < 0$

$$|x_n - x^*| < \left| \frac{f(\alpha)}{f(\gamma_1)} \right|^n \frac{\beta - \alpha}{|\lambda - 1|^n} \quad (8)$$

where x_1 defined in(3)

Case2, for $f(\beta)f(x_1) < 0$

$$|x_n - x^*| < \left| \frac{f(\beta)}{f(\gamma_2)} \right|^n \frac{\beta - \alpha}{\left| \frac{1}{\lambda} - 1 \right|^n} \quad (9)$$

where x_1 defined in(4)

Theorem 2: Let f be a continuous and monotone function defined on $[\alpha; \beta]$ with

$$\lambda = \frac{f(\beta)}{f(\alpha)} < 0 \quad (10)$$

where $f(\alpha) \leq f(x) \leq f(\beta)$ as an global extremum for all $x \in [\alpha; \beta]$ then the previous theorem becomes:

Case1, for $f(\alpha)f(x_1) < 0$

$$|x_n - x^*| < \frac{\beta - \alpha}{|\lambda - 1|^n} \quad (11)$$

where x_1 defined in(3)

Case2, for $f(\beta)f(x_1) < 0$

$$|x_n - x^*| < \frac{\beta - \alpha}{\left|\frac{1}{\lambda} - 1\right|^n} \quad (12)$$

In next study we are interesting in theorem2. Going back to the previous theorem2, it's clear that $\lambda < 0$, thus it with first case we noted that if $\lambda = -1$ we have that

$$|x_n - x^*| < \frac{\beta - \alpha}{2^n}$$

which is the error bound of the bisection method, so a new MBM tied to the value of the real number $\lambda \neq 0$ and if $\lambda \in]-1; 0[$ the new MBM is slowly than the bisection method and that's what we have to calculate the value of λ before working with a new MBM. However if $\lambda < -1$ the new MBM is better than the bisection method. Since for the second case if $\lambda = -1$ we have that $|x_n - x^*|$ becomes the error bound of the bisection method and if $\lambda \leq -1$ the new MBM is slowly than the bisection method and the contrary if $\lambda \in]-1; 0[$ the new MBM has better convergence than bisection method. To see how many iterations will be necessary, for $f(\alpha)f(x_1) < 0$ suppose we want to have

$$|x_n - x^*| < \eta_1$$

This will be satisfied if

$$\frac{\beta - \alpha}{|\lambda - 1|^n} < \eta_1$$

Taking logarithms of both sides, we can solve this to give

$$n \geq \frac{\ln\left(\frac{\beta - \alpha}{\eta_1}\right)}{\ln|\lambda - 1|} \quad (13)$$

and for the case where $f(\beta)f(x_1) < 0$

$$n \geq \frac{\ln\left(\frac{\beta - \alpha}{\eta_1}\right)}{\ln\left|\frac{1}{\lambda} - 1\right|} \quad (14)$$

2. Algorithm for a new Modified Bisection Method for multi-roots:

In this section we provide an algorithm that relies on the method of a new MBM which finding the roots of function. This algorithm is with the R software version i386 3.3.3.

The R is an integrated suite of software facilities for data manipulation, calculation and graphical display. R is very much a vehicle for newly developing methods of interactive data analysis. It has developed rapidly, and has been extended by a large collection of packages, many people use R as a statistics system. There

are about 25 packages supplied with R (called "standard" and "recommended" packages) and many more are available through the CRAN family of Internet sites (via <http://CRAN.R-project.org>) and elsewhere.

We can state an alternative a new MBM algorithm for multi-roots as the following ,

- Step1: Divide the interval into several intervals by $h = (b - a) / t$ where $t \in \mathbb{N}$ is the number of intervals for example $t = 10$.
- Step2: For $i = 1, \dots, 10$, and tolerance 1×10^{-7} ; and $\alpha_i = \alpha_i$, $\beta_i = \beta_{i+1}$ where $f(\alpha_i)f(\beta_{i+1}) < 0$.

- Step3: Compute

$$x_1 = \alpha_i - f(\alpha_i) \frac{\beta_{i+1} - \alpha_i}{f(\beta_{i+1}) - f(\alpha_i)}.$$

- Step4: If $f(\alpha_i)f(x_1) < 0$ for $n \geq 1$, compute

$$x_{n+1} = \alpha_i - f(\alpha_i) \frac{x_n - \alpha_i}{f(x_n) - f(\alpha_i)}$$

ELSE

$$x_{n+1} = \beta_{i+1} - f(\beta_{i+1}) \frac{\beta_{i+1} - x_n}{f(\beta_{i+1}) - f(x_n)}$$

- Step 5: If $|x_{n+1} - x^*| < 1 \times 10^{-7}$, then stop program (i.e $x_{n+1} = x^*$) the zero is x_{n+1} . ELSE if $f(\alpha_i)f(x_1) < 0$, $\alpha_i = \alpha_i$; $x_{n+1}^* = x_{n+1}$ else

$$\beta_{i+1} = \beta_{i+1}; x_{n+1}^* = x_{n+1}$$

and set $n = n + 1$, GOTO Step3.

2.1 A new MBM for multi-roots With The R Software:

Next we present our algorithm for multi-roots with the R software as the following ,

```
ridhaRoot<-function(f,a,b,num=10,eps=1e-07)
```

```
{h=abs(b-a)/num
i=0
j=0
a1=b1=0
while(i<=num)
{ a1=a+i*h
b1=a1+h
if(f(a1)==0){print(a1)
print(f(a1))}
else if (f(b1)==0){print(b1)
print(f(b1))}
else if (f(a1)*f(b1)<0){
repeat{ if(abs(b1-a1)<eps) break
m=(b1-a1)/(f(b1)-f(a1))
d=b1-f(b1)*m
if(f(a1)*f(d)<0) b1<-d else a1<-d }
```

```

print(j+1)
j=j+1
print(d)
print(f(d))
}
i=i+1
}
if(j==0)
print("finding root is fail")
else print("finding root is successful")
}

```

Example1: we go to finding the roots of $f(x) = x^2 - 2$ on the interval $[-2;2]$

```

f <- function(x){x^2-2}
ridhaRoot(f,-2,2)
#####
> f <- function(x){x^2-2}
> ridhaRoot(f,-2,2)
[1] 1
[1] -1.414214
[1] 4.440892e-16
[1] 2
[1] 1.414214
[1] 4.440892e-16
[1] "finding root is successful"
#####

```

Example2: We go to finding the roots of $f(x) = x^3 - 2x + 2$ on the interval $[-3;3]$

```

f <- function(x){x^3-2*x+2}
ridhaRoot(f,-3,3)
#####
> f <- function(x){x^3-2*x+2}
> ridhaRoot(f,-3,3)
[1] 1
[1] -1.769292
[1] 0
[1] "finding root is successful"
#####

```

3. Discussion

Hence, by the intermediate value theorem a modified bisection method, it works when f is a continuous function on $[\alpha, \beta]$ with the initial condition $f(\alpha)f(\beta) < 0$ existing with each iteration. And the select the number λ is an important role in credible of a new MBM because if we have $|\lambda - 1| < 2$ with case1 or $\left| \frac{1}{\lambda} - 1 \right| < 2$ with the case2 as we describe in theorem2, the new MBM become useless and the Bisection method is better than a new MBM

with this situation. Unlike this situation the new MBM is better because we fund that have $|\lambda - 1| > 2$ with case1 or $\left| \frac{1}{\lambda} - 1 \right| > 2$ with the case2 and faster convergent than bisection method. And another important point is the interval $[\alpha, \beta]$, increase on $f(\alpha)f(\beta) < 0$ supporting power the new MBM is selecting images $f(\alpha)$ and $f(\beta)$ as an global extremum and this is indicated by the theorem2 that if f is a continuous and monotone on $[\alpha, \beta]$. More than if the difference $|\lambda - 1|$ with case1 or $\left| \frac{1}{\lambda} - 1 \right|$ with the case2 is too large the error bound will become too small and little of the iteration number. Finally, in practice we're out several important points of the application this new numerical method for roots finding of nonlinear equation called Kouider Method (KM) which based on the new MBM. The important point for KM is the choose interval by take the following points, $f(\alpha)f(\beta) < 0$ with $f(\alpha)$ and $f(\beta)$ as an global extremum and more importantly it whenever the big different $|\lambda - 1|$ with case1 or $\left| \frac{1}{\lambda} - 1 \right|$ with the case2 as we define in theorem2, the new MBM become a good.

4. Conclusion

In this study, we presented a new numerical method as we called KM of looking for the roots of a zero equation. In order to apply the KM, correctly on the interval $[\alpha, \beta]$ we have to present the initial condition, is the intermediate value theorem with $f(\alpha)$ and $f(\beta)$ are the global extremum for all $x \in [\alpha, \beta]$, with the important points is to take the greatest value of $f(\beta)$ if the case1 in thoreme2 is investigated or take the smallest value of $f(\beta)$ if the case2 in thoreme2, is investigated as we mentioned to her in the previous element. More precisely, we know that the function is continuous, so to speed up the process of searching for the roots of a function we can take $\lambda < -1$ if the case1 in thoreme2 is investigated and take $-1 < \lambda < 0$ with $\lambda \neq 0$ if the case2 in thoreme2 is investigated and its easy without count the derivative of a function at the reference point, which is not always easy with the error bound of the Newton Raphsone method . In the practice, the initial solution is

really important for the Newton method. But some initial solutions can make the method Newton diverges.

Hence. by KM it works when f is a continuous function on $[\alpha, \beta]$ where $(\alpha, \beta) \in \mathbb{R}^2$ with the initial condition $f(\alpha)f(\beta) < 0$ existing for with $f(\alpha)$ and $f(\beta)$ are the global extremum for all $x \in [\alpha, \beta]$.

5. Proof of the theorem1

Let f is a continuous function on $[\alpha, \beta]$ $f(\alpha)$ and $f(\beta)$ have opposite signs i.e $f(\alpha)f(\beta) < 0$, and from (3) or (4) we have $\alpha < x_1 < \beta$. Then form the new MBM we get two cases as following:

Case1: $f(\alpha)f(x_1) < 0$ then for any $n \geq 0$ $\alpha < x^* < x_n < x_1$, so we have $x_n - x^* < x_n - \alpha < x_1 - \alpha < \beta - \alpha$. Since, we found the terms x_n by(5) with $x_0 = \beta$ as

$$\begin{aligned} x_1 &= \alpha - f(\alpha) \frac{x_0 - \alpha}{f(x_0) - f(\alpha)} \\ x_2 &= \alpha - f(\alpha) \frac{x_1 - \alpha}{f(x_1) - f(\alpha)} \\ x_3 &= \alpha - f(\alpha) \frac{x_2 - \alpha}{f(x_2) - f(\alpha)} \\ &\vdots \\ x_n &= \alpha - f(\alpha) \frac{x_{n-1} - \alpha}{f(x_{n-1}) - f(\alpha)} \end{aligned}$$

by recurrence we can write x_n as

$$x_n = \alpha - f(\alpha)^n \frac{\beta - \alpha}{\prod_{j=0}^{n-1} (f(x_j) - f(\alpha))} \quad (15)$$

then for any $n \geq 0$ and (12) we find

$$x_n - \alpha = -f(\alpha)^n \frac{\beta - \alpha}{\prod_{j=0}^{n-1} (f(x_j) - f(\alpha))}$$

Since $|x_n - x^*| < |x_n - \alpha|$ we have

$$|x_n - x^*| < |f(\alpha)|^n \frac{\beta - \alpha}{\prod_{j=0}^{n-1} |f(x_j) - f(\alpha)|} \quad (16)$$

Suppose that $f(\gamma_1) \leq f(x) \leq f(\gamma_2)$ for all $x \in [\alpha, \beta]$ and $(\gamma_1, \gamma_2) \in [\alpha, \beta]^2$ we have for $j = \{0, 1, \dots, n-1\}$

$$\frac{1}{|f(x_j) - f(\alpha)|} < \frac{1}{|f(\gamma_2) - f(\gamma_1)|} \quad (17)$$

to this end we can get

$$|x_n - x^*| < (\beta - \alpha) \left| \frac{f(\alpha)}{f(\gamma_2) - f(\gamma_1)} \right|^n$$

we potted

$$\lambda = \frac{f(\gamma_2)}{f(\gamma_1)}$$

we find

$$|x_n - x^*| < \left| \frac{f(\alpha)}{f(\gamma_1)} \right|^n \frac{(\beta - \alpha)}{|\lambda - 1|^n}$$

and if $f(\gamma_1) = f(\alpha)$ we have

$$|x_n - x^*| < \frac{(\beta - \alpha)}{|\lambda - 1|^n}$$

Case2: $f(\beta)f(x_1) < 0$ then for any $n \geq 0$

$x_1 < x_n < x^* < \beta$, so we have $x_n - x^* < x_n - \beta < x_1 - \beta < \beta - \alpha$;

Since we found the terms x_n with (6)

and $x_0 = \alpha$, by recurrence we can write x_n as

$$x_n = \beta - f(\beta)^n \frac{\beta - \alpha}{\prod_{j=0}^{n-1} (f(\beta) - f(x_j))} \quad (18)$$

then for any $n \geq 0$ and (12) we find

$$x_n - \beta = -f(\beta)^n \frac{\beta - \alpha}{\prod_{j=0}^{n-1} (f(\beta) - f(x_j))}$$

Since $|x_n - x^*| < |x_n - \alpha|$ and we suppose that

$f(\gamma_1) \leq f(x) \leq f(\gamma_2)$ for all $x \in [\alpha, \beta]$ and

$(\gamma_1, \gamma_2) \in [\alpha, \beta]^2$ we have for $j = \{0, 1, \dots, n-1\}$

$$\frac{1}{|f(\beta) - f(x_j)|} < \frac{1}{|f(\gamma_2) - f(\gamma_1)|} \quad (19)$$

to this end we can get

$$|x_n - x^*| < (\beta - \alpha) \left| \frac{f(\beta)}{f(\gamma_2) - f(\gamma_1)} \right|^n$$

we potted

$$\lambda = \frac{f(\gamma_2)}{f(\gamma_1)}$$

we find

$$|x_n - x^*| < \left| \frac{f(\beta)}{f(\gamma_2)} \right|^n \frac{(\beta - \alpha)}{\left| \frac{1}{\lambda} - 1 \right|^n}$$

and if $f(\gamma_2) = f(\beta)$ we have

$$|x_n - x^*| < \frac{(\beta - \alpha)}{\left| \frac{1}{\lambda} - 1 \right|^n}$$

Reference

- [1] Autar, K.K and Egwu, E (2008), <http://www.numericalmethods.eng.usf.edu> Retrieved on 20th February, 2014.
- [2] Emmanuel Paradis, "R for Beginners", Institut des Sciences de l'Evolution ' Universit'e Montpellier II F-34095 Montpellier c'edex 05 France, 2005 https://cran.r-project.org/doc/contrib/Paradis-rdebuts_en.pdf.
- [3] Ehiwario, J.C., Aghamie, S.O, "Comparative Study

of Bisection, Newton-Raphson and Secant Methods of Root- Finding Problems”, IOSR Journal of Engineering Vol. 04, ||V1|| PP 01-07 , 2014

- [4] http://www.efunda.com/math/num_rootfinding-cfm February,2014
- [5] Jay, Villanueva, “Approximation of equations with a hand-held calculator”, Florida Memorial University Miami, FL, jvillanu@fmuniv.edu. <http://archives.math.utk.edu/ICTCM/VOL23/C012/paper>
- [6] Pierre Lafaye de Micheaux, Remy Drouilhet, Benoit Liquet, “The R Software: Fundamentals of Programming and Statistical Analysis”, Springer. <http://web.maths.unsw.edu.au/~lafaye/textes/book> R2013
- [7] Tanakan ,S,“A New Algorithm of Modified Bisection Method for Nonlinear Equation”, Applied Mathematical Sciences 7.123, 2013, 6107-6114.
- [8] S. Muangchan and S. Sompong, “Technique of Initial Solution Finding for the Newton Method”, Int. J. Contemp. Math. Sciences, Vol. 6, no. 41, 2037-2043 , 2011
- [9] Srivastava,R.B and Srivastava, S, “Comparison of Numerical Rate of Convergence of Bisection, Newton and Secant Methods”, Journal of Chemical, Biological and Physical Sciences. Vol 2(1) pp 472- 479 , 2011.
- [10] Y. Cohen, J. Cohen ,“Statistics and Data with R: An Applied Approach Through Examples”,. Wiley, 2008.
- [11] Wikipedia: The Free Encyclopedia. Wikimedia Foundation Inc. Newton’s method, Available from [http://http://en.wikipedia.org/wiki/Newton’s method](http://http://en.wikipedia.org/wiki/Newton's_method), Retrieved 5 September 2013.