

Radiation damping of a rotating dipole

Optik Available online 9 December 2019, 164021

The article has been repeatedly rejected without consideration.

JETP Letters

Our Editorial Board has considered your paper "Radiation damping of a rotating dipole" on 3.10.2019 and come to the conclusion that it cannot be accepted by JETP Letters as not requiring rapid publication.

Yours sincerely, Managing Editor I. Podyniglazova

УФН

№ 6655

09 октября 2019 г.

Храпко Р.И.

Уважаемый Радий Игоревич!

Ваша статья «Торможение излучением вращающегося диполя» была рассмотрена членами редколлегии УФН.

Как Вы знаете, журнал УФН – обзорный, и тем самым он заметно отличается от специализированных журналов, публикующих оригинальные (новые) научные результаты. О специфических требованиях УФН Вы можете прочитать в 1-м выпуске УФН за 2013 г. («От редакционной коллегии»). Там сказано, в частности, что редколлегия может отклонять даже вполне «доброкачественные» статьи, как не прошедшие по конкурсу.

В настоящее время в УФН *очень большой портфель уже одобренных рукописей* (см. <https://ufn.ru/ru/articles/accepted/>), что вынуждает нас особо строго относиться к отбору статей, которые мы можем рекомендовать в печать, причём исключительно на конкурсной основе.

По этим причинам было принято решение отказаться от публикации Вашей статьи в журнале УФН. Рекомендуем Вам направить статью в соответствующий специализированный журнал.

С уважением,
Заместитель главного
редактора журнала
«Успехи физических наук»

академик РАН



О.В. Руденко

Physical Review Letters

LJ16511. Your manuscript has been considered. We regret to inform you that we have concluded that it is not suitable for publication in Physical Review Letters.

Yours sincerely, **Serena Dalena** Associate Editor Physical Review Letters Email: prl@aps.org

RE: From Khrapko Radi

Dear **Serena Dalena**,

Your conclusion contradicts the fact that “The Physical Review editors attempt to accept papers that are scientifically sound, important to the field, and contain significant new results in physics”, since you do not deny that my article proves that the spin of an electromagnetic radiation is actually described by the spin tensor and confirms the presence of the spin radiation by a rotating dipole, the radiation, which not seen before.

Your unsubstantiated conclusion confirms that the Physical Review Letters reviewers unable to review articles on the spin tensor.

Optics Letters

ID: 378337 Your recent submission to Optics Letters has not been received successfully. I regret to inform you that The Optical Society is unable to consider your recent submission to Optics Letters for publication.

Sincerely, Optics Letters Manuscripts Office olmss@osa.org

Journal of Optics

JOPT-106905 We have assessed your manuscript and have considered its suitability for the journal very carefully. We regret to inform you that your article will not be considered for review as it does not meet our strict publication criteria.

Editor-in-Chief: L. N. Hazra

JOSA B submission (380469)

Your recent submission to JOSA B has not been received successfully. I regret to inform you that The Optical Society is unable to consider your recent submission to JOSA B for publication.

Sincerely, JOSA B Manuscripts Office jbmss@osa.org

RE: JOSA B submission (380469)

From: Khrapko Radi

Dear JOSA B Manuscripts Office, Your letter contradicts the previous one (see below). What is the reason for the refusal?

JOSA B resubmission complete (380469)

Thank you for your resubmission to the Journal of the Optical Society of America B. Your manuscript is being checked again and prepared for the review process. Your manuscript ID appears above.

Sincerely, JOSA B Manuscripts Office




ELSEVIER

Optik

Available online 9 December 2019, 164021

[In Press, Journal Pre-proof](#) [What are Journal Pre-proof articles?](#)



Radiation damping of a rotating dipole

Author links open overlay panel [Radi. I.Khrapko](#)¹

Moscow Aviation Institute - Volokolamskoe shosse 4, 125993 Moscow, Russia

Received 30 October 2019, Accepted 6 December 2019, Available online 9 December 2019.

<https://doi.org/10.1016/j.ijleo.2019.164021>

Abstract

Jefimenko's generalization of the Coulomb and Bio-Savard laws is used to calculate the reaction of the radiation on the rotating electric dipole. It is found that the energy taken from the dipole is equal to the recognized value of the radiated energy. At the same time, it is confirmed that the angular momentum flux exceeds the generally accepted value by the spin radiation not seen before.

PACS 03.50.De **Keywords** classical spin Jefimenko's generalization spin tensor

1. Introduction

As shown in articles [1–4], the radiation of a rotating electric dipole contains, in addition to the generally recognized angular momentum flux [1 - 9]

$$dL_z / dt = \omega^3 p^2 / 6\pi\epsilon_0 c^3, \quad (1.1)$$

which is localized mainly near the equatorial plane. with the angular distribution

$$dL_z / dt d\Omega = \omega^3 p^2 \sin^2 \theta / 16\pi^2 \epsilon_0 c^3, \quad (1.2)$$

also the spin flux

$$dS_z / dt = \omega^3 p^2 / 12\pi\epsilon_0 c^3, \quad (1.3)$$

which is localized mainly near the axis of rotation of the dipole, with the angular distribution

$$dS_z / dt d\Omega = \omega^3 p^2 \cos^2 \theta / 16\pi^2 \epsilon_0 c^3 \quad (1.4)$$

($d\Omega = \sin \theta d\theta d\varphi$, and the unit system is used in which $\text{div}\mathbf{E} = \rho / \epsilon_0$).

Thus, the total angular momentum flux is

$$dJ_z / dt = dL_z / dt + dS_z / dt = \omega^3 p^2 / 4\pi\epsilon_0 c^3. \quad (1.5)$$

Meanwhile, the power radiated by a rotating dipole is recognized as

$$P = \omega^4 p^2 / 6\pi\epsilon_0 c^3, \quad (1.6)$$

and, therefore, the usual mechanics equality

$$P = \omega dJ / dt. \quad (1.7)$$

is violated

The result (1.5) was obtained in [2 - 4] by integrating the sum of the spin tensor and the moment of the Poynting vector over the surface surrounding a rotating dipole. In [1], this result was confirmed using the quantum mechanical Feynman proof and, independently, calculating the reaction of the magnetic vector potential field on the dipole.

In the present work, the energy (1.6) and angular momentum (1.5) fluxes are calculated using the same type by the use of the retarded electromagnetic Jefimenko's field [10]. It turned out that the forces acting on the dipole and responsible for the loss of energy differ from the forces responsible for the loss of angular momentum.

As a rotating electric dipole, we consider a pair of oscillating dipoles perpendicular to each other and having a quarter-period oscillation shift in time,

$$p^x = p \exp(-i\omega t), \quad p^y = ip \exp(-i\omega t), \quad (1.8)$$

2. Energy loss by a rotating dipole

The value (1.6) is calculated as the result of the influence of the electromagnetic field of the dipole on the dipole itself, according to the formula for the density of the resulting power

$$P_\wedge = -(\mathbf{j} \cdot \mathbf{E}); \quad (2.1)$$

here \mathbf{j} and \mathbf{E} are current density flowing along the dipole and the electric field in the dipole, respectively, and the index \wedge for P_\wedge means, in this case, "volume density", $dP = P_\wedge d^3x$. First, the effect of the x -dipole on itself is calculated.

In this paper, the electric field near the dipole is calculated from the known formula, taking into account the retardation [10 (6.55)]:

$$\mathbf{E}(\mathbf{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3x' \left\{ \frac{\hat{\mathbf{r}}}{r^2} [\rho(\mathbf{x}', t')_{\text{ret}}] + \frac{\hat{\mathbf{r}}}{cr} [\partial_{t'} \rho(\mathbf{x}', t')_{\text{ret}}] - \frac{1}{c^2 r} [\partial_{t'} \mathbf{j}(\mathbf{x}', t')_{\text{ret}}] \right\}, \quad (2.2)$$

and an "elementary vibrator" is considered as a dipole; the current of the dipole is the same at all points, and the charges are only at the ends (see Figure 1)




Figure 1. x -dipole

The dipole current I_x is obtained by differentiating the charge

$$p^x = ql \exp(-i\omega t), \quad (2.3)$$

$$I_x = \partial_t p^x / l = -i\omega q \exp(-i\omega t). \quad (2.4)$$

The first term of expression (2.2),

$$E_1^x = \frac{1}{4\pi\epsilon_0} \int d^3x' \left\{ \frac{1}{r^2} [\rho(\mathbf{x}', t')_{\text{ret}}] \right\},$$

is simply the retarded Coulomb field at the point x . Therefore, replacing

$$d^3x' \rho \rightarrow d\tilde{q}', \quad t \rightarrow t - (l/2 \pm x)/c, \quad r \rightarrow l/2 \pm x$$

and taking into account the direction of the electric field, we obtain the electric field strength from both charges at the point x :

$$E_1^x(x) = \frac{q}{4\pi\epsilon_0} \left[\frac{-\exp[-i\omega t + i\omega(l/2 + x)/c]}{(l/2 + x)^2} - \frac{\exp[-i\omega t + i\omega(l/2 - x)/c]}{(l/2 - x)^2} \right]. \quad (2.5)$$

The corresponding contribution of this term to the power generated by the dipole is given by formula (2.1) (we replace $jd^3x \rightarrow Idx$ from (5), and the bar means complex conjugation)

$$\begin{aligned} P_1 &= \frac{-1}{2} \int_{-l/2}^{l/2} dx \Re\{\bar{I} E_1^x\} \\ &= -\frac{\omega q^2}{8\pi\epsilon_0} \int_{-l/2}^{l/2} dx \Re\left\{ i \exp(i\omega t) \left[\frac{-\exp[-i\omega t + i\omega(l/2 + x)/c]}{(l/2 + x)^2} - \frac{\exp[-i\omega t + i\omega(l/2 - x)/c]}{(l/2 - x)^2} \right] \right\} \\ &= -\frac{\omega q^2}{8\pi\epsilon_0} \int_{-l/2}^{l/2} dx \left[\frac{\sin[\omega(l/2 + x)/c]}{(l/2 + x)^2} + \frac{\sin[\omega(l/2 - x)/c]}{(l/2 - x)^2} \right]. \end{aligned} \quad (2.6)$$

Taking into account the small size of the dipole, we consider only two terms of the expansion of the sine in a series

$$\begin{aligned} P_1 &= -\frac{\omega q^2}{8\pi\epsilon_0} \int_{-l/2}^{l/2} dx \left[\frac{\omega}{c(l/2 + x)} - \frac{\omega^3(l/2 + x)}{6c^3} + \frac{\omega}{c(l/2 - x)} - \frac{\omega^3(l/2 - x)}{6c^3} \right] \\ P_1 &= -\frac{\omega q^2}{8\pi\epsilon_0} \int_{-l/2}^{l/2} dx \left[\frac{\omega}{c(l/2 + x)} - \frac{\omega^3(l/2 + x)}{6c^3} + \frac{\omega}{c(l/2 - x)} - \frac{\omega^3(l/2 - x)}{6c^3} \right]. \end{aligned} \quad (2.7)$$

Similarly to formula (2.5), we find the electric field provided by the second term of formula (2.2)

$$E_2^x = \frac{i\omega q}{4\pi\epsilon_0 c} \left[\frac{\exp[-i\omega t + i\omega(l/2 + x)/c]}{(l/2 + x)} + \frac{\exp[-i\omega t + i\omega(l/2 - x)/c]}{(l/2 - x)} \right]. \quad (2.8)$$

In contrast to formula (2.5), this formula contains i .

Formula (2.1) gives the contribution of this term, E_2^x , to the power generated by the dipole

$$\begin{aligned} P_2 &= \frac{\omega^2 q^2}{8\pi\epsilon_0 c} \int_{-l/2}^{l/2} dx \Re\left\{ \exp(i\omega t) \left[\frac{\exp[-i\omega t + i\omega(l/2 + x)/c]}{(l/2 + x)} + \frac{\exp[-i\omega t + i\omega(l/2 - x)/c]}{(l/2 - x)} \right] \right\} \\ &= \frac{\omega^2 q^2}{8\pi\epsilon_0 c} \int_{-l/2}^{l/2} dx \left[\frac{\cos[\omega(l/2 + x)/c]}{(l/2 + x)} + \frac{\cos[\omega(l/2 - x)/c]}{(l/2 - x)} \right]. \end{aligned} \quad (2.9)$$

Restricting ourselves to two terms of the expansion of cosine in a series, we have

$$P_2 = \frac{\omega^2 q^2}{8\pi\epsilon_0 c} \int_{-l/2}^{l/2} dx \left[\frac{1}{(l/2 + x)} - \frac{\omega^2(l/2 + x)}{2c^2} + \frac{1}{(l/2 - x)} - \frac{\omega^2(l/2 - x)}{2c^2} \right]. \quad (2.10)$$

Surprisingly, the integrals diverging at the ends of the dipole are shortened upon the addition $P_1 + P_2$, and the remaining terms are constants. As $l/6 - l/2 = -l/3$, this part of the power is

$$P_1 + P_2 = -\frac{\omega^4 p^2}{24\pi\epsilon_0 c^3}. \quad (2.11)$$

The third term of formula (2.2),

$$E_3^x = \frac{1}{4\pi\epsilon_0} \int dx' \left\{ -\frac{1}{c^2 r} [\partial_{t'} I(x', t')]_{\text{ret}} \right\},$$

uses the derivative of the current

$$\partial_{t'} I_x = -\omega^2 q \exp(-i\omega t). \quad (2.12)$$

To calculate the strength at the point x , we divided the region of integration into two parts by the point x

$$E_3^x = -\frac{\omega^2 q}{4\pi\epsilon_0 c^2} \left\{ \int_{-l/2}^x dx' \frac{-\exp[-i\omega t + i\omega(x-x')/c]}{x-x'} + \int_x^{l/2} dx' \frac{-\exp[-i\omega t + i\omega(x'-x)/c]}{x'-x} \right\}. \quad (2.13)$$

Using formula (2.1), $dP = -(\mathbf{j} \cdot \mathbf{E})d^3x = -IE^x dx$, and current (2.4), we obtain the power corresponding to the third term of formula (2.2)

$$\begin{aligned} P_3 &= -\frac{\omega^3 q^2}{8\pi\epsilon_0 c^2} \int_{-l/2}^{l/2} dx \Re \left\{ i\omega \exp(i\omega t) \left[\int_{-l/2}^x dx' \frac{\exp[-i\omega t + i\omega(x-x')/c]}{x-x'} + \int_x^{l/2} dx' \frac{\exp[-i\omega t + i\omega(x'-x)/c]}{x'-x} \right] \right\} \\ &= -\frac{\omega^3 q^2}{8\pi\epsilon_0 c^2} \int_{-l/2}^{l/2} dx \left[\int_{-l/2}^x dx' \frac{-\sin[\omega(x-x')/c]}{x-x'} + \int_x^{l/2} dx' \frac{-\sin[\omega(x'-x)/c]}{x'-x} \right]. \end{aligned} \quad (2.14)$$

Restricting ourselves to one term in the expansion of the sine in a series, we easily obtain

$$P_3 = \frac{\omega^4 d^2}{8\pi\epsilon_0 c^3}. \quad (2.15)$$

Thus, the power radiated by one x -dipole is

$$P = P_1 + P_2 + P_3 = \omega^4 p^2 / (12\pi\epsilon_0 c^3). \quad (2.16)$$

Naturally, the y -dipole, acting on itself, makes the same contribution. It will be shown below that the electric field of one oscillating dipole in the territory of another oscillating dipole is perpendicular to the current, and therefore does not produce energy. So a *rotating* dipole delivers power (1.6) to the radiation:

$$P = \omega^4 p^2 / 6\pi\epsilon_0 c^3.$$

3. The torque experienced by the charges of a rotating dipole

We now calculate the electric field \mathbf{E} created by the y -dipole at the location of the charge of the x -dipole, that is, at the point $x = l/2$ (see Fig. 2)




Fig. 2. The pair of oscillating dipoles. Current elements and radii used in the formulas are indicated.

To shorten the notation, we rewrite formula (2.2) in terms of charges and currents:

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \left(\frac{\hat{\mathbf{r}}}{r^2} [q_y]_{\text{ret}} + \frac{\hat{\mathbf{r}}}{cr} [\partial_t q_y]_{\text{ret}} - \int_{-l/2}^{l/2} \frac{1}{c^2 r} [\partial_t I_y]_{\text{ret}} d\mathbf{y} \right); \quad (3.1)$$

here the charge and current belong to the y-dipole:

$$q_y = iq \exp(-i\omega t), \quad [q_y]_{\text{ret}} = iq \exp(-i\omega t + i\omega r/c), \quad [\partial_t q_y]_{\text{ret}} = \omega q \exp(-i\omega t + i\omega r/c) \quad (3.2)$$

$$I_y = \partial_t q_y = \omega q \exp(-i\omega t), \quad [I_y]_{\text{ret}} = \omega q \exp(-i\omega t + i\omega r/c), \quad [\partial_t I_y]_{\text{ret}} = -i\omega^2 q \exp(-i\omega t + i\omega r/c). \quad (3.3)$$

The electric field created by the charge consists of two terms:

$$\mathbf{E}_1 + \mathbf{E}_2 = \hat{\mathbf{r}}([q_y]_{\text{ret}}/r^2 + [\partial_t q_y]_{\text{ret}}/cr)/4\pi\epsilon_0. \quad (3.4)$$

However, the forces created by these terms on the charge $q_x = q \exp(-i\omega t)$ are mutually eliminated when the dipole size tends to zero, although they tend to infinity.

$$\begin{aligned} F_1 + F_2 &= \Re\{(E_1 + E_2)\bar{q}_x\}/2 = \Re\{i \exp(i\omega r/c)/r^2 + \omega \exp(i\omega r/c)/cr\} q^2 / 8\pi\epsilon_0 \\ &= \{-\sin(\omega r/c)/r^2 + \omega \cos(\omega r/c)/cr\} q^2 / 8\pi\epsilon_0 \rightarrow (-\omega/cr + \omega/cr) q^2 / 8\pi\epsilon_0 = 0. \end{aligned} \quad (3.5)$$

Similarly, the total interaction forces of other pairs of charges are zero. So the damping of the rotating dipole is provided only by the third term of the formula (3.1):

$$\mathbf{E}_3 = - \int_{-l/2}^{l/2} \frac{1}{c^2 r} [-i\omega^2 q \exp(-i\omega t + i\omega r/c)] d\mathbf{y} \quad (3.6)$$

The force acting on the charge q_x along the y-axis is

$$\begin{aligned} F_3 &= \Re\{E_3 \bar{q}_x\}/2 = \Re\left\{ \int_{-l/2}^{l/2} \frac{1}{c^2 r} i\omega^2 q^2 \exp(i\omega r/c) dy \right\} / 8\pi\epsilon_0 \\ &= - \int_{-l/2}^{l/2} \frac{1}{c^2 r} \omega^2 q^2 \sin(\omega r/c) dy / 8\pi\epsilon_0 \rightarrow - \int_{-l/2}^{l/2} \omega^3 q^2 dy / 8\pi\epsilon_0 c^3 = -\omega^3 q^2 l / 8\pi\epsilon_0 c^3. \end{aligned} \quad (3.7)$$

The torque acting on both charges of the x-dipole is $-\omega^3 q^2 l^2 / 8\pi\epsilon_0 c^3$. Therefore, the torque acting on the rotating dipole is directed against the rotation of the dipole and is equal to (1.5)

$$\omega^3 p^2 / 4\pi\epsilon_0 c^3.$$

4. Magnetic field torque

In addition to the electric field (2.2), Jefimenko's formulas give the magnetic field [10 (6.56)],

$$\mathbf{B}(\mathbf{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3 x' \left\{ [\mathbf{j}(\mathbf{x}', t')]_{\text{ret}} \times \frac{\hat{\mathbf{r}}}{r^2} + [\partial_t \mathbf{j}(\mathbf{x}', t')]_{\text{ret}} \times \frac{\hat{\mathbf{r}}}{cr} \right\}, \quad (4.1)$$

This field acts on dipoles (1.8) by the Lorentz force. Consider the magnetic field created by the y-dipole on the territory of the x-dipole. By analogy with (3.1), we write

$$\mathbf{B} = \frac{1}{4\pi\epsilon_0} \int \left([I_y]_{\text{ret}} / r^2 + [\partial_t I_y]_{\text{ret}} / cr \right) d\mathbf{y} \times \hat{\mathbf{r}}, \quad (4.2)$$

However, using (3.3) and (2.4), we find that the average value of the Lorentz force acting on the x-dipole is zero:

$$\begin{aligned} \mathbf{F} &= \Re\left\{ \int \bar{\mathbf{I}}_x d\mathbf{x} \times \mathbf{B} \right\} / 2 = \Re\left\{ \iint i\omega q [\omega q \exp(i\omega r/c)/r^2 - i\omega^2 q \exp(i\omega r/c)/cr] d\mathbf{x} \times (d\mathbf{y} \times \hat{\mathbf{r}}) \right\} / 8\pi\epsilon_0 \\ &= \iint [-\sin(\omega r/c)\omega^2 / r^2 + \cos(\omega r/c)\omega^3 / cr] q^2 d\mathbf{x} \times (d\mathbf{y} \times \hat{\mathbf{r}}) / 8\pi\epsilon_0 \rightarrow 0 \quad \text{if } r \rightarrow 0. \end{aligned} \quad (4.3)$$

So the total torque acting on the rotating dipole is (1.5)

5. Conclusion

This calculation confirms the presence of spin radiation by a rotating dipole, the radiation that was predicted using the spin tensor. This proves that the spin of electromagnetic radiation is actually

described by the spin tensor. The spin tensor has been successfully used to describe also plane waves and beams [11–17].

I am eternally grateful to Professor Robert Romer for the courageous publication of my question: "Does a plane wave really not carry spin?" (was submitted on 07 October, 1999) [18].

References

- [1] Khrapko R I Spin radiation from a rotating dipole. *Optik* 181 (2019) 1080-1084
- [2] Khrapko R I. Spin of dipole radiation. (2001) <http://trudymai.ru/published.php?ID=34635>
- [3] Khrapko R I "Radiation of spin by a rotator", <http://www.ma.utexas.edu/cgi-bin/mps?key=03-315> (2003)
- [4] Khrapko R I. Spin is not a moment of momentum <http://trudymai.ru/published.php?ID=28834> (2012)
- [5] Landau L D, Lifshitz E M *The Classical Theory of Fields* (Pergamon: N. Y., 1975). .
- [6] Corney A. *Atomic and Laser Spectroscopy*; (Oxford University Press, 1977).
- [7] Sommerfeld A. *Atombau und Spektrallinien* 1 Band (FR1EDR. V1EWEG & SOHN BRAUNSCHWEIO 1951)
- [8] Vul'fson K S Angular momentum of electromagnetic waves *Sov. Phys. Usp.* 30 724–728 (1987)
- [9] Barabanov A L Angular momentum in classical electrodynamics *Phys. Usp.* 36 (11) 1068–1074 (1993)
- [10] Jackson J D *Classical Electrodynamics*, (John Wiley, 1999).
- [11] Khrapko R. I. Absorption of angular momentum of a plane wave *Optik* 154 (2018) 806–810
- [12] Khrapko R. I. "Reflection of light from a moving mirror" *Optik* **136** (2017) 503–506
- [13] Khrapko R.I. "Mechanical stresses produced by a light beam" *J. Modern Optics*, **55**, 1487-1500 (2008)
- [14] Khrapko R. I. True energy-momentum tensors are unique. Electrodynamics spin tensor is not zero <https://arxiv.org/abs/physics/0102084> (2001).
- [15] Khapko R. I. Spin transferred to a mirror reflecting light. *International Conference on Electrical, Communication, Electronics, Instrumentation and Computing (ICECEIC)* during 30th- 31st January 2019. <http://khrapkori.wmsite.ru/ftpgetfile.php?id=188&module=files>
- [16] Khapko R. I. Absorption of Spin by a Conducting Medium *AASCIT Journal of Physics* Vol. 4, No. 2, Page: 59-63 (2018)
- [17] Khrapko R. I. Origin of Spin: Paradox of the classical Beth experiment. In *Unfolding the Labyrinth: Open Problems in Mathematics, Physics, Astrophysics, and other areas of science* (Hexis - Phoenix 2006), pp. 57-71 <https://arxiv.org/abs/math/0609238>
- [18] Khrapko R. I. Does plane wave not carry a spin? *Amer. J. Phys.* **69**, 405 (2001)