

Quanton based model of field interactions

Ayman Kassem (a) , (b)

(a) aeronautical engineer, Cairo , Egypt

(b) **Contact address** : ayman_kassem@egyptair.com

Abstract

The mechanism of the universe's inflation is variation of energy in

space and in time , the relationship between space and time

varying energy fields is governed by energy constraining inside

what was once a Planck scale entity : the quanton

as energy varies in space or in time, it creates associated fields

and through their interactions, inflationary momentum and the

fundamental forces are generated

this model comes in three parts : energy constraining , where the

evolution of the quanton and its different transitions are discussed

until the stable state is reached

the second part , electromagnetic waves in terms of space and

time varying energy fields and role of Maxwell equations in the

evolution of the quanton

the third part , energy fields and their interactions , while using

basic physics concepts , this model shows that the origin many of

the physical phenomena can be traced back to the quanton based

world

Key words

space and time varying fields , energy degrees of freedom

Contents

1.Physical basis of this model	8
2.Definition of the model	9
3.Mathematical brief	11
4. field variation parameters	13
5.Energy constraining	14
6.Bridging the gap between mathematics and physics in energy constraining	29
7.Energy degrees of freedom	32
8.Superposition principle inside quanton	38
9.Defining directional component of energy fields	40
10.Dimensional energy symmetry	42
11.Energy density / Degree of freedom relationship	45
12.Energy constraining and the release of thermal energy	49
13.Energy constraining: a possible origin of cosmic microwave	

back ground(CMB)	51
14. why do quantons split ?	53
15. Mechanism of quanton splitting and expansion	55
16.mathematics behind expansion / constraining inside the Quanton	61
17.wave like properties of space fabric	65
18. quanton evolution and its degrees of freedom	71
19.variation of energy fields with time	76
20.energy fields parameters	78
21.Dimensions of energy fields	85
22. field definition inside the quanton	88
23.what maintains the integrity of the quanton ?	90
24. Q/AQ pair as a wave form	91
25.a. electromagnetic waves as space and time varying fields	93
25.b. differences between quanton and electromagnetic waves	98

26. Maxwell equations of energy fields	100
27. role of Maxwell equations in the evolution of the quanton	104
28. anti quanton evolution and its degrees of freedom	105
29. Lorentz transformation of EM energy fields	109
30. some concepts behind space fabric	112
31. quanton field interactions	113
32. bound and unbound fields	114
33. types energy field interactions	117
34. quanton field interactions	122
35. generation of space fabric binding interaction	133
36. Dimensions of energy field interactions	135
38. dark energy and dark matter in terms of quanton interaction Potentials	136
38. why quanton does not achieve equilibrium	143
39. inverse relationship between wave length and energy- a possible explanation	144

40. quanton stable degrees of freedom	146
41. role of individual energy fields in the formation of space fabric	148
42. energy fields and the generation of the fundamental Interactions	149
43. degrees of freedom of normal matter quanton	152
44. gravitational mass and its relativistic effect	155
45. field parameters for the normal matter	157
46. scalarized degree of freedom	160
47. normal matter field interactions	162
48. gravitational interaction	169
49. atomic charge , electric field and binding energy	171
50. strong nuclear energy self-interaction	173
51. gravitational like attraction of dark matter	178
52. gravitational distortion of space fabric	184

53. electromagnetic waves interactions	187
54. CPT symmetry at the quanton level	193
55. conclusions	194
56. references	194

1. The physical basis of this model

This model is based on the following two concepts

a-the relationship between energy density inside the quanton

and its parameters (defined in terms of parameters : k , ω , or r_q

(quanton radius)) is an energy degree of freedom relationship

b- the complex nature of the energy expansion in the form of

space varying and time varying fields

the following points will be discussed throughout the model

1-as energy expands from a packet state (energy non varying in

space or time) , It creates associated fields that vary in space and

in time

2- the symmetric nature of this variation in space and in time

3- as a result of this symmetry , the relationship between those

Space and time varying fields is governed by : energy degrees of

Freedom

2-Definition of the model

2.a Quantons

1-quantons are an accumulation of space and time varying energy

fields , as those fields vary at periodic rate , they possess wave

like behaviour, (it will be later discussed why those fields do

not interact directly with electromagnetic waves ,and when and

how such waves leave the quanton)

each quanton is composed of two different type of energy fields

which interact to form a binding relationships

2-they exist in lattice form which constitutes the space fabric

3-quantons are spherical in shape due to equi-partition of energy

(here it will be called :dimensional energy symmetry) but may

vary in their energy content (packet energy) and in volume with

time as they expand and split

4-Quantons are held in a quasi equilibrium state under the effect

of Internal and external interactions of energy fields

5-due to the imbalance of these interactions the quantons

expand , then split up , the resulting pair share

up the original energy content




Fig. (1) Summary of the quanton states

2.b.Anti quantons

anti quantons are similar to quantons but the dominant nature

of their energy differs from that of the quanton

2.b.1 Anti quanton generation

anti quantons are generated from quantons , quantons and anti quantons exist in pairs as they become a quantum entity of the form Q+AQ

3. Mathematical brief

1-The following formulations for various energy fields inside or outside the quanton (anti quanton) ,

$$E_{sf} = \frac{\partial E}{\partial s} : \text{ free space varying field} \quad (1-3)$$

$$E_{tf} = \frac{\partial E}{\partial t} \text{ (free time varying field} \quad (2-3)$$

$$E_{sc} = \int E \, ds \quad \text{space varying constrained field} \quad (3-3)$$

$$E_{tc} = \int E \, dt \quad \text{time varying constrained field} \quad (4-3)$$

$$E_s = E_{sf} E_{sc} , \quad E_t = E_{tf} E_{tc} \quad (5,6-3)$$

2-quanton and anti quanton energy density equation is in the form

$$\text{of } E_q = E_{sf} E_{sc} E_{tf} E_{tc} \text{ and neither in the form} \quad (7-3)$$

$$E_q = \sqrt{E_{sf}^2 + E_{sc}^2 + E_{tf}^2 + E_{tc}^2} \quad \text{nor the form}$$

$$E_q = E_{sf} + E_{sc} + E_{tf} + E_{tc}$$

3- spatial energy fields are vector quantities which have

direction as well as magnitude.

4 -an energy field like free space varying energy can be defined as

$$E_{sf} = K_{sf} D_{sf} \psi_{sf} \quad (8-3)$$

where D_{sf} : energy field strength (degree of freedom parameter –

in exponential terms of the constant c),

K_{sf} : field intensity parameter which is defined in terms of the

quanton total energy divided by four degrees of freedom

and ψ_{sf} is reserved for variation parameter of space varying

energy field

5-the two types of quanton energy fields $E_{qf} = E_{sf} E_{tc}$ (9-3)

and $E_{qc} = E_{sc} E_{tf}$ can be expressed by the one-dimensional PDE

$$(E_{qf})_{tt} = c^2 (E_{qf})_{xx} \quad \text{or} \quad (E_{qc})_{tt} = c^2 (E_{qc})_{xx} \quad (10-3)$$

6- $E = E_s E_t$ (an energy packet state – energy not varying in space

or time , not associated with fields) (11-3)

which is generated by energy constraining

4. variation parameters of energy fields

quanton (or anti quanton) energy density defined as the

multiplication of field strengths and intensities of four types of

energies and two opposing natures which takes the form

$$E_q = (E_{sf} E_{tc}) (E_{sc} E_{tf}) = E_{qf} E_{qc} \quad (1-4)$$

Each of those four functions expresses the change of either

space or time as follows ,

1 – $\psi_{sf} = e^{+\frac{jr}{2r_q}}$ which defines change of free energy field in space

, $r = (x, y, z)$, (2-4)

2 – $\psi_{sc} = e^{-j\frac{jr}{2r_q}}$ defines change of constrained energy field

in space

(3-4)

3 – $\psi_{tf} = e^{+j\omega t}$: that expresses variation of free energy field

in time

4 – $\psi_{tf} = e^{-j\omega t}$: variation parameter of constrained energy

field in time

5. Energy constraining

1-Energy constraining describes evolution , interaction of energy

fields which is summarized as

a-the act of containment free energy fields (E_{sf} , E_{tf}) inside

quantons (this will be discussed in the section : Maxwell

equations role in the evolution of quantons)

b-the appearance of constrained energies ($E_{sc} E_{sc}$)

c- evolution of the quanton fields' degrees of freedom

d-energy fields expansion inside the quanton and its subsequent

splitting

e- the release of radiation energy as a product of the quanton

expansion

2-as energy expands in space in the form of space and time

varying fields , it's said to have free degrees of freedom , and it

must express this degree of freedom in space in a symmetric way

with respect to all spatial dimensions ,and this is only possible

inside a spherical structure, a quanton , so , dimensional energy

symmetry (DES) is behind the evolution of the quantons as a

spatially symmetric shape

3-as energy is released , it must expand , not only by variation

In space but by variation in time as well , hence the appearance of

energy fields E_{sf} , E_{tf} (free energy that varies in space

and free Energy that varies in time) ,such expansion takes the form

$$\frac{\partial}{\partial s} (E) = \frac{\partial}{\partial s} (E_s E_t) = \frac{\partial E}{\partial s} \frac{\partial E}{\partial t} = E_{sf} E_{tf}$$

4-energy fields cannot vary in space and time simultaneously

, so no energy field is in the form $E_{sf,tf} = fn(s, t)$

, but rather $E = E_{sf}(x, y, z) E_{tf}(t)$

and this is because the relationship between the expansion of

space varying and time varying fields is diametric , as the time

varying field (curls) the free expansion of space varying field

hence the appearance of the quanton (this point will be further

discussed in the section Maxwell equations of energy fields)

5-Energy fields can either be free in space varying

($E_{sf} = \frac{\partial E}{\partial s}$) or free in time varying field ($E_{tf} = \frac{\partial E}{\partial t}$)

or constrained in space ($E_{sc} = \int E_s ds$) or constrained in

time ($E_{tc} = \int E dt$) , while non space or time varying energy

in the form ($E = E_s E_t$) can be defined as an energy packet state

:energy that does not change in space or in time

6-the appearance of constrained energy fields inside the

quanton (anti quanton) , is due to the fact that free energies
 (E_{sf} E_{tf}) seek to form a more stable binding interactions with
 these newly appeared constrained energies (E_{sc} E_{tc}) under
 inflationary conditions rather than the less stable repulsive self
 Interactions (discussed in detail in the section : space fabric field
 interactions and why space fabric generates binding interactions)
 7-as energy expands by space varying field (E_{sf}) , it must
 have a constrained time varying field (E_{tc}) such that
 $E_{qf} = E_{sf}E_{tc}$, so the field E_{qf} is due to a predominantly free space
 varying energy field (E_{sf})
 8-as time varying energy expands (E_{tf}) , it must be expand in part
 by variation in space as well , hence the appearance of space
 varying constrained energy field (E_{sc}) such that $E_{qc} = E_{sc} E_{tf}$,
 hence the field (E_{qc}) is a predominantly space constrained energy

field (E_{sc})

9-as energy expands from a packet state ($E = E_s E_t$) , it possesses

Four degrees of freedom , and for the quanton (or anti quanton) to

exist as an independent energy entity , it must possess all of those

Four degrees of freedom (needless to say one of them varies in

time)

10- based on the previous point, inside the quanton (anti quanton)

energy fields cannot expand by free variation in space and in time

in the form $E_q = E_{sf} E_{tc}$ or of the form $E_{aq} = E_{sc} E_{tf}$ alone

the emerging fields become

$$E_q = (E_{sf} E_{tc})(E_{sc} E_{tf}) = E_{qf} E_{qc} \quad (1-5)$$

and for anti quanton

$$E_{aq} = \left(\frac{E_{sf} E_{tc}}{c} \right) (c E_{sc} E_{tf}) = \left(\frac{E_{qf}}{c} \right) (c E_{qc}) \quad (2-5)$$

This quanton energy density equation represents two fields

(discussed in the section : wave model inside the quanton)

one of them is free energy dominated or $E_{qf} = (E_{sf}E_{tc})$,and the

other is constrained energy dominated $E_{qc} = (E_{sc}E_{tf})$

the anti quanton's energy density equation is the same as the

energy density equation of quanton's , but energy levels (i.e

degrees of freedom) of various fields are different form those of the

quanton (this will be discussed later in the sections : quanton

and anti quanton evolution and their energy degrees of freedom)

11- the fields E_{qf} , E_{qc} are orthogonal (will be discussed further in

the section: Maxwell's equations of energy fields)

12- for free energy fields E_{sf} , E_{tf} , differentiation is the

mathematical expression of free energy expansion by variation in

space or time , while integration is the corresponding mathematical

expression of constraining of free fields varying in space or time

a-for space varying field , full expansion in space

$$\frac{\partial}{\partial x} \frac{\partial y}{\partial y} \frac{\partial}{\partial z} (E_s) = \frac{\partial E}{\partial s} = E_{sf} \text{ (energy expands from a packet state to$$

become a space varying field)

b-constraining of free energy fields takes the form

$$\iiint E_{sf} \, dx \, dy \, dz = E_s \text{ (free space varying field is packetized-$$

reduced into a non varying state) , (3-5)

c- for free time varying field

$$\frac{\partial}{\partial t} (E_t) = \frac{\partial E}{\partial t} = E_{tf} \text{ , and constraining } \int E_{tf} = E_t \text{ (4-5)$$

13- for constrained energy fields E_{sc} , E_{tc} , integration is the

mathematical expression of free energy expansion by variation in

space or time , and differentiation is the corresponding

mathematical expression of energy constraining in space or time

a- for constrained space varying field , expansion in space is

defined as $\iiint E_s \, dx \, dy \, dz = E_{sc}$ (expansion of constrained space

varying field)

b- while constraining takes the form

$$\frac{\partial}{\partial x} \frac{\partial y}{\partial y} \frac{\partial}{\partial z} (E_{sc}) = \frac{\partial}{\partial s} (E_{sc}) = E_s \quad (\text{reduction of constrained space})$$

varying field into a packet state-non varying in space or time),
(5-5)

c- for time varying field

expansion in time $\int E_t dt = E_{tc}$, and when being constrained

$$\frac{\partial}{\partial t} (E_{tc}) = E_t \quad (6-5)$$

14- energy field (free- constrained) expansion inside the quanton is

more or less a process of differentiating two variables

15- expansion of (free- constrained) energy fields by variation in

space or time follows differentiation of two variables rules

$$\frac{\partial}{\partial x} (f(x) g(x)) = \frac{\partial f}{\partial x} g(x) + \frac{\partial g}{\partial x} f(x)$$

results of an energy expansion process inside the quanton =

expansion of the (free -constrained fields) +

constraining of (free- constrained fields)

let's consider the case of expansion of free space varying field E_{sf}

$$\text{inside the quanton or } \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) = \frac{\partial E}{\partial s} \int E dt = E_{sf} E_{tc} \quad (7-5)$$

(this step will be further elaborated in the chapters : quanton

degrees of freedom and the role of Maxwell equations in the

evolution of the quanton)

16- similarly for the case of free time varying field

$$\frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) = \frac{\partial E}{\partial t} \int E ds = E_{sc} E_{tf} \quad (8-5)$$

Now the quanton energy density equation

$$E_q = \left(\frac{\partial E}{\partial s} \int E dt \right) \left(\int E ds \frac{\partial E}{\partial t} \right) = (E_{sf} E_{tc}) (E_{sc} E_{tf}) \quad (9-5)$$

, to summerize

process	Free energy	Constrained energy field
expansion	differentiation	integration
constraining	integration	differentiation

Table (1) Mathematical expression of energy expansion /constrained inside the quanton

17- the quanton's four degrees freedom are the sum of free

energy fields' degrees of freedom plus the constrained energy

fields' degrees of freedom or

$$Dof_q = Dof_{sf} + Dof_{sc} + Dof_{tf} + Dof_{tc} = 4 \quad (10-5)$$

18- it is understood that the space varying energy fields (free and constrained) have three degrees of freedom or

$$Dof_{sf} + Dof_{sc} = 3 \quad (11-5)$$

while time varying energy fields (free and constrained) have one

$$\text{degree of freedom or } Dof_{tf} + Dof_{tc} = \text{one} \quad (12-5)$$

19—energy fields $E_{sf}, E_{tf}, E_{sc}, E_{tc}$ do not have the dimensions of energy , but their product (E_q) does have the dimensions of

energy density which is defined as energy divided

$$\text{by three dimensional volume } [E_q] = \left[\frac{\text{energy}}{\text{volume}} \right] = M L^{-1} T^{-2}$$

(later , it will be shown that this energy density is in fact 4

dimensional that expands in 3 D space)

20- as free energy fields expand , constraining of the expanding

fields takes place

$$\frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) = \frac{\partial E}{\partial s} + \int \left(\frac{\partial E}{\partial s} \right) ds \text{ or}$$

$$\frac{\partial}{\partial s} (E_{sf}) = \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) = \frac{\partial E}{\partial s} + \text{constraining term}$$

and
$$\frac{\partial}{\partial t} (E_{tf}) = \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) = \frac{\partial E}{\partial t} + \int \left(\frac{\partial E}{\partial t} \right) dt \text{ or}$$

$$\frac{\partial}{\partial t} (E_{tf}) = \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) = \frac{\partial E}{\partial t} + \text{constraining term}$$

and so on for higher order derivatives, and this process represents

the expansion of the free energy by variation in space or time

which must be accompanied by constraining while inside the

quanton

21- as constrained energy fields expand , constraining of the

expanding fields takes place

$$\int \left(\int E ds \right) ds = \int E ds + \frac{\partial}{\partial s} \left(\int E ds \right) \text{ or}$$

$$\int \left(\int E ds \right) ds = \left(\int E_{sc} ds \right) ds = \int E ds + \text{constraining term} , \text{ and}$$

$$\int (\int E dt) dt = (\int E_{tc} dt) dt = \int E dt + \frac{\partial}{\partial t} (\int E dt) \quad \text{or}$$

$$\int (\int E dt) dt = (\int E_{tc} dt) dt = \int E dt + \text{constraining term} ,$$

this means that expansion of constrained energy fields must also

be accompanied by constraining of those constrained energy

fields

22- when energy is released from a field constraining process

For free or constrained energy field as in (20) or (21) , it is released

in the packet state $E = E_s E_t$ (energy non varying in space or

time) in other words , released energy cannot take the form of

E_s or E_t as either of those forms of energy do not exist

independently

23- a cycle of expansion and constraining is not a reversible

process due to losses and effect of entropy (irreversible process)

(will be further clarified in the section energy constraining and the

origin of entropy)

24- energy degree of freedom must be identical in spatial

dimensions for (E_{sx}, E_{sy}, E_{sz}) for each field otherwise energy

field is deemed to be unstable

25- an energy field expansion process results in an expanding

field plus energy constraining products , so it is expected that the

total energy content of the quanton (or anti quanton) to decrease

during the process of expansion (this point discussed in the

section : energy constraining and origin of entropy)

26- since the quanton is a quantum entity , its packet energy – total

energy content of the quanton – is governed solely by the Planck

–Einstein relationship so, quanton energy is determined by its

wave parameters $(k, \omega \text{ or } r_q)$, while an energy degree of

freedom- which is defined in terms of the constant (c) , is just a

mechanism of division of energy between the various space and

time varying fields

27- recalling point (7) , energy of the following forms do not exist

independently

a- $\frac{\partial E}{\partial s} \frac{\partial E}{\partial t}$ (energy field can not expand in space and in time

Simultaneously without having a constrained field)

b – $(\int E ds \int E dt)$ (energy field cannot be constrained in

space and in time simultaneously without having an free

expansion component)

28- though quanton includes both energy types (free and

constrained) energy fields , but there is a dominant type of energy

field , this is based on which field has the majority of *Dof's*

29- for the quanton , the free energy field is the dominant while

for anti quanton , the constrained field is the dominant type of

energy field

30- the packet in this model assumes two roles

a-The packet energy : total energy of the quanton which is defined

$$\text{as } E_p = \frac{h}{2\pi} \omega = \int E_q dV \quad (13-5)$$

b- packet state which is the result from constraining process

and defined as $E = E_s E_t$

(energy that does not vary in space or in time)

6. Bridging the gap between mathematics physics of energy constraining

1-While differentiation of two functions involves

differentiating only one at a time and maintaining the other

constant , in real world this is not possible since an expanding

energy field must vary either in space or in time

2-when dealing with expansion of constrained energy fields

integration is the physical equivalent to mathematically

maintaining one function as a constant

3-expansion of two energy fields of the form

$$\frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right) = \left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right) \left(\int E ds \int E dt \right)$$

Could not be in the form $\left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right) + \left(\int E ds \int E dt \right)$

Since the energy fields $\left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right)$ or $\left(\int E ds \int E dt \right)$

are unstable in this form as the quanton is in the process of

formation and free or constrained energy fields could not exist

independently

4-The quanton energy density equation

$$E_q = \left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right) \left(\int E ds \int E dt \right), \text{expresses two physical entities}$$

(free energy fields : $\left(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t} \right)$ and constrained energy fields

$\left(\int E ds \int E dt \right)$) and each of those types of fields behave as single

physical entity (ie single variable) , so four different energy

fields , are in fact ,representing only two variables instead of

four(energy field interactions will be based on this particular point)

5-recalling points (13), (14) from previous section , for complex

Energy fields (free /constrained)

$E_q = E_{sf} E_{sc} E_{tf} E_{tc}$ energy constraining which happens through

Quanton expansion in space is defined as

$$\frac{\partial}{\partial s} (E_q) = \frac{\partial}{\partial s} ((E_{sf} E_{tc})(E_{sc} E_{tf}))$$

$$(\frac{\partial E_{sf}}{\partial s} \int E_{tc} dt)(\int E_{sc} ds \frac{\partial E_{tf}}{\partial t}) + (\int E_{sf} ds \frac{\partial E_{tc}}{\partial t})(\frac{\partial E_{sc}}{\partial s} \int E_{tf} dt)$$

$$= (E_{sf} E_{tc} E_{sc} E_{tf}) + E_s E_t \quad (1-6)$$

6- when dealing with Energy field expansion inside the quanton

There would be two terms as a result of the expansion

a-expansion term : differentiating free energies *integration of

constrained energies

b-constraining term : integrating free energies * differentiating

constrained fields

7- energy expansion process inside the quanton , involves

both free and constrained energies , and to avoid confusion while

using the gradient operator (∇) for both types of field expansion

,the use of the differential / integral operators will be maintained

$\frac{\partial}{\partial S}$ for expansion of free energy fields and \int for the expansion

of constrained energy fields inside the quanton

8- when dealing with energy expansion we will use the

wave like form $E_q = (E_{sf} E_{tc})(E_{sc} E_{tf})$

while when dealing with fields and energy interactions

the form $E_q = (E_{sf} E_{tf})(E_{sc} E_{tc})$ will be used

7. Energy Degrees of freedom

1-as energy is allowed to vary in every dimension in space or in

time , it is said to have an energy degree of freedom

2- the quanton energy density is defined in terms of the degrees

of freedom of its wave parameters ($\omega, k, or r_q$)

3- E_q (quanton energy density)will be shown to be directly

proportional to ω^4 , k^4 or $\frac{1}{r_q^4}$

4- while the energy density of the quanton is defined in terms of

ω^4 , k^4 or $\frac{1}{r_q^4}$, however the energy fields are defined in terms of

Field strength or in terms of the constant (c) in the form of

$D_{sf} = c^{Dof_{sf}}$, Dof_{sf} : degrees of freedom of free space varying

field (transformation from degrees of freedom from formulation in

terms of wave parameters , to degrees of freedom in terms of (c))

5-For space varying and time varying energy fields , where the

resultant energy density is in the form

$E_q = (E_{sf} E_{tc})(E_{sc}E_{tf})$ and not in the square root form

$$E_q = \sqrt{E_{sf}^2 + E_{sc}^2 + E_{tf}^2 + E_{tc}^2}$$

this multiplier form of resultant , allowed (c) to become an energy

degree of freedom in a multiplier (exponential) form , where
 energy is divided up symmetrically , between the space and time
 varying fields , hence the symmetric division of energy across all
 dimensions

6-the constant (c) plays a bigger role than being the velocity of light
 or the velocity of transmission of the fundamental forces , as it
 plays the role of ratio between space and time varying fields , this
 is based on the following

a- the constant (c) represents the relationship between energy
 field expansion by variation in space and in time , for the wave
 parameters of the fields E_{tc} , E_{sf}

$$\psi_{tc} , \psi_{sf} \text{ where } \psi_{tc} = e^{-j\omega t} , \psi_{sf} = e^{+jk(x+y+z)}$$

$$\psi = \psi_{tc} \psi_{sf}$$

$$\frac{\partial \psi_{tc}}{\partial t} = -j\omega \psi_{tc} , \quad \frac{\partial \psi_{sf}}{\partial x} = jk \psi_{sf}$$

$$\left(\frac{\partial \Psi}{\partial t}\right) = \frac{\partial}{\partial t} (\Psi_{sf} \Psi_{tc}) = \Psi_{sf} \frac{\partial \Psi_{tc}}{\partial t} = -j\omega \Psi_{sf} \Psi_{tc}$$

$$\left(\frac{\partial \Psi}{\partial x}\right) = \frac{\partial}{\partial x} (\Psi_{sf} \Psi_{tc}) = \frac{\partial \Psi_{sf}}{\partial x} \Psi_{tc} = jk \Psi_{sf} \Psi_{tc}$$

$$-\frac{\left(\frac{\partial \Psi}{\partial t}\right)}{\left(\frac{\partial \Psi}{\partial x}\right)} = \frac{j\omega \Psi_{sf} \Psi_{tc}}{jk \Psi_{sf} \Psi_{tc}} = \frac{\omega}{k} = c \quad (1-7)$$

which is the relationship between rate of field variation in space

and in time

b- recalling the Lagrangian (L) of an action as $\frac{d}{dt} \frac{\partial L}{\partial x'} - \frac{\partial L}{\partial x} = 0$

given that momentum $P = \frac{\partial L}{\partial x'}$

we get $\frac{\partial P}{\partial t} = \frac{\partial L}{\partial x}$ or alternatively $\frac{\partial L}{\partial P} = \frac{\partial x}{\partial t} = c$

an energy degree of freedom: the rate of change of the total

energy of the system with respect to its momentum

c-the same result can be obtained directly from the energy

momentum relationship $E^2 = P^2 c^2 + m_0^2 c^4$

Differentiating both sides $2 E dE = 2 P dP$

$$\frac{dE}{dP} = \left(\frac{P c}{E} \right) c \quad , \text{ and } \frac{dE}{dP} = c$$

where for space fabric case ($m_o = \text{zero}$) , $E = P c$

which is an alternative definition of the energy degree of freedom

6-both results of (a) and (c) are equivalent , given that

$$\Psi = \Psi_{sf} \Psi_{tc}$$

using the Schrödinger equation ,for time and space derivatives

$$-\frac{\partial \Psi}{\partial t} = \frac{jE}{2\pi h} \Psi$$

$$\nabla \Psi = \frac{jP}{2\pi h} \Psi$$

$$\frac{\frac{\partial \Psi}{\partial t}}{\nabla \Psi} = \frac{E}{P} = c \quad (2-7)$$

Based on the above points ,the division of energy density between

space and time varying fields can be done where strength of space

and time varying energy fields (Dof) is expressed in terms of the

constant (c) that defines the relationship between their rate of

variation , It is worth noting that

1-energy field degrees of freedom (field strength) is not related to the total energy of the quanton , as it is only a mechanism for the division of the quanton energy density between the various space and time varying energy fields , and what differs the total energy content of any quanton from another is only the rate of variation of fields with time and space according to Planck Einstein

relationship $E_p = \frac{h}{2\pi}\omega$

2- the energy degrees of freedom can be classified as follows

a- active (actual degrees of freedom) that belong to the energy fields inside or outside the quanton

b- kinetic degree of freedom which expresses the propagation of energy fields (outside the quanton in the form of electromagnetic waves) in one direction , this kinetic degree of freedom is subtracted from the existing four degrees of energy freedom for

space and time varying fields (discussed in the section :

electromagnetic waves out of quanton) , where $Dof's = (2)+1$

instead of $(3)+(1)$

c- scalarized degrees of freedom : when a degree of freedom of an

energy field becomes part of its intensity parameter instead of its

strength parameter (discussed in the section : normal mater

quantons)

8.The superposition principle inside the quanton

1-the linear superposition of energy fields still applies inside and

Outside the quanton with a resultant field which equals to the

addition of the individual field intensities on condition that

a-those fields must be of the same type (free / constrained) and

b- have the same degree of freedom

$$E_{sfi} + E_{sfj} = K_{sfi} D_{sf} + K_{sfj} D_{sf}$$

$$= (K_{sfi} + K_{sfj}) D_{sf} \quad (1-8)$$

$$\begin{aligned}
(E_{sfi} E_{tci}) + (E_{sfj} E_{tcj}) &= (K_{sfi} D_{sf})(K_{tci} D_{tc}) + (K_{sfj} D_{sf})(K_{tcj} D_{tc}) \\
&= (K_{sfi} K_{tci})(D_{sf} D_{tc}) + (K_{sfj} K_{tcj})(D_{sf} D_{tc}) \\
&= (K_{sfi} + K_{sfj})(K_{tci} + K_{tcj})(D_{sf} D_{tc}) \quad (2-8)
\end{aligned}$$

while the superposition of fields of different nature (free /

constrained) or fields that do have different energy Dof's

the superposition is done by adding their field strength (ie

exponential degree of freedom) and multiplying their intensities

2- the exponential form of superposition applies, as energy fields

are defined in terms of energy degree of freedom (Dof) , which is

expressed as the exponent of (c^{Dof})

the resulting superposition inside the quanton will not be a linear

one instead , it is an exponential superposition where

$$\begin{aligned}
E_{sfi} E_{scj} &= (K_{sfi} D_{sf})(K_{scj} D_{sc}) \\
&= (K_{sfi} K_{scj})(D_{sf} D_{sc}) \quad (3-8)
\end{aligned}$$

$$\begin{aligned}
E_{sf} E_{tc} &= (K_{sf} D_{sf}) (K_{tc} D_{tc}) \\
&= (K_{sf} K_{tc}) (D_{sf} D_{tc}) \quad (4-8)
\end{aligned}$$

and for the quanton as a whole

$$\begin{aligned}
E_q &= (E_{sf} E_{tc}) (E_{sc} E_{tf}) = (K_{sf} D_{sf}) (K_{tc} D_{tc}) (K_{sc} D_{sc}) (K_{tf} D_{tf}) \\
&= (K_{sf} K_{tc} K_{sc} K_{tf}) c^{Dof_{sf} + Dof_{tc} + Dof_{sc} + Dof_{tf}} = (K_{sf} K_{tc} K_{sc} K_{tf}) c^4 \quad (5-8)
\end{aligned}$$

3- inside the quanton , instead of the addition of the same

type of energy , the exponential addition can be between two

different types of energy fields (space and time varying fields) and

of two different natures (free / constrained) to give a complex

energy field

the main reason behind this is that free and constrained fields

cannot be considered as an independent energy entity individually

, since neither of them does possess four degrees of freedom

and hence their individual Dof's must be added exponentially to

obtain either a complex field or the total energy density of the quanton if the addition is for all four energy fields

9. Definition of directional field directional components

For free space varying fields

$$E_{sf} = \sqrt{E_{sfx}^2 + E_{sfy}^2 + E_{sfz}^2} \quad (1-9)$$

and space varying constrained field

$$E_{sc} = \sqrt{E_{scx}^2 + E_{scy}^2 + E_{scz}^2} \quad (2-9)$$

for spatially varying fields

$$E_{sx} = E_{sfx} E_{scx} \quad , \quad E_{sy} = E_{sfy} E_{scy} \quad (3-9)$$

$$E_{sz} = E_{sfz} E_{scz} \quad , \quad (4-9)$$

And for time varying fields,

$$E_t = E_{tf} E_{tc} \quad (5-9)$$

Those are 8 components , 6 that vary in space and 2 that vary in time 3 are constrained space varying and one is constrained time

varying , and 3 are free space varying and one is free in time

, it is worth noting that

1-spatial and time varying energy fields cannot exist

independently of each other , as discussed previously

2- the quanton fields $E_{sf}, E_{sc}, E_{tf}, E_{tc}$ neither have the

dimensions of energy nor the energy density but their product

has the dimension of energy divided by three dimensional volume

10.Dimesional energy symmetry (DES)

Dimensional energy symmetry is the mechanism which ensures the

uniformity and homogeneity of energy under inflationary

conditions inside the quanton ,given that $E_q = (E_{sf}E_{tc})(E_{sc}E_{tf})$

energy as it expands in along the x- axis

$\frac{\partial}{\partial x} (E_q)$ will not only give as the result of the expansion

$$\left(\frac{\partial E_{sf}}{\partial x} \int E_{tc} dt\right) \left(\int E_{sc} dx \frac{\partial E_{tf}}{\partial t}\right) +$$

$(\int E_{sf} \, dx \frac{\partial E_{tc}}{\partial t}) (\frac{\partial E_{sc}}{\partial x} \int E_{tf} \, dt)$, but it will be of the form

$$\frac{\partial}{\partial x} (E_q) = \frac{\partial}{\partial x} (E_{sf} E_{tc} E_{sc} E_{tf}) =$$

$$\frac{\partial}{\partial x} (E_{sf} E_{sc}) \frac{1}{\partial x} \frac{\partial y}{\partial t} \frac{\partial}{\partial y} (E_{sf} E_{sc}) \frac{1}{\partial x} \frac{\partial z}{\partial t} \frac{\partial}{\partial z} (E_{sf} E_{sc}) \frac{1}{\partial x} \frac{\partial}{\partial t} (E_{tf} E_{tc})$$

$$= (\frac{\partial}{\partial x}) (\frac{1}{\partial x} \frac{\partial y}{\partial t} \frac{\partial}{\partial y}) (\frac{1}{\partial x} \frac{\partial z}{\partial t} \frac{\partial}{\partial z}) (E_{sf}) \int (E_{tc}) \, dt) *$$

$$(\int \int \int (E_{sc}) \, dx \, dy (\frac{1}{\partial y} \frac{dx}{dt}) \, dz (\frac{1}{\partial z} \frac{dx}{dt}) (\frac{1}{\partial x} \frac{\partial}{\partial t}) (E_{tf})) +$$

$$(\int \int \int (E_{sf}) \, dx \, dy (\frac{1}{\partial y} \frac{dx}{dt}) \, dz (\frac{1}{\partial z} \frac{dx}{dt})) (\frac{1}{\partial x} \frac{\partial}{\partial t}) (E_{tc}) *$$

$$(\frac{\partial}{\partial x}) (\frac{1}{\partial x} \frac{\partial y}{\partial t} \frac{\partial}{\partial y}) (\frac{1}{\partial x} \frac{\partial z}{\partial t} \frac{\partial}{\partial z}) (E_{sc}) (\int (E_{tf}) \, dt (\frac{\partial x}{\partial t}))$$

Given that $\frac{\partial x}{\partial t} = \frac{\partial y}{\partial t} = \frac{\partial z}{\partial t} = c$

$$\frac{\partial}{\partial x} (E_q) = (\frac{\partial E_{sf}}{\partial s} \int E_{tc} \, dt) (\int E_{sc} \, ds \frac{\partial E_{tf}}{\partial t}) +$$

$$(\int E_{sf} \, ds \frac{\partial E_{tc}}{\partial t}) (\frac{\partial E_{sc}}{\partial s} \int E_{tc} \, dt) = \frac{\partial}{\partial s} (E_q) \quad (1-10)$$

Note : we applied the chain rule for differentiation and

integration by change of variables

We can clearly see that as energy expands along one axis ,it must not only expand along other spatial and temporal axes but be constrained along the spatial and temporal axes as well ,

we conclude that

1-Events in one direction are immediately reflected in the other spatial and temporal directions , and by the same magnitude

2-The uniformity and the homogeneity of space fabric is ensured through the role time plays as the link between all the three spatial axis (and via the constant (c))

3-to satisfy dimensional energy symmetry for quanton , the degrees of freedom must be symmetric with respect space and time varying energy fields

Define the Dof_q , D_q (in terms of c)

Where the degree of freedom parameter

$$Dof_q = Dof_{sf} + Dof_{tf} + Dof_{sc} + Dof_{tc} = 4 \quad (2-10)$$

$$\text{Energy field strength parameter } D_q = D_{sf} D_{tf} D_{sc} D_{tc} = c^4 \quad (3-10)$$

$$D_s = c^3, \quad D_{sf} = c^{Dof_{sf}}, \quad D_{sc} = c^{Dof_{sc}} = c^{3-Dof_{sf}} \quad (4,5,6,7-10)$$

$$D_t = c, \quad D_{tf} = c^{Dof_{tf}}, \quad D_{tc} = c^{Dof_{tc}} = c^{1-Dof_{tf}} \quad (8,9,10-10)$$

$$Dof_{sf_x} = Dof_{sf_y} = Dof_{sf_z} \quad (11-10)$$

4-the degree of freedom of constrained space varying field must

be identical for spatial time varying components

$$Dof_{sc_x} = Dof_{sc_y} = Dof_{sc_z} \quad (12-10)$$

in other words for free and constrained fields the degree of

freedom must be expressed in a symmetric way across all spatial

and time varying fields




Fig (2) Role of dimensional energy symmetry in ensuring the uniformity of energy distribution inside the quanton

11. Energy density / Degree of freedom relationship

When assessing the relationship between the quanton energy density E_q and its wave parameters , two methods will be used , and as it turned out , the two are equivalent , first is the simplified method which is discussed here , and secondly , the analytical , which is dealt with in the section : energy field parameters

a-simplified method

based on two basic assumptions

1-high symmetry , the quanton volume can be

represented by a highly symmetric 3 dimensionally symmetric

shape (a sphere) given that ($2r_q = \lambda$) ,

2- the uniformity of the field inside the quanton

the total energy at any radius (a) can be defined as

$$E(a) = \frac{hc}{2r_q} \left(\frac{a}{r_q} \right), \quad (a < r_q)$$

under such assumptions , the following relations would be used as

an approximation

$$E_p = \int_{V_q} E_q dV = E_q V_q = \frac{h}{2\pi} \omega \quad (1-11)$$

where E_q is the average energy field density inside the quanton

, E_p : packet energy ,

based on assumption of uniform energy density inside the

quanton , $V_q = \frac{4}{3}\pi r_q^3$

given that $\omega = kc = \frac{\pi c}{r_q}$, $k = \frac{2\pi}{\lambda} = \frac{\pi}{r_q}$, $r_q = \frac{\lambda}{2}$ (2-11)

$V_q = \frac{4}{3}\pi r_q^3 = \frac{4}{3}\pi \left(\frac{\pi^3}{k^3}\right) = \frac{4}{3}\frac{\pi^4 c^3}{\omega^3}$ (3-11)

This shows that the quanton volume can be defined in terms of the parameters k , ω , this indicates that the relationship

$E_p = \int_{V_q} E_q dV = E_q V_q$ is not only a volumetric relationship but

an energy degree of freedom as well , now the energy density can

Be written as

$E_q = \frac{E_p}{\int_{V_q} dV} = \frac{h\omega}{(2\pi)\left(\frac{4\pi}{3}r_q^3\right)}$ ($r_q = \frac{\pi c}{\omega}$) , substituting for r_q^3

$E_q = \frac{3h\omega}{(2\pi)(4\pi)}\left(\frac{\omega^3}{c^3}\right) = \frac{3h\omega^4}{8\pi^5 c^3}$ (4-11)

and this is a very important relationship since the term

$\frac{3h}{8\pi^5 c^3} = \text{constant}$, in other words , (5-11)

Field energy density inside the quanton is linearly proportional to

the four degrees of freedom as expressed by either (ω^4, k^4 or $\frac{1}{r_q^4}$),

$$\text{define } h_q = \frac{3h}{8\pi^5 c^3} \quad (6-11)$$

$$E_q = h_q \omega^4 = h_q k^4 c^4 = h_q \frac{\pi^4 c^4}{r_q^4} \quad (7-11)$$

substituting $E_p = E_q V_q = h_q V_q \omega^4$

from (2-11) and given that $\omega = kc = \frac{\pi c}{r_q}$

$$\frac{V_{q2}}{V_{q1}} = \left(\frac{r_{q2}}{r_{q1}} \right)^3 = \frac{\omega_1^3}{\omega_2^3} \text{ and } \frac{r_{q2}}{r_{q1}} = \frac{\omega_1}{\omega_2} \quad (8-11)$$

12 -Energy constraining and the release of thermal energy

1-as the quantons expand , field constraining takes place

(transformation into a packet state – energy non varying in space

or time)

2-Energy constraining during quanton inflation as follows

a-Expansion of free energy fields ($\frac{\partial E_{sf}}{\partial s} \frac{\partial E_{tf}}{\partial t}$) must be accompanied

by constraining of part - of the expanding free energy fields in the

form $(\int E_{sf} ds \int E_{tf} dt)$

b-expansion of constrained fields $(\int E_{sc} ds \int E_{tc} dt)$ must be

accompanied by an constraining of part of the expanding field

in the form $(\frac{\partial E_{sc}}{\partial s} \frac{\partial E_{tc}}{\partial t})$

c-In both cases , the result will be the release of energy in a packet

state (non varying in space or time) of the form $E = E_s E_t$

for the free type of energy as it expands

$$\frac{\partial}{\partial s} [(E_{sf} E_{tc})(E_{sc} E_{tf})] = (\frac{\partial E_{sf}}{\partial s} \int E_{tc} dt) (\int E_{sc} ds \frac{\partial E_{tf}}{\partial t})$$

$$+(\int E_{sf} ds \frac{\partial E_{tc}}{\partial t}) (\frac{\partial E_{sc}}{\partial s} \int E_{tf} dt)$$

given that $\frac{\partial E_{sf}}{\partial s} = E_{sf}$, $\int E_{tc} dt = E_{tc}$, $\int E_{sc} ds = E_{sc}$, $\frac{\partial E_{tf}}{\partial t} = E_{tf}$

and $(\int E_{sf} ds \frac{\partial E_{tc}}{\partial t}) = E_s E_t$, $(\frac{\partial E_{sc}}{\partial s} \int E_{tf} dt) = E_s E_t$

the results of field expansion inside the quanton can be defined as

a- expansion term :

$$(\frac{\partial E_{sf}}{\partial s} \int E_{tc} dt) (\int E_{sc} ds \frac{\partial E_{tf}}{\partial t}) = (E_{sf} E_{tc})(E_{sc} E_{tf})$$

Corresponds to the expanding fields which expand through variation of energy in space and in time , the nature of expanding fields is the same as the original type of fields (though with lesser energy content)

b- The constraining term : $(\int E_{sf} ds \frac{\partial E_{tc}}{\partial t}) (\frac{\partial E_{sc}}{\partial s} \int E_{tf} dt) = E_s E_t$

represents the release of energy in a packet state which is due to the constraining of part of the free and constrained fields a part of this energy packet is released from the quanton as it expands




Fig (3). Quanton expansion cycle

13.energy constraining -a possible origin of cosmic microwave back ground (CMB)

Inflation of the universe (expansion of space fabric) is a free expansion process and is accompanied by the release of thermal energy , the idea that a free expansion process gives off heat is rather odd , since expansion is closely related to reduction in temperature , in fact any release of thermal energy is more than offset by the effects of inflation , so the net result would be a reduction in temperature (observed as thermal degradation of

CMB photons)

inflation of the universe is a free expansion process , which according to the second law of thermodynamics, is an irreversible process , this irreversibility is due to losses in the form of space fabric giving off heat during expansion process

the origin of this release of thermal energy : is energy constraining

Based on the previous results ,we can conclude that the

CMB is due to release of thermal energy during free expansion of the space fabric itself

the extraordinarily high degree of CMB homogeneity with variation of the order of (10^{-5}) , reflects the high degree of homogeneity of space fabric it self as it releases radiation during the free expansion process , and , in fact energy constraining inside the quantons is behind that release of this radiation energy

14. why do quantons split ?

to show the source of this released energy we consider the case of a quanton as it expands (discussed in the following section) from a volume (V_{q1}) to (V_{q2})

1-the quanton radius r_q and its volume V_q should change in the

Following manner $\frac{V_{q2}}{V_{q1}} = \left(\frac{r_{q2}}{r_{q1}}\right)^3$ as the quanton expands in three

Dimensional space

2- quanton energy fields change periodically with time

, this variation at the rate of ω rad/sec , and vary in space at

the rate of k ($= \frac{\pi}{r_q}$) , the total energy of the quanton (as a quantum

entity) is governed by Planck Einstein relationship (function only

in its wave parameters) , namely $E_p = hf = \frac{hkc}{2\pi} = \frac{hc}{2r_q}$

The relationship between quantons of different energy content can

be put in the form $\frac{E_{p2}}{E_{p1}} = \frac{\omega_2}{\omega_1} = \frac{k_2}{k_1} = \frac{\lambda_1}{\lambda_2} = \frac{r_{q1}}{r_{q2}}$

Which means that the quanton radius (and the wave length

of its characteristic wave behaviour are inversely proportional

to its total (packet) energy content

3-recalling here the first concept upon which this model is based

namely the Dof relationship between energy density inside the

quanton and its total energy content

energy density inside the quanton can be assessed as

$$E_p = E_q V_q \quad \text{or} \quad E_q = \frac{E_p}{V_q} \quad , \text{ then}$$

$$\frac{E_{q2}}{E_{q1}} = \left(\frac{E_{p2}}{E_{p1}} \right) \left(\frac{V_{q1}}{V_{q2}} \right)$$

Substituting for $\left(\frac{E_{p2}}{E_{p1}} \right) = \left(\frac{r_{q1}}{r_{q2}} \right)$, and $\left(\frac{V_{q1}}{V_{q2}} \right) = \left(\frac{r_{q1}}{r_{q2}} \right)^3$

$$\text{We get } \frac{E_{q2}}{E_{q1}} = \left(\frac{r_{q1}}{r_{q2}} \right)^4 \quad (1-14)$$

Which deviates from what we would expect in a classical

Volume / density relationship of the form $\frac{q_2}{q_1} = \frac{V_1}{V_2} = \left(\frac{r_1}{r_2} \right)^3$

and this is due to the fact that energy density inside the quanton is

inversely proportional to r_q^4 and not to r_q^3

this previous relationship can be obtained directly from the

equation (6-11) , namely $E_q = \pi^4 h_q \frac{c^4}{r_q^4}$ or

$$E_q = \left(\frac{\text{constant}}{r_q^4} \right)$$

Hence , as quantons expand into a three dimensional space ,

They have to release energy , in the form of radiation

But , energy release from such a process would be excessive

Instead , the quantons , as they expand , do split , to allow for

subsequent expansion , put this time with minimal release

of thermal energy

15.Mechanism of quanton splitting

There are two mechanisms that can cause the quantons to

Expand , namely

a-Splitting action of the quantons due dimensional energy

asymmetry

b-The sole release of energy from the quantons

as for the first mechanism

1-Stage (1-2): expansion under the effect of self interacting repulsive field

1-the two types of fields inside the quanton (free Dominated E_{qf}

and constrained dominated E_{qf}) interact , creating a binding

relationship but since the energy Dof's (i.e field strength) of both

fields are not the same , the field of the dominant type of energy

self-interact creating a repulsive interaction that causes the

quantons to expand , the self interacting (unbound) field is

($E_{sfu} E_{tfu}$) for quantons and ($E_{scu} E_{tcu}$) for anti quantons -

discussed in the section :quanton field interactions)

2-the unbound field is at the origin of the quanton inflationary

energy , which has a greater potential (in terms of the quanton total energy) than the quanton binding energy , as a result , it overcomes this binding and causes the quanton to the expand

3-as the quanton expands its wave parameters (ω , k) are altered

While its energy content remains the same since there's no energy release from the quantons at this stage

as a result the quanton has either to

a- to release thermal energy to maintain the relationship $E_p = \frac{h\omega}{2\pi}$ or

b- to split thereby reducing its overall energy content and allowing for further expansion

2-Stage (2-3) dimensional energy asymmetry occurs and quanton splits

Since the quanton parameters (ω , k) do not reflect its energy

content , ($\frac{E_{p2}}{E_{p1}}$ must equal to $\frac{r_{q1}}{r_{q2}} = \frac{\omega_2}{\omega_1} = \frac{k_2}{k_1}$, while E_{p2} still equals

E_{p1}) , this conflict causes quanton to split as a mechanism to

restore the relationship

$$(r_{q3}) = \left(\frac{r_{q2}}{2}\right) = r_{q1} , \quad E_{p3} = \frac{E_{p2}}{2} = \frac{E_{p1}}{2}$$

, the splitting corresponds to a quanton radius $r_{q2} = x r_{q1}$, $x > 1$

3-Stage (3- 4) quanton expands further

following quanton splitting quanton packet (total energy)

$E_{p3} = \frac{E_{p2}}{2}$, wave parameters (ω, k) also must expand further to

Satisfy the relationship $\frac{E_{p4}}{E_{p2}} = \frac{\omega_4}{\omega_1} = \frac{k_4}{k_1} = \frac{r_{q1}}{r_{q4}} = \frac{1}{2}$

as the quantons expand , they release thermal energy in the form

of CMB photons , and to arrive at the final pseudo stable state

stage	(1)	(2)	(3)	(4)
Total quanton energy : E_p	E_{p1}	E_{p1}	$\frac{E_{p1}}{2}$	$< \frac{E_{p1}}{2}$
Wave parameter ω	ω_1	$\frac{\omega_1}{x}$	$\sqrt[3]{\frac{2}{x^3}} \omega_1$	$< \frac{\omega_1}{2}$
quanton energy density : E_q	E_{q1}	$\frac{E_{q1}}{x^3}$	$\frac{E_{q1}}{x^3}$	$< \frac{E_{q1}}{16}$
Quanton radius r_q	r_{q1}	$x r_{q1}$	$\sqrt[3]{\frac{x^3}{2}} r_{q1}$	$> 2 r_{q1}$
Quanton volume V_q	V_{q1}	$x^3 V_{q1}$	$\frac{x^3 V_{q1}}{2}$	$> 8 V_{q1}$
Number of quantons	one	one	two	two

Table (2) Summary of the stages of the quanton splitting and expansion $x > 1$




Fig (4) cycle of quanton splitting and subsequent inflation

The second method is the pure release of thermal energy

followed up by a subsequent quanton expansion

this mechanism is such an inefficient one in comparison to the fore described method of quanton splitting and subsequent expansion , given the high efficiency of the splitting process as a mechanism to manage the expansion of the quanton through both inflation and multiplication while on the other hand minimizing the thermal energy release , it is clear that such quanton splitting and subsequent expansion is the actual mechanism of space fabric expansion

the release of the radiative energy during the process of expansion of the quanton is not related to the re-establishment of the wave parameter relationship with the quanton energy

a simple explanation lies in the fact that all the quanton energy fields are involved in different interactions , mainly binding ones while energy in a packet state is not involved in any of those

binding interactions , and already possesses four degrees of freedom ,as a result , small part of this energy escapes in the form of radiative energy

the previous model for quanton splitting serves as preliminary and introductory one since the CMB radiation is statistically distributed frequencies indicate that the quanton frequencies are also statistically distributed and the splitting occurs non symmetrically

16.mathematics behind constraining

1- as the quanton forms , the nature of the energy field changes (from free to constrained)

2- to perform such an operation energy fields must transit through a packet state (energy that does not vary in space or in time)

3-and as energy field strength is in terms of Dof , its operator

(integration / differentiation) has to be applied at an exponential

level , thus the exponent of field variation parameter which is

operated upon

the expansion of the free space varying energy

$$\mathbf{a} - \text{for } \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) = \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) \left(\int \left(\frac{\partial E}{\partial s} \right) ds \right) = \left(\frac{\partial E}{\partial s} \right) (E_s)$$

$$\frac{\partial}{\partial s} (K_{sf} D_{sf} \Psi_{sf}) = K_{sf} D_{sf} \frac{\partial}{\partial s} (\Psi_{sf})$$

$$= K_{sf} D_{sf} (e^{jks}) \left(e^{\frac{\partial}{\partial s}(jks)} \right) \quad (1-16)$$

$$= [K_{sf} D_{sf} (e^{jks})] [K_s D_s e^{(jk)}]$$

$$= \frac{\partial E}{\partial s} (K_s D_s e^{(jk)}) = \left(\frac{\partial E}{\partial s} \right) (E_s)$$

$$\mathbf{b} - \text{for } \frac{\partial}{\partial s} \left[\left(\frac{\partial E}{\partial s} \right) (E_s) \right] = \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) \int (E) dt = \left(\frac{\partial E}{\partial s} \right) \left(\int E dt \right)$$

$$\frac{\partial}{\partial s} \left[\left(\frac{\partial E}{\partial s} \right) (E_s) \right] = \frac{\partial E}{\partial s} (K_s D_s e^{-\int(jk) ds}) \quad (2-16)$$

We perform the following change of the integration parameters

$$\mathbf{K} = \frac{\omega}{c} , \quad ds = dx dy dz , \quad dx , dy , dz = c dt$$

$$\begin{aligned}
\frac{\partial}{\partial s} \left[\left(\frac{\partial E}{\partial s} \right) (E_s) \right] &= K_{sf} D_{sf} (e^{jks}) (K_{tc} D_{tc} e^{-\int \frac{\omega}{c} dt}) \\
&= K_{sf} D_{sf} (e^{jks}) (K_{tc} D_{tc} e^{-j\omega dt}) \quad (3-16) \\
&= \frac{\partial E}{\partial s} \int E dt
\end{aligned}$$

For the case of expansion of free time varying energy

$$\begin{aligned}
\text{a-for } \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) &= \left(\frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right) \left(\int \left(\frac{\partial E}{\partial t} \right) dt \right) = \left(\frac{\partial E}{\partial t} \right) (E_t) \\
\frac{\partial}{\partial t} (K_{tf} D_{tf} \Psi_{tf}) &= K_{tf} D_{tf} \frac{\partial}{\partial t} (\Psi_{tf}) \\
&= K_{tf} D_{tf} (e^{j\omega t}) (e^{\frac{\partial}{\partial t} (j\omega t)}) \quad (4-16) \\
&= K_{tf} D_{tf} (e^{j\omega t}) (K_t D_t e^{(j\omega)}) \\
&= \frac{\partial E}{\partial t} (K_t D_t e^{(j\omega)}) = \left(\frac{\partial E}{\partial t} \right) (E_t)
\end{aligned}$$

$$\begin{aligned}
\text{b-for } \frac{\partial}{\partial t} \left[\left(\frac{\partial E}{\partial t} \right) (E_t) \right] &= \left(\frac{\partial E}{\partial t} \right) \left(\int E ds \right) \\
= \frac{\partial}{\partial t} \left[\left(\frac{\partial E}{\partial t} \right) (E_t) \right] &= \frac{\partial E}{\partial t} (K_t D_t e^{-\int (j\omega) dt}) \quad (5-16) \\
&= K_{tf} D_{tf} (e^{j\omega t}) (K_t D_t e^{-\int (j\omega) dt})
\end{aligned}$$

again , we perform the change of the integration parameters

$$\omega = kc \quad , \quad dt = \frac{dx}{c} = \frac{dy}{c} = \frac{dz}{c} \quad , \quad ds = dx dy dz$$

$$\frac{\partial}{\partial t} \left[\left(\frac{\partial E}{\partial t} \right) (E_t) \right] = K_{tf} D_{tf} (e^{j\omega t}) (K_{sc} D_{sc} e^{-\int kc \frac{ds}{c}}) \quad (6-16)$$

$$= [K_{tf} D_{tf} (e^{j\omega t})] [K_{sc} D_{sc} (e^{-jks})]$$

$$= \frac{\partial E}{\partial t} \int E ds$$

and for the quanton expansion and constraining terms

a-Expansion part

as mentioned earlier , the expansion of constrained fields is

Handled by integration process

$$\begin{aligned} \frac{\partial}{\partial s} (E_q) &= \frac{\partial}{\partial s} ((E_{sf} E_{tc})(E_{sc} E_{tf})) \\ &= K_{sf} K_{tc} D_{sf} D_{tc} \frac{\partial}{\partial x} \psi_{sf} \int \psi_{tc} dt (K_{sc} K_{tf} D_{sc} D_{tf} \int \psi_{sc} dx \frac{\partial}{\partial t} \psi_{tf}) \quad (7-16) \end{aligned}$$

$$= K_{sf} K_{tc} D_{sf} D_{tc} \left(\frac{jk}{-j\omega} \psi_{sf} \psi_{tc} \right) (K_{sc} K_{tf} D_{sc} D_{tf}) \left(\frac{j\omega}{-jk} \psi_{sc} \psi_{tf} \right)$$

$$= K_{sf} \psi_{sf} D_{sf} (K_{tc} D_{tc} \psi_{tc}) ((K_{sc} D_{sc} \psi_{sc}) (K_{tf} D_{tf} \psi_{tf}))$$

$$= (E_{sf} E_{tc}) (E_{sc} E_{tf}) = E_q$$

b- constraining term

$$\begin{aligned} & \left(\int E_{sf} ds \frac{\partial E_{tc}}{\partial t} \right) \left(\frac{\partial E_{sc}}{\partial s} \int E_{tf} dt \right) \\ &= (K_{sf} K_{tc} D_{sf} D_{tc} e^{\frac{\partial}{\partial s}(jks)} e^{\frac{\partial}{\partial t}(-j\omega t)}) (K_{sc} K_{tf} D_{sc} D_{tf} e^{\frac{\partial}{\partial s}(-jks)} e^{\frac{\partial}{\partial t}(j\omega t)}) \\ &= K_{sf} K_{tc} D_{sf} D_{tc} e^{(jk)} e^{(-j\omega)} (K_{sc} K_{tf} D_{sc} D_{tf} e^{(-jk)} e^{(j\omega)}) \\ &= K_s D_s K_t D_t = E_s E_t \end{aligned}$$

To summarize, the exponential differentiation / integration would be applied in either of the following cases

1-change of the nature of the energy field (free / constrained)

or (space varying / time varying) and vice versa

2- change in the degrees of freedom of any energy field (Dof rearrangement between fields)

17. Wave- like properties of space fabric

Energy which varies in time and varies in space has wave like

properties as it changes at periodic rate that equals

ω rad /sec ($= 2 \pi f$) and the space varying field ,which varies

along its radius $r_q (= \frac{\pi}{k})$, such that $\omega r_q = \text{constant} = \pi c$

In fact the quanton (or anti quanton) is represented by

two (wave like) equations ,

to show how the wave equations would look like for the energy

free and constrained energy fields , first remembering

that $\psi_{sf} = e^{jkx}$, $\psi_{tc} = e^{-j\omega t}$, $\psi_{sc} = e^{-jkx}$, $\psi_{tf} = e^{j\omega t}$

the free energy dominated wave parameters

$\psi_{qf} = (\psi_{sf} \psi_{tc})$ differentiating both sides w.r.t time

$$\frac{\partial \psi_{qf}}{\partial t} = \frac{\partial \psi_{tc}}{\partial t} \psi_{sf} = -j\omega \psi_{sf} \psi_{tc}$$

$$\frac{\partial^2 \psi_{qf}}{\partial t^2} = \frac{\partial^2 \psi_{tc}}{\partial t^2} \psi_{sf} = -\omega^2 \psi_{sf} \psi_{tc}$$

while differentiating w.r.t (x)

$$\frac{\partial \psi_{qf}}{\partial x} = \frac{\partial \psi_{sf}}{\partial x} \psi_{tc}$$

$$\frac{\partial^2 \psi_{qf}}{\partial x^2} = \frac{\partial^2 \psi_{sf}}{\partial x^2} \psi_{tc} = -k^2 \psi_{sf} \psi_{tc}$$

For a wave equation $\frac{\partial^2 \psi_{qf}}{\partial t^2} = c^2 \frac{\partial^2 \psi_{qf}}{\partial x^2}$ to be satisfied

$$\frac{\partial^2 \psi_{tc}}{\partial t^2} = c^2 \left(\frac{\partial^2 \psi_{sf}}{\partial x^2} \frac{\psi_{tc}}{\psi_{sf}} \right) \quad \text{or} \quad (E_{qf})_{tt} = c^2 (E_{qf})_{xx} \quad \text{as before} \quad (1-17)$$

Which is the PDE for free energy dominated field

similarly ,for the constrained energy dominated wave

$\psi_{qc} = (\psi_{sc} \psi_{tf})$ differentiating both sides w.r.t time

$$\frac{\partial \psi_{qc}}{\partial t} = \frac{\partial \psi_{tf}}{\partial t} \psi_{sc}$$

$$\frac{\partial^2 \psi_{qc}}{\partial t^2} = \frac{\partial^2 \psi_{tf}}{\partial t^2} \psi_{sc} \quad , \quad \text{while differentiating w.r.t } x$$

$$\frac{\partial \psi_{qc}}{\partial x} = \frac{\partial \psi_{sc}}{\partial x} \psi_{tf}$$

$$\frac{\partial^2 \psi_{qc}}{\partial x^2} = \frac{\partial^2 \psi_{sc}}{\partial x^2} \psi_{tf} \quad ,$$

for a wave equation $\frac{\partial^2 \psi_{qc}}{\partial t^2} = c^2 \frac{\partial^2 \psi_{qc}}{\partial x^2}$ to be satisfied

$$\frac{\partial^2 \psi_{tf}}{\partial t^2} = c^2 \left(\frac{\partial^2 \psi_{sc}}{\partial x^2} \frac{\psi_{tf}}{\psi_{sc}} \right) \quad \text{or} \quad (E_{qc})_{tt} = c^2 (E_{qc})_{xx} \quad (2-17)$$

this PDE for the constrained energy dominated field,

which also shows how for wave equation of space and time

varying fields would look like , but does the quanton energy

density equation in its differential / integral form really represent

two wave equations ?

a-For the free energy dominated term ($\frac{\partial E}{\partial s} \int E dt$)

Differentiating with respect to time $\frac{\partial}{\partial t} (\frac{\partial E}{\partial s} \int E dt) =$

$$[(\frac{\partial x}{\partial t}) \frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \int E dt] + [\frac{\partial E}{\partial s} \frac{\partial}{\partial t} (\int E dt)]$$

$$= [c \frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \int E dt] + [\frac{\partial E}{\partial s} \frac{\partial}{\partial t} (\int E dt)]$$

Differentiating again with respect to time

$$\frac{\partial}{\partial t} [c \frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \int E dt] + \frac{\partial}{\partial t} [\frac{\partial E}{\partial s} \frac{\partial}{\partial t} (\int E dt)]$$

$$= [c (\frac{\partial x}{\partial t}) \frac{\partial^2}{\partial x^2} (\frac{\partial E}{\partial s}) \int E dt] + [c \frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \frac{\partial}{\partial t} (\int E dt)]$$

$$+ [(\frac{\partial x}{\partial t}) \frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \frac{\partial}{\partial t} (\int E dt)] + [\frac{\partial E}{\partial s} \frac{\partial^2}{\partial t^2} (\int E dt)] =$$

$$[c^2 \frac{\partial^2}{\partial x^2} (\frac{\partial E}{\partial s}) \int E dt] + 2 c [\frac{\partial}{\partial x} (\frac{\partial E}{\partial s}) \frac{\partial}{\partial t} (\int E dt)] + [\frac{\partial E}{\partial s} \frac{\partial^2}{\partial t^2} (\int E dt)]$$

For the same energy type differentiating with respect to x

$$\begin{aligned} \frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \int E dt \right) &= \left[\frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \int E dt \right] + \left[\frac{\partial E}{\partial s} \frac{1}{\frac{\partial x}{\partial t}} \frac{\partial}{\partial t} \left(\int E dt \right) \right] \\ &= \left[\frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \int E dt \right] + \left[\frac{\partial E}{\partial s} \frac{1}{c} \frac{\partial}{\partial t} \left(\int E dt \right) \right] \end{aligned}$$

differentiating again with respect to x

$$\begin{aligned} \frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \int E dt \right] + \frac{\partial}{\partial x} \left[\frac{1}{c} \frac{\partial E}{\partial s} \frac{\partial}{\partial t} \left(\int E dt \right) \right] \\ = \left[\frac{\partial^2}{\partial x^2} \left(\frac{\partial E}{\partial s} \right) \int E dt \right] + \left[\frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \frac{1}{\frac{\partial x}{\partial t}} \frac{\partial}{\partial t} \left(\int E dt \right) \right] + \\ \left[\frac{1}{c} \frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \frac{\partial}{\partial t} \left(\int E dt \right) \right] + \left[\frac{1}{c} \frac{\partial E}{\partial s} \frac{1}{\frac{\partial x}{\partial t}} \frac{\partial^2}{\partial t^2} \left(\int E dt \right) \right] \\ = \left[\frac{\partial^2}{\partial x^2} \left(\frac{\partial E}{\partial s} \right) \int E dt \right] + 2 \frac{1}{c} \left[\frac{\partial}{\partial x} \left(\frac{\partial E}{\partial s} \right) \frac{\partial}{\partial t} \left(\int E dt \right) \right] + \left(\frac{1}{c} \right)^2 \left[\frac{\partial E}{\partial s} \frac{\partial^2}{\partial t^2} \left(\int E dt \right) \right] \end{aligned}$$

by comparing the results of both double differentiation

$$\frac{\partial^2}{\partial t^2} \left(\frac{\partial E}{\partial s} \int E dt \right) = c^2 \frac{\partial^2}{\partial x^2} \left(\frac{\partial E}{\partial s} \int E dt \right)$$

which is customary form of for a wave relation

b- For the constrained energy dominated wave

$$E_{qc} = \int E ds \frac{\partial E}{\partial t} \quad , \quad \text{expanding in x direction}$$

$$\begin{aligned}
\frac{\partial}{\partial x} \left(\int E ds \frac{\partial E}{\partial t} \right) &= \left[\frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\int E ds \frac{\partial}{\partial x} \left(\frac{\partial E}{\partial t} \right) \right] \\
&= \left[\frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\int E ds \frac{1}{\frac{\partial x}{c}} \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] \\
&= \left[\frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\frac{1}{c} \int E ds \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right]
\end{aligned}$$

Differentiating again with respect to x- axis

$$\begin{aligned}
\frac{\partial}{\partial x} \left[\frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \frac{\partial}{\partial x} \left[\frac{1}{c} \int E ds \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] \\
= \left[\frac{\partial^2}{\partial x^2} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\frac{\partial}{\partial x} \left(\int E ds \right) \frac{1}{\frac{\partial x}{c}} \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] + \\
\left[\frac{1}{c} \frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] + \left[\frac{1}{c} \int E ds \frac{1}{\frac{\partial x}{c}} \frac{\partial^2}{\partial t^2} \left(\frac{\partial E}{\partial t} \right) \right] = \\
= \left[\frac{\partial^2}{\partial x^2} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + 2 \left[\frac{1}{c} \frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] + \left(\frac{1}{c} \right)^2 \left[\int E ds \frac{\partial^2}{\partial t^2} \left(\frac{\partial E}{\partial t} \right) \right]
\end{aligned}$$

when differentiating with respect to time

$$\begin{aligned}
\frac{\partial}{\partial t} \left(\int E ds \frac{\partial E}{\partial t} \right) &= \left[\left(\frac{\partial x}{\partial t} \right) \frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\int E ds \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right] \\
&= \left[c \frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \left[\int E ds \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right]
\end{aligned}$$

Differentiating again with respect to time

$$\frac{\partial}{\partial t} \left[c \frac{\partial}{\partial x} \left(\int E ds \right) \frac{\partial E}{\partial t} \right] + \frac{\partial}{\partial t} \left[\int E ds \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \right]$$

$$= [c \left(\frac{\partial x}{\partial t} \right) \frac{\partial^2}{\partial x^2} (\int E ds) \frac{\partial E}{\partial t}] + [c \frac{\partial}{\partial x} (\int E ds) \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right)] +$$

$$[\left(\frac{\partial x}{\partial t} \right) \frac{\partial}{\partial x} (\int E ds) \frac{\partial E}{\partial t}] + [\int E ds \frac{\partial^2}{\partial t^2} \left(\frac{\partial E}{\partial t} \right)] =$$

$$= [c^2 \frac{\partial^2}{\partial x^2} (\int E ds) \frac{\partial E}{\partial t}] + 2 [c \frac{\partial}{\partial x} (\int E ds) \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right)] + [(\int E ds) \frac{\partial^2}{\partial t^2} \left(\frac{\partial E}{\partial t} \right)]$$

by comparing the results of both double differentiations

$$\frac{\partial^2}{\partial t^2} \left(\int E ds \frac{\partial E}{\partial t} \right) = c^2 \frac{\partial^2}{\partial x^2} \left(\int E ds \frac{\partial E}{\partial t} \right)$$

which is the usual form of the wave equation

18. quanton evolution and degrees of freedom

evolution of the quanton takes place as both free fields

(E_{sf}) and (E_{tf}) coexist

1-as free energy field expands by variation in space It must vary in

time , so , a constrained time varying field appears

$$a - \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) = \left[\frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) \right] \left[\int \left(\frac{\partial E}{\partial s} \right) ds \right] = \left(\frac{\partial E}{\partial s} \right) (E_s) \quad (1-18)$$

expanding again in time

$$b - \frac{\partial}{\partial s} \left[\left(\frac{\partial E}{\partial s} \right) (E_s) \right] = \frac{\partial}{\partial s} \left(\frac{\partial E}{\partial s} \right) \int (E) dt = \left(\frac{\partial E}{\partial s} \right) \left(\int E dt \right) \quad (2-18)$$

2-as the time varying field (E_{tf}) expands , a part of it must vary in

space in the form of constrained space varying energy field

$$a - \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) = \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \left(\int \frac{\partial E}{\partial t} dt \right) = \left(\frac{\partial E}{\partial t} \right) (E_t) \quad (3-18)$$

when expanding in space

$$b - \frac{\partial}{\partial t} \left[\left(\frac{\partial E}{\partial t} \right) (E_t) \right] = \frac{\partial}{\partial t} \left(\frac{\partial E}{\partial t} \right) \int (E) ds = \left(\frac{\partial E}{\partial t} \right) \int E ds \quad (4-18)$$

and since non of fields possesses all four Dof's

neither field can exist independently , and for the stable

quanton energy density equation becomes

$$E_q = \left(\frac{\partial E}{\partial s} \int E dt \right) \left(\int E ds \frac{\partial E}{\partial t} \right) = (E_{sf} E_{tc}) (E_{sc} E_{tf}) = E_{qf} E_{qc} \quad (5-18)$$

which expresses two apparently separate (but otherwise linked)

fields

3-for space constrained energy field (E_{sc}),its energy Dof equals

one third of the corresponding free energy field (E_{sf})

4-for free time varying energy field(E_{tf}) , its energy degree of freedom equals one third of the corresponding time constrained energy field (E_{tc}) , the previous discussion can be summarized in the following 4 equations by solving them the quanton Dof's for the four energy fields can be obtained


$$Dof_{sf} = 3 Dof_{sc} \quad , \quad Dof_{tc} = 3 Dof_{tf}$$

$$Dof_{sf} + Dof_{sc} = 3 \quad , \quad Dof_{tf} + Dof_{tc} = 1 \quad ,$$

which gives the following results

$$Dof_{sf} = 2.25 \quad , \quad Dof_{sc} = 0.75$$

$$Dof_{tf} = 0.25 \quad , \quad Dof_{tc} = 0.75$$



Fig(5).Tree diagram for the evolution and the degrees of freedom of quanton energy fields

5- as free and constrained energy fields create corresponding field

binding interaction inside the quanton , the result of such this

interaction would be a rearrangement of the degrees of freedom in

such a way that maximizes its stability

this can be ensured by creating a symmetry of free and

constrained degrees of freedom (discussed in the section :

quanton stable Dof) , the stable energy Dof (after

rearrangement)

$$Dof_{sf} = 2 \quad , \quad Dof_{tf} = 0.5$$

$$Dof_{sc} = 1 \quad , \quad Dof_{tc} = 0.5$$

4-for the quanton system despite having a constrained energy

fields, it is dominated by the free energy field of the form $(\frac{\partial E}{\partial s} \frac{\partial E}{\partial t})$

since this energy term represents 2.5 degrees of freedom while the

constrained type $(\int E ds \int E dt)$ constitutes 1.5 energy Dof's out

of four

5-the number of (unbound) degrees of freedom of free energy

fields is equivalent to the number of free energy degrees

of freedom (space plus time varying) minus the energy

constrained degrees of freedom (space and time varying)

8- (unbound) free field is manifested in the form of quanton

inflation

$$D_{sfu} D_{tfu} \text{ (unbound field strength)} = \frac{(D_{sf} D_{tf})}{(D_{sc} D_{tc})} = \frac{c^{2.25} c^{0.25}}{c^{0.75} c^{0.75}} = c \quad (6-18)$$

unbound free Dof = (\sum (free Dof) – \sum (constrained Dof)

$$= [(Dof_{sf}) + (Dof_{tf})] - [(Dof_{sc}) + (Dof_{tc})] = (2.5-1.5) = +1$$

19.Variation of quanton energy fields with time

not only the unbound energy fields $E_{sfu} E_{sfu}$ of the quanton

(or $E_{scu} E_{scu}$ for anti quanton) which change with time as the

quanton (or anti quanton) expands , but rather all the other

energy fields , and this is so , to ensure the uniformity of energy

density inside the quanton

19.a-Variation of space varying energy with time

$$\frac{\partial E_{sf}}{\partial t} = \frac{\partial E_{sf}}{\partial x} \frac{\partial x}{\partial t} = c \frac{\partial E_{sf}}{\partial x} \quad (1-19)$$

$$\frac{\partial E_{sf}}{\partial x} = j k E_{sf} \quad (2-19)$$

$$\frac{\partial E_{sf}}{\partial t} = j k c E_{sf} \quad (3-19)$$

b- time varying energy field variation

$$\frac{\partial E_{tf}}{\partial t} = j \omega E_{tf} \quad (4-19)$$

c-Relative rate of Variation between different energy fields

$$\frac{\partial E_{sf}}{\partial E_{tf}} = \frac{\partial E_{sf}}{\partial x} \left(\frac{\partial x}{\partial t} \right) \frac{1}{\left(\frac{\partial E_{tf}}{\partial t} \right)} = (jkc E_{sf}) \left(\frac{1}{j\omega E_{tf}} \right) = \frac{E_{sf}}{E_{tf}} = \frac{D_{sf}}{D_{tf}} \quad (5-19)$$

The same results can be reached when considering the wave

parameters of energy fields

$$\frac{\partial \Psi_{tf}}{\partial \Psi_{sf}} = \left(\frac{\partial \Psi_{tf}}{\partial t} \right) \left(\frac{1}{\frac{\partial x}{\partial t}} \right) \left(\frac{1}{\frac{\partial \Psi_{sf}}{\partial x}} \right) \quad (6-19)$$

$$\text{given that } \Psi_{tf} = e^{+j\omega t} \quad , \quad \frac{\partial \Psi_{tf}}{\partial t} = j\omega \Psi_{tf}$$

$$\Psi_{sf} = e^{+jk(x+y+z)} \quad , \quad \frac{\partial \Psi_{sf}}{\partial x} = jk \Psi_{sf}$$

$$\frac{\partial \Psi_{tf}}{\partial \Psi_{sf}} = \frac{1}{c} \frac{\omega}{k} = 1 = \text{constant} \quad , \quad (7-19)$$

$$\text{While from before } \frac{\partial E_{tf}}{\partial t} = j\omega E_{tf} \quad , \quad \frac{\partial E_{sf}}{\partial x} = jk E_{sf}$$

$$\frac{\partial E_{tf}}{\partial E_{sf}} = \frac{1}{c} \frac{\omega}{k} \frac{E_{tf}}{E_{sf}} = \frac{E_{tf}}{E_{sf}} = \frac{D_{tf}}{D_{sf}} \quad , \quad \text{which means that} \quad (8-19)$$

1-the rate of variation of energy fields wave parameters

with respect to each other is constant (=1) (same rate of variation for all energy fields)

2-relative rate of variation in time of all energy fields is equal to the ratio between their degrees of freedom and this is due to the uniformity of their variation parameters

20. Energy field parameters

while the energy degrees of freedom of the quanton total energy (packet energy E_p) are in terms of the wave parameters (k , ω , r_q) , the energy degrees of freedom for the energy fields are in terms of the constant (c) as pointed out earlier , this is because the constant (c) is what determines the relationship between the rate of variation of space and time varying fields , previously , the energy density constant (h_q) was determined while using an approximative method , now the analytical method

shall be used to assess it

Recalling first that the quanton fields are infinite in range , and

The definition of the variation parameters of E_{qf} , E_{qc} fields which

corresponds to an exponentially decaying field away from the

quanton , the free and constrained fields can be put

as $E_{qf}(x) = E_{qf} e^{-j(\frac{x}{2r_q})}$ (free energy dominated field)

$E_{qc}(x) = E_{qc} e^{-j(\frac{x}{r_q})}$ (constrained energy dominated field)

and the quanton energy density is in the form

$$E_q = (E_{sf}E_{tc})(E_{sc}E_{tf}) = E_{qf} E_{qc} = E_q e^{-j(\frac{x}{r_q})} \quad (1-20)$$

E_q : represents the average energy density over time

to assess the entire energy stored in both fields , the quanton

packet energy would be equal to the volumetric integration

$$E_p = \frac{h\omega}{2\pi} = \iiint_{-\infty}^{\infty} E_q e^{-j(\frac{x+y+z}{r_q})} dx dy dz = \quad (2-20)$$

$$= (2)^3 \iiint_0^{\infty} E_q e^{-j(\frac{x+y+z}{r_q})} dx dy dz \quad (\text{ symmetric integration})$$

$$\begin{aligned}
 & \mathbf{x, y, z} = \infty \\
 & = 8 (r_q)^3 E_q e^{-j(\frac{x+y+z}{r_q})} \Big|_{\mathbf{x, y, z} = 0} = 8 (r_q)^3 E_q
 \end{aligned}$$

$$E_q = \frac{h\omega}{16\pi r_q^3} = \frac{h\omega^4}{16\pi^4 c^3}, \quad (3-20)$$

$$\text{where } \frac{h}{16\pi^4 c^3} = h_q \text{ (energy density constant)} \quad (4-20)$$

to relate the average energy density E_q to its maximum value

E_{q0} over time, we use the quanton /anti quanton wave model

$$E_q = \frac{1}{2}(E_{qf} + cE_{qc}) * \frac{1}{2}\left(\frac{E_{qf}}{c} + E_{qc}\right)$$

and since $E_{qfo} = cE_{qco}$

$$E_q = E_{qfo} \cos\left(\frac{\pi r}{2r_q} - \omega t\right) E_{qco} \cos\left(\frac{\pi r}{2r_q} - \omega t\right) = E_{q0} \cos^2\left(\frac{\pi r}{2r_q} - \omega t\right) \quad (5-20)$$

The average value of a periodic function is defined as

$$E_q = \frac{1}{T} \int_0^T E_{q0} (t) \quad (6-20)$$

$$E_q = E_{q0} \int_0^T \cos^2\left(\frac{\pi r}{2r_q} - \omega t\right) dt \quad (7-20)$$

The value of this integration equals to $\left(\frac{1}{2}\right)$

$$E_{q0} = 2E_q = \frac{h \omega^4}{8\pi^4 c^3}, \quad (8-20)$$

Here , the quanton is represented by an equivalent volume = $8 r_q^3$

The same result can be reached alternatively ,when calculating the

Vacuum energy density ρ_v at any point in space as the summation

of individual energy density contributions of (N_q) quantons

$\rho_v = \sum_i^{N_q} \rho_{vi}$, which leads to the same integration and the same

energy density constant , and in general the vacuum energy

density is equivalent to the quanton average energy density

$$\rho_v = E_q \quad (8-20)$$

While in terms of the wave parameter (k), or the quanton radius

the quanton energy density takes the form

$$E_q = \left(\frac{h}{16 \pi^4 c^3} \right) k^4 c^4 = \frac{h c}{16 r_q^4} \quad (9-20)$$

energy as it expands by variation in space and time , it has four

degrees of freedom, this can be used to define the various

space and time varying fields

$$E_q = \frac{h}{16 \pi^4 c^3} k^4 c^4 = \text{constant} * \left(\frac{2\pi}{\lambda}\right)^4 c^4 = \frac{\text{constant}}{4D \text{ volume}} * c^4 \quad (10-20)$$

This relationship does not only expresses a volumetric relationship

of energy density as it expands into a 4 D volume , but it

expresses an energy density – degree of freedom relationship as it

can be put in terms of the wave parameters ($k, \omega, \frac{1}{r_q}$)

parameter	Analytical method	Approximative method
Integration volume	Universe volume	N/A
Equivalent volume	$\frac{\text{Universe volume}}{\text{number of quantons}}$	$\frac{4\pi}{3} r_q^3$
Quanton shape	cubic	spherical
Quanton dimensions	Each side = $2 r_q$	Radius = r_q
Density constant h_q	$\frac{h}{16 \pi^4 c^3}$	$\frac{3 h}{8 \pi^5 c^3}$
Energy density inside the quanton	$E_q (= \frac{E_{q0}}{2})$	$(\frac{6}{\pi}) E_q$
Free and constrained field	Inside and outside quanton (both Propagate throughout space)	Inside quanton only
Vacuum energy density ρ_v	E_q	E_q

Table (3) why vacuum energy density is uniform throughout space no difference between the analytical and approximative method of determining the quanton energy density constant h_q

the energy degrees of freedom which can be put as

$$D_q = c^4 = D_{sf} D_{sc} D_{tf} D_{tc} = \text{the field strength parameter of energy}$$

fields

$$\text{where } D_{sf} = c^{Dof_{sf}} \quad , \quad D_{sc} = c^{Dof_{sc}} \quad (11, 12-20)$$

$$D_{tf} = c^{Dof_{tf}} \quad , \quad D_{tc} = c^{Dof_{tc}} \quad (13, 14-20)$$

$$E_q = \frac{h}{16 \pi^4 c^3} k^4 c^4 = K_q^4 c^4 \quad (15, 16-20)$$

the quantity $K_q^4 = \left(\frac{h}{16 \pi^4 c^3} k^4 \right)$ can be put as

$$K_q^4 = h_q k^4 = K_{sf} K_{sc} K_{tf} K_{tc} \quad (= \text{energy field intensity parameter}) \quad (16-20)$$

$$\text{where } K_{sf} = K_q = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} k \quad (17-20)$$

$$K_{sc} = K_q = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} k \quad , \quad K_{tf} = K_{tc} = K_q = \sqrt[4]{\left(\frac{h}{16 \pi^4 c^3}\right) \frac{\omega}{c}} \quad (18-20)$$

it must be noted that while $\frac{E_q}{\omega^4} = \left(\frac{h}{16 \pi^4 c^3}\right) = h_q = \text{constant}$,

its fourth root is not a constant , $\sqrt[4]{\frac{E_q}{\omega^4}}$ or $\sqrt[4]{\frac{E_q}{k^4}} \neq \text{constant}$

the division of the field intensity parameter does not follow the

energy degree of freedom but follows the division of field types

(free dominated and constrained dominated fields) otherwise the

Energy fields E_{sf} , E_{tc} or E_{sc} , E_{tf} could exist independently

one can be drawn to think that the division of (K_q^4) between

various energy fields such that $K_{sf} = K_q^{Dof_{sf}} = K_q^2$, or

$K_{tf} = K_q^{Dof_{tf}}$, but since there are no wave parameters in nature of

k^2 or $\omega^{0.5}$ due to the symmetry of the wave behavior between

various fields which is previously defined as $\frac{\partial \Psi_{tf}}{\partial \Psi_{sf}} = \frac{1}{c} \frac{\omega}{k} = \text{constant}$

and
$$\frac{\partial E_{tf}}{\partial E_{sf}} = \frac{1}{c} \frac{\omega}{k} \frac{E_{tf}}{E_{sf}} = \frac{E_{tf}}{E_{sf}} = \frac{D_{tf}}{D_{sf}}$$

this leads to the following result : $K_{sf} = K_{sc} = K_{tf} = K_{tc} = K_q$ (19-20)

Finally , we can write the energy fields themselves as

$$E_{sf} = E_{sfo} \quad \psi_{sf} = K_q D_q^{Dof_{sf}} \psi_{sf} = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} k c^{2.0} \psi_{sf} = \sqrt[4]{\frac{h}{16 c^3}} \frac{c^{2.0}}{r_q} \psi_{sf} \quad (20-20)$$

$$E_{sc} = E_{sco} \quad \psi_{sc} = K_q D_q^{Dof_{sc}} \psi_{sc} = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} k c^{1.0} \psi_{sc} = \sqrt[4]{\frac{h}{16 c^3}} \frac{c^{1.0}}{r_q} \psi_{sf}$$

(21-20)

$$E_{tf} = E_{tfo} \quad \psi_{tf} = K_q D_q^{Dof_{tf}} \quad \psi_{tf} = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} \frac{\omega}{c} c^{0.5} \quad \psi_{tf} = \sqrt[4]{\frac{h}{16 c^3}} \frac{c^{0.5}}{r_q} \psi_{tf}$$

(22-20)

$$E_{tc} = E_{tco} \quad \psi_{tc} = K_q D_q^{Dof_{tc}} \quad \psi_{tc} = \sqrt[4]{\frac{h}{16 \pi^4 c^3}} \frac{\omega}{c} c^{0.5} \quad \psi_{tc} = \sqrt[4]{\frac{h}{16 c^3}} \frac{c^{0.5}}{r_q} \psi_{tc}$$

(23-20)

$$\frac{E_{sf}}{E_{tc}} = \frac{K_q D_q^{Dof_{sf}} \psi_{sf}}{K_q D_q^{Dof_{tc}} \psi_{tc}} = \frac{D_q^{Dof_{sf}} \psi_{sf}}{D_q^{Dof_{tc}} \psi_{tc}} = c^{1.5} \frac{\psi_{sf}}{\psi_{tc}}$$

(24-20)

a unified value of (K_q) for all energy fields ensures that the

relationship between the different fields depends only on their the

degrees of freedom and not on the intensity of such fields

in general a field energy can be seen as the product of two terms

field energy = field intensity (defined in terms of : K_q) * field

strength (D_q : defined in terms of energy degrees of freedom)

21.Dimensions of vector energy fields

While being a scalar quantity , energy as it expands in the form of

space and time varying fields which are vector quantities

individual energy content of various fields in the form

$E_p = \int_{V_q} E_{sf} dV$ does not exist ,

and this is due to the fact that quanton energy fields are

inextricably linked to the quanton volume in a dependence

relationship , this does not make it possible to determine the

total energy of each individual field

the energy fields must be defined in terms of the quanton

dimensions , in addition to energy dimensions and degrees of

freedom for each energy field

the quanton radius (r_q) and , its volume (V_q) are not constant but

rather inversely proportional to its packet energy content , and

consequently its energy fields

while $V_q = \text{fn}(r_q^3) = \text{fn}(\lambda^3) = \text{fn}(\frac{1}{\omega^3})$

and $E_q = E_{sf} E_{sc} E_{tf} E_{tc} = \left(\frac{h}{16 \pi^4 c^3}\right) \omega^4 = \text{constant} * \omega^4$

hence $V_q = \text{fn}\left(\frac{\omega}{E_{sf} E_{sc} E_{tf} E_{tc}}\right)$ ie quanton volume is dependent on the

product of all four energy field densities

dimensions of individual energy fields are expected to be as

follows

$$(E_{sf}) = \left[\sqrt[4]{\left(\frac{h}{16\pi^4 c^3}\right)} k c^{2.0} \psi_{sf} \right]$$

$$[E_{sf}] = M^{0.25} L^{0.5-0.75-1+2.0} T^{-0.25+0.75-2.0} = M^{0.25} L^{0.75} T^{-1.50} \quad (1-21)$$

$$[E_{sc}] = \left[\sqrt[4]{\left(\frac{h}{16\pi^4 c^3}\right)} k c^{1.0} \psi_{sc} \right] = M^{0.25} L^{-0.25} T^{-0.50} , \quad (2-21)$$

$$[E_{tc}] = \left[\sqrt[4]{\left(\frac{h}{16\pi^4 c^3}\right)} \frac{w}{c} c^{0.50} \psi_{tc} \right] = M^{0.25} L^{-0.75} T^{0.00} \quad (3-21)$$

$$[E_{tf}] = \left[\sqrt[4]{\left(\frac{h}{16\pi^4 c^3}\right)} \frac{w}{c} c^{0.50} \psi_{tc} \right] = M^{0.25} L^{-0.75} T^{0.00} \quad (4-21)$$

21 b. effect of fixed relative ratio between space and time varying fields

As it had been mentioned previously , exponential degrees of

freedom while in terms of the constant (c) , provide a

mechanism for the division of energy density between the space

and time varying energy fields so as to maintain a constant ratio

between them , for space varying fields value (in magnitude)

$$E_s = E_{sf} E_{sc} = (K_q c^2) (K_q c) = K_q^2 c^3 = \sqrt[2]{\left(\frac{h}{16\pi^4 c^3}\right)} k^2$$

for time varying energy fields

$$E_t = E_{tf} E_{tc} = (K_q c^{0.5}) (K_q c^{0.5}) = K_q^2 c = \sqrt[2]{\left(\frac{h}{16\pi^4 c^3}\right)} k^2$$

the relative ratio between space and time varying energy fields

$$\frac{E_{sf} E_{sc}}{E_{tf} E_{tc}} = \text{constant} = c^2 , \text{ the ratio of the space and time varying}$$

energy fields does vary as the wave parameters change

very high values of (k , ω) corresponding to relatively high

percentage for the share of the time varying fields, as the universe

expands , this percentage drops while the percentage of the space

varying energy fields increases comparatively

22. field representation inside the quanton

While free and constrained fields extend beyond the quanton , yet

an expression for those fields inside the quanton can be provided

2- under condition of equipartition of energy in spatial axes

quanton energy fields must be at any instant symmetrically

expressed in all three dimensional space , based on this ,

each of the quanton fields is represented by concentric toroidal

solenoid

while in the proper (own) frame of reference the free dominated

field components can be defined as

$$E_{qfxi}^* = 0 \quad (1-22)$$

$$E_{qfyi}^* = E_{qfi} \sin (\omega t) \quad (2-22)$$

$$E_{qfzi}^* = E_{qfi} \cos (\omega t) \quad (3-22)$$

And the constrained energy dominated field components

$$E_{qcxi}^* = E_{qci} \cos (\omega t) \quad (4-22)$$

$$E_{qcyi}^* = E_{qci} \sin (\omega t) \quad (5-22)$$

$$E_{qfzi}^* = 0 \quad (6-22)$$

The proper frame of reference (x^*, y^*, z^*) is related to the observer frame of reference (x, y, z) via 3 dimensional transformation matrix (T)

$$\begin{vmatrix} E_{qfxi}^* \\ E_{qfyi}^* \\ E_{qfzi}^* \end{vmatrix} = [T] \begin{vmatrix} E_{qfxi} \\ E_{qfyi} \\ E_{qfzi} \end{vmatrix} \quad (7-22)$$

$$\begin{vmatrix} E_{qcxi}^* \\ E_{qcyi}^* \\ E_{qczi}^* \end{vmatrix} = [T] \begin{vmatrix} E_{qcxi} \\ E_{qcyi} \\ E_{qczi} \end{vmatrix} \quad (8-22)$$

The matrix (T) which has the angles θ, ϕ, ψ (Euler's angles)

as its elements

and the resultant fields are $E_{qfx} = \sum_i^n E_{qfxi}$ (9-22)

$$E_{qfy} = \sum_i^n E_{qfyi} \quad , \quad E_{qfz} = \sum_i^n E_{qfzi} \quad (10, 11-22)$$

$$E_{qcx} = \sum_i^n E_{qcxi} \quad , \quad E_{qcy} = \sum_i^n E_{qcyi} \quad , \quad E_{qcz} = \sum_i^n E_{qczi} \quad (12,13,14-22)$$

23. what maintains the integrity of the quanton ?

1-Free and constrained energy dominated fields E_{qf} and E_{qc}

are interacting through free / constrained energy field interaction

this interaction creates a binding relationship that maintains

the integrity of the quanton

2- any radiative energy that leaves the quanton must have four

degrees of freedom (transmission of energy through space can

only take place while fields varying in space and time have those

four Dof's)

under such a condition , no individual field (E_{qf} or E_{qc}) can leave

the quanton independently , instead , both fields can leave the

quanton conjointly in the form of electromagnetic waves

24.quanton wave form : (Q+AQ) pair

This model illustrates that the quanton -anti quanton pair

Would create a form of quanton waves , later this concept

would be used to develop a model for electromagnetic waves in

terms of space and time varying fields

quantons and anti quantons exist in pairs in the form (Q+AQ)

this linear superposition form is due to the fact that either quanton

or anti quanton is a separate but not independent energy system

as the pair is considered to be a single quantum entity .

to fulfil the wave behaviour (linear supposition of fields), the Dof

symmetry condition must be satisfied

a-For the higher degree of freedom field pair (2.5 Dof's)

$$(E_{qc})_{aq} = (E_{qf})_q \quad \text{or} \quad (\text{Dof}_{qc})_{aq} = (\text{Dof}_{qf})_q \quad (1-24)$$

b-for the lower degree of freedom pair (1.5 Dof's)

$$(E_{qf})_{aq} = (E_{qc})_q \quad \text{or} \quad (\text{Dof}_{qf})_{aq} = (\text{Dof}_{qc})_q \quad (2-24)$$

a model for the energy fields given that $E_q = E_{sf} E_{tc} E_{sc} E_{tf} = E_{qf} E_{qc}$

and $E_{aq} = \left(\frac{E_{sf} E_{tc}}{c}\right) (c E_{sc} E_{tf})$ wave form is as follows

$$\text{higher Dof } E_{wff} = \frac{1}{2} [(E_{qf})_q + (E_{qc})_{aq}] \quad (3-24)$$

$$\text{lower Dof : } E_{wcc} = \frac{1}{2} [[E_{qc}]_q + (E_{qf})_{aq}] \quad (4-24)$$

$$E_{wff}(x) = \frac{1}{2} (D_{sf} D_{tc} + c D_{sc} D_{tf}) = \frac{1}{2} K_q^2 c^{2.5} \cos\left(\frac{\pi x}{2r_q}\right) - \omega t \quad (5-24)$$

$$E_{wcc}(x) = \frac{1}{2} (D_{sc} D_{tf} + \frac{1}{c} D_{sf} D_{tc}) = \frac{1}{2} K_q^2 c^{1.5} \cos\left(\frac{\pi x}{r_q}\right) - \omega t \quad (6-24)$$

the symmetry between free and constrained fields Dof's

does not mean that (Q-AQ) would not expand or there would not be

radiative energy release from the pair as the Q/AQ pair expands

while the energy density of such a pair

$$E_q = \frac{1}{2} [(E_{qf})_q + (E_{qc})_{aq}] * \frac{1}{2} [[E_{qc}]_q + (E_{qf})_{aq}]$$

and due to the symmetry of interaction where $(E_{qf})_q = (E_{qc})_{aq}$

$$[E_{qc}]_q = (E_{qf})_{aq}$$

$$\begin{aligned} E_q &= \frac{1}{4} \frac{1}{c} E_{qf}^2 + 2 * \frac{c}{4} E_{qf} E_{qc} + \frac{c}{4} E_{qc}^2 \\ &= \frac{1}{4} \left(\frac{E_{qf}^2}{c} + 2 E_{qf} E_{qc} + c E_{qc}^2 \right) = E_{qf} E_{qc} \end{aligned}$$

25. Electromagnetic waves as space and time varying fields

The difference between quanton – anti quanton pair and electromagnetic waves lie in the fact that electromagnetic waves propagate in linear directions, and consequently, one degree of freedom is subtracted from space varying fields (free and constrained) , as it becomes a kinetic degree of freedom , this relativistic effect is split equally between free and constrained fields or each of the free and the constrained waves have one half of Dof less than the corresponding quaton fields, radiative (electromagnetic) energy is released from the quanton in the following one dimensional form

Propagation of electromagnetic energy long the x- direction

The formulation of electromagnetic waves in terms of energy fields

Depends on the system of units

Under the (Esu) system U (volumetric electromagnetic energy

$$\text{density) } = E^2 = c^2 B^2$$

$$(\epsilon)= 1 \quad , \mu = \frac{1}{c^2} \quad , \text{ under such system}$$

electric and the magnetic fields are defined as follows

$$E_f(x) = \frac{E_{qf}(x)}{\sqrt{c}} = \frac{E_{sf}(x) E_{tc}}{\sqrt{c}} \quad , \quad B_c(x) = \frac{E_{qc}(x)}{\sqrt{c}} = \frac{E_{sc}(x) E_{tf}}{\sqrt{c}} \quad (1-25)$$

where $E_f(x)$ is the electric field due to the free energy dominated

wave , $B_c(x)$ is the magnetic field due to the constrained energy

dominated wave which propagate along x- axis

$$\text{given that } \cos(kx-\omega t) = \frac{1}{2} (e^{j(kx-\omega t)} + e^{-j(kx-\omega t)})$$

define the electromagnetic (sinusoidal waves) as $E(x)$, $B(x)$

$$E(x) = \frac{1}{2} (E_f(x) + c B_c(x)) = \frac{1}{2} \left(\frac{E_{sf}(x) E_{tc}}{\sqrt{c}} + \sqrt{c} E_{sc}(x) E_{tf} \right) \quad (2-25)$$

$$B(x) = \frac{1}{2} (B_c(x) + \frac{1}{c} E_f(x)) = \frac{1}{2} \left(\frac{E_{sc}(x) E_{tf}}{\sqrt{c}} + \frac{1}{c} \frac{E_{sf}(x) E_{tc}}{\sqrt{c}} \right) \quad (3-25)$$

for the (si) system of units

$$U = \epsilon_0 E^2 = \frac{1}{\mu_0} B^2$$

$$E_f(x) = \frac{E_{qf}(x)}{\sqrt{c}} = \frac{E_{sf}(x) E_{tc}}{\sqrt{c}} \quad , \quad B_c(x) = \frac{E_{qc}(x)}{\sqrt{c}} = \frac{E_{sc}(x) E_{tf}}{\sqrt{c}} \quad (4-25)$$

define the electromagnetic (sinusoidal waves) as $E(x)$, $B(x)$

$$E(x) = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} (E_f(x) + c B_c(x))$$

$$= \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sf}(x) E_{tc}}{\sqrt{c}} + \sqrt{c} E_{sc}(x) E_{tf} \right) \quad (5-25)$$

$$B(x) = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} (B_c(x) + \frac{1}{c} E_f(x)) = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sc}(x) E_{tf}}{\sqrt{c}} + \frac{1}{c} \frac{E_{sf}(x) E_{tc}}{\sqrt{c}} \right) \quad (6-25)$$

$$B(x) = \frac{1}{2} \left(\frac{E_{qc}(x)}{\sqrt{\epsilon_0} \sqrt{c}} + \sqrt{\mu_0} \frac{E_{qf}(x)}{\sqrt{c}} \right) \quad (7-25)$$

and as a magnitude, $E_o(x) = \left(\frac{1}{\sqrt{\epsilon_0}} \sqrt{\frac{h}{16 \pi^4 c^3}} \right) (k^2 c^2) (Dof = 2) \quad (8-25)$

$$B_o(x) = \left(\frac{1}{\sqrt{\epsilon_0}} \sqrt{\frac{h}{16 \pi c^3}} \right) (k^2 c) (Dof = one) \quad (9-25)$$

To note that

1-space and time varying energy fields leave the quanton in the

form of electromagnetic (radiation) energy , where there is no

field component along the direction of the wave propagation , this

absence of fields in the wave propagation direction is translated

into a kinetic degree of freedom which is subtracted

from the free and constrained dominated fields Dof's ,in other

$$\text{words } Dof_{\text{ electric field }} + Dof_{\text{ magnetic field }} + Dof_{\text{ kinetic }} = 3+1 = 4 \quad (10-25)$$

2 -energy leaves the quanton in the form of an energy packet

$E = E_s E_t$, and the expansion of this energy packet in space

is different from that inside the quanton

energy expansion inside the quanton is in the form

$$\frac{\partial}{\partial s} (E_s E_t) = E_q = E_{sf} E_{tf} E_{sc} E_{tc} , \text{ while outside the quanton}$$

$$\text{it takes the form } E_q = \frac{\partial}{\partial s} (E_s E_t) = c\epsilon_0 \left(\frac{E_{sf} E_{tc}}{\sqrt{\epsilon_0 \sqrt{c}}} \right) \left(\frac{E_{sc} E_{tf}}{\sqrt{\epsilon_0 \sqrt{c}}} \right) = c\epsilon_0 E B$$

and as energy has to be ejected from the quanton , one degree of

energy freedom became a kinetic degree of freedom , and so the

quanton instead of being stationary becomes relativistic quanton

anti quanton pair

3- the difference between the two cases of energy expansion , is

due to the absence of the field component in the relativistic
 (Q/AQ) pair propagation direction which means that this quanton
 pair is a two dimensional one and must substitute this lost Dof
 with a relativistic Dof to maintain dimensional energy symmetry
 4-electromagnetic waves leave quanton under two constraints
 a-Integrity of the energy is maintained (no dispersion)
 b-free and constrained fields (E_{qf} , E_{qc}) cannot leave the
 quanton independently , as the electromagnetic waves are the
 mechanism of transmission of energy through 3D space , they
 must have energy fields which are varying in space and time
 whose energy Dof = 4 (one of them a kinetic Dof)
 this is achieved by cross linking free and constrained fields in the
 form for sinusoidal waves

$$E(x) = \frac{1}{2} (E_f + c B_c) , \quad B(x) = \frac{1}{2} (B_c + \frac{1}{c} E_f)$$

5- electromagnetic waves in the form

$$\mathbf{E} = \frac{1}{2} \left(\frac{E_{qf}}{\sqrt{c}} + c \frac{E_{qc}}{\sqrt{c}} \right), \quad \mathbf{B} = \frac{1}{2} \left(\frac{E_{qf}}{c\sqrt{c}} + \frac{E_{qc}}{\sqrt{c}} \right) \text{ can be seen as a relativistic}$$

Two dimensional quanton anti quanton pairs , where one energy degree of freedom is replaced by a kinetic energy degree of freedom as the waves are formed ,

dimensional analysis ,

based on free and constrained energy field dimensions , the dimensions of electromagnetic field can be determined

$$\text{the electric field } [\mathbf{E}] = \left[\frac{E_{sf} E_{tc}}{\sqrt{c}} \right] = M^{+0.5} L^{-0.5} T^{-1} \quad (11-25)$$

$$\text{and the magnetic field } [\mathbf{B}] = \left[\frac{E_{sf} E_{tc}}{c\sqrt{c}} \right] = M^{+0.5} L^{+1.5} T^{00.0} \quad (12-25)$$

$$[\mathbf{U}] = \text{electromagnetic energy density} = \left[\frac{E}{V} \right] = [\epsilon E^2] = M L^{-1} T^{-2}$$

(ϵ : can be chosen according to a system of units to be = 1)

$$\mathbf{U} = (\mathbf{E}_f + c \mathbf{B}_c)^2 = \frac{1}{4} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} + \sqrt{c} E_{sc} E_{tf} \right)^2$$

$$[U] = \frac{1}{4} * 4 \left(\sqrt{\frac{h}{16 \pi^4 c^3}} \right)^2 (k^2 c^2)^2 = \left(\frac{hc}{16 r_q^4} \right) = \frac{hc}{\lambda^4} \quad (13-25)$$

$= \left[\frac{E}{V} \right] = M L^{-1} T^{-2}$, this is the generic form (non statistical) of

Electromagnetic energy density while in terms of the magnetic

$$\text{field } \left[\frac{E}{V} \right] = \left[\frac{B^2}{\mu} \right] = M L^{-1} T^{-2}$$

$$[U] = c^2 \left(B_c + \frac{1}{c} E_f \right)^2 = c^2 \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} + \frac{1}{c} \frac{E_{sf} E_{tc}}{\sqrt{c}} \right)^2$$

(μ : chosen according to a system of units to be $= \frac{1}{c^2}$)

$$U = \left(\sqrt{\frac{h}{16 \pi^4 c^3}} \right)^2 (k^2 c)^2 = \frac{hc}{16 \pi^4} k^4 = \left[\frac{E}{V} \right] = M L^{-1} T^{-2}$$

	<i>Dof_x</i> (kinetic)	<i>Dof_{yz}</i>	<i>Dof_t</i>	<i>Total Dof</i>
<i>E</i> (x)	0.5	<i>E_{sfyz}</i> (x)=1.50	<i>E_{tc}</i> =0.5	2.50
<i>B</i> (x)	0.5	<i>E_{scyz}</i> (x)=0.50	<i>E_{tf}</i> =0.5	1.50
total	1.00	2.00	1.00	4.00

Table (4) How degrees of freedom of freedom are shared among the different energy fields for the case of electromagnetic waves

25.b. Differences between quanton and electromagnetic waves

	Quanton waves	electromagnetic
Kinetic degrees of freedom	none	one
Nature of fields	three dimensional	Two dimensional
Dof_{sf} , Dof_{sc}	2, 1	1.5 , 0.5
Field energy density	4-Dimensional	3D+relativistic Dof
Wave vector (pointing vector)	static (rotational)	one directional translation
Viewed as	Static three dimensional (Q+AQ) pair	Relativistic two dimensional (Q+AQ) pair

Table (5) Comparison between quanton (free /constrained) waves and electromagnetic waves

26.Maxwell equations of energy fields

As energy variation in space and time creates dynamic fields, so we can relate the four Maxwell equations for electromagnetism to their original form for energy fields we have defined the electromagnetic waves as the relativistic expansion of quantons / anti quantons pair that is travelling through space at velocity (c) in the form

$$\mathbf{E} = \frac{1}{2} \left(\left(\frac{\mathbf{E}_{sf} \mathbf{E}_{tc}}{\sqrt{\epsilon_0} \sqrt{c}} \right)_q + \left(\frac{\mathbf{E}_{sc} \mathbf{E}_{tf}}{\sqrt{c} \sqrt{\epsilon_0}} \right)_{aq} \right)$$

$$\mathbf{B} = \frac{1}{2} \left(\left(\frac{\mathbf{E}_{sc} \mathbf{E}_{tf}}{\sqrt{\epsilon_0} \sqrt{c}} \right)_q + \left(\frac{\mathbf{E}_{sf} \mathbf{E}_{tc}}{\sqrt{\epsilon_0} \sqrt{c}} \right)_{aq} \right)$$

substituting in the four Maxwell equations with the constituent energy fields corresponding to the electric and magnetic fields

1-Gauss law of electric field

$$\nabla \cdot \mathbf{E} = \frac{\rho_c}{\epsilon_0}$$

ρ_c : charge density

$$\nabla \cdot \mathbf{E} = \nabla \cdot \left(\left(\frac{\mathbf{E}_{sf} \mathbf{E}_{tc}}{\sqrt{\epsilon_0} \sqrt{c}} \right)_q + \left(\frac{\mathbf{E}_{sc} \mathbf{E}_{tf}}{\sqrt{c} \sqrt{\epsilon_0}} \right)_{aq} \right) = 2 \left(\frac{\rho_c}{\epsilon_0} \right) \quad (1-26)$$

$(\mathbf{E}_{tc} \nabla \cdot \mathbf{E}_{sf})_q + (\mathbf{E}_{tf} \nabla \cdot \mathbf{E}_{sc})_{aq} = 0$ (for electromagnetic waves and space fabric case)

Where $\nabla \cdot \mathbf{E}_{tf} = 0$, $\nabla \cdot \mathbf{E}_{tc} = 0$ (\mathbf{E}_{tf} , \mathbf{E}_{tc} are function of time only)

$$\text{Or } (\mathbf{E}_{tc} \nabla \cdot \mathbf{E}_{sf})_q = - (\mathbf{E}_{tf} \nabla \cdot \mathbf{E}_{sc})_{aq} \quad (2-26)$$

2-Gauss law of magnetic field

$$\nabla \cdot \mathbf{B} = 0$$

$$\left(\frac{\mathbf{E}_{tf}}{\sqrt{\epsilon_0}\sqrt{c}} \nabla \cdot \mathbf{E}_{sc}\right)_q + \left(\frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0}\sqrt{c}} \nabla \cdot \mathbf{E}_{sf}\right)_{aq} = 0 \quad (3-26)$$

$$\left(\mathbf{E}_{tc} \nabla \cdot \mathbf{E}_{sf}\right)_{aq} = -\left(\mathbf{E}_{tf} \nabla \cdot \mathbf{E}_{sc}\right)_q \quad (4-26)$$

3-faraday's law for electric field

$$\nabla \times \mathbf{E} = -\frac{\partial B}{\partial t}$$

$$\nabla \times \mathbf{E} = \left(\frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0}\sqrt{c}} \nabla \times \mathbf{E}_{sf}\right)_q + \left(\frac{\mathbf{E}_{tf}}{\sqrt{c}\sqrt{\epsilon_0}} \nabla \times \mathbf{E}_{sc}\right)_{aq} \quad (5-26)$$

$$= \left(\frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0}\sqrt{c}} \nabla \times \mathbf{E}_{sf}\right)_q + \left(\frac{\mathbf{E}_{tf}}{\sqrt{c}\sqrt{\epsilon_0}} \nabla \times \mathbf{E}_{sc}\right)_{aq} \quad (6-26)$$

$$-\frac{\partial B}{\partial t} = -\frac{\partial}{\partial t} \left(\mathbf{E}_{sc} \frac{\mathbf{E}_{tf}}{\sqrt{\epsilon_0}\sqrt{c}}\right)_q - \left(\mathbf{E}_{sf} \frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0}\sqrt{c}}\right)_{aq} \quad (7-26)$$

$$= -\left(\frac{\mathbf{E}_{sc}}{\sqrt{\epsilon_0}\sqrt{c}} \frac{\partial \mathbf{E}_{tf}}{\partial t}\right)_q - \left(\frac{\mathbf{E}_{sf}}{\sqrt{\epsilon_0}\sqrt{c}} \frac{\partial \mathbf{E}_{tc}}{\partial t}\right)_{aq}$$

By comparing eq 6 , 7 We get

$$\left(\nabla \times \mathbf{E}_{sf} \frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0}\sqrt{c}}\right)_q = -\left(\frac{\mathbf{E}_{sc}}{\sqrt{\epsilon_0}\sqrt{c}} \frac{\partial \mathbf{E}_{tf}}{\partial t}\right)_q \quad \text{or}$$

$$\left(\mathbf{E}_{tc} \nabla \times \mathbf{E}_{sf}\right)_q = -\left(\mathbf{E}_{sc} \frac{\partial \mathbf{E}_{tf}}{\partial t}\right)_q \quad \text{and} \quad (8-26)$$

$$\left(\frac{\mathbf{E}_{tf}}{\sqrt{c}\sqrt{\epsilon_0}} \nabla \times \mathbf{E}_{sc}\right)_{aq} = -\left(\frac{\mathbf{E}_{sf}}{\sqrt{\epsilon_0}\sqrt{c}} \frac{\partial \mathbf{E}_{tc}}{\partial t}\right)_{aq} \quad \text{or}$$

$$(\mathbf{E}_{tf} \nabla \times \mathbf{E}_{sc})_{aq} = - (\mathbf{E}_{sf} \frac{\partial \mathbf{E}_{tc}}{\partial t})_{aq} \quad (9-26)$$

where $\frac{\partial}{\partial t} (\mathbf{E}_{sf}) = 0$, $\frac{\partial}{\partial t} (\mathbf{E}_{sc}) = 0$

(\mathbf{E}_{sf} , \mathbf{E}_{sc} are function of space only)

4-ampere's law for magnetic field

$$\nabla \times \mathbf{B} = \mu_0 (\mathbf{j} + \epsilon_0 \frac{\partial \mathbf{E}}{\partial t})$$

Where $\mu_0 \epsilon_0 = \frac{1}{c^2}$

$$\begin{aligned} \nabla \times \mathbf{B} &= \nabla \times \left(\frac{\mathbf{E}_{sc}}{\sqrt{\epsilon_0} \sqrt{c}} \mathbf{E}_{tf} \right)_q + \nabla \times \left(\frac{\mathbf{E}_{sf}}{\sqrt{\epsilon_0} \sqrt{c}} \mathbf{E}_{tc} \right)_{aq} \\ &= \left(\frac{\mathbf{E}_{tf}}{\sqrt{\epsilon_0} \sqrt{c}} \nabla \times \mathbf{E}_{sc} \right)_q + \left(\frac{\mathbf{E}_{tc}}{\sqrt{\epsilon_0} \sqrt{c}} \nabla \times \mathbf{E}_{sf} \right)_{aq} \end{aligned} \quad (10-26)$$

$$\frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t} = \frac{1}{c^2} \frac{\partial}{\partial t} \left[\left(\frac{\mathbf{E}_{sf}}{\sqrt{\epsilon_0} \sqrt{c}} \mathbf{E}_{tc} \right)_q + \left(\frac{\mathbf{E}_{sc}}{\sqrt{\epsilon_0} \sqrt{c}} \mathbf{E}_{tf} \right)_{aq} \right]$$

$$= \frac{1}{c^2} \left[\left(\frac{\mathbf{E}_{sf}}{\sqrt{\epsilon_0} \sqrt{c}} \frac{\partial \mathbf{E}_{tc}}{\partial t} \right)_q + \left(\frac{\mathbf{E}_{sc}}{\sqrt{c} \sqrt{\epsilon_0}} \frac{\partial \mathbf{E}_{tf}}{\partial t} \right)_{aq} \right] \quad (11-26)$$

By comparing eq 10 , 11 we get

$$(\mathbf{E}_{tf} \nabla \times \mathbf{E}_{sc})_q = \frac{1}{c^2} (\mathbf{E}_{sf} \frac{\partial \mathbf{E}_{tc}}{\partial t})_q \quad \text{and}$$

$$(\mathbf{E}_{tc} \nabla \times \mathbf{E}_{sf})_{aq} = \frac{1}{c^2} (\mathbf{E}_{sc} \frac{\partial \mathbf{E}_{tf}}{\partial t})_{aq} \quad (12-26)$$

It is worth noting that

1- the equations (2 , 3) can be put in the following form

$$\text{For quantons : } \frac{\nabla_x E_{sf}}{\frac{\partial E_{tf}}{\partial t}} = - \frac{E_{sc}}{E_{tc}} \quad (13-26)$$

$$\text{For anti quantons : } \frac{\nabla_x E_{sc}}{\frac{\partial E_{tc}}{\partial t}} = - \frac{E_{sf}}{E_{tf}} \quad (14-26)$$

2- Maxwell equations remain invariant under relativistic effects

as this effect is split equally between two fields

27.Role of Maxwell equations in the evolution of the quanton

Based on the previous results of Maxwell's equations which link the

fields of both the quanton and the anti quanton together , the

quanton 's own form of Maxwell equations can be deduced

1-the basic fields during the primordial time were in the form

E_{sf} , E_{tf} (free energy field that varies in space and free energy field

that varies in time) as the formation of the quanton took the path

of the coexistence of both fields

2-as energy expands by varying in time (E_{tf}), its rate of variation

Induces a curl in the space varying field such that

$$\nabla_{\mathbf{x}} E_{sf} = - \frac{E_{sc}}{E_{tc}} \frac{\partial E_{tf}}{\partial t} \text{ in other words , the rate of variation of } E_{tf}$$

causes E_{sf} to curl into the quanton as it is formed hence , the

energy fields E_{sf} E_{tc} are contained into a quanton formation

5- the rate of variation of the time varying field E_{tc} induces a

formation of a curl in the constrained space varying field E_{sc} ,

$$\text{such that } \nabla_{\mathbf{x}} E_{sc} = \frac{1}{c^2} \frac{E_{sf}}{E_{tf}} \frac{\partial E_{tc}}{\partial t} \text{ , such that fields } E_{sc} , E_{tf}$$

are also contained in the quanton as it formed

28. Anti quanton evolution and its degrees of freedom

Initially , this model proposed anti quanton to have evolved from

free time varying energy (E_{tf}) , however, through later work , many

changes had to be made to match a more refined version of space

fabric evolution from a single quanton , the existence of anti

quanton as a stable part of space fabric may seem to be

problematic ,

however , other evidence still weighs in its favour , namely

1-its role in the electromagnetic wave generation

(already discussed in electromagnetic section)

2-its role in the formation of the negatively charged particles

(electrons , down quarks)

3-anti quanton is stable under expansion conditions

(no degeneration)

4-the interactions generated by anti quanton energy fields are

symmetric to those of the quanton ,hence , it can not affect the

space fabric homogeneity and integrity




Fig (6).hypothetical tree diagram for the evolution and the degrees of freedom of anti quanton energy fields, and why the independent evolution of the anti quanton now seems to be problematic

From the above degree of freedom evolution diagram , the anti quanton would have evolved from energy fields E_{sc} , E_{tc} such space and time varying fields could not evolve independently under inflationary conditions, an alternative scenario is offered, which is the evolution of the anti quanton from the quanton itself and through the pathways of Maxwell's equations , which are in its generalized form , links the variation of both space and time

a-free dominated field splits , reduced into a packet state

$$\int \mathbf{E}_{qf} \, ds = \int \mathbf{E}_{sf} \, ds \frac{\partial \mathbf{E}_{tc}}{\partial t} = \mathbf{E}_s \mathbf{E}_t \quad (1-28)$$

then expands as a constrained space varying field

$$\int (\mathbf{E}_s) \, ds \frac{\partial}{\partial t} (\mathbf{E}_t) = \int \mathbf{E}_s \, ds \frac{\partial \mathbf{E}_{tf}}{\partial t} = (\mathbf{E}_{qc})_{aq} \quad (2-28)$$

b-for the quanton's constrained space dominated field

$$\int \mathbf{E}_{qc} \, ds = \frac{\partial}{\partial s} (\mathbf{E}_{sc}) \int (\mathbf{E}_{tf}) \, dt = \mathbf{E}_s \mathbf{E}_t \quad (3-28)$$

then expands as a free space varying field

$$\frac{\partial}{\partial s} (\mathbf{E}_s) \int (\mathbf{E}_t) \, dt = \frac{\partial \mathbf{E}_s}{\partial t} \int \mathbf{E}_t \, dt = (\mathbf{E}_{qf})_{aq} \quad (4-28)$$

Anti quanton is the mirror image of the quanton's Dof's

5-For the energy degrees of freedom inside anti quanton ,

They are governed by the Maxwell's equations

rate of variation of \mathbf{E}_{tc} curls \mathbf{E}_{sc} such that

$$(\mathbf{E}_{tf} \nabla \times \mathbf{E}_{sc})_{aq} = -(\mathbf{E}_{sf} \frac{\partial \mathbf{E}_{tc}}{\partial t})_{aq}$$

rate of variation of E_{tf} curls E_{sf} such that

$$(E_{tc} \nabla \times E_{sf})_{aq} = \frac{1}{c^2} (E_{sc} \frac{\partial E_{tf}}{\partial t})_{aq}$$

The relationship between Q/AQ pair is governed by

$$(E_{tf} \nabla \cdot E_{sc})_{aq} = - (E_{tc} \nabla \cdot E_{sf})_q$$

$$(E_{tc} \nabla \cdot E_{sf})_{aq} = - (E_{tf} \nabla \cdot E_{sc})_q$$

6-the dominant energy of the anti quanton system is constrained

$$D_{net} (\text{unbound}) = \frac{\text{constrained fields Dof}}{\text{free fields Dof}}$$

$$D_{scu} D_{tcu} (\text{unbound}) = \frac{(D_{tc} D_{sc})}{(D_{sf} D_{tf})} = \frac{c^2 c^{0.5}}{c c^{0.5}} = c \quad (5-28)$$

$$(\text{unbound})_{\text{constrained Dof}} = (\sum(\text{constrained Dof}))$$

$$- \sum(\text{free Dof}) = [(Dof_{sc}) + (Dof_{tc})] - [(Dof_{sf}) + (Dof_{tf})]$$

$$= (2.5-1.5) = +1$$

29.Lorentz transformation of energy fields

In the previous chapters we have discussed the concept of a

relativistic quanton and how it is represented electromagnetic

waves in the form of space and time varying fields

Here , the Lorentz transformation will be discussed , for the electromagnetic waves (this time in terms of the quanton

Energy fields)

Considering the case when energy fields are seen by an observer traveling at relativistic velocity along x axis

2-for Lorentz transformation of electromagnetic waves, and while denoting (') for the case of a moving frame of reference , the transformation takes the form

$$E_x' = E_x , E_y' = \gamma (E_y + \beta c B_z)$$

$$E_z' = \gamma (E_z + \beta c B_y) , B_x' = B_x$$

$$B_y' = \gamma (B_y - \frac{v E_z}{c^2}) , B_z' = \gamma (B_z - \frac{v E_y}{c^2})$$

In this case the electric field is represented by the field $E_y(x)$

and the magnetic field is represented by the field $B_z(x)$

Using the same transformation for the case of free and constrained energy dominated system , where

$$E = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} (E_f + c B_c) = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} + c \frac{E_{sc} E_{tf}}{\sqrt{c}} \right)$$

$$B = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} (B_c + \frac{1}{c} E_f) = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} + \frac{1}{c} \frac{E_{sf} E_{tc}}{\sqrt{c}} \right)$$

after substitution, we get for E and B

$$E_y' = \frac{\gamma}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) + c \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) + v \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) + \frac{v}{c} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) \right) \quad (1-29)$$

$$= \frac{\gamma}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) \left(1 + \frac{v}{c} \right) + c \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) \left(1 + \frac{v}{c} \right) \quad (2-29)$$

$$E_y' = \frac{\gamma}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) + c \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) \right) \left(1 + \frac{v}{c} \right) = \sqrt{\frac{\left(1 + \frac{v}{c} \right)}{\left(1 - \frac{v}{c} \right)}} E_y \quad (3-29)$$

$$\text{Where } \gamma \left(1 + \frac{v}{c} \right) = \frac{\sqrt{\left(1 + \frac{v}{c} \right)} \sqrt{\left(1 + \frac{v}{c} \right)}}{\sqrt{\left(1 + \frac{v}{c} \right)} \sqrt{\left(1 - \frac{v}{c} \right)}} = \sqrt{\frac{\left(1 + \frac{v}{c} \right)}{\left(1 - \frac{v}{c} \right)}} \quad (4-29)$$

$$B_z' = \frac{\gamma}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) + \frac{1}{c} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) \right) - \frac{v}{c^2} \left(\left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) + c \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) \right) \quad (5-29)$$

$$B_z' = \frac{\gamma}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sc} E_{tf}}{\sqrt{c}} \right) \left(1 - \frac{v}{c} \right) + \frac{1}{c} \left(\frac{E_{sf} E_{tc}}{\sqrt{c}} \right) \left(1 - \frac{v}{c} \right) = \sqrt{\frac{\left(1 - \frac{v}{c} \right)}{\left(1 + \frac{v}{c} \right)}} B_z \quad (6-29)$$

$$\text{Where } \gamma \left(1 - \frac{v}{c}\right) = \frac{\sqrt{\left(1 - \frac{v}{c}\right)} \sqrt{\left(1 - \frac{v}{c}\right)}}{\sqrt{\left(1 + \frac{v}{c}\right)} \sqrt{\left(1 - \frac{v}{c}\right)}} = \sqrt{\frac{\left(1 - \frac{v}{c}\right)}{\left(1 + \frac{v}{c}\right)}} \quad (7-29)$$

For a comoving frame of reference at v where $\beta = \frac{v}{c}$ and $\gamma = \frac{1}{\sqrt{1-\beta^2}}$

The electromagnetic fields as viewed by moving observer

$$\text{are } E' = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sf}' E_{tc}'}{\sqrt{c}} + c \frac{E_{sc}' E_{tf}'}{\sqrt{c}} \right) = \frac{1}{\sqrt{\epsilon_0}} \sqrt{\frac{1+\beta}{1-\beta}} K_q^2 c^2 \cos(k'r' - \omega't') \quad (8-29)$$

$$B' = \frac{1}{2} \frac{1}{\sqrt{\epsilon_0}} \left(\frac{E_{sc}' E_{tf}'}{\sqrt{c}} + \frac{1}{c} \frac{E_{sf}' E_{tc}'}{\sqrt{c}} \right) = \frac{1}{\sqrt{\epsilon_0}} \sqrt{\frac{1-\beta}{1+\beta}} K_q^2 c \cos(k'r' - \omega't') \quad (9-29)$$

$$\text{Where } k' = \sqrt{\frac{1-\beta}{1+\beta}} k, \quad r' = \sqrt{\frac{1-\beta}{1+\beta}} r$$

$$\omega' = \sqrt{\frac{1-\beta}{1+\beta}} \omega, \quad t' = \sqrt{\frac{1-\beta}{1+\beta}} t$$

$$\text{to note that the product } E_y' B_z' = \sqrt{\frac{\left(1 + \frac{v}{c}\right)}{\left(1 - \frac{v}{c}\right)}} E_y \sqrt{\frac{\left(1 + \frac{v}{c}\right)}{\left(1 + \frac{v}{c}\right)}} B_z$$

$= E_y B_z = \text{constant}$, irrespective of the frame of reference

30. some concepts behind space fabric

1-equipartition of energy or dimensional energy symmetry (with

respect to time and space variation of energy)

2- field interaction , no silent energy field , energy fields

of different types (free / constrained) interact with other energy

fields of different or similar nature to create a binding or repulsive

interactions

3-Preservation of space fabric integrity (in the form of space fabric

Binding and retaining interactions)

4-energy field interactions are expressed at all the scales (energy

fields are infinite in range)

31.Interactions of energy fields

1-Energy as it varies in space or time creates associated

dynamic fields that exist inside as well as outside the quantons

2-the nature of the field interactions depends on the type of the

energy field (free or constrained energy dominated)

3- energy field interaction is according to following manner

a-Interaction of energy fields of similar type (free or constrained)

is repulsive in nature

b-interaction energy fields of different type creates a binding interaction

4-an energy field can interact with another energy field only if

they have the same field strength (they both have the same *Dof*'s)

(necessity condition)

5- same energy field can self-interact to generate a repulsive reaction

6-though energy fields are infinite in their range of action but this range can still be divided into 3 main zones

a-inside quantons b- outside quantons :short range

c-outside quantons : long range

32.Bound and unbound fields

1-inside the quanton , interaction between energy fields of different

nature (free- constrained) generates a binding interaction and those energy fields which are involved in such an interaction are said to be bound fields , while energy fields that do not generate such interactions are said to be unbound fields

2-for quantons free energy fields are split into two parts :bound and unbound part $E_{sf} = K_{sf} (D_{sfb} D_{sfu})$, $E_{tf} = K_{tf} (D_{tfb} D_{tfu})$

3-fields generated by free energy (E_{sfb} E_{tfb}) fully interact with fields generated by constrained energy (E_{sc} E_{tc}) in a binding interaction , while for anti quanton $E_{sc} = K_{sc} (D_{scb} D_{scu})$,

$E_{tc} = K_{tc} (D_{tcb} D_{tcu})$, and the binding interaction is between fields (E_{scb} E_{tcb}) and (E_{sf} E_{tf})

4-Fields created by free and constrained energies are best described as having flux lines , and the number of those flux lines is not indefinite in number

5- unbound energy fields are repulsive in nature due to their

self-interaction

6-for the space fabric case , binding interaction expresses a state

of equilibrium due to the symmetry interacting energy fields

(equal in strength and intensity)

7- for an energy system to be under equilibrium , all its energy

fields must be tied in a binding relationships (with other energy

fields) at all the scales (absence of unbound fields)

8-bound energy fields create binding interactions necessary

for the integrity of the space fabric (later they will be called

quanton binding (E_b) and retaining (E_t) interactions)

9-all remaining unbound energy fields and through the self

interaction give rise to quanton inflation , splitting and on larger

scale inflationary momentum

33. types of field interactions

33.a.single interactions

1-Single Interactions of the type $E_{binding\ ij} = \frac{(E_{sfi})(E_{scj})}{(\Delta r_{ij})}$

Do not exist in nature since space varying fields cannot exist

independently of time varying fields

2-simple interactions between different energy fields inside and

around the quanton do not generate four dimensional potential

energies , so we use the term interaction ($E_{ij\ binding}$) to describe

the simple binding between energy fields of the type

($E_{sfi}E_{tfi}$), ($E_{scj}E_{scj}$).

3-the dimensions of any interaction depend on its degrees of

freedom

the interactions between energy fields ($E_{sfi}E_{tfi}$) and ($E_{scj}E_{tcj}$) can

be assessed as follows :

the binding interaction($E_{binding-ij}$) between fields (E_{sfi},E_{tfi})

and $(E_{scj} E_{tcj})$ (represented by shared flux lines between the fields)

is proportional to the generated flux (φ_{ij}) between the two energy

fields, the flux itself is proportional to the product of the Dof's and

intensities of those two fields , and follows the same guidelines

outlined in the section : superposition principle inside the qunton ,

namely

1-the generated interaction Dof's equal to the summation of

energy degrees of freedom of both fields (proportional to the

product of field strength of both fields)-for example

$$D_{binding-ij} = (D_{sfi} D_{tfi})(D_{scj} D_{tcj}) = c^{Dof_{sf}+Dof_{tf}+Dof_{sc}+Dof_{tc}} \quad (1-33)$$

2- the interaction intensity must be proportional to the product of

intensity of both fields as defined by the parameter K_q

$$(\text{for example } (K_{sfi} K_{tfi})(K_{scj} K_{tcj}) = K_q^4)$$

3- the interaction must be related to true energy , so dimensions of

the energy fields intensities must always represent the real binding energy , in other words interactions must be always in terms of

$$K_q^4 \quad (K_{ij \text{ binding}} = (K_{sf} K_{tf}) (K_{sc} K_{tc}) = K_q^4)$$

as the term K_q^4 represents an energy density divided by c^4

4 – the binding relationship for the case of two fields

$$E_{\text{binding } ij} = \frac{\varphi_{ij}}{(\Delta r_{ij})} = \frac{(E_{sfb} E_{tbi})(E_{scj} E_{tcj})}{(\Delta r_{ij})} \quad (2-33)$$

$$= \frac{(K_{sfi} K_{scj}) (D_{sfb} D_{sc}) (K_{tbi} K_{tcj}) (D_{tfb} D_{tc})}{(\Delta r_{ij})}$$

$$= \frac{(K_{sfi} K_{tbi}) (K_{scj} K_{tcj}) (D_{sfb} D_{tfb}) (D_{sf} D_{sc})}{(\Delta r_{ij})}$$

$$= \frac{(K_q^4) (D_{sfb} D_{tfb}) (D_{sf} D_{sc})}{(\Delta r_{ij})} \quad (3-33)$$

$$= \sqrt{\alpha_b} \frac{h}{2r_q v_q} c^{Dof_{sfb} + Dof_{tfb} + Dof_{sc} + Dof_{tc}} \quad (4-33)$$

α_b : parameter of interaction , Δr_{ij} : effective distance between

Two fields , while E_{sfi} is defined as being equal to $K_{sfi} D_{sf}$ (which

expresses the energy field as the product of its strength (Dof) and

intensity ,

the dimensions of such an interaction would be $\frac{\text{Energy}}{c^{4-(Dof_{total})} (3D \text{ volume})}$

where $Dof_{total} = Dof_{free} + Dof_{constrained}$

so only interactions which have four degrees of freedom are

able of generating a binding that has the true dimensions of energy

density

33.b multiple fields interactions

1- Energy fields tend to form higher order interactions whenever

possible (multiple field interactions) (this is true up to $Dof = 4$)

2- hyper interactions (summation of Dof of constituent fields

greater than 4) are inhibited inside and outside quanton .

for real interactions , Dofs must be equal or less than (4) whether

it is a single or multiple interaction

(in real spaces only real interactions can be generated)

3- simpler interactions can combine to form a multiple interaction

with higher degrees of freedom (up to 4)

so ,multiple complex field interactions are generated as a result of

two simple binding interactions of the type $(E_{sfbi} E_{tfbi})(E_{scj} E_{tcj})$

that can combine with another simple interaction

$(E_{sfbj} E_{tfbj})(E_{sci} E_{tcj})$ to form a complex one of the type

$$E_{binding\ ij} = \frac{(E_{sfbi} E_{tfbi})(E_{sci} E_{tci})(E_{sfbj} E_{tfbj})(E_{scj} E_{tcj})}{(\Delta r_{ij})} \quad (5-33)$$

which is the case of gravitation

33.d. nonbinding (repulsive) interactions

while inside the quanton , the unbound field $E_{sfu} E_{tfu}$ (or

$E_{scu} E_{tcu}$ for the case of anti quanton) generates self-interaction

that gives rise only to simple repulsive interactions inside the

quanton , while outside the quanton (anti quanton) the generated

self-interacting field can be involved in a repulsive interaction

as well with another energy field of the same nature (free or

constrained) and the generated interaction would always be a

repulsive one ,

as this energy field cannot create a binding interaction with

another field with opposing type due to this repulsive self

interacting nature even if they share the same Dof's

$$\begin{aligned}
 E_{rij} &= (E_{sfui}E_{tfui})(E_{sfuj}E_{tfuj}) \frac{1}{(\Delta r_{ij})} \\
 &= (K_{qi}^2 D_{sfu}D_{tfu}) (K_{qj}^2 D_{sfu}D_{tfu}) \frac{1}{(\Delta r_{ij})} \quad (6-33)
 \end{aligned}$$

$$= K_q^4 (D_{sfu} D_{tfu})^2 \frac{1}{(\Delta r_{ij})}$$

$$= \sqrt{\alpha_r} \frac{h}{2r_q v_q} c^{Dof_{sfu}+Dof_{tfu}+Dof_{sfu}+Dof_{tfu}} \quad (7-33)$$

and once outside the quanton , the fields behave as complex ones

so , they must interact with another field (simple or complex) of

the same energy nature to generate a nonbinding (repulsive)

interaction in both cases

34. quanton field interactions

34.a-inside quantons

34.a.1The quanton retaining interaction (E_t)

the free and constrained energy fields interact with the energy of

an opposite nature inside the quanton to create the quanton

retaining interaction (E_t)

This is interaction is between (the bound part) of the free

energy field ($E_{sfb}E_{tfb}$) and constrained energy field ($E_{sc}E_{tc}$)

the binding part of the free energy field that participates in this

interaction has to have the same degrees of freedom as

constrained field (due to the symmetry of Dof's of the interaction)

and is expressed as

$$E_{sf}E_{tf} = (K_{sf}K_{tf}) (D_{sfb}D_{tfb}) (D_{sfu}D_{tfu})$$

$$(D_{sf}D_{tf})_{binding} = (D_{sfb}D_{tfb}) = D_{sc}D_{tc} \quad \text{or} \quad (1-34)$$

$$(D_{sfb} D_{tfb}) = c^{1.50} \quad (2-34)$$

$$(D_{sfu}D_{tfu}) = \frac{E_{sf}E_{tf}}{E_{sc}E_{tc}} = \frac{K_q^2 D_{sf}D_{tf}}{K_q^2 D_{sc}D_{tc}} = \frac{D_{sf}D_{tf}}{D_{sc}D_{tc}} = \frac{c^{2.50}}{c^{1.50}} = c \quad (3-34)$$

the generated retaining interaction (E_t) that maintains the quanton's integrity and prevents it from disintegration, the retaining interaction (E_t) is binding energy type since it is developed between two fields of different nature

this interaction takes the following form for a single quanton

$$(E_t)_q = (E_{sf}E_{tf})_{binding} (E_{sc}E_{tc}) \quad (4-34)$$

$$= [K_q^2(D_{sfb} D_{tfb})] [K_q^2(D_{sc}D_{tc})]$$

$$(E_t)_q = K_q^4 c^3 = \frac{\sqrt{\alpha_t h} k^4}{16\pi^4} = \frac{\sqrt{\alpha_t h}}{16 r_q^4} \quad (5-34)$$

Where the term $(E_{sf}E_{tf})_{binding}$ represents the binding part of the free energy ($E_{sf}E_{tf}$) that interacts with constrained energy

$(E_{sc}E_{tc})$, (r_q) is the quanton radius

α_t : retaining interaction parameter

while for anti quanton case the retaining interaction would be

$$(E_t)_{aq} = (E_{sc} E_{tc})_{bound} (E_{sf} E_{tf}) = \quad (6-34)$$

$$= [K_q^2 (D_{scb} D_{tcb})] [K_q^2 (D_{sf} D_{tf})]$$

$$(E_t)_{aq} = K_q^4 c^3 = \frac{\sqrt{\alpha_t} h}{16 r_q^4} \quad (7-34)$$

as for the dimensions of such interaction , which has three Dof's , it

should be $[\frac{energy}{volume * c}] = M L^{-2} T^{-1}$

34.a.2.quanton inflationary interaction (E_i)

Type : simple nonbinding(repulsive)

Inflationary interaction can be thought of as the result of the

self-interaction of the unbound part of free energy field which is

not involved in the retaining interaction (E_t)

the result of this self-interaction is the appearance of a repulsive

interaction (E_i) that causes quanton to expand ,

the generated quanton inflationary interaction would be in the form

$$(E_i)_q = \left(\sqrt{(E_{sf} E_{tf})_{unbound}} \right)^2 \quad (8-34)$$

$$= (K_q^2 \sqrt{(D_{sfu} D_{tfu})}) (K_q^2 \sqrt{(D_{sfu} D_{tfu})})$$

$$(E_i)_q = K_q^4 c = \frac{\sqrt{\alpha_i h}}{16 \pi c^2 r_q^4} \quad (9-34)$$

α_i : inflationary interaction parameter

the inflationary interaction is at the origin of the quanton's inflation

and subsequent division , which is a synonym with space fabric

expansion , this self-interaction can be thought of as energy field

of a strength $\sqrt{D_{sfu} D_{tfu}}$ that is interacting with another energy

field of similar magnitude creating this repulsive interaction

the dimensions of such a energy-like interaction , which has

one Dof , it should be $[\frac{energy}{volume * c^3}] = M L^{-4} T^{+1}$

While for the case of anti quanton , the inflationary energy

$$(E_i)_{aq} = (K_q^2 \sqrt{(D_{scu} D_{tcu})})^2 K_q^2 \sqrt{(D_{scu} D_{tcu})}$$

$$(E_i)_{aq} = K_q^4 c = \frac{\sqrt{\alpha_i h}}{16 \pi c^2 r_q^4} \quad (10-34)$$

34.b-outside quanton

34.b.1-Space fabric binding interaction (E_b)

Type : multiple binding

as energy fields are not limited in range to inside the quanton ,
the fields of the free energy outside the quanton interact with the
fields of the constrained energies of other quantons to generate
the binding interaction (E_b) and vice versa

the generated binding interaction (E_b) is responsible for
maintaining the space fabric integrity , it is represented by two
contributions due to quantons and anti quantons ,

where (E_{bi})_q is the binding interaction developed between the
quanton (q_i) and other quantons (q_j) or anti quantons (aq_j) ,

a-For the quantons case

$$E_{bfi} = E_b (E_{sfbi} E_{tfbi})_q = \left\{ [(E_{sfbi} E_{tfbi})_q \sum_j^n (E_{scj} E_{tcj})_q] \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right) \right. \\ \left. + [(E_{sfbi} E_{tfbi})_q \sum_j^n (E_{scbj} E_{tcbj})_{aq}] \right\} \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right) \quad (12-34)$$

$$\begin{aligned}
&= \{ [K_{qi}^2 (D_{sfb} D_{tfb})_q \sum_j^n K_{qj}^2 (D_{sc} D_{tc})_q] \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right) + \right. \\
& \left. [K_{qi}^2 (D_{sfb} D_{tfb})_q \sum_j^n K_{qj}^2 (D_{scb} D_{tcb})_{aq}] \right\} \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right) \\
E_{bfi} &= K_q^4 c^3 \left(\sum_j^n \left(\frac{r_q}{(r_i - r_j)_{q-q}} \right) + \left(\sum_j^n \frac{r_q}{(r_i - r_j)_{q-aa}} \right) \right) \\
&= \frac{\sqrt{\alpha_b} h}{2} \frac{1}{8 r_q^3} \left(\sum_j^n \left(\frac{1}{(r_i - r_j)_{q-q}} \right) + \left(\sum_j^n \frac{1}{(r_i - r_j)_{q-aa}} \right) \right) \quad (13-34)
\end{aligned}$$

Where the term $(D_{sfb} D_{tfb})_q$ represents the bound part of the free energy $(E_{sfb} E_{tfb})$ that interacts with constrained energy $(E_{sc} E_{tc})$,

$(r_i - r_j)$: the distance between quantons (q_i) and (q_j) or

anti quantons (aq_j) , $(i \neq j)$,

$\sqrt{\alpha_b}$: binding interaction parameter

while the binding interaction due to the constrained field $E_{sc} E_{tc}$

will be in the form

$$E_{bci} = E_{bi} (E_{sci} E_{tci})_q = \{ [(E_{sci} E_{tci})_q \sum_j^n (E_{sfbj} E_{tfbj})_q] \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right) \}$$

$$+ \{ [E_{sci}E_{tci})_q (E_{sfj}E_{tfj})_{aq}] \left(\frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)} \right) \} \quad (14-34)$$

$$= \{ K_q^4 [(D_{sc}D_{tc})_q \sum_j^n (D_{sf}D_{tf})_{aq}] \left(\frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)} \right) +$$

$$[K_q^4 (D_{sc}D_{tc})_q \sum_j^n (D_{sf}D_{tf})_{aq}] \left(\frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)} \right) \}$$

$$= K_q^4 c^3 \left(\sum_j^n \left(\frac{r_q}{(r_i-r_j)_{q-q}} \right) + \left(\sum_j^n \frac{r_q}{(r_i-r_j)_{q-aa}} \right) \right)$$

$$E_{bci} = \frac{\sqrt{\alpha_b} h}{2\pi} \frac{1}{V_q} \left(\sum_j^n \left(\frac{1}{(r_i-r_j)_{q-q}} \right) + \left(\sum_j^n \frac{1}{(r_i-r_j)_{q-aa}} \right) \right) \quad (15-34)$$

which is the same expression as before or $E_{bf} (E_{sfi}E_{tfi}) =$

$E_{bc} (E_{sci}E_{tci})_q$ and this is due to the symmetry of interactions

Later a single expression for both interactions will be developed

Which will be of a multiple binding type ,

of course there would be no counting of any quantons , as the

summation can be handled by assessing energy density over an

integration volume

as for the factor $\frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)}$, while for a single quanton of a radius r_q

it has a total binding energy between bound free energy fields

$(E_{sfb} E_{tfb})$ and constrained energy fields $(E_{sc} E_{tc})$ that is

equivalent To $E_{tp} = \int_{V_q} E_t dV = \int_{V_q} (E_{sfb} E_{tfb}) (E_{sc} E_{tc}) dV$

$$= \frac{h}{16(\pi)^4} k^4 V_q$$

$$= \frac{\sqrt{\alpha_t} h}{2} \frac{1}{8 r_q^3} \frac{1}{r_q} V_q = \sqrt{\alpha_t} \frac{h}{2 r_q} \frac{1}{V_q} V_q = \sqrt{\alpha_t} \frac{h}{2 r_q}$$

which says that the binding energy is directly proportional to $(\frac{1}{r_q})$,

now for the case of a virtual quanton whose radius now becomes

$(r_i - r_j)$ instead of r_q , the binding energy between the two energy

fields inside two separate quantons q_i , q_j becomes

$$E_{bp} = (\int_{V_{qi}} (E_{sfb} E_{tfb}) dV \int_{V_{qj}} (E_{sc} E_{tc}) dV) \frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)}$$

$$= K_{qi}^2 (D_{sfb} D_{tfb}) K_{qj}^2 (D_{sc} D_{tc}) V_q \frac{\sqrt{r_{qi}r_{qj}}}{(r_i-r_j)}$$

$$\begin{aligned}
&= \sqrt{\alpha_b} c^3 \sqrt[2]{\frac{h}{2\pi c^3 V_{qi} r_{qi}}} \sqrt[2]{\frac{h}{2\pi c^3 V_{qj} r_{qj}}} V_q \frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \\
&= \frac{\sqrt{\alpha_b} h}{2(r_i - r_j)}
\end{aligned}$$

this factor $\left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)}\right)$ acts as a conversion factor for the calculation of

the binding between any energy fields regardless whether they

belong to the same quanton or not

34.b.2-Quanton repulsive interaction (E_r)

Type : repulsive

Out side the quanton , the unbound free energy field $(E_{sfui}E_{tfui})_q$

generates a repulsive interaction with other quantons ' unbound

free energy $(E_{sfuj}E_{tfuj})_q$

for quanton (q_i)

$$E_r((E_{sfui} E_{tfui})_q) = [(E_{sfui}E_{tfui})_q \sum_j^n (E_{sfuj}E_{tfuj})_q \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)}\right)] \quad (16-34)$$

$$= [K_{qi}^2 D_{sfu}D_{tfu})_q \sum_j^n (K_{qj}^2 D_{sfu}D_{tfu})_q \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)}\right)]$$

$$= \sqrt{\alpha_r} c^2 \sqrt[2]{\frac{h}{16 c^3 r_{qi}^4}} \sum_j^n \sqrt[2]{\frac{h}{16 c^3 r_{qj}^4}} \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)_{q-q}} \right)$$

$$E_r = \frac{\sqrt{\alpha_r} h}{16 c r_q^3} \sum_j^n \left(\frac{1}{(r_i - r_j)_{q-q}} \right) \quad (17-34)$$

α_r : repulsive interaction parameter

The dimensions of such a energy density interaction , which has

two Dof's , it should be $\left[\frac{\text{energy}}{\text{volume} \cdot c} \right] (= M L^{-3} T^{+00})$

For anti quanton (aq_i)

generated interaction due to unbound field ($E_{scui} E_{tcui})_{aq}$) outside

the anti quanton is also a repulsive in nature in nature since this

energy interacts with the surrounding anti quantons' unbound

constrained energy field ($E_{scuj} E_{tcuj})_{aq}$ to generate an repulsive

interaction

$$E_r((E_{scui})_{aq}) = [(E_{scui} E_{tcui})_{aq} \sum_j^n (E_{scui} E_{tcui})_{aq}] \frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \quad (18-34)$$

$$= [K_{qi}^2 \sqrt{D_{tfu} D_{tfu}}]_{aq} \sum_j^n [K_{qj}^2 \sqrt{D_{tfu} D_{tfu}}]_{aq} \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right)$$

$$\begin{aligned}
&= K_q^4 c^2 \sum_j^n \frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)_{aq-aq}} \\
&= \sqrt{\alpha_r} c^2 \sum_j^n \sqrt{\frac{h}{16 c^3 r_{qi}^4}} \sqrt{\frac{h}{16 \pi c^3 r_{qj}^4}} \left(\frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)_{aq-aq}} \right) \\
E_{ri} &= \frac{\sqrt{\alpha_r} h}{16 c r_q^3} \sum_j^n \left(\frac{1}{(r_i - r_j)_{aq-aq}} \right) \tag{19-34}
\end{aligned}$$

35. generation of space fabric binding interaction (E_b)

1- energy fields out of the quanton , which generate the quanton binding interaction are also at the origin of dark matter gravitation

like effect as well as at the origin of gravitation

if inter-quanton binding were not present , there would have

been no gravitational like effect of dark matter , nor gravitation

for normal matter

2-The generated free energy fields out of the quanton are not in

the form of $(E_{sf} E_{tf})$ (2.5 Dof's) , instead the free energy field

out of the quanton is divided into two parts :

first part which is the binding part which forms the retaining

interaction (E_t) or $(E_{sfb}E_{tfb})_q = K_q^2(D_{sfb}D_{tfb})_q$ and has

(1.5 Dof's) , and the second part which generates the quanton

inflationary interaction (E_i) namely the unbound part

$((E_{sfu}E_{tfu})_q = K_q^2(D_{sfu}D_{tfu})_q$ which has one degree of freedom,

so we can summarize the energy fields as they leave the quanton

as follows

a - $E_{sc}E_{tc}$ (1.5 Dof's) (bound constrained energy)

b- $(E_{sfb}E_{tfb})$ (1.5 Dof's) (bound free energy)

c - $(E_{sfu}E_{tfu})$ (one Dof) (unbound self-interacting

free energy) , and for anti quanton case

a - $E_{sf}E_{tf}$ (1.5 Dof's) (bound free field)

b- $(E_{scb}E_{tcb})$ (1.5 Dof's) (bound constrained field)

c - $(E_{scu}E_{tcu})$ (0.5+0.5 Dof) (unbound constrained field)

3-each energy field can only interact with an energy field which

has the similar degrees of freedom

4-the free energy fields $(E_{sf}E_{tf})_{bound}$ of the quanton or $(E_{sf}E_{tf})$

of the anti quantons create in an interaction with the constrained

energy field $(E_{sc}E_{tc})$ of the other quantons or $(E_{sc}E_{tc})_{bound}$ of the

anti quantons which generates a more stable binding energy rather

than the less stable repulsive interaction with an energy field of the

same nature

5-binding energy fields out of the quanton are symmetric to those

out of the anti quanton (1.5 Dof's of each energy type), and they

all the generate a binding interaction (E_b)

36.Dimensions of energy field interactions

While interactions that generate real energy density have 4 Dof's ,

interactions that involve space fabric , have different dimensions

generally , the number of energy Dof's involved in an interaction is

what determines its dimensions

From the previous discussion , we can deduce some rules

regarding the dimensionality of an interaction (E_i) that involves

($Dof_i = x$) degrees of freedom

dimensions of interaction $[E_i] = \left(\frac{\text{energy}}{\text{volume}} \right) \left(\frac{1}{c^{4-x}} \right) =$

$$= M L^{2-3-4+x} T^{-2+4-x} = M L^{x-5} T^{2-x} = \left[\frac{M L^{x-2} T^{2-x}}{\text{volume}} \right]$$

$$= \frac{\text{energy}}{\text{volume}} \left(\frac{T^x}{L^x} \right)$$

For the special case of $x=4$, $[E_{D_4}] = M L^{-1} T^{-2} = \left(\frac{\text{energy}}{\text{volume}} \right)$

37-dark energy and dark matter in terms of quanton interaction potentials

Previously the quanton interactions were discussed in terms of

energy density , alternatively , those interactions can be assessed

in terms of the quanton packet energy via volumetric integration

$$E_{tp} = \left(\int_{V_q} (E_{sf} E_{tf})_{bound} (E_{sc} E_{tc}) \right) dV$$

$$= [(K_q^2 (D_{sfb} D_{tfb}) (K_q^2 (D_{sc} D_{tc}))] V_q \quad (1-37)$$

$$= K_q^4 c^3 V_q = \alpha_t \frac{h k^4}{16 \pi} = \frac{\sqrt{\alpha_t} h}{2\pi} \frac{1}{8 r_q^3 r_q} V_q$$

$$E_{tp} = \sqrt{\alpha_t} \frac{h}{2 r_q} \quad (2-37)$$

for the inflationary interaction

$$E_{ip} = \int_{V_q} K_q^2 (\sqrt{D_{sfu} D_{tfu}}) (K_q^2 \sqrt{D_{sfu} D_{tfu}}) dV$$

$$= [(K_q^4 (D_{sfu} D_{tfu}))] V_q$$


$$= \sqrt{\alpha_i} \frac{h}{2 V_q r_q c^2} V_q = \sqrt{\alpha_i} \frac{h}{2 r_q c^2} \quad (3-37)$$

for the repulsive interaction

$$E_{rp} = \int_{V_q} K_{qi}^2 (D_{sfu} D_{tfu}) (K_{qj}^2 D_{sfu} D_{tfu}) dV$$

$$= [(K_q^4 (D_{sc} D_{tc})^2] V_q$$

$$= \sqrt{\alpha_r} \frac{h}{2 V_q r_q c} V_q = \sqrt{\alpha_r} \frac{h}{2 r_q c^2} \quad (3-37)$$



Fig(7).the relationship between quanton packet energy and the Energy of various interactions

37b.multiple form of quanton interactions

the interactions of the Q/AQ pair combine to form higher order

interactions (of Dof = four)

this particular point addresses the question why the quanton

evolved to become a Q/AQ pair

37b.1.binding interaction

for a multiple interaction which combines both binding of

$$= \frac{\alpha_b hc}{2} \left(\sum_j^n \left(\frac{1}{(r_i - r_j)_{q-q}} \right) + \left(\sum_j^n \frac{1}{(r_i - r_j)_{q-aq}} \right) \right) \quad (7-37)$$

$$E_{ref} = \frac{hc}{2 r_{ref}}, \quad r_{ref} = \sqrt{r_{qi} r_{qj}} \quad (8-37)$$

E_{bt} = total binding potential of quanton / anti quanton pair $(= \frac{c^2 E_{bp}^2}{E_{ref}})$

37b.2. retaining interaction

for the retaining interaction that combines both bindings of Q /AQ

pair , given that $E_t = (E_{sfb} E_{tfb})(E_{sc} E_{tc})$

$$\frac{c^2 E_{tp}^2}{E_{ref}} = \frac{c^2}{E_{ref}} \int_{V_q} (E_{sfb} E_{tfb})_q (E_{sb} E_{tc})_q dV \int_{V_{aq}} (E_{sf} E_{tf})_{aq} (E_{sfb} E_{tcb})_{aq} dV \quad (9-37)$$

$$= \frac{2 r_{ref} c^2}{hc} (K_q)^4 (D_{sfb} D_{tfb} D_{sc} D_{tc})_q V_q (D_{sf} D_{tf} D_{scb} D_{tcb})_{aq} V_{aq} \quad (10-37)$$

$$\frac{2 \alpha_t r_{ref} c^2}{hc} \left(\frac{h}{16 r_q^4} c^3 V_q \right) \left(\frac{h}{16 r_q^4} c^3 V_{aq} \right)$$

$$\text{total retaining interaction potential } \frac{c^2 E_{tp}^2}{E_{ref}} = \frac{\alpha_t hc}{2 r_q} \quad (11-37)$$

The summation of both the binding and retaining interactions

For the total number of quantons N_q represents the dark matter

with its largely gravitational effects

$$E_u * f_{DM} = \frac{1}{4} \left[N_q \frac{2r_q c^2 E_{tp}^2}{2h} + \frac{1}{2} \sum_i^m \sum_j^n \frac{2r_q c^2 E_{tb}^2}{2h} \right] \quad (12-37)$$

$$= \frac{1}{4} \left[N_q \frac{\alpha_i h}{2r_q} + \frac{1}{2} \sum_i^m \sum_j^n \frac{\alpha_b}{2((r_i - r_j))} \right]$$

Where f_{DM} represents the dark matter fraction of the total energy

of the universe (0.26) , E_u : total energy in the universe

and the summation for $n = N_q$, $m = N_q - 1$, $i \neq j$

37b.3. inflationary and repulsive interactions in multiple form

While for the combined inflationary interaction due to unbound

fields (a, b) of both the Q/Q pair

$$\frac{c^6 E_{ip}^2}{E_{ref}} = \frac{c^6}{E_{ref}} \int_{V_q} (E_{sfu} E_{tfu})_q \, dV \int_{V_{aq}} (E_{scu} E_{tcu})_{aq} \, dV \quad (13-37)$$

$$= \frac{c^6}{E_{ref}} \left[K_q^4 (D_{sfu} D_{tfu})_q V_q (K_q^4 (D_{scu} D_{tcu})_q) V_{aq} \right] \quad (14-37)$$

$$= \frac{2 r_{ref} c^5}{h} \left(\frac{h}{16 c^3 r_q^4} c V_q \right) \left(\frac{h}{16 c^3 r_q^4} c V_{aq} \right)$$

$$\frac{c^6 E_{ip}^2}{E_{ref}} = \alpha_i \frac{hc}{2V_q r_q} V_q = \alpha_i \frac{hc}{2 r_q} \quad (15-37)$$

the combined repulsive interaction of the Q/AQ pair

$$\frac{c^4 E_{rp}^2}{E_{ref}} = \frac{c^4}{E_{ref}}$$

$$\left[\left(\int_{V_{qi}} (E_{sfui} E_{tfui})_q (E_{scui} E_{tcui})_{aq} dV \sum_j^n \int_{V_{qj}} (E_{sfuj} E_{tfuj})_q (E_{scuj} E_{tcuj})_{aq} dV \right) \frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \right] \quad (16-37)$$

$$= \frac{2 r_{ref} c^3}{h}$$

$$\left[(K_{qi})^4 (D_{sfui} D_{tfui} D_{scui} D_{tcui})_q V_{qi} \sum_j^n K_{qj}^4 (D_{sfuj} D_{tfuj} D_{scj} D_{tcj})_{aq} V_{qj} \right] \frac{\sqrt{r_{qi} r_{qj}}}{(r_i - r_j)} \quad (17-37)$$

$$\frac{c^4 E_{rp}^2}{E_{ref}} = \frac{2\alpha_b c}{h} \left[\frac{h}{2V_q c^3 r_{qi}} c^3 V_{qi} \left(\sum_j^n \frac{h}{2V_q c^3 r_{qj}} c^3 V_{qj} \frac{r_{qi} r_{qj}}{(r_i - r_j)_{q-q}} \right) \right]$$

$$= \frac{\alpha_r hc}{2} \left(\sum_j^n \left(\frac{1}{(r_i - r_j)_{q-q}} \right) \right) \quad (18-37)$$

The summation of both the inflationary and the repulsive

Interactions for the total number of quantons N_q in the universe

represents the dark energy with its largely inflationary effects

$$E_u * f_{DE} = \frac{1}{4} \left[N_q \frac{2\alpha_i r_q c^6 E_{ip}^2}{2h} + \frac{1}{2} \sum_i^m \sum_j^n \frac{2\alpha_b r_q c^4 E_{rp}^2}{2h} \right] \quad (19-37)$$

$$= \frac{1}{4} \left[N_q \frac{\alpha_i h}{2r_q} + \frac{1}{2} \sum_i^m \sum_j^n \frac{\alpha_r}{2((r_i - r_j))} \right]$$

Where f_{DM} represents the dark energy fraction of the total energy of the universe (0.68) , E_u : total energy in the universe

38. Why quanton does not achieve equilibrium

Energy fields inside the quanton try to achieve stability in the form of binding Interaction which has the maximum number of degrees of freedom , the rearrangement the quanton Dof's to satisfy the condition would be as follows : $Dof_{tf} = Dof_{tc} = 0.5$

$$Dof_{sf} = Dof_{sc} = 1.5 \quad , \quad Dof_{sf} Dof_{tf} = Dof_{sc} Dof_{tc} = 2$$

This binding interaction here has all four Dof's

under such conditions the quanton is in equilibrium ,no unbound fields exist to cause quanton inflation or splitting ,

But this will not happen as such a condition would entail that there would be no inflation of the universe beyond the single quanton , which would remain in this state indefinitely

this scenario is not possible as energy has to expand , by variation in space and variation in time since the repulsive self interaction (represented by the dark energy) has the majority of the quanton potential in comparison to the biding potential (represented by the dark matter)

39.The inverse relationship between wave length / energy – a possible explanation

The quanton retaining (binding) interaction took the form

$E_t = (E_{sfb}E_{tfb})(E_{sc}E_{tc})$, unlike any other potentials Like

$U_g = G \frac{Mm}{r}$ or $U_e = K \frac{Q_i Q_j}{r}$, term $(\frac{1}{\Delta r})$ does not appear in this

binding potential

In fact , the quanton , like any other quantum system has its energy

which is defined as $E_p = \frac{hkc}{2\pi}$, can alternatively be written as

$$E_p(\text{ packet energy }) = \frac{hkc}{2\pi} = \frac{hc}{2r_q} \quad (\text{ where } k = \frac{\pi}{r_q})$$

While $E_{tp} = \int_{V_q} E_t \, dv = E_t V_q = \frac{\sqrt{\alpha_t} h}{2 r_q}$

(E_{tp} : total retaining energy inside quanton)

this shows that the quanton radius is inversely proportional to

retaining energy (a binding type interaction) , which already

satisfies the inverse proportionality law

as the quanton energy E_p decreases , its retaining energy

decreases and consequently quanton radius and its wave length

increases , this shows that the term $\frac{1}{r_q}$ is inherently present in the

retaining interaction as well as all forms of quanton interactions

and for the particular case of electromagnetic waves , the inverse

relationship between the wavelength and the energy of the wave is

an expression of an increased binding energy which leads to

a corresponding change in the relativistic quanton dimensions or

its wave length

40. Quanton stable degrees of freedom

The condition of interaction between free and constrained fields

$$(E_{sf} E_{tf})_{binding} = (E_{sc} E_{tc}) \text{ or } (D_{sfb} D_{tfb}) = (D_{sc} D_{tc})$$

, however in addition to generalized matching between the binding

Dof's , a stable interaction requires the symmetry of the binding

Dof's of both sides

To achieve this , a rearrangement of the space and time varying

fields Dof takes place such that

For quations

$$Dof_{tf} = Dof_{tc} \text{ , } Dof_{sfb} = Dof_{sc} \quad (1-40)$$

Energy field	Transient Dof's	Stable Dof's
E_{sf}	2.25	2.00
$E_{sfb} E_{tfb}$ (bound)	1.5	1.50
$E_{sfu} E_{tfu}$ (unbound)	1.00	1.00
E_{sc} (bound)	0.75	1.00
E_{tf}	0.25	0.50
E_{tc} (bound)	0.75	0.50

Table (6) transient and stable quanton degrees of freedom

and for anti quanton

$$Dof_{tf} = Dof_{tc} \quad , \quad Dof_{scb} = Dof_{sf} \quad (2-40)$$

Energy field	Transient Dof's	Stable Dof's
E_{sc}	2.25	2.00
$E_{scb} E_{tcb}$ (bound)	1.5	1.50
$E_{scu} E_{tcu}$ (unbound)	1.00	1.00
E_{sf} (bound)	0.75	1.00
E_{tc}	0.25	0.50
E_{tf} (bound)	0.75	0.50

Table(7) anti quanton transient and stable degrees of freedom

The unbound field remains of the form

$$E_{unbound} = E_{sfu} E_{tfu} = K_q^2 D_{sfu} D_{sfu} \quad \text{and this specific symmetry}$$


Is not translated into the unbound field of the nature

$$E_{unbound} = E_{sfu} = K_q D_{sfu}$$

41. Role of individual energy fields in the formation of space fabric interactions

Energy field	Role inside quanton	Role outside quanton (short range)	interaction at cosmological scale
$E_{sfb} E_{tfb}$ (bound)	Quanton retaining interaction E_t	Quanton binding interaction E_b	Dark matter gravitational like effect
$E_{sc} E_{tc}$ (bound)	Quanton retaining interaction E_t	Quanton binding interaction E_b	Dark matter gravitational like effect
$E_{sfu} E_{tfu}$ (unbound)	Quanton inflationary interaction E_i	Quanton repulsive interaction E_r	Matter distortion of space fabric

Table(8)summary of the role of individual energy fields and their interactions at Planck and cosmological scale for the quantons of space fabric



Fig(8).Summary of the quanton energy fields and the generated interactions

	structure	quanton			Anti quanton		
structure	Energy field	$(E_{sfbj}E_{tfbj})$ (bound)	$E_{scj}E_{tcj}$ (bound)	$E_{sfnj}E_{tfnj}$ unbound	$E_{scbj}E_{tcbj}$ (bound)	$E_{sfj}E_{tfj}$ (bound)	$E_{scui}E_{tcuj}$ (unbound)
quanton	$(E_{sfbi}E_{tfbi})$ (bound)	N/A	E_b	N/A	E_b	N/A	N/A
	$E_{sci}E_{tci}$ (bound)	E_b	N/A	N/A	N/A	E_b	N/A
	$(E_{sfui}E_{tfui})$ unbound)	N/A	N/A	E_r	N/A	N/A	N/A
Anti quanton	$(E_{scbi}E_{tcbi})$ (bound)	E_b	N/A	N/A	N/A	E_b	N/A
	$E_{sfi}E_{tfi}$ (bound)	N/A	E_b	N/A	E_b	N/A	N/A
	$(E_{scui}E_{tcui})$ (unbound)	N/A	N/A	N/A	N/A	N/A	E_r

Table(9) Summary of the generated interactions outside quanton / anti quanton due to different energy fields

42. energy fields' role in the generation of the fundamental forces

1-ordinary matter evolved from quanton / anti quanton pair as they

split in a process that led to the rearrangement of energy fields

Degrees of freedom which became different compared space fabric

case

2-normal matter quantons are quantized , but not a quantum entity

and can be regarded as at the origin of gravitational mass

in addition to gravitational mass ,normal matter is composed of associated fields

3-normal matter quantons are comprise only two degrees of freedom as the remaining two become scalarized (transformed from being part of the field strength to being part of its intensity)

4-for the case of space fabric qunaton system is not under Equilibrium of interactions (equilibrium: absence of the repulsive self interacting fields) ,as it expands and splits , for the case of normal matter quntons , they are under an actual equilibrium of interactions due to the complete symmetry between free and constrained fields , where no inflation or splitting

5-under such conditions , normal matter quantons and anti quantons became identical

6-for space fabric , unbound fields inside the quanton , give

rise to quanton inflation , for the normal matter , the unbound energy fields (associated fields) gave rise to fundamental forces through their interactions with other fields (except gravitation where it is originated from bound energy fields inside the quanton) a model for this rearrangement in the structure of the quanton / anti quanton for the normal matter is as follows

1- the normal matter quantons are formed from space varying fields (E_{sfb} , E_{tfb} E_{scb} E_{tcb}) for quantons and for anti quantons and due to fact that they both have the same Dof for both cases , normal matter quantons and anti quantons are identical

2-unbound fields (E_{sfu} , E_{tfu}), or (E_{sfu} , E_{tfu}) have the following roles ,

a-for the gluons_: they gave rise to part of the strong nuclear force

b-for the electrically charged particles : they are at the origin of

the atomic electric field

3-Difference in the degree of freedom between bound and unbound energy fields prevented the interaction between both groups

43.Degrees of freedom of quantons of normal matter
a-gravitational mass

We recall that the normal matter quantons have only two Dof's and for normal matter both quantons and anti quantons are identical

1-since normal matter quantons are under equilibrium of interactions the bound fields now can reflect the space time symmetry such that

$$Dof_{sfb} = Dof_{scb} = 0.75 \quad , \quad Dof_{tfb} = Dof_{tcb} = 0.25 \quad (1-43)$$

$$(Dof_{sfb} + Dof_{scb}) = 1.5 \quad , \quad \sum Dof_m = 2 \quad (2-43)$$

Energy of the gravitational mass take the non-relativistic form

$$E_m = \sum_i^m \int_{V_p} E_{sfb} E_{tfb} E_{scb} E_{tcb} dV , \quad (3-43)$$

The volumetric integration represents bound fields that are involved in formation of gravitational mass

b-charged atomic fields

a-the unbound fields (E_{sfu} , E_{scu}) have the same Dof

$$(Dof_{sfu} = Dof_{scu} = 1.5 \text{ Dof's}) \quad (4-43)$$

b-unbound fields (E_{tfu} , E_{tcu}) also have the same Dof

$$(Dof_{tfu} = Dof_{tcu} = 0.5) \quad (5-43)$$

3-for positively charged particles : the atomic field is represented by the unbound field (E_{sfu} E_{tfu})

While for the negatively charged particles , the associated atomic Field is represented by the unbound field (E_{scu} E_{tcu})

4-for normal matter , the active degrees of freedom are four

: two for the normal matter quanton , and two for the associated

fields

4- Due to absence of the curl (point source) , the atomic electric

Field becomes invariant (but its denotation is maintained)

c-gluon field

unbound free based field (E_{tfu} E_{tfu}) , and constrained based

field (E_{scu} E_{tcu}) , each field is split into main fields representing

(1.5) Dof's and auxiliary field representing (0.5) Dof's

parameter	Space fabric quantons	Normal matter quantons
Nature of the quanton	Quantum entity	<u>Not</u> a quantum entity
Bound fields	<u>Q</u> : $(E_{sfb} E_{tfb}) (E_{sc} E_{tc})$ <u>AQ</u> : $(E_{sf} E_{tf})(E_{scb} E_{tcb})$	<u>Q or AQ</u> : $E_{sfb} E_{tfb} E_{scb} E_{tcb}$ (both are identical)
unbound fields	<u>unbound fields</u> : <u>Q</u> : $(E_{sfu} E_{tfu})$ <u>AQ</u> : $(E_{scu} E_{tcu})$	<u>unbound fields</u> <u>Gluons</u> <u>Q</u> : $E_{sfu} E_{tfu}$ (main and auxiliary) <u>AQ</u> : $E_{scu} E_{tcu}$ (main and auxiliary) <u>Positive particles</u> $E_{sfu} E_{tfu}$ <u>negative particles</u> $E_{scu} E_{tcu}$
Wave behaviour	Q+AQ pair has wave properties	Inside quantons : No wave behaviour , only binding energy fields
Degrees of freedom	Four	Dof_q : two Associated unbound fields : two
Nature of fields E_{qf} , E_{qc}	orthogonal	parallel
Quanton Expansion , splitting	Quantons Expand , and split ,	No expansion or splitting (quantons are under actual equilibrium)
Variation (under static conditions)	Varying in space and in time	invariant

Table (10) Summary of the differences between space fabric and normal matter quantons

44. Gravitational mass and its relativistic effect

The reduced quanton of the normal matter is composed of a pair

Of coplanar space varying fields (free /constrained) which

$$\mathbf{E}_{qf} = \mathbf{E}_{sfb} \mathbf{E}_{tcb} , \quad \mathbf{E}_{qc} = \mathbf{E}_{scb} \mathbf{E}_{tfb} \quad (1-44)$$

$$\mathbf{E}_m = \sum_j^n \int_{V_p} \mathbf{E}_{qf} \mathbf{E}_{qcbj} \, dV = \sum_j^n \mathbf{E}_{qf} \mathbf{E}_{qcbj} V_{pj} \quad (2-44)$$

unlike the case of quanton fields or electromagnetic waves

(where the free dominated field \mathbf{E}_{qf} and the constrained

dominated \mathbf{E}_{qc} are orthogonal to each other) ,for the normal

matter the free and the constrained energy dominated fields

are coplanar (exist in one plane) , the magnitude of the energy

density is represented by dot product of both fields , this would

lead to the development of field equations of gravitational mass

$$\mathbf{E}_{qf} \times \mathbf{E}_{qc} = 0 \quad (\text{fields are coplanar , their cross product equals} \\ \text{zero}) \quad (3-44)$$

$$\nabla \cdot \mathbf{E}_{qf} = - \nabla \cdot \mathbf{E}_{qc} \quad (\text{completely asymmetric fields}) \quad (4-44)$$

44.b For the relativistic effects of the gravitational matter

as the inertial body moves along a certain direction (x) , the fields

E_{qf} , E_{qc} undergo a restriction , from being 3 dimensional , to

becoming two dimensional (y, z) which is orthogonal to the

movement direction

the main driving force behind this change is to maintain the

integrity of the matter , under such conditions , we would expect

there would be no energy fields along the direction of motion

the relativistic mass under Lorentz transform of transverse

energy fields now becomes $E_{mo}' = (E_{qf}' E_{qc}' V_p)$ (5-44)

$$E_{mo}' = \frac{(E_{qf} E_{qc} V_p)}{\sqrt{1 - (\frac{v}{c})^2}} = \frac{E_{mo}}{\sqrt{1 - \beta^2}} \quad (6-44)$$

and the same results can be obtained via the energy momentum

relationship where $Pc = \frac{E}{c} v = \frac{(E_{qf} E_{qc} V_p)}{c} v$ (7-44)

$$E_m^2 = m^2 c^4 = P^2 c^2 + m_0^2 c^4$$

$$(E_{qf}' E_{qc}' V_p)^2 = (E_{qf}' E_{qc}' V_p)^2 \frac{v^2}{c^2} + (E_{qf} E_{qc} V_p)^2 \quad (8-44)$$

$$(E_{qf}' E_{qc}' V_p)^2 \left(1 - \frac{v^2}{c^2}\right) = (E_{qf} E_{qc} V_p)^2 \quad (9-44)$$

$$(E_{qf}' E_{qc}' V_p) = E_m = \frac{E_{qf} E_{qc} V_p}{\sqrt{1-\beta^2}} = \frac{E_{m0}}{\sqrt{1-\beta^2}} \quad (10-44)$$

45. Energy field parameters for normal matter

Normal matter quanton which is composed of bound energy fields

$(E_{sfb} E_{tfb}) (E_{sch} E_{sch})$ is not a quantum entity (as it possesses only

two degrees of freedom) ,no splitting or expansion , yet it can be

quantized form using the relationship $E_p = \frac{\alpha_m h c}{2 r_p}$

where r_p (particle radius) = fixed

$$\begin{aligned} E_m = M c^2 &= \sum_j^n \frac{m_j}{c^2} c^4 = \sum_j^n \frac{\alpha_m h}{2 c^3 r_{pj}} c^4 \\ &= n \frac{\alpha_m h}{2 c^3 r_p} c^4 \end{aligned} \quad (1-45)$$

where $\frac{\alpha_m h}{2 c^3 r_p} = \text{constant}$ (2-45)

this is quantized energy relationships and not a quantum

relationship since the Planck Einstein relationship is not applicable

namely $E_m \neq \text{fn} \left(\frac{1}{r_p} \right)$, r_p represents the radius of gravitational

mass's quanton ,

normal matter energy is presented in this quantized form

as it will serve two main purposes

1-to define field interactions in terms of the constant (c)

2-to facilitate studying interactions with quantum based fields.

the parameters ω , k for the quanton are now replaced by the

alternative characteristic length (r_p)

now the energy of the gravitational mass

$$E_m = \sum_i^n \int_{V_p} E_{sfbi} E_{scbi} E_{tfb} E_{tcb} dV = (E_{sfb} E_{scb} E_{tfb} E_{tcb}) V_p \quad (3-45)$$

$$E_m = n E_{sfb} E_{scb} E_{tfb} E_{tcb} V_p = n E_m V_p = n \frac{\alpha_m hc}{2 r_p} \quad (4-45)$$

and as an energy density

$$E_{pm} = \frac{n\alpha_m hc}{2r_p (V_p)} = \frac{n\alpha_m hc}{2r_q (8r_p^3)} = n \frac{\alpha_m hc}{16r_p^4} \quad (5-45)$$

45.a-NM Bound energy fields

These degrees of freedom here become part of the intensity parameter as the NM quanton has two Dof's only

(scalarized degrees of freedom)

$$E_{sfb} = \sqrt[4]{\frac{\alpha_m h c^2}{16\pi^4 c^3} \frac{c^{0.75}}{r_p}} = K_p c^{0.75} = K_{sfb} D_{sfb} \quad (6-45)$$

$$K_{sfb} = K_p = \sqrt[4]{\frac{\alpha_m h c^2}{16\pi^4 c^3} \frac{1}{r_p}}, \quad D_{sfb} = c^{0.75} \quad (7-45)$$

$$E_{tfb} = \sqrt[4]{\frac{\alpha_m h c^2}{16\pi^4 c^3} \frac{c^{0.25}}{r_p}} = K_p c^{0.25} = K_{tfb} D_{tfb} \quad (8-45)$$

$$E_{scb} = K_{scb} D_{scb} = \sqrt[4]{\frac{\alpha_m h c^2}{16\pi^4 c^3} \frac{c}{r_p^2}} = K_p c^{0.75} = K_{scb} D_{scb} \quad (9-45)$$

$$E_{tcb} = \sqrt[4]{\frac{\alpha_m h c^2}{16\pi^4 c^3} \frac{c^{0.25}}{r_p}} = K_p c^{0.25} = K_{tcb} D_{sfb} \quad (10-45)$$

where $K_{sfb} = K_{tfb} = K_{scb} = K_{tcb} = K_p^4$

45.b- unbound energy fields

45.b.1-positively charged particles

$$E_{sfu} = \sqrt[4]{\frac{\alpha_e h}{16 \pi^4 c^3}} \frac{c^{1.5}}{r_p} = K_{sfu} D_{sfu} = K_p c^{1.5} \quad (11-45)$$

$$E_{tfu} = \sqrt[4]{\frac{\alpha_e h}{16 \pi^4 c^3}} \frac{c^{0.5}}{r_p} = K_{tfu} D_{tfu} = K_p c^{0.5} \quad (12-45)$$

$$K_{sfu} K_{tfu} = K_p^2, \quad \alpha_e = \frac{1}{137} \quad (13-45)$$

45.b.2-negatively charged particles

$$E_{scu} = \sqrt[4]{\frac{\alpha_e h}{16 \pi^4 c^3}} \frac{c^{1.5}}{r_p} = K_{scu} D_{scu} = K_p c^{1.5} \quad (14-45)$$

$$E_{tcu} = K_{tcu} D_{tcu} = K_p c^{0.5}, \quad K_{scu} K_{tcu} = K_p^2 \quad (15-45)$$

The energy content of gravitational mass system as the product of

$$E_m = \sum_j^n V_{pj} E_{qmj} = (\sum_j^n E_{sfbj} E_{scbj} V_{pj})$$

given that $V_p = \text{constant}$, $\sum_j^n V_{pj} = n V_p$

$$E_m = n V_p (\sum_j^n E_{sfbj} E_{tfbj}) (E_{scbj} E_{tcbj})$$

Where the dimension of the bound fields $E_{sfb}E_{tfb}E_{scb}E_{tcb}$ is

$$\left[\frac{hc}{c^2 r_p^4} \right] = M^1 L^{-3} T^{00} = \frac{\text{energy}}{\text{volume} * c^2} = \frac{\text{mass}}{\text{volume}}$$

46. scalarized degrees of freedom, a possible origin of gravitational mass,

Gravitational mass density is represented by the product of the normal matter field intensities

the field intensity parameter for normal matter quantons

$$K_{sfb} K_{tfb} K_{scb} K_{tcb} = K_p^4 = \left(\sqrt[4]{\frac{\alpha_m h}{16 \pi^4 c}} \right)^4 \left(\frac{1}{r_p^4} \right)$$

$$= \frac{\alpha_m h}{16 \pi^4 c r_p^4} = \frac{mass}{volume}$$

for normal matter the intensity parameter became

$$K_{sfb} = K_{tfb} = K_{scb} = K_{tcb} = \sqrt[4]{\frac{h c^2}{16 \pi^4 c^3}} \frac{1}{r_p} = \sqrt[4]{\frac{h}{16 \pi^4 c}} \frac{1}{r_p} \text{ instead of}$$

$$\sqrt[4]{\frac{h}{16 \pi c^3}} \frac{1}{r_p} \text{ for the space fabric quantons}$$

while energy density equation of normal matter quanton is in the

form $E_q = E_{sfb} E_{scb} E_{tfb} E_{tcb}$, with a reduction of over all degree

of freedom from four to two due to the fact that two degrees of

freedom now transformed from belonging to the field strength

parameter to become a part of the field intensity

as a result of this reduction of degrees of freedom the normal

matter, Dof of quantons representing the gravitational mass

become of the form (1.5+0.5) instead of (3+1)

gauge theory prevents the gauge particles from acquiring mass ,

however , under low dimensions conditions , photons , gluons

can acquire a dynamic mass under Schwinger model of reduced

dimensions , here we suggest a generalization which proposes

that reduction in the energy degree of freedom is possibly at

the origin of mass generation (rest / dynamic)

47. field interactions of normal matter

47.a-Quanton retaining interaction (Type : single binding)

$$(E_t) = (E_{sfb} E_{tfb}) (E_{scb} E_{tcb}) \quad (1-47)$$

$$= (K_p^2 D_{sfb} D_{tfb}) (K_p^2 D_{scb} D_{tcb})$$

$$= \int_{V_p} (K_p^2 \frac{c}{r_p^2}) (K_p^2 \frac{c}{r_p^2}) dV$$

$$E_t = \alpha_t \left(\frac{h c^2}{16 \pi^4 c^3} \right) \frac{c^2}{r_p^4} (8 r_p^3) = \alpha_t \frac{h c}{2 r_p} \quad (2-47)$$

Where $K_p = \sqrt[4]{\frac{h}{16 \pi c}}$

This interaction has two degrees of freedom (compared to three

For the case of space fabric) , the dimensions of energy= $M^1 L^2 T^{-2}$

47.b-quanton's gravitational-like binding

type : multiple binding

the normal matter particles would develop a gravitational type of

binding as energy fields tend to form higher order interactions up

to four degrees of freedom

bound energy fields of each quanton form a gravitational

binding interaction with bound energy fields of other quantons

of the form $(E_{sfbi} E_{tfbi}) (E_{scbj} E_{tcbj})$ and $(E_{scbi} E_{tcbi}) (E_{sfbj} E_{tfbj})$

to generate the gravitational binding energy E_{gb} between particle

p_i and other particles p_j

formulation of the gravitational binding energy of the normal matter differs from all other interactions due to the following reasons

1-for normal matter space and time varying fields

$(E_{sfb} E_{tfb})(E_{scb} E_{tcb})$ the intensity parameter is of nature (K_p^4)

2-the gravitational binding interaction is based on two binding interactions for particles p_i, p_j , which are

a-between $(E_{sfb_i} E_{tfb_i})$ and $(E_{scb_j} E_{tcb_j})$

b-between $(E_{sfb_j} E_{tfb_j})$ and $(E_{scb_i} E_{tcb_i})$

those two simple interactions combine to form gravitational binding since each one of those interactions has only two degrees of freedom (combinations of interactions allowed up to 4 Dof's)

3-The resulting interaction has would be in the form

$$E_g = K_g (K_{pi}^4 c^2) (K_{pj}^4 c^2) \frac{r_{pi} r_{pj}}{(r_i - r_j)} \quad (3-47)$$

intensity term is becomes $(K_p^4)^2$ instead of (K_p^4) which is

required for true energy generated by the interaction

as E_g has the dimensions of energy $M^1L^+2T^{-2}$, the constant K_g

appears as a dimensional correction since each of the parameters

$$[K_{pi}^4 V_{pi}][K_{pj}^4 V_{pj}] = \left(\frac{h}{c\pi r_p}\right)^2 = \left[\frac{\text{energy}}{c^2}\right]*\left[\frac{\text{energy}}{c^2}\right]$$

To obtain a truly binding interaction E_g (in terms of energy

With dimensions $M^1L^+2T^{-2}$) the constant K_g should be equivalent

$$\text{to } \frac{c^4}{E_{ref}} \text{ where for normal matter } E_{ref} = \frac{hc}{2r_p} \quad (4-47)$$

$$E_{gbi} = \frac{c^4}{E_{ref}} \left[\left(\int_{V_{pi}} E_{sfbi} E_{tfbi} E_{scbi} E_{tcbi} dV \right) \sum_j^n \left(\int_{V_{pj}} E_{sfbj} E_{tfbj} E_{scbj} E_{tcbj} dV \right) \left(\frac{\sqrt{r_{pi}r_{pj}}}{(r_i-r_j)} \right) \right] \quad (5-47)$$

$$= \frac{c^4}{E_{ref}} [K_{pi}^4 D_{sfbi} D_{scbi} D_{sfbi} D_{scbi} V_{qi}] \sum_j^n K_{pj}^4 D_{sfbj} D_{tfbj} D_{scbj} D_{tcbj} V_{qj} \left(\frac{\sqrt{r_{pi}r_{pj}}}{(r_i-r_j)} \right)$$

$$(E_{gbi}) = \frac{c^4 r_p^2}{E_{ref}} \left[(K_{pi}^4 V_{pi} c) \left(\sum_j^n K_{pj}^4 V_{pj} c \right) \left(\frac{1}{(r_i-r_j)} \right) \right]$$

Which is a summation for particles (j)

$$\text{Given that } r_{pi}=r_{pj} = r_p , K_{pi} = K_{pj} = K_p = \sqrt[4]{\frac{h}{16\pi^4 c}}$$

$$\int_{V_p} E_{sfb} E_{tfb} E_{scb} E_{tc} dV = E_{sfb} E_{tfb} E_{scb} E_{tcb} V_p$$

$$V_{pi} = V_{pj} = V_p = 8 r_p^3$$

$$K_p^4 c = \frac{h}{16 c} \left(\frac{1}{r_p}\right)^4 c = \frac{h}{2 c r_p} \frac{1}{V_p}$$

$$\begin{aligned} E_{gbi} &= \frac{2\alpha_g c^4 r_p^2}{hc} \left[\frac{h}{2 c V_{pi}} \frac{1}{r_p} V_{pi} \right] \sum_j^n \left(\frac{h}{2 c V_{pj}} \frac{1}{r_p} V_{qj} \right) \left(\frac{1}{(r_i - r_j)} \right) \\ &= \left(\frac{\alpha_g c^4}{hc} \right) \left(\frac{h}{c} \right) \sum_j^n \left(\frac{h}{2c} \right) \left(\frac{1}{(r_i - r_j)} \right) \end{aligned} \quad (6-47)$$

$$= (\alpha_g c^2) \sum_j^n \left(\frac{h}{2c} \right) \left(\frac{1}{(r_i - r_j)} \right)$$

$$(E_{gbi}) = \frac{\alpha_g h c}{2} \sum_j^n \left(\frac{1}{(r_i - r_j)} \right) \quad (7-47)$$

$$\text{Where } G \text{ can be defined in terms of } \left(\frac{2\alpha_g c^3 r_p^2}{h} \right) \quad (8-47)$$

$$\text{And } r_p = \sqrt{\frac{Gh}{2\alpha_g c^3}}, \quad (9-47)$$

$$r = \sqrt{\frac{Gh}{2\pi c^3}} \text{ is nothing other than the Planck length}$$

it is worth noting that while the gravitational constant G remains


invariant with time as the normal matter particle radius $r_p =$

constant , the binding parameter for space fabric

$K_g = \frac{2\alpha_g c^2 r_q^2}{h}$ is a variable with time as the quanton radius itself

r_q varies with time

47.b.Role of normal matter energy fields in the generation of the fundamental forces



Fig(9).Summary of the interactions of bound and unbound fields for gluons

Energy field	Role at short range(inside quanton)	Role outside quanton	interactions at long range
$E_{sfb} E_{tfb}$ (bound)	1-Quanton retaining interaction E_t	Quanton binding E_b (gravitational binding)	1-gravitation 2- space fabric distortion
$E_{scb} E_{tcb}$ (bound)	1-Quanton retaining interaction E_t	Quanton binding E_b (gravitational binding)	1-gravitation 2- space fabric distortion
$E_{sfu} E_{tfu}$ (unbound)	1-Strong nuclear force (attraction and repulsion part) 2-dark matter gravitation like effect	1-Strong nuclear force (attraction and repulsion part) 2-dark matter gravitation like effect	N/A
$E_{scu}E_{tcu}(unbound)$	Strong nuclear force (attraction and repulsion part	Strong nuclear force (attraction and repulsion part	N/A

Table(11) Summary of the role of the interactions developed by each energy field at Planck and cosmological scales for gluons




Fig (10).Summary of developed interactions due to bound and unbound fields for positively charged particles

Energy field	Role at short range(inside quanton)	interactions outside quanton	Long range interactions
$E_{sfb}E_{tfb}$ (bound)	1-Quanton retaining interaction E_t 2-inertial retaining energy	Quanton binding interaction E_b (gravitational binding)	1-gravitation 2- space fabric distortion
$E_{scb}E_{tcb}$ (bound)	1-Quanton retaining interaction E_t 2-inertial retaining energy	Quanton binding interaction E_b (gravitational binding)	1-gravitation 2- space fabric distortion
$E_{sfu} E_{tfu}$ (unbound)	Atomic electric field	Atomic Electric field	Atomic Electric field
$E_{scu}E_{tcu}$ (unbound)	N/A	N/A	N/A

Table(12) Summary of the interactions developed by each energy fields at different scales for positively charged particles

48.Gravitation binding interaction

Type : multiple binding

outside quantons , energy field interactions are involved in

maintaining inter-quanton integrity via the gravitational binding

interaction , but as pointed out earlier that energy fields are infinite

in range , there is a residual amount that is left untied in any

binding interaction which gives rise to gravitation , defined as the

summation of interactions due to this residual free and

constrained field outside of the quanton between two bodies (i ,j)

E_g : gravitational binding energy

The term ($E_{sfbi} E_{tfbi} E_{scbi} E_{tcbi} V_{qi}$) represents the mass(= $\frac{E_{mi}}{c^2}$) ,

$$\text{where } E_g = G \left(\frac{E_{mi}}{c^2} \right) \left(\frac{E_{mj}}{c^2} \right) \frac{1}{(r_i - r_j)} = G \frac{m_i m_j}{(r_i - r_j)} \quad (G = \frac{2\alpha_g c^3 r_p^2}{h})$$

$$= \frac{2}{hc} [\sum_i^m \int_{V_{pi}} E_{sfbi} E_{tfbi} E_{scbi} E_{tcbi} dV] [\sum_j^n \int_{V_{pj}} E_{sfbj} E_{tfbj} E_{scbj} E_{tcbj} dV] \frac{r_{pi} r_{pj}}{(r_i - r_j)} \quad (1-48)$$

$$= \frac{2 r_p^2}{hc} (\sum_i^m K_{pi}^4 D_{sfbi} D_{tfbi} D_{scbi} D_{tcbi} V_{pi}) (\sum_j^n K_{pj}^4 D_{sfbj} D_{tfbj} D_{scbj} D_{tcbj} V_{pj}) \frac{1}{(r_i - r_j)}$$

$$= \frac{2\alpha_g r_p^2}{hc} \left(\sum_i^m \frac{h}{16 c r_{pi}^3} \frac{c^2}{r_{pi}} V_{pi} \sum_j^n \frac{h}{16 c r_{pj}^3} \frac{c^2}{r_{pj}} V_{pj} \frac{1}{(r_i - r_j)} \right)$$

$$E_g = \frac{2\alpha_g r_p^2}{hc} \left(\sum_i^m \frac{h c}{2 (8 r_{pi}^3)} \frac{V_{pi}}{r_{pi}} \sum_j^n \frac{h c}{2 (8 r_{pj}^3)} \frac{V_{pj}}{r_{pj}} \frac{1}{(r_i - r_j)} \right)$$

$$= \alpha_g c r_p^2 \left(\sum_i^m \left(\frac{h}{2 r_{pi}} \right) \sum_j^n \frac{1}{r_{pj}} \frac{1}{(r_i - r_j)} \right)$$

$$E_g = \frac{\alpha_g h c}{2} \left(\sum_i^m \sum_j^n \frac{1}{(r_i - r_j)} \right) \quad (2-48)$$

to note that the gravitation is the only force due to residual of

fields between two bound energy fields $(E_{sfb} , E_{tfb}) (E_{scb} , E_{tcb})$

those energy field form the retaining interaction E_t first , then the

gravitational like binding interaction E_{gb} and gravitation at last and

this is one of the reasons behind the weakness of gravitation in

comparison to other forces

49.Atomic electric charge and field

unbound fields for the case of charged particles are expressed in

the form of atomic electric field and ensuing electric charge ,

those unbound energy fields must now be defined in terms of

dimensions the new particle structure rather than the quanton

dimensions

energy stored in the positive atomic electric field is in the form

$$E_e = \sum_i^m \int_{V_p} E_{qi} dV = \int_{V_p} (E_{sfu}E_{tfu})^2 dV = \sum_i^m E_{qi} V_{pi} \quad (1-49)$$

$$\text{Or } \alpha_e \frac{hc}{2 r_p} = \frac{Q^2}{4\pi\epsilon_0 r_p} \quad (2-49)$$

$\alpha_e =$ coupling constant for atomic electric field , V_p : particle

Volume , for the case of positively charged particles (free energy dominated), the atomic charge can be assessed using Gauss law

, where $\int E_{sfu} E_{tfu} dA = \frac{Q}{\epsilon_0}$

$Q =$ charge density , $E_{sfu} E_{tfu}$ are the unbound now invariant atomic (static) electric field

$E(+)$ = $E_{sfu} E_{tfu} = \frac{Q}{4\pi \epsilon_0 r_p^2}$, r_p : estimated radius of the particle

$Q(+)$ = $4 \pi \epsilon_0 r_p^2 (E_{sfu} E_{tfu})$

$$= 4 \pi \epsilon_0 r_p^2 K_p^2 D_{sfu} D_{tfu} \quad (3-49)$$

Which has the dimensions of $M^{0.5} L^{+1.5} T^{-1}$

The accompanying electric field at any point (r_0) becomes

$$E (+) = \frac{Q}{4 \pi \epsilon_0 (\Delta r_0)^2} = \frac{r_p^2 E_{sfu} E_{tfu}}{(\Delta r_0)^2} = \sqrt{\frac{\alpha_e h c}{2 V_p r_p}} \frac{r_p^2}{(\Delta r_0)^2} \quad (4-49)$$

Which has the dimensions of $M^{0.5} L^{-0.5} T^{-1}$

for negatively charged particles (anti quantons dominant)

$$Q(-) = 4 \pi \epsilon_0 r_p^2 E_{scu} E_{tcu} \quad (5-49)$$

Where E_{scu} E_{tcu} are the unbound invariant constrained fields

49b. Electric binding energy

$$E_e = K_e \frac{Q_i Q_j}{(\Delta r_{ij})}$$

$$= K_e (4 \pi \epsilon_0 r_p^2) (E_{sfui} E_{tfui}) (4 \pi \epsilon_0 r_p^2) (E_{scuj} E_{tcuj}) \frac{\sqrt{r_{pi} r_{pj}}}{(r_i - r_j)} \quad (6-49)$$

$$E_e = \frac{4 \pi \epsilon_0 \alpha_e hc}{(r_i - r_j)} \quad (7-49)$$

K_e : Coulomb Constant ($=4 \pi \epsilon_0$) ,

50. Strong nuclear binding / repulsive interaction

1-It is represented by self-interaction of the unbound free and

constrained energy fields

2-real energies (which have the dimension of $ML^{+2}T^{-2}$) must be

generated by interactions which have four degrees of freedom

(terms of c^4) , so we should expect the strong self interaction

also to be to have four degrees of freedom

3-gluons are based equitably on both free and constrained fields

so as to provide for the symmetry of the self interaction

free energy field based flux tube V_{fi} of the form $(E_{sfu} E_{tfu})$

this field which has two Dof's is complex in nature , as it is split to

main and auxiliary fields such that

$$E_{sfu} E_{tfu} = K_p^2 (D_{sfu} D_{tfu}) \quad (1-50)$$

$$= K_p^2 (D_{sfum} D_{tfum}) (D_{sfua} D_{tfua}) \quad (2-50)$$

$$= (E_{sfum} E_{tfum}) (E_{sfua} E_{tfua}) \quad (3-50)$$

where $D_{sfum} D_{tfum} = c^{1.5}$, $D_{sfua} D_{tfua} = c^{0.5}$

constrained energy field based flux tube V_{fj} in the form

$(E_{scu} E_{tcu})$, which has two Dof's and is split also into main and

auxiliary fields $E_{scu} E_{tcu} = K_p^2 (D_{scu} D_{tcu}) \quad (4-50)$

$$= K_p^2 (D_{scum} D_{tfum}) (D_{scua} D_{tcua}) \quad (5-50)$$

$$= (E_{scum} E_{tcum}) (E_{scua} E_{tcua}) \quad (6-50)$$

where $D_{scum} D_{tcum} = c^{1.5}$, $D_{scua} D_{tcua} = c^{0.5}$

4-energy stored in the flux tubes

$$E_s = \int_{V_f} (E_{sfu} E_{tfu})^2 dV \quad \text{and} \quad (7-50)$$

$$E_s = \int_{V_f} (E_{scu} E_{tcu})^2 dV \quad (8-50)$$

, V_f : flux tube volume

5- strong force self interaction degrees of freedom are in the form

(1.5+0.5) Dof's for free or constrained based gluons , and this is

since the energy fields can achieve full merger only while inside

quanton

a-Repulsive part (self interaction) type : simple nonbinding

the repulsive part of strong nuclear force is a self interaction

based gluon flux tubes with free energy fields $(E_{sfu} E_{tfu})$ in

addition to self interaction of the constrained energy field based

flux tubes or $(E_{scu}E_{tcu})$ and generating the repulsive part of the strong binding energy , the interaction takes the form

$$E_{sr} = \left(\int_{V_f} K_p^4 [D_{sfu}D_{tfu}]^2 dV + \int_{V_f} K_p^4 [D_{scu}D_{tcu}]^2 dV \right) \left(\frac{r_p}{\Delta r_p} \right) \quad (9-50)$$

Δr_p : characteristic length : distance between two quarks ,

the first term describes the contribution of free fields , while the

second term describes the contribution of constrained fields

$$E_{sr} = K_p^4 \left(\sum_j^n [D_{sfu}D_{tfu}]^2 V_{fj} + \sum_j^n [D_{scu}D_{tcu}]^2 V_f \right) \left(\frac{r_p}{\Delta r_p} \right) \quad (10-50)$$

$$= \alpha_s \left(\sqrt{\left(\frac{h}{2 c^3 V_p r_p} \right)} \right)^2 (c^2)^2 \sum_j^n V_{fj} \left(\frac{r_p}{\Delta r_p} \right)$$

$$E_{sr} = \alpha_s \frac{hc}{2 \Delta r_p} \sum_j^n \left(\frac{V_{fj}}{V_p} \right) \quad (11-50)$$

α_s : strong coupling constant

This repulsive interaction has four degrees of freedom and the

dimensions of $M^1 L^{+2} T^{-2}$

50b- the binding part type : simple binding

the attraction part is generated by the interaction between free

energy dominated flux tubes and constrained dominated gluon

flux tubes

$$\begin{aligned}
 E_{sb} &= \left(\int_{V_f} (E_{sfu} E_{tfu} \ E_{scu} E_{tcu} dV) \left(\frac{r_p}{\Delta r_f} \right) \right. \\
 E_{sb} &= K_p^4 \left(\int_{V_f} (D_{sfu} D_{tfu} \ D_{scu} E_{tcu} dV) \left(\frac{r_p}{\Delta r_f} \right) \right. & (12-50) \\
 &= K_p^4 \sum_j^n (D_{sfu} D_{tfu}) (D_{scu} D_{tcu}) V_{fj} \left(\frac{r_p}{\Delta r_f} \right) \\
 &= \alpha_s \left(\sqrt{\left(\frac{h}{2 c^3 V_p r_p} \right)} \right)^2 (c^2)^2 \sum_j^n V_{fj} \left(\frac{r_p}{\Delta r_f} \right) \\
 &= \alpha_s \frac{h c}{2 V_p r_p} V_f \left(\frac{r_p}{\Delta r_f} \right) = \alpha_s \frac{h c}{2 \Delta r_f} \sum_j^n \frac{V_{fj}}{V_p} & (12-50)
 \end{aligned}$$

Δr_f = average distance between the flux tubes (n)

it is noted that the distance (Δr_f) between flux tubes = constant

as the distance between quarks increases , V_f increases linearly

as more energy is being added to the flux tubes , so the potential

for the attraction energy increases linearly with the distance ,

unlike the case of repulsive interaction where (Δr_p) (distance

between quarks) changes and the value of the interaction

changes accordingly , while energy content of the flux tubes

remains the same

51.Gravitational like attraction of dark matter

Type : multiple binding

1-The interaction that generates the gravitational like attraction of

the dark matter is between fields of the bound fields $(E_{sfb} E_{tfb})_s$

or $(E_{scb} E_{tcb})_s$ of space fabric and the unbound main free field

$(E_{sfum} E_{tfum})_m$ or main constrained field $(E_{scum} E_{tcum})_m$ of the

normal matter's gluons which are responsible for the nuclear

strong force

2-space fabric bound fields $(E_{sfb} E_{tfb})_s$ or $(E_{scb} E_{tcb})_s$

have 1.5 Dof's each and could not create a gravitational binding

interaction with the galactic normal matter binding fields

$(E_{sfb} E_{tfb})_m$, $(E_{scb} E_{tcb})_m$ (two Dof's each) of normal matter

quantons , those same energy fields which generate the

gravitation binding and the reason behind this is due to field Dof

mismatch

3-gravitation like attraction of the dark matter is one-sided only In

the sense that there is no gravitational like attraction of the normal

matter on the space fabric , and this is because there is no long

range action of the unbound gluon fields ($E_{sfum} E_{tfum}$)

or ($E_{scum} E_{tcum}$) of the nuclear strong force (their fields are in the

form of flux tubes which is short acting range)

51.b.Gravitational like effect on normal matter free energy field

While in its density interaction form $E_{gs} =$

$$[(E_{sfumi} E_{tfumi} E_{scumi} E_{tcumi})_m \sum_j^n (E_{sfbj} E_{tfbj} E_{scj} E_{tcj})_s \left(\frac{\sqrt{r_{pi} r_{qj}}}{(r_i - r_j)} \right)] (1-51)$$

This binding interaction has three Dof's and the dimensions

of $\left(\frac{\text{energy}}{c}\right)$, while in its complex form

$$E_{gs} = \frac{c^2}{E_{ref}}$$

$$\left[\int_{V_f} (E_{sfumi} E_{tfumi} E_{scumi} E_{sfumi})_m \, dV \sum_j^n \int_{V_{qs}} (E_{sfbj} E_{tfbj} E_{scj} E_{tcj})_{qs} \, dV \right] \left(\frac{\sqrt{r_{pi} r_{qj}}}{(r_i - r_j)} \right) \quad (2-51)$$

$$= \frac{c^2}{E_{ref}} \sum_i^m K_{pi}^4 (D_{sfum} D_{tfum} D_{tcum} D_{tcum})_m V_{fi} \sum_j^n K_{qj}^4 (D_{sfbj} D_{tfbj} D_{sc} D_{tc})_s V_{qj} \left(\frac{\sqrt{r_{pi} r_{qj}}}{(r_i - r_j)} \right)$$

(3-51)

$$= \frac{2\sqrt{\alpha_s \alpha_b} r_{ref} c^2}{hc} \left[\left(c^3 \left(\frac{h}{2 c^3 V_{pi} r_{pi}} \right) V_{fi} \sum_j^n c^3 \left(\frac{h}{2 c^3 V_{qj} r_{qj}} \right) V_{qj} \left(\frac{\sqrt{r_{pi} r_{qj}}}{(r_i - r_j)} \right) \right) \right]_{m-s}$$

$$E_{gs} = \frac{\sqrt{\alpha_s \alpha_b} h}{2} \frac{V_{fi}}{V_{pi}} \sum_j^n \left(\frac{1}{(r_i - r_j)} \right)_{m-qs} \quad (4-51)$$

this binding interaction between free and constrained energy

fields of gluon (i) of the normal matter with bound free constrained

energy fields of the space fabric quantons or anti quantons (j)

$$E_{ref} = \frac{hc}{2\sqrt{r_p r_q}} \text{ , and since the quanton radius of space fabric is}$$

varying with time , it is expected that the gravitational parameter

of this interaction to be varying also with time as well

51.c. Why normal matter gravitation generates space fabric distortion

It had been proposed that bound energy fields $(E_{sfb} E_{tfb})_s$ or

$(E_{scb} E_{tcb})_s$ of space fabric which generate the space fabric

binding interaction E_b , would also generate the gravitational

attraction of the dark matter on the normal matter E_{gs} , since the

binding interaction is more stable than the repulsive alternative

,but why this is not the case for the normal matter, which, based

on the fore-mentioned discussion, the bound fields of normal

matter $(E_{sfb} E_{tfb})$, $(E_{scb} E_{tcb})$ which generate gravitation should

have generated another binding interaction to the space fabric,

reasons are as follows

1-the unbound fields of space fabric $(E_{sfu} E_{tfu})_q$ of quanton

and $(E_{scu}E_{tcu})_{aq}$ of anti quanton generate self-interacting fields those fields , which are at the origin of the quanton expansion , splitting and the inflationary momentum in general , are repulsive in nature , which means that they are complex repulsive fields (have combined Dof that is equal $0.5 +0.5$)those repulsive fields do not merge to generate a resultant field of Dof strength = 1) as those fields are of the form :

$$(K_q^2 \sqrt{(D_{sfu} D_{tfu})_q}) (K_q^2 \sqrt{(D_{sfu} D_{tfu})_q}) \text{ or}$$

$$(K_q^2 \sqrt{(D_{scu} D_{tcu})_{aq}}) (K_q^2 \sqrt{(D_{scu} D_{tcu})_{aq}}) \text{ and not of the form}$$

$K_q^2 (D_{sfu} D_{tfu})_q$ or $K_q^2 (D_{scu} D_{tcu})_{aq}$, this means that those

unbound energy fields of space fabric can only be involved in repulsive interaction with bound fields of normal matter

$(E_{sfb} E_{tfb})_m$, $(E_{scb} E_{tcb})_m$ (which are generating gravitation

force) ,and this repulsive interaction is at the origin of normal

matter distortion of space fabric

2-a gravitational binding interactions between galactic normal

matter and space fabric bound fields could not be generated , since

for normal matter binding energy fields of $(E_{sfb} E_{scb})_q$ each type

of energy fields has two Dof's , while for space fabric bound fields

$(E_{sfb} E_{tfb})_s$ or $(E_{scb} E_{tcb})$, each complex field has 1.5 Dof's

(remembering that fields leave space fabric quanton as complex

fields of free or constrained nature) , as a result , there is both Dof

mismatch as well as field type mismatch

52.Gravitational distortion of space fabric

52.a.1-the repulsive interaction on a quanton (i) of space fabric

$$E_{di} = \frac{c^4}{E_{ref}}$$

$$\left[\int_{V_{qs}} [(E_{sfui} E_{tfui})_{qs} (E_{scui} E_{tcui})_{aqs}] dV \right] \left(\sum_j^n \int_{V_p} (E_{sfbj} E_{sfbj})_m (E_{scbj} E_{scbj})_m dV \right) \frac{\sqrt{r_{qi} r_{pj}}}{(r_i - r_j)}$$

(1-52)

$$= \frac{2 r_{ref} c^4}{hc}$$

$$[K_{qi}^4 (D_{sfu} D_{tfu})_{qs} (D_{scui} D_{tcui})_{aqs} V_{qi} (\sum_j^n K_{pj}^4 (D_{sfbj} D_{tfbj})_m (D_{scbj} D_{tcbj})_m V_{pj}] \left(\frac{\sqrt{r_{qi} r_{pj}}}{(r_i - r_j)} \right)$$

(2-52)

$$= \frac{2\sqrt{\alpha_r \alpha_g} r_{qi} r_{pj} c^4}{h} \frac{h c^2}{2c^3 V_{qi} r_{qi}} V_{qi} \sum_j^n \frac{h c^2}{2c V_{qi} r_{qi}} V_{pj} \left(\frac{1}{(r_i - r_j)} \right)$$

(3-52)

$$E_{di} = \sqrt{\alpha_r \alpha_g} \left(\frac{h}{2} \right) \sum_j^n \left(\frac{1}{(r_i - r_j)} \right)$$

(4-52)

for quanton and anti quanton pair (i) of space fabric the summation for 1 to n for *j* *quantons* of the stellar matter bound fields , this interaction has four degrees of freedom

52.b.Evidence of space fabric distortion : case of abnormal galagtic rotational curves

1-the contribution of the dark matter to the rotation curves of galaxies is increasing away from the galactic bulge

this is suggestive of a presence of a repulsive effect of normal

galactic mass near the bulge which causes

a-reduced space fabric quanton energy density near the bulge

(which leads to near Keplerian pattern of rotational velocities)

b-an increased space fabric quanton energy density away from the galactic bulge and consequently ,an increased effect of the gravitational like effect of dark matter and increased rotation curve speeds away from the galactic bulge


2- a localized drop in the rotational curve of spiral galaxies was observed , this localized drop coincides with the spiral arms of the spiral galaxies , an interpretation of such a phenomena can be put as follows , an accumulation of galactic mass in the spiral arms causes a distortion in the nearby region of the space fabric , and as a result of this distortion a drop in the gravitational like effect of the dark matter takes place , and thus causing this characteristic localized drop of rotational curves of spiral galaxies

examples : rotational curve of the milky way , localized bottoming coincides with and scutum –centaurus and orion - Cygnus arms ,

for other spiral galaxies : NGC 2590, NGC 1620 , NGC 7674 , NGC


7217, NGC 2998 , NGC 801 , they all show the same localized

rotation curve bottoming characteristic of spiral galaxies




Fig(11).Rotational speed of the milky display characteristic localized bottoming which coincides with spiral arms

Source <https://web.njit.edu/~gary/202/Lecture25.html>



Fig(12). Source : <http://hyperphysics.phy-astr.gsu.edu/hbase/Astro/darmat.html>



Copyright © Addison Wesley.

Fig (13). Source :

<http://ircamera.as.arizona.edu/Astr2016/lectures/darkmatter.htm>

rotational curves of various spiral galaxies display characteristic localized bottoming which coincides with spiral arms : an evidence of mass distortion space fabric and gravitational pattern

53.Electromagnetic field interactions

the rearranged degrees of freedom are

$$Dof_{sf} Dof_{tf} = 2.5 - 0.5 = 2.00 \quad , \quad (Dof_{sf} Dof_{tf})_{bound} = 1.00$$

$$(Dof_{sc} Dof_{tc}) = 1.5 - 0.5 = 1 \quad , \quad (Dof_{sf} Dof_{tf})_{unbound} = 1.00$$

53.a-Retaining energy interaction

While in its density form

$$\underline{E}_t = \epsilon_0 \mathbf{E} \mathbf{B}$$

$$E_t = \left[\frac{1}{2} \left(\left(\frac{E_{sfb} E_{tc}}{\sqrt{\epsilon_0} \sqrt{c}} \right) + \left(\frac{\sqrt{c} E_{sc} E_{tfb}}{\sqrt{\epsilon_0}} \right) \right) * \frac{1}{2} \left(\left(\frac{E_{sc} E_{tfb}}{\sqrt{\epsilon_0} \sqrt{c}} \right) + \left(\frac{E_{sf} E_{tcb}}{c \sqrt{\epsilon_0} \sqrt{c}} \right) \right) \right] \quad (1-53)$$

$$= \frac{1}{4} \left(\frac{(E_{sfb} E_{tc})^2}{c^2} + 2 \frac{(E_{sfb} E_{tc})(E_{sc} E_{tfb})}{c} + (E_{sc} E_{tfb})^2 \right)$$

And due to symmetry $E_{sfb} E_{tfb} = c E_{sc} E_{tc}$

$$E_t = \frac{(E_{sfb} E_{tc})^2}{c^2} = (E_{sc} E_{tfb})^2 = \frac{(E_{sc} E_{tc})(E_{sfb} E_{tfb})}{c}$$

And as a potential of the total photon energy

$$E_{tp} = \int_{V_q} (E_{sfb} E_{tfb}) (E_{sc} E_{tc}) \, dV \quad (2-53)$$

$$= [K_{qs}^2 (D_{sfb} D_{tfb})] [K_{qs}^2 (D_{sfb} D_{tfb})] V_q$$

$$E_{tp} = K_{qs}^4 c^2 V_q = \alpha_t \frac{hk^4}{16c\pi^4} V_q = \alpha_t \frac{h}{16 c r_q^4} V_q$$

$$E_{tp} = \alpha_t \frac{h}{2 c r_q} \quad (3-53)$$

Which has two degrees of freedom compared to the case of space

Fabric's retaining interaction which has 3 Dof's due to the

relativistic degree of freedom

53.b inflationary energy interaction , repulsive energy interaction

Same as space fabric

52.c-Gravitational like interaction of electromagnetic waves

the conversion of one degree of freedom into relativistic one

allows for a Dof match between the bound fields of the

electromagnetic waves and the bound fields of the normal matter ,

which was not possible for the case of space fabric

the binding interaction $E_{gbi} =$

$$= \frac{c^4}{E_{ref}} \left[\int_{V_{qe}} E_{sfbi} E_{tfbi} E_{sci} E_{tci} dV \right] \left(\sum_j^n \int_{V_p} \frac{E_{sfbj} E_{tfbj} E_{scbj} E_{tcbj}}{c^2} dV \right) \left(\frac{\sqrt{r_{qi} r_{pj}}}{(r_i - r_j)} \right) \quad (4-53)$$

$$= \frac{2c^4 r_{qi} r_{pj}}{hc} (K_{qi}^4 D_{sfb} D_{tfb} D_{sc} D_{tc} V_{qi}) \left(\sum_j^n K_{pj}^4 D_{scb} D_{tfb} D_{scb} D_{tcb} V_{pj} \right) \left(\frac{1}{(r_i - r_j)} \right)$$

$$= \frac{2c^3 r_q r_p}{h} (K_{qi}^4 c^2 V_{qi}) \left(\sum_j^n K_{pj}^4 c^2 V_{pj} \right) \left(\frac{1}{(r_i - r_j)} \right) \quad (4-53)$$

$$= \frac{2c^3 r_q r_p \sqrt{\alpha_b \alpha_g}}{h} \left(\frac{h c^2}{2\pi c^3 V_{qi} r_{qi}} \right) V_{qi} \sum_j^n \left(\frac{h c^2}{2\pi c V_{pj} r_{pj}} \right) V_{pj} \left(\frac{1}{(r_i - r_j)} \right)$$

$$E_{gbi} = \frac{h c \sqrt{\alpha_b \alpha_g}}{2} \sum_j^n \left(\frac{1}{(r_i - r_j)} \right) \quad (5-53)$$

noting that this gravitational like binding can exist between two different electromagnetic waves

2-now the parameter $K_g = \frac{c^4}{E_{ref}} = \frac{2\sqrt{r_{qi}r_{pj}}c^3}{h}$ (previously it was

defined as $K_g = \frac{2\pi r_p c^3}{h}$ for the case of normal mater gravitation)

53 d.dark matter distortion of electromagnetic waves

unbound fields of space fabric interact with electromagnetic waves , keeping in mind that those unbound fields can only create a repulsive interaction

$$E_{rei} = \frac{c^4}{E_{ref}}$$

$$[(\int_{V_{qe}} (E_{sfui}E_{tfui})_{qe} (E_{scui}E_{tcui})_{aqe} dV)(\sum_j^n \int_{V_{qs}} (E_{sfuj}E_{tfuj})_{qs} (E_{scuj}E_{tcuj})_{aq_s} dV)](\frac{\sqrt{r_{qi}r_{pj}}}{(r_i-r_j)}) \quad (6-53)$$

$$= \frac{2r_{qi}r_{pj}c^3}{h}$$

$$[K_{qi}^4 (E_{sfui}E_{tfui})_{q-e} (E_{scui}E_{tcui})_{aq-e} V_{qi}] [\sum_j^n K_{pj}^4 (E_{sfuj}E_{tfuj})_{q-s} (E_{scuj}E_{tcuj})_{aq-s} V_{qj}] (\frac{1}{(r_i-r_j)}) \quad (7-53)$$

$$= \frac{2r_{qi}r_{pj}c^3}{h} (K_{qi}^4 c^2 V_{qi}) (\sum_j^n K_{pj}^4 c^2 V_{qj}) (\frac{1}{(r_i-r_j)})$$

$$= \frac{2\alpha_r r_{qi} r_{pj} c^3}{h} \left(\frac{h}{2\pi c^3 V_{qi} r_{qi}} \right) c^2 V_{qi} \sum_j^n \left(\frac{h}{2\pi c^3 V_{qj} r_{qi}} \right) c^2 V_{qj} \left(\frac{1}{(r_i - r_j)} \right)$$

$$E_{gbi} = \frac{\alpha_r h c}{2} \sum_j^n \frac{1}{(r_i - r_j)} \quad (8-53)$$

This interaction which has four Dof's and between free and constrained unbound fields (i) of electromagnetic wave and unbound free and constrained fields of quanton and anti quanton pair (j)

interaction	free energy field	constrained energy field	Dof	Interaction type
For space fabric 1- E_t : quanton retaining 2- E_b : quanton binding	$E_{sfb}E_{tfb}$	$E_{scb}E_{tcb}$	three	multiple binding
For normal matter E_t : quanton retaining	$E_{sfb} E_{tfb}$	$E_{scb} E_{tcb}$	two	Single binding
E_i : quanton inflationary	$E_{sfu}E_{tfu}$	$E_{scu} E_{tcu}$	one	repulsive
E_r : quanton repulsive	$E_{sfu}E_{tfu}$	$E_{scu} E_{tcu}$	two	repulsive
1-gravitation binding 2- Gravitation	$E_{sfb} E_{tfb}$	$E_{scb} E_{tcb}$	four	Multiple binding
Electric force	$E_{sfu} E_{tfu}$	$E_{scu}E_{tcu}$	four	a-single binding or b-repulsive
Strong nuclear	$E_{sfu}E_{tfu}$	$E_{scu} E_{tcu}$	four	a-single binding b-repulsive
Dark matter gravitation like effect	$(E_{sfb}E_{tfb})_s$ $(E_{sfu}E_{tfu})_m$	$(E_{scb}E_{tcb})_s$ $(E_{scu}E_{tcu})_m$	three	Multiple binding
gravitation distortion of space fabric	$(E_{sfu} E_{tfu})_s$, $(E_{sfb})_m$,	$(E_{scu} E_{tcu})_s$, $(E_{scb})_m$,	four	repulsive
gravitation like binding of EM waves	$(E_{sfb} E_{tfb})_e$ $(E_{sfb} E_{tfb})_m$	$(E_{sc} E_{tc})_e$ $(E_{scb} E_{tcb})_m$	four	multiple binding
Dark matter distortion of electromagnetic waves	$(E_{sf} E_{tf})_e$ $(E_{sfu} E_{tfu})_s$	$(E_{sc} E_{tc})_e$ $(E_{scu} E_{tcu})_s$	four	repulsive

Table(11) Summary of interactions , their types and their energy field source

54.CPT symmetry at the Quanton scale

CPT symmetry has its origins at the quanton scale ,as it reflects symmetries created due to energy constraining ,as the degrees of freedom of anti quanton free and constrained fields are mirror symmetric to those of the quanton's

	Quanton	Anti quanton
Nature of dominant energy	free	constrained
Main-space-varying	E_{sf}	E_{sc}
Auxiliary-space varying	E_{sc}	E_{sf}
Main time varying	E_{tf}	E_{tc}
Auxiliary time varying	E_{tc}	E_{tf}

Table(12) Mirror symmetry between quanton and anti quanton

Dominant energy	free	constrained
time	Positive configuration for the time varying fields	negative configuration for the time varying fields
Spatial configuration	Positive position vector configuration for the space varying fields	negative position vector configuration for the space varying fields
charge	positive atomic fields and charges due to unbound fields $E_{sfu}E_{tfu}$	negative atomic fields and charges due to unbound fields $E_{scu}E_{tcu}$

Table(13) CPT symmetry and its link to quanton / anti quanton mirror symmetry

55. Conclusions

Uniformity and homogeneity of CMB testifies to its origin which is the release of radiation from the space fabric as a direct result of the process of free expansion of the universe (second law of Thermodynamics) , this gives a gate way for further understanding of the quanton interactions .

56. References

Basic physics .

