

A PROOF OF TWIN PRIME CONJECTURE

T. AGAMA

ABSTRACT. In this paper we prove the twin prime conjecture by showing that

$$\sum_{n \leq x} \Lambda(n)\Lambda(n+2) \geq (1 + o(1)) \frac{x}{2\mathcal{C}(2)}$$

for some $\mathcal{C} := \mathcal{C}(2) > 0$. We start by developing a general method for estimating correlations of the form

$$\sum_{n \leq x} G(n)G(n+l)$$

for a fixed $1 \leq l \leq x$ and where $G : \mathbb{N} \rightarrow \mathbb{R}^+$.

1. Introduction and statement

Consider the sum

$$\sum_{n \leq x} G(n)G(x-n)$$

and

$$\sum_{n \leq x} G(n)G(n+l)$$

where $1 \leq l \leq x$. It is generally not easy to control sums of these forms, and unfortunately many of the open problems in number theory can be phrased in this manner. The twin prime conjecture conjectured by De polignac, which is one of the important open problems can be expressed in this form as

$$\sum_{n \leq x} \Lambda(n)\Lambda(n+2)$$

and it is the case that obtaining a non-trivial lower bound for this correlation solves the twin prime conjecture. There are a good number of techniques in the literature for studying such sums, like the circle method of Hardy and littlewood, the sieve method and many others.

In this paper, we introduce the area method. This method can also be used to control correlated sums of the form above. The novelty of this method is that it allows us to write any of these correlated sums as a double sum, which is much easier to estimate using existing tools such as the summation by part formula. As an application we obtain the result:

Date: October 7, 2019.

2000 Mathematics Subject Classification. Primary 54C40, 14E20; Secondary 46E25, 20C20.

Theorem 1.1. *There exist some constant $\mathcal{C} := \mathcal{C}(2) > 0$, such that*

$$\sum_{n \leq x} \Lambda(n)\Lambda(n+2) \geq (1 + o(1)) \frac{x}{2\mathcal{C}(2)}.$$

2. The area method

In this section we introduce and develop a fundamental method for solving problems related to correlations of arithmetic functions. This method is fundamental in the sense that it uses the properties of four main geometric shapes, namely the triangle, the trapezium, the rectangle and the square. The basic identity we will derive is an outgrowth of exploiting the areas of these shapes and putting them together in a unified manner.

Theorem 2.1. *Let $\{r_j\}_{j=1}^n$ and $\{h_j\}_{j=1}^n$ be any sequence of real numbers, and let r and h be any real numbers satisfying $\sum_{j=1}^n r_j = r$ and $\sum_{j=1}^n h_j = h$, and*

$$(r^2 + h^2)^{1/2} = \sum_{j=1}^n (r_j^2 + h_j^2)^{1/2},$$

then

$$\sum_{j=2}^n r_j h_j = \sum_{j=2}^n h_j \left(\sum_{i=1}^j r_i + \sum_{i=1}^{j-1} r_i \right) - 2 \sum_{j=1}^{n-1} r_j \sum_{k=1}^{n-j} h_{j+k}.$$

Proof. Consider a right angled triangle, say $\triangle ABC$ in a plane, with height h and base r . Next, let us partition the height of the triangle into n parts, not necessarily equal. Now, we link those partitions along the height to the hypotenuse, with the aid of a parallel line. At the point of contact of each line to the hypotenuse, we drop down a vertical line to the next line connecting the last point of the previous partition, thereby forming another right-angled triangle, say $\triangle A_1 B_1 C_1$ with base and height r_1 and h_1 respectively. We remark that this triangle is covered by the triangle $\triangle ABC$, with hypotenuse constituting a proportion of the hypotenuse of triangle $\triangle ABC$. We continue this process until we obtain n right-angled triangles $\triangle A_j B_j C_j$, each with base and height r_j and h_j for $j = 1, 2, \dots, n$. This construction satisfies

$$h = \sum_{j=1}^n h_j \text{ and } r = \sum_{j=1}^n r_j$$

and

$$(r^2 + h^2)^{1/2} = \sum_{j=1}^n (r_j^2 + h_j^2)^{1/2}.$$

Now, let us deform the original triangle $\triangle ABC$ by removing the smaller triangles $\triangle A_j B_j C_j$ for $j = 1, 2, \dots, n$. Essentially we are left with rectangles and squares piled on each other with each end poking out a bit further than the one just above,

and we observe that the total area of this portrait is given by the relation

$$\begin{aligned}\mathcal{A}_1 &= r_1 h_2 + (r_1 + r_2) h_3 + \cdots + (r_1 + r_2 + \cdots + r_{n-2}) h_{n-1} + (r_1 + r_2 + \cdots + r_{n-1}) h_n \\ &= r_1 (h_2 + h_3 + \cdots + h_n) + r_2 (h_3 + h_4 + \cdots + h_n) + \cdots + r_{n-2} (h_{n-1} + h_n) + r_{n-1} h_n \\ &= \sum_{j=1}^{n-1} r_j \sum_{k=1}^{n-j} h_{j+k}.\end{aligned}$$

On the other hand, we observe that the area of this portrait is the same as the difference of the area of triangle ΔABC and the sum of the areas of triangles $\Delta A_j B_j C_j$ for $j = 1, 2, \dots, n$. That is

$$\mathcal{A}_1 = \frac{1}{2} r h - \frac{1}{2} \sum_{j=1}^n r_j h_j.$$

This completes the first part of the argument. For the second part, along the hypotenuse, let us construct small pieces of triangle, each of base and height (r_i, h_i) ($i = 1, 2, \dots, n$) so that the trapezoid and the one triangle formed by partitioning becomes rectangles and squares. We observe also that this construction satisfies the relation

$$(r^2 + h^2)^{1/2} = \sum_{i=1}^n (r_i^2 + h_i^2)^{1/2},$$

Now, we compute the area of the triangle in two different ways. By direct strategy, we have that the area of the triangle, denoted \mathcal{A} , is given by

$$\mathcal{A} = 1/2 \left(\sum_{i=1}^n r_i \right) \left(\sum_{i=1}^n h_i \right).$$

On the other hand, we compute the area of the triangle by computing the area of each trapezium and the one remaining triangle and sum them together. That is,

$$\mathcal{A} = h_n/2 \left(\sum_{i=1}^n r_i + \sum_{i=1}^{n-1} r_i \right) + h_{n-1}/2 \left(\sum_{i=1}^{n-1} r_i + \sum_{i=1}^{n-2} r_i \right) + \cdots + 1/2 r_1 h_1.$$

By comparing the area of the second argument, and linking this to the first argument, the result follows immediately. \square

Remark 2.2. Next we state a result for a general lower bound for any two-point correlation that captures all real arithmetic function.

Theorem 2.3. *Let $f : \mathbb{N} \rightarrow \mathbb{R}^+$, a real-valued function. If*

$$\sum_{n \leq x} f(n) f(n + l_0) > 0$$

then there exist some constant $\mathcal{C} := \mathcal{C}(l_0) > 0$ such that

$$\sum_{n \leq x} f(n) f(n + l_0) \geq \frac{1}{\mathcal{C}(l_0) x} \sum_{2 \leq n \leq x} f(n) \sum_{m \leq n-1} f(m).$$

Proof. By Theorem 2.1, we obtain the identity by taking $f(j) = r_j = h_j$

$$\sum_{n \leq x-1} \sum_{j \leq x-n} f(n) f(n+j) = \sum_{2 \leq n \leq x} f(n) \sum_{m \leq n-1} f(m).$$

It follows that

$$\begin{aligned}
\sum_{n \leq x-1} \sum_{j \leq x-n} f(n)f(n+j) &\leq \sum_{n \leq x-1} \sum_{j < x} f(n)f(n+j) \\
&= \sum_{n \leq x} f(n)f(n+1) + \sum_{n \leq x} f(n)f(n+2) \\
&\quad + \cdots + \sum_{n \leq x} f(n)f(n+l_0) + \cdots + \sum_{n \leq x} f(n)f(n+x) \\
&\leq |\mathcal{M}(l_0)| \sum_{n \leq x} f(n)f(n+l_0) \\
&\quad + |\mathcal{N}(l_0)| \sum_{n \leq x} f(n)f(n+l_0) \\
&\quad + \cdots + \sum_{n \leq x} f(n)f(n+l_0) + \cdots + |\mathcal{R}(l_0)| \sum_{n \leq x} f(n)f(n+l_0) \\
&= \left(|\mathcal{M}(l_0)| + |\mathcal{N}(l_0)| + \cdots + 1 \right. \\
&\quad \left. + \cdots + |\mathcal{R}(l_0)| \right) \sum_{n \leq x} f(n)f(n+l_0) \\
&\leq \mathcal{C}(l_0)x \sum_{n < x} f(n)f(n+l_0).
\end{aligned}$$

where $\max\{|\mathcal{M}(l_0)|, |\mathcal{N}(l_0)|, \dots, |\mathcal{R}(l_0)|\} = \mathcal{C}(l_0)$. By inverting this inequality, the result follows immediately. \square

The nature of the implicit constant $\mathcal{C}(l_0)$ could also depend on the structure of the function we are being given. The von mangoldt function, contrary to many class of arithmetic functions, has a relatively small such constant. This behaviour stems from the fact that the Von-mangoldt function is defined on the prime powers. Thus one would expect most terms of sums of the form

$$\sum_{n \leq x-1} \sum_{j \leq x-n} \Lambda(n)\Lambda(n+j)$$

to fall off when j is odd for any prime power $n = p^k$ such that $j + p^k \neq 2^s$.

3. Proof of the twin prime conjecture

We are now ready to prove the twin prime conjecture. We assemble the tools we have developed thus far to solve the problem.

Theorem 3.1. *There exist some constant $\mathcal{C} := \mathcal{C}(2) > 0$, such that*

$$\sum_{n \leq x} \Lambda(n)\Lambda(n+2) \geq (1 + o(1)) \frac{x}{2\mathcal{C}(2)}.$$

Proof. By invoking Theorem 2.3, we can write

$$\sum_{n \leq x} \Lambda(n)\Lambda(n+2) \geq \frac{1}{\mathcal{C}(2)x} \sum_{2 \leq n \leq x} \Lambda(n) \sum_{m \leq n-1} \Lambda(m).$$

Using the prime number theorem [1] of the form

$$\sum_{n \leq x} \Lambda(n) = (1 + o(1))x,$$

the result follows immediately by using partial summation. \square

Remark 3.2. It is important to remark that with the lower bound in Theorem 3.1, we have solved the twin prime conjecture. This method not only does it solve the twin prime conjecture, but is good in terms of its generality, for it can be used to obtain lower bounds for a general class of correlated sums of the form

$$\sum_{n \leq x} f(n)f(n+k)$$

for a uniform $1 \leq k \leq x$.

4. Conclusion

The method adopted in this paper to prove the twin prime conjecture is simple and very elegant. In the spirit of solving the Goldbach conjecture, this method can also be exploited to develop an estimate for general sums of the form

$$\sum_{n \leq x} G(n)G(x-n)$$

which we do not pursue in this paper. ¹.

REFERENCES

1. Montgomery, H.L, and Vaughan, R.C, *Multiplicative number theory 1:Classical theory*. vol.97, Cambridge university press, 2006.

DEPARTMENT OF MATHEMATICS, AFRICAN INSTITUTE FOR MATHEMATICAL SCIENCE, GHANA
E-mail address: theophilus@aims.edu.gh/emperordagama@yahoo.com