

I think that it is possible the quantization of the gravity using the quantum field theory (I use the Lewis Ryder book) and the classical theory of fields (I use the Landau-Lifshits book).

It is evident, from the Lifshits book, that the Euler-Lagrange equation for the gravitational field is the Einstein field equation obtained from the Lagrangian of the gravitational field (I write the vacuum solution):

$$G = \mathcal{L} = g^{ik} (\Gamma_{il}^m \Gamma_{km}^l - \Gamma_{ik}^l \Gamma_{lm}^m)$$

the generalized variables, used in the Euler-Lagrange equation are:

$$\mathcal{L} \left(g^{ik}, \frac{\partial g^{lm}}{\partial x^r} \right)$$

as is evident from the variation of the action:

$$\delta S_g = -\frac{c^3}{16\pi k} \int \left\{ \frac{\partial(G\sqrt{-g})}{\partial g^{ik}} - \frac{\partial}{\partial x^l} \frac{\partial(G\sqrt{-g})}{\partial \frac{\partial g^{ik}}{\partial x^l}} \right\} \delta g^{ik} d\Omega$$

so that it is simple to obtain the Hamiltonian from this action if the Christoffel symbols are expressed in the variables subject to variation (how I got it):

$$\Gamma_{kl}^i = -\frac{1}{2} \left(g_{pk} \frac{\partial g^{ip}}{\partial x^l} + g_{lp} \frac{\partial g^{ip}}{\partial x^k} - g^{ip} g_{kw} g_{lz} \frac{\partial g^{wz}}{\partial x^p} \right)$$

then, the Hamiltonian is

$$\mathcal{H} = \frac{\partial g^{ij}}{\partial x^0} \frac{\partial \mathcal{L}}{\partial \frac{\partial g^{ij}}{\partial x^0}} - \mathcal{L}$$

the Lagrangian is a quadratic function in the derivative of the metric tensor, so that the Hamiltonian is just a quadratic function of the metric tensor; there are no linear terms of the time derivative of the metric tensor:

$$\mathcal{H} = \frac{1}{2} \frac{\partial \mathcal{L}}{\partial \frac{\partial g^{ik}}{\partial x^0} \partial \frac{\partial g^{ik}}{\partial x^0}} (\partial_0 g^{ik})^2 - \frac{1}{2} \frac{\partial \mathcal{L}}{\partial \frac{\partial g^{ij}}{\partial x^\mu} \partial \frac{\partial g^{lm}}{\partial x^\nu}} \partial_\mu g^{ik} \partial_\nu g^{lm}$$

where $\mu \in 1, 2, 3$ and so for each greek letter.

It is evident that the classical quantization is possible, if it is possible to write the time derivative of the metric tensor like a function of the generalized moment (I don't use the Heisenberg picture):

$$\begin{aligned} \hat{H} &= \int \mathcal{H} d^3x \\ i\hbar \frac{\partial \Psi(g^{ij}, \partial_\mu g^{kl})}{\partial t} &= H\Psi(g^{ij}, \partial_\mu g^{kl}) \\ \hat{\pi}_{rs}\Psi(g^{ik}, \partial_\mu g^{lm}) &= -i\hbar \frac{\partial}{\partial \partial_0 g^{rs}} \Psi(g^{ik}, \partial_\mu g^{lm}) \\ \hat{g}^{rs}\Psi(g^{ik}, \partial_\mu g^{lm}) &= g^{rs}\Psi(g^{ik}, \partial_\mu g^{lm}) \end{aligned}$$

I write the Hamiltonian, and the momentum of the Lagrangian of the vacuum starting from the second derivative of the Lagrangian:

$$\begin{aligned} \frac{\partial^2 \mathcal{L}}{\partial g^{ab} \partial g^{cd}} &= g^{ik} \left(\frac{\partial \Gamma_{il}^m}{\partial g^{ab}} \frac{\partial \Gamma_{km}^l}{\partial g^{cd}} + \frac{\partial \Gamma_{il}^m}{\partial g^{cd}} \frac{\partial \Gamma_{km}^l}{\partial g^{ab}} - \frac{\partial \Gamma_{ik}^l}{\partial g^{ab}} \frac{\partial \Gamma_{lm}^m}{\partial g^{cd}} - \frac{\partial \Gamma_{ik}^l}{\partial g^{cd}} \frac{\partial \Gamma_{lm}^m}{\partial g^{ab}} \right) = \\ &= \delta_a^s \delta_c^r g_{bd} + \delta_a^s \delta_d^r g_{bc} + \delta_b^s \delta_c^r g_{ad} + \delta_b^s \delta_d^r g_{ac} - g_{ac} g_{bd} g^{rs} - g_{ad} g_{bc} g^{rs} - \delta_a^s \delta_b^r g_{cd} - \delta_a^r \delta_b^s g_{cd} - \delta_c^r \delta_d^s g_{ab} - \delta_c^r \delta_d^s g_{ab} + 2g_{ab} g_{cd} g^{rs} \end{aligned}$$

the calculations are simpler if a new terms Υ_0^a is introduced to evaluate the remaining components of the metric tensor:

$$\begin{aligned}\Lambda_{a0} &= (\delta_a^0 g_{c0} + g_{ac}) \partial_0 g^{c0} - (\delta_a^0 g_{cd} + g^{00} g_{ac} g_{d0}) \partial_0 g^{cd} = (\delta_a^0 g_{00} + g_{a0}) \partial_0 g^{00} + (\delta_a^0 g_{\alpha0} + g_{a\alpha}) \partial_0 g^{\alpha0} - (\delta_a^0 g_{00} + g^{00} g_{a0} g_{00}) \partial_0 g^{00} + \\ &- (\delta_a^0 g_{\alpha0} + g^{00} g_{a\alpha} g_{00}) \partial_0 g^{\alpha0} - (\delta_a^0 g_{\alpha0} + g^{00} g_{a0} g_{\alpha0}) \partial_0 g^{\alpha0} - (\delta_a^0 g_{\alpha\beta} + g^{00} g_{a\alpha} g_{\beta0}) \partial_0 g^{\alpha\beta} = g_{a0}(1 - g^{00} g_{00}) \partial_0 g^{00} + \\ &+ (g_{a\alpha} - \delta_a^0 g_{\alpha0} - g^{00} g_{a\alpha} g_{00} - g^{00} g_{a0} g_{\alpha0}) \partial_0 g^{\alpha0} - (\delta_a^0 g_{\alpha\beta} + g^{00} g_{a\alpha} g_{\beta0}) \partial_0 g^{\alpha\beta} \\ \Upsilon_{a0} &= \Lambda_{a0} + (\delta_a^0 g_{\alpha\beta} + g^{00} g_{a\alpha} g_{\beta0}) \partial_0 g^{\alpha\beta} = \Lambda_{a0} - \left(\frac{\delta_a^0 g_{\alpha\beta}}{g^{00}} + g_{a\alpha} g_{\beta0} \right) g^{\alpha\mu} g^{\beta\nu} \Lambda_{\mu\nu} = \\ &= g_{a0}(1 - g^{00} g_{00}) \partial_0 g^{00} + (g_{a\alpha} - \delta_a^0 g_{\alpha0} - g^{00} g_{00} g_{a\alpha} - g^{00} g_{a0} g_{\alpha0}) \partial_0 g^{\alpha0} \\ \Upsilon_{\kappa0} &= \Lambda_{\kappa0} - g_{\kappa\alpha} g_{\beta0} g^{\alpha\mu} g^{\beta\nu} \Lambda_{\mu\nu} = g_{\kappa0}(1 - g^{00} g_{00}) \partial_0 g^{00} + [g_{\kappa\alpha}(1 - g^{00} g_{00}) - g^{00} g_{\kappa0} g_{\alpha0}] \partial_0 g^{\alpha0} \\ \Upsilon_{00} &= \Lambda_{00} - \left(\frac{g_{\alpha\beta}}{g^{00}} + g_{a\alpha} g_{\beta0} \right) g^{\alpha\mu} g^{\beta\nu} \Lambda_{\mu\nu} = g_{00}(1 - g^{00} g_{00}) \partial_0 g^{00} - 2g^{00} g_{00} g_{\alpha0} \partial_0 g^{\alpha0}\end{aligned}$$

then it is possible to obtain the $\partial_0 g^{a0}$ as a $\Upsilon(\Lambda(\Theta(\pi^{lm})))$ function:

$$\begin{pmatrix} \Upsilon_{10} - \frac{g_{10}}{g_{00}} \Upsilon_{00} \\ \Upsilon_{20} - \frac{g_{20}}{g_{00}} \Upsilon_{00} \\ \Upsilon_{30} - \frac{g_{30}}{g_{00}} \Upsilon_{00} \end{pmatrix} = \begin{pmatrix} g_{11}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{10} & g_{12}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{20} & g_{13}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{30} \\ g_{21}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{10} & g_{22}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{20} & g_{23}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{30} \\ g_{31}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{10} & g_{32}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{20} & g_{33}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{30} \end{pmatrix} \begin{pmatrix} \partial_0 g^{10} \\ \partial_0 g^{20} \\ \partial_0 g^{30} \end{pmatrix}$$

the determinant of the linear system is in general not null:

$$\begin{vmatrix} g_{11}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{10} & g_{12}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{20} & g_{13}(1 - g^{00} g_{00}) + g^{00} g_{10} g_{30} \\ g_{21}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{10} & g_{22}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{20} & g_{23}(1 - g^{00} g_{00}) + g^{00} g_{20} g_{30} \\ g_{31}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{10} & g_{32}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{20} & g_{33}(1 - g^{00} g_{00}) + g^{00} g_{30} g_{30} \end{vmatrix} = (1 - g^{00} g_{00})^2 (\gamma - g^{00} g) \geq 0$$

where $g = |g_{ab}|$ and $\gamma = |g_{\alpha\beta}|$, then it is possible to obtain the transformation between time derivatives of the metric tensor and generalized moments $\partial_0 g^{ab} = P^{ab} + Q^{ab,cd} \pi_{cd}$.

$$\begin{aligned}\partial_0 g^{\kappa0} &= \frac{\sum_{\alpha_1 \alpha_2 \alpha_3 \beta_1 \beta_2} \epsilon_{\kappa \beta_1 \beta_2} \epsilon_{\alpha_1 \alpha_2 \alpha_3} [g_{\alpha_1 \beta_1}(1 - g^{00} g_{00}) + g^{00} g_{\alpha_1 0} g_{\beta_1 0}] [g_{\alpha_2 \beta_2}(1 - g^{00} g_{00}) + g^{00} g_{\alpha_2 0} g_{\beta_2 0}] [\Upsilon_{\alpha_3 0} - \frac{g_{\alpha_3 0}}{g_{00}} \Upsilon_{00}]}{2(1 - g^{00} g_{00})^2 (\gamma - g^{00} g)} = \\ &= \frac{\sum_{\alpha_1 \alpha_2 \alpha_3 \beta_1 \beta_2} \epsilon_{\kappa \beta_1 \beta_2} \epsilon_{\alpha_1 \alpha_2 \alpha_3} [g_{\alpha_1 \beta_1} g_{\alpha_2 \beta_2}(1 - g^{00} g_{00})^2 + 2g^{00} g_{\alpha_1 0} g_{\beta_1 0} g_{\alpha_2 0} g_{\beta_2 0}] [\Upsilon_{\alpha_3 0} - \frac{g_{\alpha_3 0}}{g_{00}} \Upsilon_{00}]}{2(1 - g^{00} g_{00})^2 (\gamma - g^{00} g)} = \\ &= \frac{\sum_{\alpha_1 \alpha_2 \alpha_3 \beta_1 \beta_2} \epsilon_{\kappa \beta_1 \beta_2} \epsilon_{\alpha_1 \alpha_2 \alpha_3} g_{\alpha_2 \beta_2} [g_{\alpha_1 \beta_1}(1 - g^{00} g_{00}) + 2g^{00} g_{\alpha_1 0} g_{\beta_1 0}] [\Upsilon_{\alpha_3 0} - \frac{g_{\alpha_3 0}}{g_{00}} \Upsilon_{00}]}{2(1 - g^{00} g_{00})(\gamma - g^{00} g)}\end{aligned}$$

It is simple to obtain the Hamiltonian like a π_{ab} function:

$$\begin{aligned}\mathcal{H} &= 2g^{00}(g_{\alpha0} g_{\alpha0} - g_{\alpha\alpha} g_{00}) \partial_0 g^{\alpha0} \partial_0 g^{00} + 2g^{00}(g_{\alpha\beta} g_{\alpha\beta} - g_{\alpha\alpha} g_{\beta\beta}) \partial_0 g^{\alpha\beta} \partial_0 g^{\beta\alpha} + \\ &- 2g_{cd} \partial_\beta g^{ad} \partial_\alpha g^{\beta c} + 2g_{cd} \partial_\beta g^{cd} \partial_\alpha g^{\alpha\beta} + g_{ac} g_{bd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd} - g_{ab} g_{cd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd} = \\ &= 2g^{00}(g_{\alpha0} g_{\alpha0} - g_{\alpha\alpha} g_{00}) (P^{\alpha0} + Q^{\alpha0,cd} \pi_{cd})^2 + \\ &= 2g^{00}(g_{\alpha\beta} g_{\alpha\beta} - g_{\alpha\alpha} g_{\beta\beta}) (P^{\alpha\beta} + Q^{\alpha\beta,cd} \pi_{cd})^2 + \\ &- 2g_{cd} \partial_\beta g^{ad} \partial_\alpha g^{\beta c} + 2g_{cd} \partial_\beta g^{cd} \partial_\alpha g^{\alpha\beta} + g_{ac} g_{bd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd} - g_{ab} g_{cd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd} \\ \hat{\mathcal{H}} &= 2g^{00}(g_{\alpha0} g_{\alpha0} - g_{\alpha\alpha} g_{00}) (P^{\alpha0} P^{\alpha0} - 2i\hbar P^{\alpha0} Q^{\alpha0,cd} \frac{\partial}{\partial g^{cd}} - \hbar^2 Q^{\alpha0,ab} Q^{\alpha0,cd} \frac{\partial^2}{\partial g^{ab} \partial g^{cd}}) + \\ &2g^{00}(g_{\alpha\beta} g_{\alpha\beta} - g_{\alpha\alpha} g_{\beta\beta}) (P^{\alpha\beta} P^{\alpha\beta} - 2i\hbar P^{\alpha\beta} Q^{\alpha\beta,cd} \frac{\partial}{\partial g^{cd}} - \hbar^2 Q^{\alpha\beta,ab} Q^{\alpha\beta,cd} \frac{\partial^2}{\partial g^{ab} \partial g^{cd}}) + \\ &- 2g_{cd} \partial_\beta g^{ad} \partial_\alpha g^{\beta c} + 2g_{cd} \partial_\beta g^{cd} \partial_\alpha g^{\alpha\beta} + g_{ac} g_{bd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd} - g_{ab} g_{cd} g^{\alpha\beta} \partial_\alpha g^{ab} \partial_\beta g^{cd}\end{aligned}$$

this is the Hamiltonian of the gravitational field, that it is quantizable (even if not uniquely), and it is similar to the Wheeler-De Witt equation.