

The electromagnetic origin of the 3 GeV peak in the flux profiles of Cosmic Rays protons.

Oswaldo F. Schilling

Departamento de Física, UFSC, 88040-900, Florianópolis, SC. Brazil

Abstract:

In two recent publications we developed a field theoretical model in which the origin of baryons is associated with the stabilization of charge vortices from vacuum. According to this model the vortices should condense from a state 2.7 GeV higher in energy than the proton rest energy. In support of such a picture, all number- flux and energy-flux profiles taken from protons cosmic rays (CR) display a peak in the range between 2.5 and 4 GeV(the form of this peak may suffer the influence on particles of the Sun magnetic field). In the present work we demonstrate through a simple calculation that the observed peak at ~ 3 GeV would correspond to the “elastic” limit of a vortex-proton structure. Any excess energy should then be released through radiation to avoid “melting” of the vortices, so that particles mostly concentrate at values of energy lower than 3 GeV.

Introduction.

In recent years we have worked extensively on heuristic models for baryons, which resulted in three publications [1-3]. In the two most recent publications[1,2] we developed a fieldtheoretical model which supplemented the essentially phenomenological treatment initially presented [3]. The model in [3] actually extends similar work of Barut[4] and Post[5] on the subject, reobtaining Barut's main final results which relate mass of baryons to the inverse of the alpha-constant [4]. In particular, Barut's n (a number of action quanta from the Bohr-Sommerfeld conditions) reappears in our treatment as a number of magnetic flux-quanta [1-3].

Perhaps the most relevant (and surprising!) result of this work has been the deduction that baryons should be the result of a process of stabilization(or condensation) of vortices of charge, generated from a parent state located at 3.7 GeV[1]. Such state might be associated with the EM vacuum and the baryons would come as a result of instabilities of EM origin taking place in that vacuum. Support for such an explanation comes from the energy distribution of protons and other particles in cosmic rays (CR)[6], which peak in the range 2.5-4 GeV, indicating a heavy concentration of particles below the peak energy. The purpose of the present paper is to supplement the brief analysis of this experimental observation advanced in [2]. As we show below the association of the peak position with the limiting "elastic" strength of a vortex of charge is rather straightforward, as well as the demonstration that at the 3.7 GeV level some form of melting might indeed might happen as energy increases.

Strained charge vortices.

At some point in their history particles in CR were subject to extreme electric and magnetic forces, and the measured flux distributions we detect now were established under the influence of such forces. Let's concentrate on the case of protons, which represent by far the majority of the contents of CR. Protons are composed of entangled constituents of different charge values, and probably different topological properties

associated with these charge values[5]. Under extreme electromagnetic forces, the proton structure is subject to huge internal stresses to avoid breakdown. We now argue that the observed peak at 3GeV energy in the spectra of CR directly probes the proton structure elastic response to such forces.

It is well known that a spring-mass oscillating system of mass m and elastic spring constant k will spontaneously oscillate at its fundamental frequency $\omega = (k/m)^{1/2}$ when subject to a constant force F along the spring- mass direction. Let's assume that the peak kinetic energy of 3 GeV corresponds to the ground state energy of a 3DIM quantum harmonic oscillator, so that $3/2\hbar\omega \approx 3 \text{ GeV}$.

With m the proton mass, one might immediately estimate the elastic constant k and natural frequency of this vortex structure, $\hbar\omega = \hbar (k/m)^{1/2} = 2 \text{ GeV} = 3.2 \times 10^{-10} \text{ J}$. We obtain the natural frequency $\omega = 3.1 \times 10^{24} \text{ rad/s}$ (we recognize that since the proton is a multi-component system other normal modes of vibration theoretically exist, but we will concentrate on the most conspicuous one, that at about 3 GeV).

The intense forces mentioned in the previous paragraphs would excite the vibration mode of frequency ω , and the internal constituents of the proton would acquire additional oscillating displacements as a result. The maximum amplitude δ related to such additional oscillating motion is given by the maximum elastic energy expression (m is the proton mass) :

$$3/2 m\omega^2\delta^2 = 3/2 \hbar\omega \quad (1)$$

One immediately obtains $\delta = 1.4 \times 10^{-16} = 0.14 \text{ fm}$. This is the magnitude of the vibrational displacements(here considered elastic) of the proton structure when excited 3 GeV above the rest state. It remains to be discussed whether the elastic regime is still applicable. The value for δ is indeed deeply revealing as far as the process taking place at 3.7 GeV is concerned, as we discuss below.

Analysis and Conclusions.

As argued in ref. [2], protons of energy around 3 GeV in theory reach their stability limits and lose the energy advantage over the parent state at 3.7 GeV. One might speculate that the proton would then “melt” into its parent phase.

In the previous section we have worked out the parameter δ , which will be used to put such rather abstract concept of melting into a quantitative form. A practical way of doing this is through the Lindemann criterion (LC) for the melting of a solid.

According to the LC a solid loses cohesion towards a disordered (“molten”) phase when atomic oscillations reach about 5 to 10% of the interatomic distance. Here we must replace the term interatomic by “inter-constituent” spacings. One might immediately evaluate if this is the case with the protons in CR. The size of a proton is estimated from about 0.6 (from the calculated profile of charge distribution) up to 0.9 fm. The inter-constituent spacing cannot be greater than the size of the particle itself. It is possible to conclude then that the amplitude $\delta = 0.14$ fm is about 15 to 23 % of the experimental inter-constituent spacing, and 30% of the calculated [1] vortex size L of 0.46 fm. The Lindemann criterion for melting(if applicable in its usual form here) is therefore exceeded at such high kinetic energies.

It seems clear that the peak exists exactly because it indicates structural instability at about 3 GeV kinetic energies. Is there any actual indication that protons disappear into vacuum or break apart? There exists ample evidence that protons profusely radiate electromagnetic energy as their speed increases in accelerators, so that energy is released before breakdown(decay!). The result is the observed concentration of CR protons up to and below about 3 GeV energies. Energy is released until strains come back to the elastic regime and the structure is stable. Such threshold can be regarded also as a fingerprint of the importance of electromagnetic forces in the formation and in the inner constitution of complex particles like the proton[see 1].

The same explanation might be attempted to the electron case, in which inner topological features [5] might play the role of the baryon constituents in the analysis.

We wish to thank Dr Indranu Suhendro for his interest on this research.

References

1. O F Schilling, Progress in Physics (U. of New Mexico) 15,185 (2019).
2. _____ 15, 197 (2019).
3. O F Schilling, Annales Louis de Broglie 43-1, 1 (2018).
4. A.O.Barut, Appl. Phys (Lasers and Optics) **B60**, 123 (1995).
5. E.J.Post, Phys. Lett. 119**A**, 47 (1986).
6. T.K.Gaisser et al., Cosmic Rays and Particle Physics, Cambridge (2016).