

# The Illusion of Quantum Mechanical Probability Waves

Wim Vegt

Abstract

In the New Theory, it will be demonstrated that because of a mistake in the Maxwell Equations, Confined Electromagnetic waves could not be considered to be the material waves expressed in Schrödinger's wave equation. The New Theory presents a new equation describing electromagnetic field configurations which are also solutions of the Schrödinger's wave equation and the relativistic quantum mechanical Dirac Equation.

## 1.1 The Creation of Probability

The physical concept of quantum mechanical probability waves has been created during the famous [1927 5<sup>th</sup> Solvay Conference](#). During that period there were several circumstances which came just together and made it possible to create an unique idea of material waves being complex (partly real and partly imaginary) and describing the probability of the appearance of a physical object (elementary particle).

The idea of complex probability waves was completely new in the beginning of the 20<sup>th</sup> century and it is hard to believe that this strange non- scientific concept of probability waves has been created at a famous scientific Conference like the [1927 5<sup>th</sup> Solvay Conference](#). Since then the New Concept has been protected carefully within the [Copenhagen Interpretation](#).

When Schrödinger published his famous material wave equation in 1926, he found spherical and elliptical solutions for the presence of the electron within the atom. With that outcome Bohr's model of the atom completely fell apart. Because in Bohr's model of the atom an electron as a particle can only exist at one place at one time. But according to the solutions of the wave equations, for a spherical solution the electron is everywhere at the same time dived equally along a sphere.

And now lines in history come together. The first idea of the material waves in Schrödinger's wave equation was the concept of confined Electromagnetic Waves. But according to Maxwell this was impossible. According to Maxwell's equations Light (Electromagnetic Waves) can only propagate along straight lines and it is impossible that Light

(Electromagnetic Waves) could confine with the surface of a sphere or an ellipse.

For that reason these material waves in Schrödinger's wave equation could only be of a different origin than Electromagnetic Waves. And that conclusion opened for Niels Bohr the opportunity to save his model of the atom.

Niels Bohr introduced the unusual concept of "Probability Waves" as the origin of the material waves in Schrödinger's wave equation. And defined the New Concept that the electron was still a particle but the physical presence of the electron in the Atom was equally divided by a spherical probability function. And like Maxwell also Niels Bohr chose for the problem-solving approach.

Niels Bohr solved two problems at the same time. He found an answer for the origin of the material waves in Schrödinger's wave equation and he could keep his model of the atom.

## **1.2 Back to the roots**

To change the nowadays popular concept of "Problem Solving Physics" into "Fundamental Physics" we have to go back in time for over 300 years. Back to the time when the vast areas of Science, Religion and Magic met each other and often collided towards each other in an unknown challenging world.

Back to the time of [Isaac Newton](#) who published in 1687 in the "Philosophiae Naturalis Principia Mathematica" a Universal Fundamental Principle in Physics in Harmony with Religion. The Universal Path, the Leitmotiv, the Universal Concept in Physics which was fundamental in science and not in conflict with the Catholic Church. Newton found the

concept of “Universal Equilibrium” which he formulated in his famous third equation Action = - Reaction. In nowadays math the concept of “Universal Equilibrium” has been formulated as:

$$\sum_{i=0}^{i=n} \overline{F}_i = 0 \quad (1)$$

Because the Inertia Force is a Reaction Force, the Inertia Force appears in the equation with a minus sign.

$$\sum_{i=0}^{i=n} \overline{F}_i - m \overline{a} = 0 \quad (2)$$

Equation (2) is a general presentation of Newton’s famous second law of motion. In a fundamental way, Newton’s second law of motion describes the required electromagnetic equation for the Gravitational-Electromagnetic Interaction in general terms, including Maxwell’s theory of Electrodynamics published in 1865 in the article: “A Dynamic Theory of the Electromagnetic Field” and Einstein’s theory of General Relativity published in 1911 the article: “On the Influence of Gravitation on the Propagation of Light”.

Because Maxwell’s 4 equations are not part of one whole uniform understanding of the universe like the fundamental equation of Newton’s second law of motion represents, Maxwell’s theory is missing a fundamental foundation.

Newton's second law of motion has been based on a profound understanding of the universe which is based on the fundamental principle of Harmony and Equilibrium, expressed in equation (2).

To describe the interaction between light and gravity and to understand electromagnetic waves and their interaction and to understand the concept of "photons" it is important to define the fundamental equation for the electromagnetic field based on the fundamental principle of Harmony and Equilibrium formulated by Newton in 1687 and published in his famous work: "[Philosophiae Naturalis Principia Mathematica](#) (Mathematical Principles of Natural Philosophy)".

To realize this, Newton's second law of motion will be the Ground, the Leitmotiv, the Universal Concept in Physics on which the New Theory will be built. The fundamental Electromagnetic force density equation has been based integral on Newton's second law of motion and has been divided into 5 separate terms (B-1 – B-5), each one describing a part of the electromagnetic and inertia force densities.

$$\sum_{i=0}^{i=5} B_i = 0 \quad (3)$$

The first term B-1 represents the inertia of the mass density of light (Electromagnetic Radiation). The terms B-2 and B-3 represent the electric force densities within the Electromagnetic Radiation (Beam of Light) and the terms B-4 and B-5 represent the magnetic force densities within the Electromagnetic Radiation (Beam of Light).

Fundamental in the New Theory is the outcome of (3) which always has to be zero according Newton's fundamental principle of "Universal Equilibrium".

To apply the concept of "Universal Equilibrium" within an electromagnetic field, the electric forces  $F_{\text{Electric}}$ , the magnetic forces  $F_{\text{Magnetic}}$  and the inertia forces will be presented separately in equation (3):

$$\sum_{i=0, j=0}^{i=n, j=m} \left( \overline{F_{\text{Electric}-i}} + \overline{F_{\text{Magnetic}-j}} - m \overline{a} \right) = 0 \quad (4)$$

### 1.3 The Inertia of Light (Term B-1)

Reducing Equation (2) to one single Force  $\overline{F}$ , equation (2) will be written in the well-known presentation:

$$\overline{F} = m \overline{a} \quad (5)$$

The right and the left term of Newton's law of motion in equation (5) has to be divided by the Volume "V" to find an equation for the force density  $\overline{f}$  related to the mass density "ρ".

$$\begin{aligned} \overline{F} &= m \overline{a} \\ \left( \frac{\overline{F}}{V} \right) &= \left( \frac{m}{V} \right) \overline{a} \\ \overline{f} &= \rho \overline{a} \end{aligned} \quad (6)$$

The Inertia Force  $\overline{F}_{\text{Inertia}}$  for Electromagnetic Radiation will be derived from Newton's second law of motion, using the relationship between the momentum vector  $\overline{p}$  for radiation expressed by the Poynting vector  $\overline{S}$  :

$$\overline{F}_{\text{INERTIA}} = -m \overline{a} = -m \frac{\Delta \overline{v}}{\Delta t} = -\frac{\Delta(m\overline{v})}{\Delta t} = -\frac{\Delta \overline{p}}{\Delta t} = -\left(\frac{V}{c^2}\right) \frac{\Delta \overline{S}}{\Delta t} \quad (7)$$

Dividing the right and the left term in equation (7) by the volume  $V$  results in the inertia force density  $\overline{f}_{\text{Inertia}}$  :

$$\begin{aligned} \overline{F}_{\text{INERTIA}} &= -m \overline{a} = -m \frac{\Delta \overline{v}}{\Delta t} = -\frac{\Delta(m\overline{v})}{\Delta t} = -\frac{\Delta \overline{p}}{\Delta t} = -\left(\frac{V}{c^2}\right) \frac{\Delta \overline{S}}{\Delta t} \\ \frac{\overline{F}_{\text{INERTIA}}}{V} &= -\frac{m}{V} \overline{a} = -\frac{m}{V} \frac{\Delta \overline{v}}{\Delta t} = -\frac{1}{V} \frac{\Delta \overline{p}}{\Delta t} = -\left(\frac{1}{c^2}\right) \frac{\Delta \overline{S}}{\Delta t} \\ \overline{f}_{\text{INERTIA}} &= -\rho \overline{a} = -\left(\frac{1}{c^2}\right) \frac{\Delta \overline{S}}{\Delta t} \quad [\text{N}/\text{m}^3] \end{aligned} \quad (8)$$

The Poynting vector  $\overline{S}$  represents the total energy transport of the electromagnetic radiation per unit surface per unit time [ $\text{J} / \text{m}^2 \text{ s}$ ]. Which can be written as the cross product of the Electric Field intensity  $\overline{E}$  and the magnetic Field intensity  $\overline{H}$  .

$$\begin{aligned} \overline{f}_{\text{INERTIA}} &= -\rho \overline{a} = -\left(\frac{1}{c^2}\right) \frac{\Delta \overline{S}}{\Delta t} = -\left(\frac{1}{c^2}\right) \frac{\Delta (\overline{E} \times \overline{H})}{\Delta t} \quad [\text{N}/\text{m}^3] \\ \overline{f}_{\text{INERTIA}} &= -\left(\frac{1}{c^2}\right) \frac{\partial (\overline{E} \times \overline{H})}{\partial t} \quad [\text{N}/\text{m}^3] \end{aligned} \quad (9)$$

## 1.4 Coulomb's Law (Colomb Force) for Electromagnetic Radiation (Term B-2 and B-4)

An example of the Coulomb Force is the Electric Force  $\overline{F}_{\text{Coulomb}}$  acting on an electric charge  $Q$  placed in an electric field  $\overline{E}$ . The equation for the Coulomb Force equals:

$$\overline{F}_{\text{Coulomb}} = \overline{E} Q \quad [\text{N}] \quad (10)$$

Dividing the right and the left term in equation (10) by the volume  $V$  results in the Electric force density  $\overline{f}_{\text{Coulomb}}$  :

$$\begin{aligned} \overline{F}_{\text{COULOMB}} &= \overline{E} Q \quad [\text{N}] \\ \frac{\overline{F}_{\text{COULOMB}}}{V} &= \overline{E} \frac{Q}{V} \quad [\text{N}/\text{m}^3] \\ \overline{f}_{\text{COULOMB}} &= \overline{E} \rho_E \quad [\text{N}/\text{m}^3] \end{aligned} \quad (11)$$

Substituting [Gauss's law in differential form](#) in (11) results in Coulombs Law for Electromagnetic Radiation for the Electric force density  $\overline{f}_{\text{Coulomb}}$  :

$$\begin{aligned} \overline{f}_{\text{COULOMB}} &= \overline{E} \rho_E \\ \overline{f}_{\text{COULOMB}} &= \overline{E} \rho_E = \overline{E} (\nabla \cdot \overline{D}) \\ \overline{f}_{\text{COULOMB}} &= \overline{E} (\nabla \cdot \overline{D}) = \varepsilon \overline{E} (\nabla \cdot \overline{E}) \quad [\text{N}/\text{m}^3] \end{aligned} \quad (12)$$


In Electromagnetic Field Configurations, there is in general no preference for the electric force densities or the magnetic force densities. In general the equations for the electric field densities are universally exchangeable with the magnetic field densities.

For the magnetic field densities, equation (12) can be written as:

$$\begin{aligned}\bar{f}_{\text{Coulomb - Electric}} &= \bar{E}(\nabla \cdot \bar{D}) = \varepsilon \bar{E}(\nabla \cdot \bar{E}) \left[ \text{N/m}^3 \right] \text{ (Term B-2)} \\ \bar{f}_{\text{Coulomb - Magnetic}} &= \bar{H}(\nabla \cdot \bar{B}) = \mu \bar{H}(\nabla \cdot \bar{H}) \left[ \text{N/m}^3 \right] \text{ (Term B-4)}\end{aligned} \quad (13)$$

## 1.5 Lorentz's Law (Lorentz Force) for Electromagnetic Radiation (Term B-3 and B-5)

An example of the Lorentz Force is the Magnetic Force  $F_{\text{Lorentz}}$  acting on an electric charge  $Q$  moving with a velocity  $v$  within a magnetic field with magnetic field intensity  $B$  (magnetic induction).



**Fig. 1. The Lorentz Force equals the cross product of the Magnetic Induction  $B$  and the velocity  $v$  of the charge  $q$  moving within the magnetic field times the value of the electric charge**

The equation for the Lorentz Force equals:

$$\vec{F}_{\text{LORENTZ}} = Q \vec{v} \times \vec{B} \quad [\text{N}] \quad (14)$$

Dividing the right and the left term in equation (14) by the volume  $V$  results in the Lorentz force density  $\vec{f}_{\text{Lorentz}}$

$$\begin{aligned}\bar{\mathbf{F}}_{\text{LORENTZ}} &= Q \bar{\mathbf{v}} \times \bar{\mathbf{B}} \quad [\text{N}] \\ \frac{\bar{\mathbf{F}}_{\text{LORENTZ}}}{V} &= - \bar{\mathbf{B}} \times \frac{Q \bar{\mathbf{v}}}{V} \quad [\text{N/ m}^3] \\ \bar{\mathbf{f}}_{\text{LORENTZ}} &= - \bar{\mathbf{B}} \times \frac{Q \bar{\mathbf{v}}}{V} = - \bar{\mathbf{B}} \times \bar{\mathbf{j}} = - \mu \bar{\mathbf{H}} \times \bar{\mathbf{j}} \quad [\text{N/ m}^3]\end{aligned} \quad (15)$$

In which  $q$  is the electric charge,  $v$  the velocity of the electric charge,  $B$  the magnetic induction and  $j$  the electric current density. Substituting Ampère's law in differential form in (15) results in Lorentz's Law for Electromagnetic Radiation

for the Electric force density  $\bar{\mathbf{f}}_{\text{Lorentz}}$  :

$$\begin{aligned}\bar{\mathbf{f}}_{\text{LORENTZ}} &= - \mu \bar{\mathbf{H}} \times (\bar{\mathbf{j}}) \\ \bar{\mathbf{f}}_{\text{LORENTZ}} &= - \mu \bar{\mathbf{H}} \times (\bar{\mathbf{j}}) = - \mu \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) \quad [\text{N/ m}^3]\end{aligned} \quad (16)$$

In Electromagnetic Field Configurations, there is in general no preference for the electric force densities or the magnetic force densities. In general the equations for the electric field densities are universally exchangeable with the magnetic field densities. For the electric field densities, equation (16) can be written as:

$$\begin{aligned}\bar{\mathbf{f}}_{\text{Coulomb - Electric}} &= - \epsilon \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) \quad [\text{N/ m}^3] \quad (\text{Term B-3}) \\ \bar{\mathbf{f}}_{\text{Coulomb - Magnetic}} &= - \mu \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) \quad [\text{N/ m}^3] \quad (\text{Term B-5})\end{aligned} \quad (17)$$

## 1.6 The Fundamental Universal Equation for the Electromagnetic field (Term B-1 + Term B-2 + Term B-3 + Term B-4 + Term B-5)

Newton's second law of motion applied within any arbitrary electromagnetic field configuration results in the fundamental equation (23) for any arbitrary electromagnetic field configuration (a beam of light):

$$\begin{aligned}
 & \text{NEWTON: } \mathbf{F}_{\text{TOTAAL}} = m \mathbf{a} \text{ represents: } \mathbf{f}_{\text{TOTAAL}} = \rho \mathbf{a} \\
 & -\rho \mathbf{a} + \mathbf{f}_{\text{TOTAAL}} = 0 \\
 & -\rho \mathbf{a} + \mathbf{f}_{\text{ELEKTRISCH}} + \mathbf{f}_{\text{MAGNETISCH}} = 0 \quad (23) \\
 & -\rho \mathbf{a} + \mathbf{F}_{\text{COULOMB}} + \mathbf{F}_{\text{LORENTZ}} + \mathbf{F}_{\text{COULOMB}} + \mathbf{F}_{\text{LORENTZ}} = 0 \\
 & -\frac{1}{c^2} \frac{\partial (\vec{\mathbf{E}} \times \vec{\mathbf{H}})}{\partial t} + \epsilon_0 \vec{\mathbf{E}} (\nabla \cdot \vec{\mathbf{E}}) - \epsilon_0 \vec{\mathbf{E}} \times (\nabla \times \vec{\mathbf{E}}) + \mu_0 \vec{\mathbf{H}} (\nabla \cdot \vec{\mathbf{H}}) - \mu_0 \vec{\mathbf{H}} \times (\nabla \times \vec{\mathbf{H}}) = 0 \\
 & \quad \text{B-1} \qquad \text{B-2} \qquad \text{B-3} \qquad \text{B-4} \qquad \text{B-5}
 \end{aligned}$$

Term B-4 is the magnetic equivalent of the (electric) Coulomb's law B-2 and Term B-3 is the electric equivalent of the (magnetic) Lorentz's law B-5.

The universal equation for the electromagnetic field (free electromagnetic waves and confined electromagnetic fields) has been presented in (24) and expresses the perfect equilibrium between the inertia forces (B-1), the electric forces (B-2 and B-3) and the magnetic forces (B-4 and B-5) in any arbitrary electromagnetic field configuration.

$$\begin{aligned}
 & -\frac{1}{c^2} \frac{\partial (\vec{\mathbf{E}} \times \vec{\mathbf{H}})}{\partial t} + \epsilon_0 \vec{\mathbf{E}} (\nabla \cdot \vec{\mathbf{E}}) - \epsilon_0 \vec{\mathbf{E}} \times (\nabla \times \vec{\mathbf{E}}) + \mu_0 \vec{\mathbf{H}} (\nabla \cdot \vec{\mathbf{H}}) - \mu_0 \vec{\mathbf{H}} \times (\nabla \times \vec{\mathbf{H}}) = 0 \quad (24) \\
 & \quad \text{B-1} \qquad \text{B-2} \qquad \text{B-3} \qquad \text{B-4} \qquad \text{B-5}
 \end{aligned}$$

## 1.7 The Universal Integration of Maxwell's Theory of Electrodynamics:

The universal equation (24) for any arbitrary electromagnetic field configuration can be written in the form:

$$\begin{aligned}
 & -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) = 0 \\
 & -\epsilon_0 \mu_0 \left( \bar{\mathbf{E}} \times \frac{\partial (\bar{\mathbf{H}})}{\partial t} + \bar{\mathbf{H}} \times \frac{\partial (\bar{\mathbf{E}})}{\partial t} \right) + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) = 0 \quad (25) \\
 & - \left( \underbrace{\epsilon_0 \bar{\mathbf{E}} \times \frac{\partial (\bar{\mathbf{B}})}{\partial t}}_{\text{M-3}} + \underbrace{\mu_0 \bar{\mathbf{H}} \times \frac{\partial (\bar{\mathbf{D}})}{\partial t}}_{\text{M-4}} \right) + \underbrace{\bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{D}})}_{\text{M-1}} - \underbrace{\epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}})}_{\text{M-3}} + \underbrace{\bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{B}})}_{\text{M-2}} - \underbrace{\mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}})}_{\text{M-4}} = 0
 \end{aligned}$$

The Maxwell Equations are presented in (26):

$$\begin{aligned}
 \nabla \cdot \bar{\mathbf{D}} &= \rho \quad (\text{M-1}) & \nabla \times \bar{\mathbf{E}} &= -\frac{\partial \mathbf{B}}{\partial t} \quad (\text{M-3}) \\
 \nabla \cdot \bar{\mathbf{B}} &= 0 \quad (\text{M-2}) & \nabla \times \bar{\mathbf{H}} &= \frac{\partial \mathbf{D}}{\partial t} \quad (\text{M-4})
 \end{aligned} \quad (26)$$

In vacuum in the absence of any charge density, it follows from (26) that all the solutions for the Maxwell's Equations are also solutions for the separate parts of the Universal Equation (25) for the Electromagnetic field.

Universal Equation for the Electromagnetic Field.

$$\begin{aligned}
 & - \left( \underbrace{\epsilon_0 \bar{\mathbf{E}} \times \frac{\partial (\bar{\mathbf{B}})}{\partial t}}_{\text{M-3}} + \underbrace{\mu_0 \bar{\mathbf{H}} \times \frac{\partial (\bar{\mathbf{D}})}{\partial t}}_{\text{M-4}} \right) + \underbrace{\bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{D}})}_{\text{M-1}} - \underbrace{\epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}})}_{\text{M-3}} + \underbrace{\bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{B}})}_{\text{M-2}} - \underbrace{\mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}})}_{\text{M-4}} = 0 \\
 & 4 \text{ Maxwell's Equations} \\
 \nabla \cdot \bar{\mathbf{D}} &= \rho \quad (\text{M-1}) & \nabla \times \bar{\mathbf{E}} &= -\frac{\partial \mathbf{B}}{\partial t} \quad (\text{M-3}) \\
 \nabla \cdot \bar{\mathbf{B}} &= 0 \quad (\text{M-2}) & \nabla \times \bar{\mathbf{H}} &= \frac{\partial \mathbf{D}}{\partial t} \quad (\text{M-4})
 \end{aligned} \quad (27)$$

Comparing the 4 Maxwell Equations (26) with the Universal Equation (24) we conclude that the 4 Maxwell equations show only the 4 parts of the Universal Dynamic Equilibrium in 4 separate terms and the 4 Maxwell equations are missing the fundamental term for inertia. For that reason it is impossible to calculate the interaction between light and gravity with the 4 Maxwell equations. To find the interaction terms between light and gravity the inertia term (B-1 in 24) is necessary.

## 1.8 Interaction between Gravity and Light (Electromagnetic Radiation).

To define the Fundamental Equation for the Interaction between Gravity and Light, an extra term (B-6) has been introduced in equation (24). The term B-6 represents the force density of the gravitational field acting on the electromagnetic mass density.

$$F_{\text{GRAVITY}} = m \bar{g} \text{ [N]}$$

Dividing both terms by the Volume V:

$$\frac{F_{\text{GRAVITY}}}{V} = \frac{m}{V} \bar{g} \text{ [N/ m}^3\text{]} \quad (28)$$

Results in the force density:

$$f_{\text{GRAVITY}} = \rho \bar{g} \text{ [N/ m}^3\text{]}$$

The specific mass “ $\rho$ ” of a beam of light follows from Einstein’s equation:

$$W = m c^2$$

Dividing both terms by the Volume V results in:

$$\frac{W}{V} = \frac{m}{V} c^2 \quad (29)$$

which represents the energy density "w" and the specific mass " $\rho$ " of the electromagnetic radiation:

$$w = \rho c^2$$

which results for an expression of the specific mass  $\rho$ :

$$\rho = \frac{1}{c^2} w = \varepsilon \mu w$$

The energy density “w” follows from the electric and the magnetic field intensities:

$$w = \frac{1}{2} \varepsilon E^2 + \frac{1}{2} \mu H^2$$

$$w = \frac{1}{2} (\varepsilon E^2 + \mu H^2) = \frac{1}{2} (\varepsilon (\vec{E} \cdot \vec{E}) + \mu (\vec{H} \cdot \vec{H})) \quad (30)$$

Substituting equation (30) in equation (29) results in the gravitational force density  $f_{\text{GRAVITY}}$  acting on an arbitrary electromagnetic field configuration (a beam of light) with mass density  $\rho$ .

$$f_{\text{GRAVITY}} = \rho \vec{g}$$

$$f_{\text{GRAVITY}} = \rho \vec{g} = \varepsilon \mu w \vec{g} = \frac{1}{2} (\varepsilon^2 \mu (\vec{E} \cdot \vec{E}) + \varepsilon \mu^2 (\vec{H} \cdot \vec{H})) \vec{g} \quad (31)$$

Substituting equation (31) in equation (24) results in the fundamental equation describing the Electromagnetic-Gravitational interaction for any arbitrary electromagnetic field configuration (a beam of light):

$$\begin{aligned} & \text{NEWTON: } \vec{F}_{\text{TOTAAL}} = m \vec{a} \text{ [N]} \\ & \text{NEWTON: Expressed in force densities: } \vec{f}_{\text{TOTAAL}} = \rho \vec{a} \text{ [N/m}^3 \text{]} \\ & -\rho \vec{a} + \vec{f}_{\text{TOTAAL}} = \vec{0} \quad (32) \\ & -\rho \vec{a} + \vec{f}_{\text{ELEKTRISCH}} + \vec{f}_{\text{MAGNETISCH}} + \vec{f}_{\text{GRAVITY}} = \vec{0} \\ & \vec{f}_{\text{INERTIA}} + \vec{f}_{\text{COULOMB}} + \vec{f}_{\text{LORENTZ}} + \vec{f}_{\text{COULOMB}} + \vec{f}_{\text{LORENTZ}} + \vec{f}_{\text{GRAVITY}} = \vec{0} \\ & -\frac{1}{c^2} \frac{\partial (\vec{E} \times \vec{H})}{\partial t} + \varepsilon_0 \vec{E} (\nabla \cdot \vec{E}) - \varepsilon_0 \vec{E} \times (\nabla \times \vec{E}) + \mu_0 \vec{H} (\nabla \cdot \vec{H}) - \mu_0 \vec{H} \times (\nabla \times \vec{H}) + \frac{1}{2} (\varepsilon^2 \mu (\vec{E} \cdot \vec{E}) + \varepsilon \mu^2 (\vec{H} \cdot \vec{H})) \vec{g} = \vec{0} \\ & \quad \text{B-1} \quad \text{B-2} \quad \text{B-3} \quad \text{B-4} \quad \text{B-5} \quad \text{B-6} \end{aligned}$$

Term B-1 represents the inertia term of the electromagnetic radiation. Term B-4 is the magnetic representation of the (electric) Coulomb’s Force B-2 and Term B-3 is the electric representation of the (magnetic) Lorentz Force B-5. Term B-6 represents the Electromagnetic-Gravitational interaction of

a gravitational field with field acceleration  $\bar{g}$  acting on an arbitrary electromagnetic field configuration ( a beam of light) with specific mass  $\rho$ .

The universal equation for the electromagnetic field (free electromagnetic waves and confined electromagnetic fields) within a gravitational field with gravity field intensity  $\bar{g}$  has been presented in (33) and expresses the perfect equilibrium between the inertia forces (B-1), the electric forces (B-2 and B-3), the magnetic forces (B-4 and B-5) and the gravitational force (B-6) in any arbitrary electromagnetic field configuration.

$$\begin{aligned}
 & - \frac{1}{c^2} \frac{\partial(\bar{E} \times \bar{H})}{\partial t} + \epsilon_0 \bar{E} (\nabla \cdot \bar{E}) - \epsilon_0 \bar{E} \times (\nabla \times \bar{E}) + \mu_0 \bar{H} (\nabla \cdot \bar{H}) - \\
 & \qquad \qquad \qquad \text{B-1} \qquad \qquad \qquad \text{B-2} \qquad \qquad \qquad \text{B-3} \qquad \qquad \qquad \text{B-4} \\
 & - \mu_0 \bar{H} \times (\nabla \times \bar{H}) + \frac{1}{2} \left( \epsilon^2 \mu (\bar{E} \cdot \bar{E}) + \epsilon \mu^2 (\bar{H} \cdot \bar{H}) \right) \bar{g} = \bar{0} \\
 & \qquad \qquad \qquad \text{B-5} \qquad \qquad \qquad \text{B-6}
 \end{aligned} \tag{33}$$

## 1.9 The Confinement of Light (Electromagnetic Radiation).

When a beam of light is approaching a strong gravitational field in the direction of the gravitational field, generated by a Black Hole, the confinement has been called a Longitudinal Black Hole. The direction of propagation of the beam of light is in the same direction (or in the opposite direction) of the gravitational field. According the first term in (33), the beam of light will be accelerated or decelerated. However, the speed of light is a universal constant and for that reason the speed of light cannot increase or decrease. Instead the intensity of the electromagnetic radiation will increase when the beam of light approaches (propagates in the opposite direction as the direction of the gravitational field) the Black Hole. And the intensity of the electromagnetic radiation will decrease when the beam of light leaves (propagates in the same direction as the direction of the gravitational field) the Black Hole.

The Gravitational-Electromagnetic Confinement for the elementary structure beyond the “[superstring](#)” / “[Black Hole](#)” is presented in equation (34).

3-Dimensional Space Domain

$$\begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} - \frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \varepsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \varepsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \quad (34) \\ + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \frac{1}{2} (\varepsilon^2 \mu (\bar{\mathbf{E}} \cdot \bar{\mathbf{E}}) + \varepsilon \mu^2 (\bar{\mathbf{H}} \cdot \bar{\mathbf{H}})) \bar{\mathbf{g}} = \bar{\mathbf{0}}$$

In which  $\bar{g}$  represents the gravitational acceleration acting on the electromagnetic mass density of the confined electromagnetic radiation.

A possible solution for equation (34) describing an Electromagnetic-Gravitational confinement within a radial gravitational field with acceleration  $\bar{g}$  has been represented in (35).

$$\begin{pmatrix} e_r \\ e_\theta \\ e_\phi \end{pmatrix} = \begin{pmatrix} 0 \\ f(r) \sin(\omega t) \\ -f(r) \cos(\omega t) \end{pmatrix} \quad \begin{pmatrix} m_r \\ m_\theta \\ m_\phi \end{pmatrix} = \begin{pmatrix} 0 \\ f(r) \cos(\omega t) \\ f(r) \sin(\omega t) \end{pmatrix} \quad \bar{g} = \begin{pmatrix} \frac{G_1}{4 \pi r^2} \\ 0 \\ 0 \end{pmatrix} \quad (35)$$

$$w_{em} = \left( \frac{\mu_0}{2} (\bar{m} \cdot \bar{m}) + \frac{\epsilon_0}{2} (\bar{e} \cdot \bar{e}) \right) = \epsilon_0 f(r)^2$$

In which the radial function  $f(r)$  equals:

$$f[r] = K e^{-\frac{G_1 \epsilon_0 \mu_0}{r} + 8 \pi \log[r]} \quad (36)$$


The solution has been calculated according [Newton's Shell Theorem](#).

**1.10 Confinement of Light (Electromagnetic Radiation) in the region smaller than “SuperStrings” with an electromagnetic mass of  $emm = 1.6726 \times 10^{-27}$  [kg] and the radius =  $3 \times 10^{-58}$  [m]**

For an electromagnetic mass of the confinement:  $emm = 1.6726 \times 10^{-27}$  [kg] (mass of proton), the radius of the confinement equals approximately  $3 \times 10^{-58}$  [m]. This is far beyond the order of [Planck's Length](#),


The [Plot graph](#) of the Electric Field Intensity  $f(r)$  of the confinement has been presented as a function of the radius in figure (2) and figure (3):

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^{-59}, 10^{-55}\} \right]$$



**Figure 2** PlotGraph of the Electric Field Intensity  $f(r)$  [V/ m] for the region  $10^{-59} < r < 10^{-55}$  [m] in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $1.6\ 726 \times 10^{-27}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#).

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^{-59}, 10^{-57}\} \right]$$



**Figure 3** PlotGraph of the Electric Field Intensity  $f(r)$  [V/m] for the region  $10^{-59} < r < 10^{-57}$  [m] in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $1.6726 \times 10^{-27}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#).

The fundamental question is: How it is possible to create confinements from “visible light” (with a wave length between  $3.9 \times 10^{-7}$  [m] until  $7 \times 10^{-7}$  [m]) within dimensions smaller than Planck's Length?

This is only possible when the wave length of the confined radiation is smaller than the dimensions of the confinement. This requires extreme high frequencies. The transformation in frequency from visible light into the extreme high frequency of the confinement is possible because of the Lorentz/ Doppler transformation during the collapse of the radiation when the confinement has been formed (implosion of visible light).

**1.10.1 Confinement of Light (Electromagnetic Radiation) in the region of “Superstrings” with dimensions in the order of Planck’s Length with an electromagnetic mass of  $10^{-4}$  [kg] and a radius =  $2 \times 10^{-35}$  [m]):**

Figure 4 and Fig. 5 represent the electromagnetic field density (along the vertical axis) as a function of the distance (along the horizontal axis) of the center of the Confinement of Light with dimensions in the order of Planck’s length. The chosen values equal:

$$f[r] = K e^{-\frac{G1 \text{ emm } \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}$$

$$G1 = 6.6740810^{-11} \quad (37)$$

$$\text{emm} = 10^{-4} \text{ [kg]}$$

$$\varepsilon_0 = 8.8510^{-12}$$


$$\mu_0 = 1.256637061435917210^{-6}$$

In which “emm” represents the electromagnetic mass of the confinement located at the center according [Newton’s Shell Theorem](#).

For an electromagnetic mass  $\text{emm} = 10^{-4}$  [kg] of the Confinement of Light, the radius of the confinement equals approximately  $2 \times 10^{-35}$  [m] and the first harmonic frequency equals  $1.5 \times 10^{27}$  [Hz].


[The Plot graph](#) of the Electric Field Intensity  $f(r)$  of the confinement has been presented as a function of the radius in figure (4) and figure (5):

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^{-36}, 10^{-25}\} \right]$$



**Figure 4** PlotGraph of the Electric Field Intensity  $f(r)$  [V/ m] for the region  $10^{-36} < r < 10^{-25}$  [m] in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $10^{-4}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#).

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^{-36}, 10^{-35}\} \right]$$



**Figure 5** PlotGraph of the Electric Field Intensity  $f(r)$  [V/ m] for the region  $10^{-36} < r < 10^{-35}$  [m] in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $10^{-4}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#).

It follows from Figure 5 that the radius of the stable gravitational electromagnetic confinement equals approximately  $2 \times 10^{-35}$  [m], which is the size of the [Planck length](#). According the theory of [superstrings](#), the fundamental constituents of reality are [strings](#) of the [Planck length](#) (about  $1.62 \times 10^{-35}$  [m]) that vibrate at resonant frequencies.

**1.10.2 Confinement of Light (Electromagnetic Radiation) in the region of a Longitudinal Black Hole with an electromagnetic mass of  $10^{40}$  [kg], a radius =  $1.5 \times 10^9$  [m] at a frequency of 0.2 [Hz].**

To realize a Gravitational-Electromagnetic confinement for a conventional Black Hole with an Electro Magnetic Mass:  $emm = 10^{40}$  [kg], the solution (30) and (31) for the Gravitational Electromagnetic Equilibrium Equation (34) results in a Gravitational Electromagnetic Confinement radius  $r = 1.5 \times 10^9$  [m] (Figure 6 and Figure 7).

$$f[r] = K e^{-\frac{G1 emm \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}$$

$$G1 = 6.6740810^{-11}$$

$$emm = 10^{40} \text{ [kg]} \tag{38}$$


$$\varepsilon_0 = 8.8510^{-12}$$

$$\mu_0 = 1.256637061435917210^{-6}$$

In which “emm” equals the electromagnetic mass of the Single Harmonic Black Hole located at the center according [Newton’s Shell Theorem](#). For an electromagnetic mass of the Single Harmonic Black Hole (SHBH), the value for the electromagnetic mass (emm) equals:  $emm = 10^{40}$  [kg], the radius of the confinement equals approximately  $1.5 \times 10^9$  [m] and the first harmonic frequency equals 0.2 [Hz].


[The Plot graph](#) of the Electric Field Intensity  $f(r)$  of the SHBH has been presented as a function of the radius in figure (6) and figure (7):

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^5, 10^{12}\} \right]$$



**Fig. 6. PlotGraph of the Electric Field Intensity  $f(r)$  [V/ m] (vertical axis) for the region  $10^5 < r < 10^{12}$  [m] (horizontal axis) in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $10^{40}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#).**

$$\text{Plot} \left[ e^{-\frac{G1 \varepsilon_0 \mu_0}{r} + 8 \pi \log[r]}, \{r, 10^9, 4 \times 10^9\} \right]$$




**Fig. 7. PlotGraph of the Electric Field Intensity  $f(r)$  [V/m] (vertical axis) for the region  $10^9 < r < 4 \cdot 10^9$  [m] (horizontal axis) in which the gravitational field acceleration has been chosen accordingly an electromagnetic mass of  $10^{40}$  [kg] located at the center of the confinement, according [Newton's Shell Theorem](#). And a corresponding Single Harmonic frequency of 0.2 [Hz].**

It follows from Figure 7 that the radius of the stable gravitational electromagnetic confinement of the SHBH equals approximately  $1.5 \cdot 10^9$  [m].

### 1.11 The Transversal Black Hole.

We consider a beam of light passing a strong gravitational field, generated by a Black Hole. According the first term in (1) the beam of light will follow a circular orbit around the Black Hole. The required Equilibrium will exist at the radius where the centrifugal electromagnetic inertia forces will be equal and opposite directed to the centripetal oriented gravitational forces on the electromagnetic mass. Figure 12 represents the orbit (colored red) of a LASER beam around a uniform intense gravitational field (Black Hole)



**Fig. 8. A LASER beam around a Black Hole captured in a circular orbit around the Black Hole in a Transversal Modus by the Gravitational interaction of the Black Hole with the mass (inertia) of the Laser Beam.**

In general, Newtons second law of motion has been presented as:

$$F = m a \quad (39)$$

In which “a” represents the acceleration which equals the difference of the velocity  $\Delta v$  divided by the time interval  $\Delta t$ .

$$a = \frac{\Delta v}{\Delta t} \quad (40)$$

The momentum  $p$  of a mechanical mass equals:

$$p = m v \quad (41)$$

Then Newton’s second law of motion can be presented as:

$$F_{\text{INERTIA}} = m a = m \frac{\Delta v}{\Delta t} = \frac{\Delta (m v)}{\Delta t} = \frac{\Delta (p)}{\Delta t} \quad (42)$$

Like a mechanical mass expresses the property of inertia, also a beam of light expressed the property of inertia. When the sun shines on the earth, the radiation of the sun presses on the earth with thousands of Newton.

Like a mechanical mass, also a beam of light has momentum. The momentum of a beam of light has been expressed by the Poynting vector  $S$  and equals the mechanical momentum vector  $p$  multiplied by the square of the speed of light  $c$  divided by the Volume.

$$F_{\text{INERTIA}} = m a = m \frac{\Delta v}{\Delta t} = \frac{\Delta (m v)}{\Delta t} = \frac{\Delta (p)}{\Delta t} = \frac{V \Delta (S)}{c^2 \Delta t} \quad (43)$$

(mechanical mass)                      (beam of light)

The inertia force density “f” equals the inertia force “F” divided by the Volume “V”.

$$f_{\text{INERTIA}} = \left( \frac{m}{V} \right) a = \rho \frac{\Delta v}{\Delta t} = \frac{1}{c^2} \frac{\Delta (S)}{\Delta t} \quad (44)$$

(mechanical mass)    (beam of light)

The well-known equation of Einstein equals:

$$W = m c^2$$

$$w = \frac{W}{V} = \frac{m}{V} c^2 = \rho c^2 \quad (45)$$

In which “w” represents the electromagnetic energy density and equals:

$$w = \frac{1}{2} \epsilon E^2 + \frac{1}{2} \mu H^2 \quad (46)$$

For electromagnetic radiation the electromagnetic impedance  $Z_0$  equals:

$$\begin{aligned}
Z_0 &= \frac{E}{H} = \sqrt{\frac{\mu}{\epsilon}} \\
H &= E \sqrt{\frac{\epsilon}{\mu}} \\
w &= \frac{1}{2} \epsilon E^2 + \frac{1}{2} \mu H^2 = \frac{1}{2} \epsilon E^2 + \frac{1}{2} \epsilon E^2 = \epsilon E^2
\end{aligned} \tag{47}$$

Substituting equation (47) in (45) and (46) results in:

$$\begin{aligned}
w &= \rho c^2 = \epsilon E^2 \\
\rho &= \frac{\epsilon}{c^2} E^2
\end{aligned} \tag{48}$$

Because the beam of light has been confined in the radial direction, it demonstrates in the radial direction the property of inertia (electromagnetic mass) and interacts with a gravitational field according Newton's second law of motion. The whole Universe is in a perfect Equilibrium. Also at the "Event Horizon" of a Black Hole does exist a perfect equilibrium between the confining gravitational force of the Black Hole and the radial directed inertia force density of the confined electromagnetic radiation (Laser Beam confined by a Black Hole at the "Event Horizon").

To determine the "Event Horizon" of the Black Hole (Radius of the circular orbit of the Laser Beam), we have to find the perfect equilibrium between the inertia force densities of the electromagnetic energy densities of the Laser Beam and the confining gravitational force acting on the electromagnetic energy densities of the Laser Beam.

$$f_{\text{GRAVITY}} = f_{\text{INERTIA}}$$

$$\rho g = \frac{1}{c^2} \frac{\Delta (S)}{\Delta t} \quad (49)$$

From Figure (8) follows the relationship between de changing in the Poynting vector  $\Delta S$  during the time interval  $\Delta t$  :

$$\text{Tan}(\alpha) = \frac{c \Delta t}{R} = \frac{\Delta S}{S}$$

$$\frac{\Delta S}{\Delta t} = \frac{c S}{R} \quad (50)$$

$$\rho g = \frac{1}{c^2} \frac{\Delta (S)}{\Delta t} = \frac{1}{c^2} \frac{c S}{R} = \frac{1}{c} \frac{S}{R}$$

The Poynting vector  $\vec{S}$  represents the total energy transport of the electromagnetic radiation per unit surface per unit time [J / m<sup>2</sup> s]. Which can be written as the cross product of the Electric Field intensity  $\vec{E}$  and the magnetic Field intensity  $\vec{H}$  .

$$\vec{S} = \vec{E} \times \vec{H}$$

$$S = E H \text{Sin}(90^\circ) = E H \quad (51)$$

$$S = E^2 \sqrt{\frac{\epsilon}{\mu}}$$

Substituting equation (49) and (51) in (50) results in an equation for the Event Horizon at radius “R” of a Transversal Black Hole.

$$\begin{aligned}
\rho g &= \frac{1}{c} \frac{S}{R} \\
R &= \frac{S}{\rho c g} \\
R &= \frac{E^2 \sqrt{\frac{\epsilon}{\mu}}}{\rho c g} = \frac{E^2 \sqrt{\frac{\epsilon}{\mu}}}{\frac{\epsilon}{c} E^2 g} \quad (52) \\
R &= \frac{c^2}{g} \approx \frac{9 \cdot 10^{16}}{g} \quad [\text{m}]
\end{aligned}$$

Equation (52) represents the perfect equilibrium between the inertia force densities of the electromagnetic mass  $\frac{1}{c^2} \frac{\Delta S}{\Delta t}$  and the centripetal oriented gravitational force density  $\frac{w}{c^2} g$  acting on the electromagnetic mass. The perfect equilibrium direction [9,10,12,13] where the inertia forces due to the circular orbit of the beam of light are in a perfect balance with the attractive gravitational forces, exists at one defined radius “R” of the beam of light (LASER Beam), independent of the intensity of the beam of light and independent of the frequency of the beam of light. Only the acceleration “g” of the gravitational field determines the radius of equilibrium “R”

$$R \approx \frac{9 \cdot 10^{16}}{g} \quad (53)$$

In which “R” is the radius of the beam of light and “g” the acceleration of the gravitational field of the “Black Hole”.

The x-y plane is oriented perpendicular on the z-direction. The speed of light towards the positive z-direction equals the speed of light (the constant "c = 300.000 km/s"). But the speed of light in the x-y plane has to be exactly zero [9,14,15]. Else the diameter of the laser beam would become larger and larger during the propagation along the positive z-direction. This is only possible because the Electromagnetic confining forces B-2, B-3, B-4 and B-5 in equation (34) compensate exactly the outward oriented radiation pressure towards the x-direction and the y-direction. The Radial Radiation Pressure has been compensated by the Coulomb Force Densities and the Lorentz Force Densities within the Laser Beam.

## 1.12 The Origin of Electric Charge and Magnetic Spin in discrete values (The introduction of Quantum Numbers)

The Gravitational-Electromagnetic Confinement for the elementary structure of the Confined Electromagnetic Radiation has been presented in equation (34).

3-Dimensional Space Domain

$$\begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} - \frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \frac{1}{2} (\epsilon^2 \mu (\bar{\mathbf{E}} \cdot \bar{\mathbf{E}}) + \epsilon \mu^2 (\bar{\mathbf{H}} \cdot \bar{\mathbf{H}})) \bar{\mathbf{g}} = \bar{\mathbf{0}} \quad (54)$$

In which  $\bar{\mathbf{g}}$  represents the (radial oriented) gravitational acceleration caused by the electromagnetic mass density of the confined electromagnetic radiation.

To find the origin of Electric Charge and Magnetic Spin we choose as an example a solution for (54) which equals:

$$\begin{pmatrix} e_r \\ e_\theta \\ e_\varphi \end{pmatrix} = \begin{pmatrix} 0 \\ f_1(r, \theta, \varphi, t) \sin(\omega t) \\ -f_2(r, \theta, \varphi, t) \cos(\omega t) \end{pmatrix} \quad \begin{pmatrix} m_r \\ m_\theta \\ m_\varphi \end{pmatrix} = \begin{pmatrix} 0 \\ f_2(r, \theta, \varphi, t) \cos(\omega t) \\ f_1(r, \theta, \varphi, t) \sin(\omega t) \end{pmatrix} \quad (55)$$

$$w_{em} = \left( \frac{\mu_0}{2} (\bar{\mathbf{m}} \cdot \bar{\mathbf{m}}) + \frac{\epsilon_0}{2} (\bar{\mathbf{e}} \cdot \bar{\mathbf{e}}) \right) = \epsilon_0 f(r)^2$$

In which  $f[r]$ ,  $f_1[r, \theta, \varphi, t]$ ,  $f_2[r, \theta, \varphi, t]$  equals:

$$\begin{aligned}
 f[r] &= K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} \\
 f_1[r, \theta, \varphi, t] &= K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} g_1[\theta, \varphi, t] \\
 f_2[r, \theta, \varphi, t] &= \frac{K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}}}{\sqrt{2}} \sqrt{-g_1[\theta, \varphi, t]^2 + \cos[2 \omega t] g_1[\theta, \varphi, t]^2 + 2h[\theta, \varphi]}
 \end{aligned} \tag{56}$$

In which  $g_1[\theta, \varphi, t]$  and  $h[\theta, \varphi]$  are arbitrary function. The Electromagnetic Confinement has been described for the electric field intensity:

$$\begin{pmatrix} e_r \\ e_\theta \\ e_\varphi \end{pmatrix} = \begin{pmatrix} 0 \\ K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} g_1[\theta, \varphi, t] \sin[\omega t] \\ -\frac{K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}}}{\sqrt{2}} \sqrt{-g_1[\theta, \varphi, t]^2 + \cos[2 \omega t] g_1[\theta, \varphi, t]^2 + 2h[\theta, \varphi]} \end{pmatrix} \tag{57}$$

The Electromagnetic confinement has been described for the magnetic field intensity:

$$\begin{pmatrix} m_r \\ m_\theta \\ m_\varphi \end{pmatrix} = \sqrt{\frac{\varepsilon_0}{\mu_0}} \begin{pmatrix} 0 \\ K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} \frac{\sqrt{-g_1[\theta, \varphi, t]^2 + \cos[2 \omega t] g_1[\theta, \varphi, t]^2 + 2h[\theta, \varphi]}}{\sqrt{2}} \\ K e^{-\frac{G1 \varepsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} g_1[\theta, \varphi, t] \sin[\omega t] \end{pmatrix} \tag{58}$$

The following functions with the quantum numbers  $\{m_1, n_1, p_1, q_1\}$  have been chosen:

$$\begin{aligned}
 f[r] &= K e^{-\frac{G_1 \epsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} \\
 g_1(\theta, \varphi, t) &= \sin(\omega t) (\sin(\pi \theta m_1) \sin(n_1 2 \pi \varphi) + 1) \\
 h(\theta, \varphi) &= \sin(\pi \theta p_1) \sin(q_1 2 \pi \varphi) + 1 \\
 g_2(\theta, \varphi, t) &= \frac{\sec(\omega t) \sqrt{\cos(2\omega t) g_1(\theta, \varphi, t)^2 - g_1(\theta, \varphi, t)^2 + 2 h(\theta, \varphi)}}{\sqrt{2}} \\
 f_1[r, \theta, \varphi, t] &= e^{-\frac{G_1 \epsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} K g_1[\theta, \varphi, t] \\
 f_2[r, \theta, \varphi, t] &= \frac{e^{-\frac{G_1 \epsilon_0 \mu_0 + 8 \pi \log[r]}{r} \frac{1}{8 \pi}} K \sqrt{-g_1[\theta, \varphi, t]^2 + \cos[2\omega t] g_1[\theta, \varphi, t]^2 + 2 h[\theta, \varphi]}}{\sqrt{2}}
 \end{aligned} \tag{59}$$

### 1.13 Electromagnetic Confinement with Electric- and Magnetic Dipoles (Electric- and Magnetic Spin) { $m_1=0, n_1=0, p_1=0, q_1=0$ }

The divergence of the electric field intensity (electric charge density  $\rho$ ) equals:

$$\rho = \nabla \cdot \begin{pmatrix} e_r \\ e_\theta \\ e_\varphi \end{pmatrix} = \frac{\sqrt{2} K_1 \cot(\theta) \sin^2(\omega t) \sqrt{1 - \sin^4(\omega t)} e^{\frac{G_1 \varepsilon_0 \mu_0}{8 \pi r}}}{r^2 \sqrt{2 - 2 \sin^4(\omega t)}} \quad (60)$$

$$\rho = \nabla \cdot \begin{pmatrix} e_r \\ e_\theta \\ e_\varphi \end{pmatrix} = \frac{\frac{1}{2} K_1 \cot(\theta) e^{\frac{G_1 \varepsilon_0 \mu_0}{8 \pi r}}}{r^2} \quad (\text{averaged over 1 period of time})$$

The divergence of the magnetic field intensity (magnetic flux density  $\phi$ ) equals:

$$\phi = \nabla \cdot \begin{pmatrix} m_r \\ m_\theta \\ m_\varphi \end{pmatrix} = \frac{K_1 \sqrt{\varepsilon_0} \cot(\theta) \sqrt{2 - 2 \sin^4(\omega t)} e^{\frac{G_1 \varepsilon_0 \mu_0}{8 \pi r}}}{\sqrt{2} \sqrt{\mu_0} r^2} \quad (61)$$

$$\phi = \nabla \cdot \begin{pmatrix} m_r \\ m_\theta \\ m_\varphi \end{pmatrix} = \frac{K_1 \sqrt{\varepsilon_0} \cot(\theta) \sqrt{\frac{3}{4}} e^{\frac{G_1 \varepsilon_0 \mu_0}{8 \pi r}}}{\sqrt{\mu_0} r^2} \quad (\text{averaged over 1 period of time})$$

In which  $K_1$  is an arbitrary variable. Because of the  $\text{Cot}(\theta)$  function, the electric divergence as well as the magnetic divergence changes from sign when the angle  $\theta$  varies between  $0^0$  until  $360^0$  forming electric dipoles (+ versus -) and magnetic dipoles (N versus S).



## 2. The Illusion of Quantum Mechanical Probability Waves.

The physical concept of quantum mechanical probability waves has been created during the famous [1927 5<sup>th</sup> Solvay Conference](#). During that period there were several circumstances which came just together and made it possible to create a unique idea of material waves being complex (partly real and partly imaginary) and describing the probability of the appearance of a physical object (elementary particle).

The idea of complex (probability) waves is directly related to the concept of confined ([standing](#)) waves. Characteristic for any [standing](#) wave is the fact that the velocity and the pressure (electric field and magnetic field) are always shifted over 90 degrees. The same principle does exist for the [standing \(confined\) electromagnetic waves](#),

For that reason every confined (standing) Electromagnetic wave can be described by a complex sum vector  $\bar{\phi}$  of the Electric Field Vector  $\bar{E}$  and the Magnetic Field Vector  $\bar{B}$  ( $\bar{E}$  has 90 degrees phase shift compared to  $\bar{B}$ ).

The vector functions  $\bar{\phi}$  and the complex conjugated vector function  $\bar{\phi}^*$  will be written as:

$$\bar{\phi} = \frac{1}{\sqrt{2} \mu} \left( \bar{B} + i \frac{\bar{E}}{c} \right) \quad (62)$$

$\bar{B}$  equals the magnetic induction,  $\bar{E}$  the electric field intensity ( $\bar{E}$  has + 90 degrees phase shift compared to  $\bar{B}$ ) and  $c$  the speed of light.

The complex conjugated vector function equals:

$$\bar{\phi}^* = \frac{1}{\sqrt{2\mu}} \left( \bar{\mathbf{B}} - i \frac{\bar{\mathbf{E}}}{c} \right) \quad (63)$$

The dot product equals the electromagnetic energy density w:

$$\bar{\phi} \cdot \bar{\phi}^* = \frac{1}{2\mu} \left( \bar{\mathbf{B}} + i \frac{\bar{\mathbf{E}}}{c} \right) \cdot \left( \bar{\mathbf{B}} - i \frac{\bar{\mathbf{E}}}{c} \right) = \frac{1}{2} \mu H^2 + \frac{1}{2} \varepsilon E^2 = w \quad (64)$$

Using Einstein's equation  $W = m c^2$ , the dot product equals the electromagnetic mass density w

$$\bar{\phi} \cdot \bar{\phi}^* \frac{1}{c^2} = \frac{\varepsilon}{2} \left( \bar{\mathbf{B}} + i \frac{\bar{\mathbf{E}}}{c} \right) \cdot \left( \bar{\mathbf{B}} - i \frac{\bar{\mathbf{E}}}{c} \right) = \frac{1}{2} \varepsilon \mu^2 H^2 + \frac{1}{2} \varepsilon^2 E^2 = \rho \text{ [kg/m}^3] \quad (65)$$

The cross product is proportional to the Poynting vector (ref. 3, page 202, equation 15).

$$\bar{\phi} \times \bar{\phi}^* = \frac{1}{2\mu} \left( \bar{\mathbf{B}} + i \frac{\bar{\mathbf{E}}}{c} \right) \times \left( \bar{\mathbf{B}} - i \frac{\bar{\mathbf{E}}}{c} \right) = i \sqrt{\varepsilon\mu} \bar{\mathbf{E}} \times \bar{\mathbf{H}} = i \sqrt{\varepsilon\mu} \bar{\mathbf{S}} \quad (66)$$

Newton's second law of motion has been described in 3 spatial dimensions, resulting in the fundamental equation for the electromagnetic field.

3-Dimensional Space Domain

$$\begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} + \begin{matrix} \text{B-1} & \text{B-2} & \text{B-3} \\ -\frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \varepsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \varepsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \\ \text{B-4} & \text{B-5} & \text{B-6} \\ + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) + \frac{1}{2} (\varepsilon^2 \mu (\bar{\mathbf{E}} \cdot \bar{\mathbf{E}}) + \varepsilon \mu^2 (\bar{\mathbf{H}} \cdot \bar{\mathbf{H}})) \bar{\mathbf{g}} = \bar{\mathbf{0}} \end{matrix} \quad (67)$$

The formal mathematical way to describe the force density results from the 4-dimensional divergence of the 4-dimensional energy momentum tensor, resulting in a 4-dimensional Force vector. Dividing the 4-dimensional Force vector by the Volume results in the 4-dimensional force density vector.

The 4-dimensional Electromagnetic Vector Potential has been defined by:

$$\varphi^{-4} = \begin{pmatrix} \varphi_4 \\ \varphi_3 \\ \varphi_2 \\ \varphi_1 \end{pmatrix} \xrightarrow{\text{CartesianCoordinateSystem}} \begin{pmatrix} \varphi_t \\ \varphi_z \\ \varphi_y \\ \varphi_x \end{pmatrix} \quad (68)$$

In which the term  $\varphi_a$  represents the 4-dimensional electromagnetic vector potential in the “a” direction while the indice “a” varies from 1 to 4. In a cartesian coordinate system the indices are chosen varying from the x,y,z and t direction. In which the indice “t” represents the time direction which has been considered to be the 4<sup>th</sup> dimension. The 4-dimensional Electromagnetic “[Maxwell Tensor](#)” has been defined by:

$$F_{ab} = \partial_b \varphi_a - \partial_a \varphi_b \quad (69)$$

Where the indices “a” and “b” vary from 1 to 4.

The 4-dimensional Electromagnetic “[Energy Momentum Tensor](#)” has been defined by:

$$T^{ab} = \frac{I}{\mu_0} \left[ F_{ac} F^{cb} + \frac{I}{4} \delta_{ab} F_{cd} F^{cd} \right] \quad (70)$$

The 4-dimensional divergence of the 4-dimensional Energy Momentum Tensor equals the 4-dimensional Force Density

4-vector  $f^a$  :

$$f^a = \partial_b T^{ab} \quad (71)$$

Substituting the electromagnetic values for the electric field intensity “E” and the magnetic field intensity “H” in (71) results in the 4-dimensional representation of Newton’s second law of motion:

$$\begin{aligned} & \text{Energy-Time Domain} \\ & \text{B-7} \\ (f_4) \quad & \nabla \cdot (\bar{E} \times \bar{H}) + \frac{1}{2} \frac{\partial (\epsilon_0 (\bar{E} \cdot \bar{E}) + \mu_0 (\bar{H} \cdot \bar{H}))}{\partial t} = 0 \\ & \text{3-Dimensional Space Domain} \\ & \text{B-1} \qquad \qquad \text{B-2} \qquad \qquad \text{B-3} \qquad \qquad (72) \\ \left( \begin{array}{c} f_3 \\ f_2 \\ f_1 \end{array} \right) & - \frac{1}{c^2} \frac{\partial (\bar{E} \times \bar{H})}{\partial t} + \epsilon_0 \bar{E} (\nabla \cdot \bar{E}) - \epsilon_0 \bar{E} \times (\nabla \times \bar{E}) + \\ & \qquad \qquad \text{B-4} \qquad \qquad \text{B-5} \\ & + \mu_0 \bar{H} (\nabla \cdot \bar{H}) - \mu_0 \bar{H} \times (\nabla \times \bar{H}) = \bar{0} \end{aligned}$$

In which  $f_1, f_2, f_3$ , represent the force densities in the 3 spatial dimensions and  $f_4$  represent the force density (energy flow) in the time dimension (4<sup>th</sup> dimension).

The 4<sup>th</sup> term in equation (72) can be written in the terms of the Poynting vector “S” and the energy density “w” representing the electromagnetic law for the conservation of energy.

$$\begin{array}{c} \text{Energy-Time Domain} \\ \text{Inner Energy} \\ \text{B-7} \end{array} \quad \left( f_4 \right) \quad \nabla \cdot \bar{S} + \frac{\partial w}{\partial t} = 0 \quad (73.1)$$

$$\begin{array}{c} \text{3-Dimensional Space Domain} \\ \text{B-1} \quad \text{B-2} \quad \text{B-3} \\ \text{B-4} \quad \text{B-5} \end{array} \quad \left( \begin{array}{c} f_3 \\ f_2 \\ f_1 \end{array} \right) \quad - \frac{1}{c^2} \frac{\partial (\bar{E} \times \bar{H})}{\partial t} + \epsilon_0 \bar{E} (\nabla \cdot \bar{E}) - \epsilon_0 \bar{E} \times (\nabla \times \bar{E}) + \mu_0 \bar{H} (\nabla \cdot \bar{H}) - \mu_0 \bar{H} \times (\nabla \times \bar{H}) = \bar{0} \quad (73.2)$$

Substituting (64) and (66) in Equation (73) results in The 4-Dimensional Equilibrium Equation (74):

$$\left( x_4 \right) \quad - \frac{i}{\sqrt{\epsilon_0 \mu_0}} \nabla \cdot (\bar{\phi} \times \bar{\phi}) = - \frac{\partial \bar{\phi} \cdot \bar{\phi}^*}{\partial t} \quad (5.1.5)$$

$$\left( \begin{array}{c} x_3 \\ x_2 \\ x_1 \end{array} \right) \quad - \frac{1}{c^2} \frac{\partial (\bar{E} \times \bar{H})}{\partial t} + \epsilon_0 \bar{E} (\nabla \cdot \bar{E}) - \epsilon_0 \bar{E} \times (\nabla \times \bar{E}) + \mu_0 \bar{H} (\nabla \cdot \bar{H}) - \mu_0 \bar{H} \times (\nabla \times \bar{H}) = \bar{0} \quad (74) \quad (5.2)$$

To transform the electromagnetic vector wave function  $\bar{\phi}$  into a scalar (spinor or one-dimensional matrix

representation), the Pauli spin matrices  $\sigma$  and the following matrices (Ref. 3 page 213, equation 99) are introduced:

$$\bar{\alpha} = \begin{bmatrix} 0 & \sigma \\ \sigma & 0 \end{bmatrix} \quad \text{and} \quad \bar{\beta} = \begin{bmatrix} \delta_{ab} & 0 \\ 0 & -\delta_{ab} \end{bmatrix} \quad (75)$$

Then equation (74) can be written as the 4-Dimensional Hyperspace Equilibrium Dirac Equation:

$$(x_4) \quad \left( \frac{i m c}{h} \bar{\beta} + \bar{\alpha} \cdot \nabla \right) \psi = - \frac{1}{c} \frac{\partial \psi}{\partial t} \quad (76.1)$$

$$\begin{pmatrix} x_3 \\ x_2 \\ x_1 \end{pmatrix} \quad - \frac{1}{c^2} \frac{\partial (\bar{\mathbf{E}} \times \bar{\mathbf{H}})}{\partial t} + \epsilon_0 \bar{\mathbf{E}} (\nabla \cdot \bar{\mathbf{E}}) - \epsilon_0 \bar{\mathbf{E}} \times (\nabla \times \bar{\mathbf{E}}) + \quad (76) \\ + \mu_0 \bar{\mathbf{H}} (\nabla \cdot \bar{\mathbf{H}}) - \mu_0 \bar{\mathbf{H}} \times (\nabla \times \bar{\mathbf{H}}) = \bar{0}$$

The fourth term (x<sub>4</sub>) equals the [relativistic Dirac equation](#) (76.1) which equals equation (102) page 213 in [Ref.3](#).

Equation (76.1) represents the relativistic quantum mechanical Dirac Equation where  $\psi$  represents the quantum mechanical probability wave function. The mathematical evidence for the equivalent for (76.1) has been published in 1995 in the article: [“A Continuous Model of Matter based on AEONs”](#). Equation (1) page 201 to Equation (102) page 213. (Doi: [10.31219/osf.io/ra7ng](https://doi.org/10.31219/osf.io/ra7ng))

The Electromagnetic Law for the conservation of Energy (73.1) and the Relativistic Dirac Equation (76.1) are identical but written in a different form.

The law of conservation of Electromagnetic Energy can be written in an electromagnetic form (73.1) or in an identical way in a quantum mechanical form (76.1):

Energy-Time Domain

Inner Energy

B-7

$$\begin{matrix} (f_4) & \nabla \cdot \bar{S} + \frac{\partial w}{\partial t} & = 0 & (73.1) \end{matrix} \quad (77)$$

$$\begin{matrix} (x_4) & \left( \frac{i m c}{h} \bar{\beta} + \bar{\alpha} \cdot \nabla \right) \psi & = - \frac{1}{c} \frac{\partial \psi}{\partial t} & (76.1) \end{matrix}$$

## 2.1 Conclusions:

It follows from equation (23) that Newton's second law of motion applied to an electromagnetic field results in a force density equation which describes the electromagnetic field configuration completely.

From equation (37) follows that confined electromagnetic radiation carries mass

From equation (60) and (61) follows that confined electromagnetic radiation carries electric charge and magnetic spin.

These 3 facts result in the conclusion that quantum mechanical probability waves predicting the location of mass, charge and spin do not exist. Mass, electric charge and magnetic spin are carried by confined electromagnetic waves and are not positioned in local points but divided by the confined electromagnetic wave energy density (mass density), charge density and spin density.

Copenhagen Interpretation	New Theory
The Universe has been built out of elementary particles	The Universe has been built out of Confined Electromagnetic Field Configurations
Elementary Particles are the fundamental building elements in the Universe	Confined Electromagnetic Field Configurations are the fundamental building elements in the Universe
Fundamental properties of matter like mass, charge and spin are carried by elementary particles	Fundamental properties of matter like mass, charge and spin are carried by Confined Electromagnetic Field Configurations
Probability waves describe the location of particles	Probability Waves do not exist. Confined electromagnetic radiation carries electric charge and magnetic spin.
Probability waves are complex waves	Confined Electromagnetic waves are not complex. The phase shift of 90 degrees between the electric standing wave and the magnetic standing wave can be written in a complex function describing simultaneously the electric field and the magnetic field.
The product of the probability function $\psi$ and the complex conjugated function $\psi^*$ equals the probability	The dot product $\bar{\phi} \cdot \bar{\phi}^* = \rho$ in which $\rho$ equals the mass density of the confined electromagnetic radiation

## 1.10 Data Availability

All the Data and all the Calculations to provide evidence to this ‘New Theory about Light’ have been published in the ‘Open Source Framework(OSF)’: <https://osf.io/gbn4p/>

DOI: [10.31219/osf.io/gbn4p](https://doi.org/10.31219/osf.io/gbn4p) (<https://doi.org/10.31219/osf.io/gbn4p> )

(Calculations in Mathematica 11.0)’ , Page 1 – 33).

## 1.11 References

- [1] Oki Gunawana and Yudistira Virgus; The one-dimensional camelback potential in the parallel dipole line trap: Stability conditions and finite size effect; Journal of Applied Physics 121, 133902 (2017); <https://doi.org/10.1063/1.4978876>
- [2] IBM Blog Research; A new effect in electromagnetism discovered – 150 years later; <https://phys.org/news/2017-10-effect-electromagnetism-years.html>
- [3] J. W. Vejt, [A Continuous Model of Matter based on AEONs](#), Physics Essays ,1995, Volume 8, Number 2, 201-224 [A Continuous model of Matter](#) ( <https://doi.org/10.31219/osf.io/ra7ng> )
- [4] Z. K. Minev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R.J. Schoelkopf, N. Mirrahimi, H.J. Carmichael and M. H. Devoret; To catch and reverse a quantum jump mid-flight; Nature; 03 June 2019 equilibrium beyond Einstein 4-Dimensional, Kaluza-Klein 5-Dimensional and Superstring 10- and 11 Dimensional Curved Hyperspaces. <https://www.nature.com/articles/s41586-019-1287-z>