

Stéphane Wojnow

wojnow.stephane@gmail.com

19 October 2021

### Abstract

*We examine a quantum point of view about the origin of the universe. We propose a simple approach to cosmology based on the Planck mass flow rate. We try to provide a perspective and simple model on the evolution of the universe from Planck time in accordance with the  $\Lambda$ CDM model.*

Keywords : origin of universe, Planck time, Planck mass flow rate, evolution of universe.

### Introduction

The theory of quantum mechanics and the theory of general relativity always discuss the initial conditions of the universe and its evolution. Recently, Bruno Valeixo Bento and Stav Zalel in their paper « If time had no beginning », have provided new insights into this issue. We will try to add some simple considerations to this question through the mass of universe and of its density and to develop this with a simple cosmological model based on the Hubble time, Planck mass flow rate and a variable coefficient  $\alpha_H$ . Finally we examine the growth process of this model in accordance with the  $\Lambda$ CDM model.

#### 1) Quantum point of view on the origin of the universe.

It is remarkable that the energy density of the quantum matter resulting from  $m_p$ ,

$$m_p c^2 / l_p^3 = 4,63 * 10^{113} \text{ J/m}^3$$

is extremely close to the energy of the quantum vacuum of the quantum field theory :

$$l_p^{-2} \approx 3,83 * 10^{69} \text{ m}^{-2}$$

Indeed, with the Planck force,  $F_p (=c^4/G)$ , and the quantum vacuum energy of the quantum field theory, we obtain this quantum vacuum energy density :

$$F_p l_p^{-2} \approx 4,63 * 10^{113} \text{ J/m}^3$$

It would thus seem that the matter of the universe seen by the observer, would emerge naturally at its instant  $t_p$ , from a fluctuation of the energy of the quantum vacuum and reciprocally in a unit of Planck sphere volume. This could be a quantum solution to the origin of the universe. We would have, in the literal and mathematical sense, a division between the "mass" of the universe and its volume, i.e. the vacuum of the universe, from the Planck time of the observer.

#### 2) Beginning of a toy cosmological model under $\Lambda$ CDM model conditions after the recombination,

It seems possible to obtain the total mass of the universe from the  $\Lambda$ CDM model otherwise. This could eventually lead to the development of a simple toy cosmological model unknown to the author, built around the Hubble constant, the Hubble time,  $t_H = 1 / H$ , the Planck mass flow rate and a variable coefficient  $\alpha_H$ .

$\alpha_H =$  **radius of the observable universe** (from calculation of the  $\Lambda$ CDM model for example) divided by the **Hubble radius** at time  $t_H$  for a flat universe :

$$\alpha_H = \frac{c}{H_0} \int_{a=0}^{a=1} \frac{da}{a^2 \sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_\Lambda}} / \frac{c}{H_0}$$

$$\alpha_H = \int_{a=0}^{a=1} \frac{da}{a^2 \sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_k a^{-2} + \Omega_\Lambda}}$$

$$\delta = \frac{c^3}{G} = \frac{m_{Pl}}{t_{Pl}} \text{ is the Planck mass flow rate}$$

$t_H = 1/H$  is the Hubble time ( $\approx 4,56 \cdot 10^{17}$  s = 14,45 billion light years today)

$R_H = c / H = c t_H$  is the Hubble radius

The increase of the total "mass" (=energy) of the universe in the sense of the  $\Lambda$ CDM model is determined by the relation :

$$M_H = \rho_c V_H$$

$$M_H = \frac{3}{8\pi G t_H^2} \frac{4\pi}{3} (c t_H \alpha_H)^3$$

$$M_H = \frac{1}{2} \frac{c^3}{G} t_H \alpha_H^3$$

$$M_H = \frac{1}{2} \frac{m_{Pl}}{t_{Pl}} t_H \alpha_H^3$$

$$M_H = \frac{1}{2} \delta t_H \alpha_H^3$$

$\alpha_H \approx 46.12$  billion light years / 14.45 billion light years  $\approx 3.19$  today if  $H_0 = 67,66$  km/s/Mpc,  $\Omega_\Lambda = 0,6889$ .

i.e. for  $H = 67.66$  km/s/Mpc and  $\Omega_\Lambda = 0.6889$  :

$$M_H \approx 2,99 \cdot 10^{54} \text{ kg}$$

in other words, the total "mass" of the universe  $\Lambda$ CDM today.

### 3) Value of $\alpha_H$ before the recombination in the cosmological toy model and consequences.

The author hypothesises that, before the recombination, the radius of the observable universe was equal to the Hubble radius. The ratio  $\alpha_H$  was then equal to 1.

Thus, the mass of the universe at Planck time would be determined by :

$$M_H = \frac{1}{2} \frac{m_{Pl}}{t_{Pl}} t_{Pl}$$

$$M_U \text{ at } t_{Pl} = \frac{m_{Pl}}{2}$$

At Planck time, and at each subsequent "Planck time grain", the other half of the mass of the universe could be, as proposed in §1, the "quantum vacuum mass" from the QFT... equal to the mass of quantum matter, in a unit Planck sphere volume.

Indeed, the energy density of matter resulting from the mass  $m_{pl}$  is equal to the energy density of the quantum vacuum of the QFT, which can be calculated from the inverse square of the Planck length and  $c^4/G$ . They can therefore have an equal share in the mass of the universe at time  $t_p$ . i.e., a Planck sphere volume is composed of half a mass of Planck matter and half a mass of Planck vacuum.

Following the reasoning of figure 2 in "If time had no beginning", of Bruno Valeixo Bento and Stav Zalel , except that we consider that we have our « Planck time grain » instead an empty set at the beginning of set, with each passing unit of Planck time, the corresponding mass is added to the mass of the universe. In our cosmological toy model the "mass" of the universe at the Hubble radius, before and after decoupling, at time  $t_H$ , grows simply following the summation :

$$M_{U \text{ Hubble at } t_H} = \sum_1^{t_H/t_p} \frac{m_{Pl}}{2}$$

i.e.

$$M_{U \text{ Hubble at } t_H} = \frac{m_{Pl}}{2} \frac{t_H}{t_{Pl}}$$

$t_H = 1/H$  is the Hubble time.  $H_0 \approx 67,66 \text{ km/s/Mpc} \approx 4,56 \cdot 10^{17}$  seconds today, so  $M_{U \text{ Hubble}} \approx 9,21 \cdot 10^{52} \text{ kg}$

Note : ... and with datas of §2 ,  $M_{U \text{ Observable}}$  becomes  $\approx 3,19^3 M_{U \text{ Hubble}} \approx 2,99 \cdot 10^{54} \text{ kg}$ .

#### 4) Determination of the mass of the universe at Planck time in the $\Lambda$ CDM model if it can be apply.

We assume a flat universe, i.e. with zero curvature. For an observer, whose universe origin is at time  $t_p$ , the radius of its observable universe before the recombination in the  $\Lambda$ CDM model is  $= l_p (= c t_p)$ , hence its volume  $V_{p1}$  :

$$\frac{4\pi}{3} (l_p)^3 = 1,768 \cdot 10^{-104} m^3$$

Its critical density  $\rho_c$  expressed in  $\text{kg/m}^3$  is at time  $t_p$  :

$$\rho_c = \frac{3(Ht_p)^2}{8\pi G} = \frac{3}{8\pi G t_p^2} = 6,153 \cdot 10^{95} \text{ kg/m}^3$$

where  $H(t_p)$  is the Hubble constant at Planck time  $t_p = 1/H(t_p)$  and  $G$  the gravitational constant.

Under these conditions, the mass of the observable universe, at Planck time  $t_p$ , is also and of course, exactly  $1/2 m_{Pl}$ . This means that half of the universe is missing. This could be a solution to the problem of the disappearance of antimatter in the formation process of the universe.

## Conclusion

We have highlighted a succinct quantum approach to the origin of the universe.

We have tried to lay the foundations of a simple toy cosmology model allowing the calculation of a total "mass" of the universe that exactly matches that of the  $\Lambda$ CDM model at the same time  $t_H = 1 / H$ . This toy model is characterised by time, creating, at each Planck time of an observer, a specific observable universe, itself probably included in a larger infinite and eternal universe, which could have no beginning and no end, as proposed by Bruno Valeixo Bento and Stav Zalel.

We proposed a determination of the mass of the universe at Planck time in the framework of general relativity in agreement with the quantum density of matter and vacuum [  $= 1 / (G t_p^2)$  kg/m<sup>3</sup> ]. Then we proposed a solution to the problem of the disappearance of antimatter in the formation process of the universe.

## References :

Bruno Valeixo Bento and Stav Zalel, If time had no beginning, Sept 27, 2021 <https://arxiv.org/pdf/2109.11953.pdf>

For the value  $l_p^{-2} = 3,83 * 10^{69}$  m<sup>-2</sup> from the QFT

Lucas Lombriser, universit  de Gen ve, communiqu  de presse ,

<https://www.unige.ch/communication/communiqu s/2019/cosmologie-une-solution-a-la-pire-prediction-en-physique/>

David W. Hogg , Distance measures in cosmology (14) and (15), 2000 December,

<https://arxiv.org/pdf/astro-ph/9905116v4.pdf>

<https://www.dcode.fr/summation-calculator>