

# ON THE GENERAL ERDŐS-TURÁN ADDITIVE BASE CONJECTURE

T. AGAMA

ABSTRACT. In this paper we introduce a multivariate version of circles of partition introduced and studied in [1]. As an application we prove a weaker general version of the Erdős-Turán additive base conjecture. The actual Erdős-Turán additive base conjecture follows from this general version as a consequence.

## 1. Introduction

In [1] we have developed a method which we feel might be a valuable resource and a recipe for studying problems concerning partition of numbers in specified subsets of  $\mathbb{N}$ . The method of circles of partition is very vast and rich and has yet unexplored applications. This method is easy to use given its combinatorial affinity. It is very elementary in nature and has parallels with configurations of points on the geometric circle.

Let us suppose that for any  $n \in \mathbb{N}$  we can write  $n = u+v$  where  $u, v \in \mathbb{M} \subset \mathbb{N}$  then the new method associate each of this summands to points on the circle generated in a certain manner by  $n > 2$  and a line joining any such associated points on the circle. This geometric correspondence turns out to useful in our development, as the results obtained in this setting are then transformed back to results concerning the partition of integers.

In this paper we study a multivariate version of the method, where we allow our base regulators to be the direct product  $\otimes$  of subsets of the natural numbers  $\mathbb{N}$ . With the goal of studying a general version of the Erdős-Turán additive base conjecture, we introduce and study the notion of the **axial** potential of the multivariate circle of partition.

**Notations.** We denote by  $\mathbb{N}_n = \{m \in \mathbb{N} \mid m \leq n\}$  the **sequence** of the first  $n$  natural numbers

## 2. Multivariate circles of partition

In this section we introduce and study the notion of **multivariate** circles of partitions. We launch the following language.

---

*Date:* December 12, 2021.

*2010 Mathematics Subject Classification.* Primary 11P32 11A41,; Secondary 11B13, 11H99.

*Key words and phrases.* circle of partition, axes; generalized circles of partition; generalized density.

**Definition 2.1.** Let  $\mathbb{A} \subseteq \mathbb{N}$ . Then we denote with

$$\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i) = \left\{ [x_1], [x_2], \dots, [x_h] \mid x_i \in \mathbb{A}_i, n = \sum_{i=1}^h x_i \right\}$$

a **multivariate** circle of partition generated by  $n \in \mathbb{N}$  with base **regulators**  $\bigotimes_{i=1}^h \mathbb{A}_i$ . We call members of the multivariate CoP multivariate points.

**Definition 2.2.** We denote the line  $\mathbb{L}_{[x_1], [x_2], \dots, [x_h]}$  joining the points  $[x_1], [x_2], \dots, [x_h]$  as a **axis** of the multivariate CoP  $\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  if and only if  $x_i \in \mathbb{A}_i$  for each  $1 \leq i \leq h$  and  $n = \sum_{i=1}^h x_i$ . We say the axis points  $[x_i]$  for each  $1 \leq i \leq h$  are **axis residents**. We do not view the axis as any different among other axis  $\mathbb{L}_{[x_1], [x_2], \dots, [x_h]}$  upto the rearrangements of its residents points. In special cases where the points  $[x_k] \in \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  such that  $hx_k = n$  then we call  $[x_i]$  the **center** of the multivariate CoP. If it exists, then we call it as a **degenerated axis**  $\mathbb{L}_{[x_k]}$  in comparison to the **real axes**  $\mathbb{L}_{[x_1], [x_2], \dots, [x_h]}$ , where not all of the weights  $x_i$  can be equal. We denote the assignment of an axis  $\mathbb{L}_{[x_1], [x_2], \dots, [x_h]}$  to the multivariate CoP  $\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  as

$$\mathbb{L}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i) \text{ which means } [x_1], [x_2], \dots, [x_h] \in \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$$

with

$$n = \sum_{i=1}^h x_i$$

for a fixed  $n \in \mathbb{N}$  with  $x_i \in \mathbb{A}_i$  for each  $1 \leq i \leq h$  or vice versa and the number of real axes of the generalized CoP as

$$\nu(\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)) := \#\{\mathbb{L}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i) \mid x_i \neq x_j\}.$$

for all  $1 < i < j \leq h$ . The lines  $\mathcal{L}_{[x_1], [x_2], \dots, [x_h]}$  joining any  $h$  arbitrary points  $[x_1], [x_2], \dots, [x_h] \in \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  which are not resident points in the multivariate CoP will be referred to as a **graph** induced by the multivariate CoP.

Throughout this paper we will denote for simplicity the multivariate circle of partition in simple wording as m-CoP. The notion of a multivariate axis is not technically convenient to work with; nonetheless it is fairly manageable if we confine ourselves to a certain class of axis of a typical CoP. As it will prove very useful in the sequel and will feature very greatly in our results in the sequel, we find it more prudent exploiting the notion of representative axis.

**Definition 2.3.** Let  $\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  be a multivariate CoP and let  $[x_1] \in \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  be a fixed point. Then we say the axis  $\mathbb{L}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A}_i)$  belongs to the class  $m$  axis of the multivariate CoP if

$$x_2 + \dots + x_h = m.$$

**Proposition 2.4.** *Let  $\mathcal{C}(n, \otimes_{i=1}^h \mathbb{N})$  be a multivariate CoP. Then there are*

$$\left\lfloor \frac{n-1}{h} \right\rfloor$$

*axis-classes of the multivariate CoP.*

Throughout this paper we will work within the axis-classes and use their representatives. For any  $s$  axis-class of a multivariate CoP  $\mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)$

$$\mathcal{C}_s := \{\mathbb{L}_{[x_1], [x_2], \dots, [x_h]}^s \hat{\in} \mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)\}$$

we denote the representative axis of the class as  $\mathfrak{R}(\mathcal{C}_s)$ . Henceforth in counting the number of axis of a typical CoP we will only count the number of representative axis or simply the number of axis-classes. We denote more generally the set of all representative axis of the axis-classes as

$$\{\mathfrak{RL}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)\}$$

and the number of all representative axis in the m-CoP as

$$\#\{\mathfrak{RL}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)\}.$$

It has been observed that for a CoP with a natural number  $\otimes_{i=1}^h \mathbb{N}$  base regulator the number of representative axis is basically the quantity

$$\left\lfloor \frac{n-1}{h} \right\rfloor.$$

### 3. Axial potential of multivariate circles of partition

In this section we introduce and study the notion of the **axial potential** of an m-CoP. We launch the following language.

**Definition 3.1.** Let  $\mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)$  be a m-CoP. Then by the  $k$  th **axial potential** denoted,  $[\mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)]^k$ , we mean the infinite sum

$$[\mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)]^k = \sum_{n=1}^{\infty} \frac{\#\left\{\mathfrak{RL}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i)\right\}^k}{\#\left\{\mathfrak{RL}_{[x_1], [x_2], \dots, [x_h]} \hat{\in} \mathcal{C}(n, \otimes_{i=1}^h \mathbb{A}_i \cup \otimes_{i=1}^h \mathbb{N})\right\}^k}.$$

We say the  $k$  th axial potential is finite if the series converges; otherwise, we say it diverges.

It is known that for any additive base  $\mathbb{A}$  of order  $h$  with  $h \geq 2$  the quantity

$$\#\{n \leq x \mid n \in \mathbb{A}\} \geq x^{\frac{1}{h}}.$$

Using this fact, we then obtain a proof of the weaker variant of the general Erdős-Turán additive base conjecture in the form below

**Theorem 3.2.** *Let  $\mathbb{A} \subset \mathbb{N}$  and suppose  $\#\left\{\mathbb{L}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\} > 0$  for all sufficiently large values of  $n$ . If  $|\mathbb{A} \cap \mathbb{N}_n| \geq n^{1-\epsilon}$  for any  $0 < \epsilon \leq \frac{1}{h}$  then*

$$\lim_{n \rightarrow \infty} \#\left\{\sum_{i=1}^h x_i \mid \sum_{i=1}^h x_i = n, x_i \in \mathbb{A}, 1 \leq i \leq h\right\} = \infty.$$

*Proof.* Under the requirement  $\mathbb{A} \subset \mathbb{N}$  and  $\#\left\{\mathbb{L}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\} > 0$  for all sufficiently large values of  $n$  then it implies that

$$\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\} > 0$$

for all sufficiently large values of  $n$  so that there exists some constant  $\mathcal{P} := \mathcal{P}(k) > 0$  such that we can write

$$\begin{aligned} \sum_{n=1}^k \frac{\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\}^h}{\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{N})\right\}^h} &= \mathcal{P} \sum_{n=1}^k \frac{\lfloor \frac{n^{1-\epsilon}-1}{h} \rfloor^h}{\lfloor \frac{n-1}{h} \rfloor^h} \\ &\gg_k \sum_{n=1}^k \frac{\frac{n^{h-h\epsilon}}{h^h}}{\lfloor \frac{n-1}{h} \rfloor^h} \end{aligned}$$

since  $\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{N})\right\} = \lfloor \frac{n-1}{h} \rfloor$ , so that we can compute the  $h^{\text{th}}$  axial potential

$$\begin{aligned} [\mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})]^h &= \sum_{n=1}^{\infty} \frac{\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\}^h}{\#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{N})\right\}^h} \\ &\gg \sum_{n=1}^{\infty} \frac{\frac{n^{h-h\epsilon}}{h^h}}{\lfloor \frac{n-1}{h} \rfloor^h} \\ &\gg \sum_{n=1}^{\infty} \frac{1}{n^{h\epsilon}} = \infty \end{aligned}$$

since  $0 \leq \epsilon \leq \frac{1}{h}$ . It follows that

$$\lim_{n \rightarrow \infty} \#\left\{\mathfrak{RL}_{[x_1],[x_2],\dots,[x_h]} \hat{\in} \mathcal{C}(n, \bigotimes_{i=1}^h \mathbb{A})\right\} = \infty$$

and it implies that

$$\lim_{n \rightarrow \infty} \#\left\{\sum_{i=1}^h x_i \mid \sum_{i=1}^h x_i = n, x_i \in \mathbb{A}, 1 \leq i \leq h\right\} = \infty.$$

This completes the proof.  $\square$

**Corollary 3.3.** *Let  $\mathbb{A} \subset \mathbb{N}$  and suppose  $\#\left\{\mathbb{L}_{[x_1],[x_2]} \in \mathcal{C}(n, \otimes_{i=1}^2 \mathbb{A})\right\} > 0$  for all sufficiently large values of  $n$ . If  $|\mathbb{A} \cap \mathbb{N}_n| \geq n^{1-\epsilon}$  for any  $0 < \epsilon \leq \frac{1}{2}$  then*

$$\lim_{n \rightarrow \infty} \#\left\{\sum_{i=1}^2 x_i \mid \sum_{i=1}^2 x_i = n, x_i \in \mathbb{A}, 1 \leq i \leq 2\right\} = \infty.$$

*Proof.* This follows by taking  $h = 2$  in Theorem 3.2.  $\square$

*Remark 3.4.* Theorem 3.2 does indicate that notion of the axial potential cannot be readily harnessed to tackle the actual general version of the Erdős-Turán additive base conjecture, since we are constraint by the inequality  $1 \leq \epsilon \leq \frac{1}{h}$ . We would rather need the larger domain  $1 \leq \epsilon \leq 1 - \frac{1}{h}$  as a requirement but then we will certainly ran into a deadlock since the series

$$\sum_{n=1}^{\infty} \frac{1}{n^h}$$

converges for  $h > 1$  so that we cannot decide on the limiting value of the quantity

$$\#\left\{\sum_{i=1}^h x_i \mid \sum_{i=1}^h x_i = n, x_i \in \mathbb{A}, 1 \leq i \leq h\right\}$$

---

#### REFERENCES

1. Agama, Theophilus and Gensel, Berndt *Studies in Additive Number Theory by Circles of Partition*, arXiv:2012.01329, 2020.

DEPARTMENT OF MATHEMATICS, AFRICAN INSTITUTE FOR MATHEMATICAL SCIENCES, GHANA.  
*E-mail address:* Theophilus@aims.edu.gh/emperordagama@yahoo.com