

# Proof of Riemann hypothesis

By Toshihiko ISHIWATA

Jan. 14, 2023

**Abstract.** This paper is a trial to prove Riemann hypothesis according to the following process. 1. We make one identity regarding  $x$  from one equation that gives Riemann zeta function  $\zeta(s)$  analytic continuation and 2 formulas  $(1/2 + a \pm bi, 1/2 - a \pm bi)$  that show non-trivial zero point of  $\zeta(s)$ . 2. We find that the above identity holds only at  $a = 0$ . 3. Therefore non-trivial zero points of  $\zeta(s)$  must be  $1/2 \pm bi$  because  $a$  cannot have any value but zero.

## 1. Introduction

The following (1) gives Riemann zeta function  $\zeta(s)$  analytic continuation to  $0 < Re(s)$ . “+ . . . . .” means infinite series in all equations in this paper.

$$1 - 2^{-s} + 3^{-s} - 4^{-s} + 5^{-s} - 6^{-s} + \dots = (1 - 2^{1-s})\zeta(s) \quad (1)$$

The following (2) shows the zero point of the left side of (1) and also non-trivial zero point of  $\zeta(s)$ .  $i$  is  $\sqrt{-1}$ .

$$S_0 = 1/2 + a \pm bi \quad (2)$$

The following (3) also shows non-trivial zero point of  $\zeta(s)$  by the functional equation of  $\zeta(s)$ .

$$S_1 = 1 - S_0 = 1/2 - a \mp bi \quad (3)$$

We define the range of  $a$  and  $b$  as  $0 \leq a < 1/2$  and  $14 < b$  respectively. Then we can show all non-trivial zero points of  $\zeta(s)$  by the above (2) and (3). Because non-trivial zero points of  $\zeta(s)$  exist in the critical strip of  $\zeta(s)$  ( $0 < Re(s) < 1$ ) and non-trivial zero points of  $\zeta(s)$  found until now exist in the range of  $14 < b$ .

We have the following (4) and (5) by substituting  $S_0$  for  $s$  in the left side of (1) and putting both the real part and the imaginary part of the left side of (1) at zero respectively.

$$1 = \frac{\cos(b \log 2)}{2^{1/2+a}} - \frac{\cos(b \log 3)}{3^{1/2+a}} + \frac{\cos(b \log 4)}{4^{1/2+a}} - \frac{\cos(b \log 5)}{5^{1/2+a}} + \dots \quad (4)$$

$$0 = \frac{\sin(b \log 2)}{2^{1/2+a}} - \frac{\sin(b \log 3)}{3^{1/2+a}} + \frac{\sin(b \log 4)}{4^{1/2+a}} - \frac{\sin(b \log 5)}{5^{1/2+a}} + \dots \quad (5)$$

We also have the following (6) and (7) by substituting  $S_1$  for  $s$  in the left side of (1) and putting both the real part and the imaginary part of the left side of (1) at zero

respectively.

$$1 = \frac{\cos(b \log 2)}{2^{1/2-a}} - \frac{\cos(b \log 3)}{3^{1/2-a}} + \frac{\cos(b \log 4)}{4^{1/2-a}} - \frac{\cos(b \log 5)}{5^{1/2-a}} + \dots \quad (6)$$

$$0 = \frac{\sin(b \log 2)}{2^{1/2-a}} - \frac{\sin(b \log 3)}{3^{1/2-a}} + \frac{\sin(b \log 4)}{4^{1/2-a}} - \frac{\sin(b \log 5)}{5^{1/2-a}} + \dots \quad (7)$$

## 2. The identity regarding $x$

We define  $f(n)$  as follows.

$$f(n) = \frac{1}{n^{1/2-a}} - \frac{1}{n^{1/2+a}} \geq 0 \quad (n = 2, 3, 4, 5, \dots) \quad (8)$$

We have the following (9) from the above (4) and (6) with the method shown in item 1.1 of [Appendix 1: Equation construction].

$$0 = f(2) \cos(b \log 2) - f(3) \cos(b \log 3) + f(4) \cos(b \log 4) - f(5) \cos(b \log 5) + \dots \quad (9)$$

We also have the following (10) from the above (5) and (7) with the method shown in item 1.2 of [Appendix 1].

$$0 = f(2) \sin(b \log 2) - f(3) \sin(b \log 3) + f(4) \sin(b \log 4) - f(5) \sin(b \log 5) + \dots \quad (10)$$

We can have the following (11) regarding real number  $x$  from the above (9) and (10) with the method shown in item 1.3 of [Appendix 1]. And the value of (11) is always zero at any value of  $x$ .

$$\begin{aligned} 0 &\equiv \cos x \{\text{the right side of (9)}\} + \sin x \{\text{the right side of (10)}\} \\ &= \cos x \{f(2) \cos(b \log 2) - f(3) \cos(b \log 3) + f(4) \cos(b \log 4) - \dots\} \\ &\quad + \sin x \{f(2) \sin(b \log 2) - f(3) \sin(b \log 3) + f(4) \sin(b \log 4) - \dots\} \\ &= f(2) \cos(b \log 2 - x) - f(3) \cos(b \log 3 - x) + f(4) \cos(b \log 4 - x) \\ &\quad - f(5) \cos(b \log 5 - x) + f(6) \cos(b \log 6 - x) - \dots \end{aligned} \quad (11)$$

At  $a = 0$  we have the following (8-1) and the above (11) holds at  $a = 0$ .

$$f(n) = \frac{1}{n^{1/2-a}} - \frac{1}{n^{1/2+a}} \equiv 0 \quad (n = 2, 3, 4, 5, \dots) \quad (8-1)$$

We have the following (12-1) by substituting  $b \log 1$  for  $x$  in (11).

$$\begin{aligned} 0 &= f(2) \cos(b \log 2 - b \log 1) - f(3) \cos(b \log 3 - b \log 1) + f(4) \cos(b \log 4 - b \log 1) \\ &\quad - f(5) \cos(b \log 5 - b \log 1) + f(6) \cos(b \log 6 - b \log 1) - \dots \end{aligned} \quad (12-1)$$

We have the following (12-2) by substituting  $b \log 2$  for  $x$  in (11).

$$\begin{aligned} 0 &= f(2) \cos(b \log 2 - b \log 2) - f(3) \cos(b \log 3 - b \log 2) + f(4) \cos(b \log 4 - b \log 2) \\ &\quad - f(5) \cos(b \log 5 - b \log 2) + f(6) \cos(b \log 6 - b \log 2) - \dots \end{aligned} \quad (12-2)$$

We have the following (12-3) by substituting  $b \log 3$  for  $x$  in (11).

$$0 = f(2) \cos(b \log 2 - b \log 3) - f(3) \cos(b \log 3 - b \log 3) + f(4) \cos(b \log 4 - b \log 3) \\ - f(5) \cos(b \log 5 - b \log 3) + f(6) \cos(b \log 6 - b \log 3) - \dots \quad (12-3)$$

In the same way as above we can have the following (12-N) by substituting  $b \log N$  for  $x$  in (11). ( $N = 4, 5, 6, 7, \dots$ )

$$0 = f(2) \cos(b \log 2 - b \log N) - f(3) \cos(b \log 3 - b \log N) + f(4) \cos(b \log 4 - b \log N) \\ - f(5) \cos(b \log 5 - b \log N) + f(6) \cos(b \log 6 - b \log N) - \dots \quad (12-N)$$

### 3. The solution for the identity of (11)

We define  $g(k, N)$  as follows. ( $k = 2, 3, 4, 5, \dots$   $N = 1, 2, 3, 4, \dots$ )

$$g(k, N) = \cos(b \log k - b \log 1) + \cos(b \log k - b \log 2) + \cos(b \log k - b \log 3) + \dots + \cos(b \log k - b \log N) \\ = \cos(b \log 1 - b \log k) + \cos(b \log 2 - b \log k) + \cos(b \log 3 - b \log k) + \dots + \cos(b \log N - b \log k) \\ = \cos(b \log 1/k) + \cos(b \log 2/k) + \cos(b \log 3/k) + \dots + \cos(b \log N/k) \quad (13)$$

We can have the following (14) from the equations of (12-1), (12-2), (12-3),  $\dots$ , (12-N) with the method shown in item 1.4 of [Appendix 1].

$$0 = f(2)\{\cos(b \log 2 - b \log 1) + \cos(b \log 2 - b \log 2) + \cos(b \log 2 - b \log 3) + \dots + \cos(b \log 2 - b \log N)\} \\ - f(3)\{\cos(b \log 3 - b \log 1) + \cos(b \log 3 - b \log 2) + \cos(b \log 3 - b \log 3) + \dots + \cos(b \log 3 - b \log N)\} \\ + f(4)\{\cos(b \log 4 - b \log 1) + \cos(b \log 4 - b \log 2) + \cos(b \log 4 - b \log 3) + \dots + \cos(b \log 4 - b \log N)\} \\ - f(5)\{\cos(b \log 5 - b \log 1) + \cos(b \log 5 - b \log 2) + \cos(b \log 5 - b \log 3) + \dots + \cos(b \log 5 - b \log N)\} \\ + \dots \\ = f(2)g(2, N) - f(3)g(3, N) + f(4)g(4, N) - f(5)g(5, N) + \dots \quad (14)$$

If (11) holds, the sum of the right sides of infinite number equations of (12-1), (12-2), (12-3), (12-4), (12-5),  $\dots$  becomes zero. The rightmost side of (14) is the sum of the right sides of  $N$  equations of (12-1), (12-2), (12-3),  $\dots$ , (12-N) as shown in item 1.4 of [Appendix 1]. Therefore if (11) holds,  $\lim_{N \rightarrow \infty} \{\text{the rightmost side of (14)}\} = 0$  must hold. Here we define  $F(a)$  as follows.

$$F(a) = f(2) - f(3) + f(4) - f(5) + \dots \quad (15)$$

We have the following (25) in [Appendix 2 : Investigation of  $g(k, N)$ ].

$$g(k, N) \sim \frac{N \cos(b \log N)}{\sqrt{1 + b^2}} \quad (N \rightarrow \infty \quad k = 2, 3, 4, 5, \dots) \quad (25)$$

From the above (15) and (25) we have the following (16).

$$\begin{aligned}
& \text{The rightmost side of (14)} \\
&= f(2)g(2, N) - f(3)g(3, N) + f(4)g(4, N) - f(5)g(5, N) + \dots \\
&\sim f(2)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} - f(3)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} + f(4)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} \\
&\quad - f(5)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} + \dots \\
&= \frac{N \cos(b \log N)}{\sqrt{1+b^2}} \{f(2) - f(3) + f(4) - f(5) + \dots\} \\
&= F(a)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} \quad (N \rightarrow \infty) \tag{16}
\end{aligned}$$

We have the following (17) by summarizing the above (16).

$$\text{The rightmost side of (14)} \sim F(a)\frac{N \cos(b \log N)}{\sqrt{1+b^2}} \quad (N \rightarrow \infty) \tag{17}$$

$\lim_{N \rightarrow \infty} \frac{N \cos(b \log N)}{\sqrt{1+b^2}}$  diverges to  $\pm\infty$ .  $0 < F(a)$  holds in  $0 < a < 1/2$  as shown in [Appendix 3 : Investigation of  $F(a)$ ]. Then  $\lim_{N \rightarrow \infty} \{\text{the rightmost side of (14)}\}$  diverges to  $\pm\infty$  in  $0 < a < 1/2$  from the above (17) i.e. (11) does not hold in  $0 < a < 1/2$ . (11) holds at  $a = 0$  as shown in item 2. Therefore the solution for the identity of (11) is only  $a = 0$ .

#### 4. Conclusion

$a$  has the range of  $0 \leq a < 1/2$  by the critical strip of  $\zeta(s)$ . However,  $a$  cannot have any value but zero as shown in the above item 3. Therefore non-trivial zero point of Riemann zeta function  $\zeta(s)$  shown by (2) and (3) must be  $1/2 \pm bi$ .

**Appendix 1. : Equation construction**

We can construct (9), (10), (11) and (14) by applying the following Theorem 1[1].

Theorem 1

If the following (Series 1) and (Series 2) converge respectively, the following (Series 3) and (Series 4) hold.

$$\text{(Series 1)} = a_1 + a_2 + a_3 + a_4 + a_5 + \dots = A$$

$$\text{(Series 2)} = b_1 + b_2 + b_3 + b_4 + b_5 + \dots = B$$

$$\text{(Series 3)} = (a_1 + b_1) + (a_2 + b_2) + (a_3 + b_3) + (a_4 + b_4) + \dots = A + B$$

$$\text{(Series 4)} = (a_1 - b_1) + (a_2 - b_2) + (a_3 - b_3) + (a_4 - b_4) + \dots = A - B$$

**1.1. Construction of (9)**

We can have (9) as (Series 4) by regarding (6) and (4) as (Series 1) and (Series 2) respectively.

**1.2. Construction of (10)**

We can have (10) as (Series 4) by regarding (7) and (5) as (Series 1) and (Series 2) respectively.

**1.3. Construction of (11)**

We can have (11) as (Series 3) by regarding the following (11-1) and (11-2) as (Series 1) and (Series 2) respectively.

$$\text{(Series 1)} = \cos x\{\text{the right side of (9)}\} \equiv 0 \tag{11-1}$$

$$\text{(Series 2)} = \sin x\{\text{the right side of (10)}\} \equiv 0 \tag{11-2}$$

**1.4. Construction of (14)**

1.4.1 We can have the following (12-1\*2) as (Series 3) by regarding the following (12-1) and (12-2) as (Series 1) and (Series 2) respectively.

$$\begin{aligned} \text{(Series 1)} = & f(2) \cos(b \log 2 - b \log 1) - f(3) \cos(b \log 3 - b \log 1) \\ & + f(4) \cos(b \log 4 - b \log 1) - f(5) \cos(b \log 5 - b \log 1) \\ & + f(6) \cos(b \log 6 - b \log 1) - \dots = 0 \end{aligned} \tag{12-1}$$

$$\begin{aligned} \text{(Series 2)} = & f(2) \cos(b \log 2 - b \log 2) - f(3) \cos(b \log 3 - b \log 2) \\ & + f(4) \cos(b \log 4 - b \log 2) - f(5) \cos(b \log 5 - b \log 2) \\ & + f(6) \cos(b \log 6 - b \log 2) - \dots = 0 \end{aligned} \tag{12-2}$$

$$\begin{aligned} \text{(Series 3)} = & f(2)\{\cos(b \log 2 - b \log 1) + \cos(b \log 2 - b \log 2)\} \\ & - f(3)\{\cos(b \log 3 - b \log 1) + \cos(b \log 3 - b \log 2)\} \\ & + f(4)\{\cos(b \log 4 - b \log 1) + \cos(b \log 4 - b \log 2)\} \\ & - f(5)\{\cos(b \log 5 - b \log 1) + \cos(b \log 5 - b \log 2)\} \\ & + \dots = 0 + 0 \end{aligned} \tag{12-1*2}$$

1.4.2 We can have the following (12-1\*3) as (Series 3) by regarding the above (12-1\*2) and the following (12-3) as (Series 1) and (Series 2) respectively.

$$\begin{aligned} \text{(Series 2)} &= f(2) \cos(b \log 2 - b \log 3) - f(3) \cos(b \log 3 - b \log 3) \\ &\quad + f(4) \cos(b \log 4 - b \log 3) - f(5) \cos(b \log 5 - b \log 3) \\ &\quad + f(6) \cos(b \log 6 - b \log 3) - \dots = 0 \end{aligned} \quad (12-3)$$

$$\begin{aligned} \text{(Series 3)} &= f(2) \{ \cos(b \log 2 - b \log 1) + \cos(b \log 2 - b \log 2) + \cos(b \log 2 - b \log 3) \} \\ &\quad - f(3) \{ \cos(b \log 3 - b \log 1) + \cos(b \log 3 - b \log 2) + \cos(b \log 3 - b \log 3) \} \\ &\quad + f(4) \{ \cos(b \log 4 - b \log 1) + \cos(b \log 4 - b \log 2) + \cos(b \log 4 - b \log 3) \} \\ &\quad - f(5) \{ \cos(b \log 5 - b \log 1) + \cos(b \log 5 - b \log 2) + \cos(b \log 5 - b \log 3) \} \\ &\quad + \dots = 0 + 0 \end{aligned} \quad (12-1*3)$$

1.4.3 We can have the following (12-1\*4) as (Series 3) by regarding the above (12-1\*3) and the following (12-4) as (Series 1) and (Series 2) respectively.

$$\begin{aligned} \text{(Series 2)} &= f(2) \cos(b \log 2 - b \log 4) - f(3) \cos(b \log 3 - b \log 4) \\ &\quad + f(4) \cos(b \log 4 - b \log 4) - f(5) \cos(b \log 5 - b \log 4) \\ &\quad + f(6) \cos(b \log 6 - b \log 4) - \dots = 0 \end{aligned} \quad (12-4)$$

$$\begin{aligned} \text{(Series 3)} &= f(2) \{ \cos(b \log 2 - b \log 1) + \cos(b \log 2 - b \log 2) + \cos(b \log 2 - b \log 3) + \cos(b \log 2 - b \log 4) \} \\ &\quad - f(3) \{ \cos(b \log 3 - b \log 1) + \cos(b \log 3 - b \log 2) + \cos(b \log 3 - b \log 3) + \cos(b \log 3 - b \log 4) \} \\ &\quad + f(4) \{ \cos(b \log 4 - b \log 1) + \cos(b \log 4 - b \log 2) + \cos(b \log 4 - b \log 3) + \cos(b \log 4 - b \log 4) \} \\ &\quad - f(5) \{ \cos(b \log 5 - b \log 1) + \cos(b \log 5 - b \log 2) + \cos(b \log 5 - b \log 3) + \cos(b \log 5 - b \log 4) \} \\ &\quad + \dots = 0 + 0 \end{aligned} \quad (12-1*4)$$

1.4.4 In the same way as above we can have the following (12-1\*N)=(14) as (Series 3) by regarding (12-1\*N-1) and (12-N) as (Series 1) and (Series 2) respectively. ( $N = 5, 6, 7, 8, \dots$ )  $g(k, N)$  is defined in page 3. ( $k = 2, 3, 4, 5, \dots$ )

$$\begin{aligned} \text{(Series 3)} &= \\ &f(2) \{ \cos(b \log 2 - b \log 1) + \cos(b \log 2 - b \log 2) + \cos(b \log 2 - b \log 3) + \dots + \cos(b \log 2 - b \log N) \} \\ &- f(3) \{ \cos(b \log 3 - b \log 1) + \cos(b \log 3 - b \log 2) + \cos(b \log 3 - b \log 3) + \dots + \cos(b \log 3 - b \log N) \} \\ &+ f(4) \{ \cos(b \log 4 - b \log 1) + \cos(b \log 4 - b \log 2) + \cos(b \log 4 - b \log 3) + \dots + \cos(b \log 4 - b \log N) \} \\ &- f(5) \{ \cos(b \log 5 - b \log 1) + \cos(b \log 5 - b \log 2) + \cos(b \log 5 - b \log 3) + \dots + \cos(b \log 5 - b \log N) \} \\ &+ \dots \\ &= f(2)g(2, N) - f(3)g(3, N) + f(4)g(4, N) - f(5)g(5, N) + \dots \\ &= 0 + 0 \end{aligned} \quad (12-1*N)$$

**Appendix 2. : Investigation of  $g(k, N)$** 

2.1 We define  $G$  and  $H$  as follows. ( $N = 1, 2, 3, 4, \dots$ )

$$\begin{aligned} G &= \lim_{N \rightarrow \infty} \frac{1}{N} \left\{ \cos(b \log \frac{1}{N}) + \cos(b \log \frac{2}{N}) + \cos(b \log \frac{3}{N}) + \dots + \cos(b \log \frac{N}{N}) \right\} \\ &= \int_0^1 \cos(b \log x) dx \end{aligned} \quad (20-1)$$

$$\begin{aligned} H &= \lim_{N \rightarrow \infty} \frac{1}{N} \left\{ \sin(b \log \frac{1}{N}) + \sin(b \log \frac{2}{N}) + \sin(b \log \frac{3}{N}) + \dots + \sin(b \log \frac{N}{N}) \right\} \\ &= \int_0^1 \sin(b \log x) dx \end{aligned} \quad (20-2)$$

We calculate  $G$  and  $H$  by Integration by parts.

$$\begin{aligned} G &= [x \cos(b \log x)]_0^1 + bH = 1 + bH \\ H &= [x \sin(b \log x)]_0^1 - bG = -bG \end{aligned}$$

Then we can have the values of  $G$  and  $H$  from the above equations as follows.

$$G = \frac{1}{1 + b^2} \quad H = \frac{-b}{1 + b^2} \quad (21)$$

2.2 We define  $E_c(N)$  and  $E_s(N)$  as follows.

$$\frac{\cos(b \log \frac{1}{N}) + \cos(b \log \frac{2}{N}) + \cos(b \log \frac{3}{N}) + \dots + \cos(b \log \frac{N}{N})}{N} - G = E_c(N) \quad (22-1)$$

$$\frac{\sin(b \log \frac{1}{N}) + \sin(b \log \frac{2}{N}) + \sin(b \log \frac{3}{N}) + \dots + \sin(b \log \frac{N}{N})}{N} - H = E_s(N) \quad (22-2)$$

From (20-1), (20-2), (22-1) and (22-2) we have the following (23).

$$\lim_{N \rightarrow \infty} E_c(N) = 0 \quad \lim_{N \rightarrow \infty} E_s(N) = 0 \quad (23)$$

2.3 From (13) we can calculate  $g(k, N)$  as follows. ( $N = 1, 2, 3, 4, \dots$ )

$$\begin{aligned} g(k, N) &= \cos(b \log 1/k) + \cos(b \log 2/k) + \cos(b \log 3/k) + \dots + \cos(b \log N/k) \\ &= N \frac{1}{N} \left\{ \cos(b \log \frac{1}{N} \frac{N}{k}) + \cos(b \log \frac{2}{N} \frac{N}{k}) + \cos(b \log \frac{3}{N} \frac{N}{k}) + \dots + \cos(b \log \frac{N}{N} \frac{N}{k}) \right\} \\ &= N \frac{1}{N} \left\{ \cos(b \log \frac{1}{N} + b \log \frac{N}{k}) + \cos(b \log \frac{2}{N} + b \log \frac{N}{k}) \right. \\ &\quad \left. + \cos(b \log \frac{3}{N} + b \log \frac{N}{k}) + \dots + \cos(b \log \frac{N}{N} + b \log \frac{N}{k}) \right\} \\ &= N \frac{1}{N} \cos(b \log \frac{N}{k}) \left\{ \cos(b \log \frac{1}{N}) + \cos(b \log \frac{2}{N}) + \cos(b \log \frac{3}{N}) + \dots + \cos(b \log \frac{N}{N}) \right\} \\ &\quad - N \frac{1}{N} \sin(b \log \frac{N}{k}) \left\{ \sin(b \log \frac{1}{N}) + \sin(b \log \frac{2}{N}) + \sin(b \log \frac{3}{N}) + \dots + \sin(b \log \frac{N}{N}) \right\} \\ &= N \cos(b \log \frac{N}{k}) G \end{aligned}$$

$$\begin{aligned}
& +N \cos(b \log \frac{N}{k}) \left\{ \frac{\cos(b \log 1/N) + \cos(b \log 2/N) + \cos(b \log 3/N) + \cdots + \cos(b \log N/N)}{N} - G \right\} \\
& -N \sin(b \log \frac{N}{k}) H \\
& -N \sin(b \log \frac{N}{k}) \left\{ \frac{\sin(b \log 1/N) + \sin(b \log 2/N) + \sin(b \log 3/N) + \cdots + \sin(b \log N/N)}{N} - H \right\} \quad (24-1)
\end{aligned}$$

$$\begin{aligned}
& = N \cos(b \log \frac{N}{k}) G + N \cos(b \log \frac{N}{k}) E_c(N) - N \sin(b \log \frac{N}{k}) H \\
& \quad - N \sin(b \log \frac{N}{k}) E_s(N) \quad (24-2)
\end{aligned}$$

$$\begin{aligned}
& = N \cos(b \log \frac{N}{k}) \frac{1}{1+b^2} + N \cos(b \log \frac{N}{k}) E_c(N) \\
& \quad + N \sin(b \log \frac{N}{k}) \frac{b}{1+b^2} - N \sin(b \log \frac{N}{k}) E_s(N) \quad (24-3)
\end{aligned}$$

$$\begin{aligned}
& = \frac{N}{\sqrt{1+b^2}} \left\{ \cos(b \log \frac{N}{k}) \frac{1}{\sqrt{1+b^2}} + \sin(b \log \frac{N}{k}) \frac{b}{\sqrt{1+b^2}} \right\} \\
& \quad + N \cos(b \log \frac{N}{k}) E_c(N) - N \sin(b \log \frac{N}{k}) E_s(N) \quad (24-4)
\end{aligned}$$

$$\begin{aligned}
& = N \left\{ \frac{\cos(b \log N/k - \tan^{-1} b)}{\sqrt{1+b^2}} \right. \\
& \quad \left. + \cos(b \log \frac{N}{k}) E_c(N) - \sin(b \log \frac{N}{k}) E_s(N) \right\} \quad (24-5)
\end{aligned}$$

$$\begin{aligned}
& = N \left[ \frac{1}{\sqrt{1+b^2}} \cos \left\{ b \log N \left( 1 - \frac{\log k}{\log N} - \frac{\tan^{-1} b}{b \log N} \right) \right\} \right. \\
& \quad \left. + \cos(b \log \frac{N}{k}) E_c(N) - \sin(b \log \frac{N}{k}) E_s(N) \right] \quad (24-6)
\end{aligned}$$

From (22-1), (22-2) and (24-1) we have (24-2). From (21) and (24-2) we have (24-3).

2.4 From (23) and the above (24-6) we have the following (25).

$$g(k, N) \sim \frac{N \cos(b \log N)}{\sqrt{1+b^2}} \quad (N \rightarrow \infty \quad k = 2, 3, 4, 5, \dots) \quad (25)$$

### Appendix 3. : Investigation of $F(a)$

#### 3.1. Investigation of $f(n)$

We have the following (8) and (15) in the text.

$$f(n) = \frac{1}{n^{1/2-a}} - \frac{1}{n^{1/2+a}} \geq 0 \quad (n = 2, 3, 4, 5, \dots \quad 0 \leq a < 1/2) \quad (8)$$

$$F(a) = f(2) - f(3) + f(4) - f(5) + f(6) - \dots \quad (15)$$

$a = 0$  is the solution for  $F(a) = 0$  due to  $f(n) \equiv 0$  at  $a = 0$ . The alternating series  $F(a)$  converges due to  $\lim_{n \rightarrow \infty} f(n) = 0$ .

We define the following (31) from the above (8) and we have the following (32) from (31).

$$f(r) = \frac{1}{r^{1/2-a}} - \frac{1}{r^{1/2+a}} \geq 0 \quad (r : \text{real number} \quad 2 \leq r) \quad (31)$$

$$\frac{df(r)}{dr} = f'(r) = \frac{1/2+a}{r^{a+3/2}} - \frac{1/2-a}{r^{3/2-a}} = \frac{1/2+a}{r^{a+3/2}} \left\{ 1 - \left( \frac{1/2-a}{1/2+a} \right) r^{2a} \right\} \quad (32)$$


The value of  $f(r)$  increases with increase of  $r$  and reaches the maximum value  $f(r_{max})$  at  $r = r_{max} = \left( \frac{1/2+a}{1/2-a} \right)^{1/(2a)}$ . Afterward  $f(r)$  decreases to zero with  $r \rightarrow \infty$ .  $f(n)$  also has the maximum value  $f(n_{max})$  at  $n = n_{max}$  and  $n_{max}$  is either of  $[r_{max}]$  and  $[r_{max}] + 1$ . Then we can have the following (34).

$$\begin{aligned} r_{max} &= \left( \frac{1/2+a}{1/2-a} \right)^{1/(2a)} = (1 + 4a + 8a^2 + \dots)^{1/(2a)} \\ &\sim (1 + 4a)^{1/(2a)} = \{(1 + 4a)^{1/(4a)}\}^2 \\ &\sim e^2 = 7.39 \quad (a \rightarrow +0) \end{aligned} \quad (34)$$

From the above (34) we have the following (35).

$$7 \leq n_{max} \quad (0 < a < 1/2) \quad (35)$$

The following (Graph 1) shows  $f(n)$  in various value of  $a$ .

Graph 1 :  $f(n)$  in various  $a$ 

We have the following (36) from (32).

$$\begin{aligned} f''(r) &= \frac{df'(r)}{dr} = \frac{(1/2 - a)(3/2 - a)}{r^{5/2-a}} - \frac{(1/2 + a)(3/2 + a)}{r^{5/2+a}} \\ &= \frac{(1/2 - a)(3/2 - a)}{r^{5/2-a}} \left\{ 1 - \frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)} r^{-2a} \right\} \end{aligned} \quad (36)$$

We have the following (37) from  $f''(r_0) = 0$ .

$$r_0 = \left\{ \frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)} \right\}^{1/(2a)} = \left( 1 + \frac{16}{3}a + \frac{128}{9}a^2 + \dots \right)^{1/(2a)} \quad (37)$$

Then we can have the following (37-1).

$$\begin{aligned} r_0 &= \left( 1 + \frac{16}{3}a + \frac{128}{9}a^2 + \dots \right)^{1/(2a)} \\ &\sim \left( 1 + \frac{16}{3}a \right)^{1/(2a)} = \left\{ \left( 1 + \frac{16}{3}a \right)^{3/(16a)} \right\}^{8/3} \\ &\sim e^{8/3} = 14.39 \quad (a \rightarrow +0) \end{aligned} \quad (37-1)$$

We can confirm the property of  $f(r)$  and  $f'(r)$  from (32), (36) and (Graph 1) as shown in the following (Table 1).

Item	Range of $r$	$f(r)$	$f'(r)$	The maximum value of $ f'(r) $
3.1.1	$2 \leq r \leq r_{max}$	Positive value. Monotonically increasing and strictly concave function. The maximum value at $r=r_{max}$ .	Positive value. Monotonically decreasing function. $f'(r)=0$ at $r=r_{max}$ .	$f'(2)$
3.1.2	$r_{max} < r \leq r_0$	Positive value. Monotonically decreasing and strictly concave function.	Negative value. Monotonically decreasing function. The minimum value at $r=r_0$ .	$-f'(r_0)$
3.1.3	$r_0 \leq r$	Positive value. Monotonically decreasing and strictly convex function. Converges to zero with $r \rightarrow \infty$ .	Negative value. Monotonically increasing function. Converges to zero with $r \rightarrow \infty$ .	$-f'(r_0)$

Table 1 : The property of  $f(r)$  and  $f'(r)$

**3.2. Verification method for  $0 < F(a)$**


We define  $F(a, n)$  as the following (38) and we have the following (39) from (38).

$$F(a, n) = f(2) - f(3) + f(4) - f(5) + \dots + (-1)^n f(n) \tag{38}$$

$$\lim_{n \rightarrow \infty} F(a, n) = F(a) \tag{39}$$

$F(a)$  is an alternating series. So  $F(a, n)$  repeats increase and decrease by  $f(n)$  with increase of  $n$  as shown in the following (Graph 2). In (Graph 2) upper points mean  $F(a, 2m)$  ( $m = 1, 2, 3, \dots$ ) and lower points mean  $F(a, 2m + 1)$ .  $F(a, 2m)$  decreases with increase of  $n$  in  $n_{max} \leq n$  and converges to  $F(a)$  with  $m \rightarrow \infty$ .  $F(a, 2m + 1)$  increases with increase of  $n$  in  $n_{max} \leq n$  and also converges to  $F(a)$  with  $m \rightarrow \infty$  due to  $\lim_{n \rightarrow \infty} f(n) = 0$ . From the above (39) we have the following (40).

$$\lim_{m \rightarrow \infty} F(a, 2m) = \lim_{m \rightarrow \infty} F(a, 2m + 1) = F(a) \tag{40}$$

Graph 2 :  $F(0.1, n)$  from  $n = 2$  to  $n = 100$ 

We define  $F1(a)$  and  $F1(a, 2m + 1)$  as follows.

$$F1(a) = \{f(2) - f(3)\} + \{f(4) - f(5)\} + \{f(6) - f(7)\} + \dots \quad (41)$$

$$\begin{aligned} F1(a, 2m + 1) &= \{f(2) - f(3)\} + \{f(4) - f(5)\} + \dots + \{f(2m) - f(2m + 1)\} \\ &= f(2) - f(3) + f(4) - f(5) + \dots + f(2m) - f(2m + 1) = F(a, 2m + 1) \end{aligned} \quad (42)$$

We have the following (43) from the above (40), (41) and (42).

$$F1(a) = \lim_{m \rightarrow \infty} F1(a, 2m + 1) = \lim_{m \rightarrow \infty} F(a, 2m + 1) = F(a) \quad (43)$$

We can use  $F1(a)$  instead of  $F(a)$  to verify  $0 < F(a)$ .

We enclose 2 terms of  $F(a)$  each from the first term with  $\{ \}$  as follows. If  $n_{max}$  is  $p$  or  $p + 1$  ( $p$ : odd number), the inside sum of  $\{ \}$  from  $f(2)$  to  $f(p)$  has negative value and the inside sum of  $\{ \}$  after  $f(p + 1)$  has positive value.

$$\begin{aligned} F(a) &= f(2) - f(3) + f(4) - f(5) + f(6) - f(7) + \dots \\ &= \{f(2) - f(3)\} + \{f(4) - f(5)\} + \dots + \{f(p-1) - f(p)\} + \{f(p+1) - f(p+2)\} + \dots \end{aligned}$$

$$(\text{inside sum of } \{ \}) < 0 \leftarrow \rightarrow (\text{inside sum of } \{ \}) > 0$$

$$(\text{total sum of } \{ \}) = -B \leftarrow \rightarrow (\text{total sum of } \{ \}) = A$$

We define as follows.

$$\{f(2) - f(3)\} + \{f(4) - f(5)\} + \dots + \{f(p-1) - f(p)\} = -B < 0$$

$$\{f(p+1) - f(p+2)\} + \{f(p+3) - f(p+4)\} + \dots = A > 0$$

We have the following (44) from the above definition.

$$F(a) = A - B \quad (44)$$

So we can verify  $0 < F(a)$  by verifying  $B < A$ .

**3.3. Investigation of  $f(n) - f(n + 1)$**

3.3.1 We have the following (45-1) from (31).


$$f(r) - f(r + 1) = \left(\frac{1}{r^{1/2-a}} - \frac{1}{r^{1/2+a}}\right) - \left\{\frac{1}{(r + 1)^{1/2-a}} - \frac{1}{(r + 1)^{1/2+a}}\right\} \quad (45-1)$$

We have the following (45-2) by differentiating  $f(r) - f(r + 1)$  regarding  $r$ .


$$\begin{aligned} \frac{df(r)}{dr} - \frac{df(r + 1)}{dr} &= \frac{1/2 + a}{r^{3/2+a}} \left\{1 - \left(\frac{r}{r + 1}\right)^{3/2+a}\right\} - \frac{1/2 - a}{r^{3/2-a}} \left\{1 - \left(\frac{r}{r + 1}\right)^{3/2-a}\right\} \\ &= C(r) - D(r) \end{aligned} \quad (45-2)$$

When  $r$  is small the value of  $f(r) - f(r + 1)$  increases with increase of  $r$  due to  $D(r) < C(r)$ . With increase of  $r$  the value reaches the maximum value  $f(r_1) - f(r_1 + 1)$  at  $r = r_1$ . Afterward the situation changes to  $C(r) < D(r)$  and the value decreases to zero with  $r \rightarrow \infty$ .

$f(n) - f(n + 1)$  also has the maximum value  $f(n_1) - f(n_1 + 1) = \{q_{max}\}$  at  $n = n_1$ .  $n_1$  is either of  $[r_1]$  and  $[r_1] + 1$ . The following (Graph 3) shows the value of  $f(n) - f(n + 1)$  in various value of  $a$ . The following (Graph 4) shows the value of  $f(n) - f(n + 1)$  at  $a = 0.1$ .



Graph 3 :  $f(n) - f(n + 1)$  in various  $a$

Graph 4 :  $f(n) - f(n + 1)$  at  $a = 0.1$ 

3.3.2 When  $n_{max}$  is even(odd) number the sign of  $f(n) - f(n + 1)$  changes from minus to plus with increase of  $n$  at  $n = n_{max}(n = n_{max} + 1)$  as shown in (Graph 4). Afterward the value reaches the maximum value  $\{q_{max}\}$  at  $n = n_1$  and the value decreases to zero with  $n \rightarrow \infty$ .

3.3.3 We can have the following (46) and (47) from (Table 1).

$$0 < f(n + 1) - f(n) = \int_n^{n+1} f'(r)dr \leq \int_2^3 f'(r)dr = f(3) - f(2) \quad (46)$$

$$(2 \leq r \leq n_{max} \quad n + 1 \leq n_{max})$$

$$0 < f(n) - f(n + 1) = \int_n^{n+1} \{-f'(r)\}dr < \int_n^{n+1} \{-f'(r_0)\}dr = -f'(r_0) \quad (47)$$

$$(n_{max} \leq r \quad n_{max} \leq n)$$


We can have the following (48) as shown in the following item 3.3.4, 3.3.5 and 3.3.6.

$$0 < -f'(r_0) < f(3) - f(2) \quad (0 < a < 1/2) \quad (48)$$

Then we can have the following (49) at the same value of  $a$  from the above (46), (47) and (48).

$$|f(n) - f(n + 1)| < f(3) - f(2) \quad (n = 3, 4, 5, \dots) \quad (49)$$

3.3.4 The following (Graph 5) is plotted by calculating  $f(3) - f(2)$  for  $a$  every 0.01.



Graph 5 :  $f(3) - f(2)$  regarding  $a$

$a$	0	0.05	0.1	0.15	0.2	0.25	0.3	0.35	0.4	0.45	0.5
$f(3)-f(2)$	0	0.014438	0.029008	0.043844	0.05908	0.074851	0.091297	0.108555	0.126771	0.146091	0.166667

Table 2 : The values of  $f(3) - f(2)$

If  $f(3) - f(2)$  has a convex or a concave in  $a_0 < a < a_0 + 0.01$ , such a convex or a concave is not displayed in the above (Graph 5). ( $a_0=0, 0.01, 0.02, \dots, 0.48, 0.49$ ) We define “The function does not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$ .” as either of the following 3 items.

- 3.3.4.1 The function does not have a local maximum value or a local minimum value in  $a_0 \leq a \leq a_0 + 0.01$ .
- 3.3.4.2 When the function has a local maximum value in  $a_0 \leq a < a_0 + 0.01$  the function is districtly concave regarding  $a$  in  $a_0 - 0.02 \leq a \leq a_0 + 0.03$ .
- 3.3.4.3 When the function has a local minimum value in  $a_0 \leq a < a_0 + 0.01$  the function is districtly convex regarding  $a$  in  $a_0 - 0.02 \leq a \leq a_0 + 0.03$ .

If the function has the property shown in the above 3 items, the graph can display the function correctly i.e. we can imagine the shape of the function easily from the graph although the graph is plotted for  $a$  every 0.01.


$f(n)$  is a monotonically increasing and districtly convex function regarding  $a$  in  $0 < a \leq 1/2$  from the following (50) and (51).  $f(n)$  meets the above item 3.3.4.1.

$$\frac{df(n)}{da} = \log n \left( \frac{1}{n^{1/2-a}} + \frac{1}{n^{a+1/2}} \right) > 0 \tag{50}$$

$$\frac{d^2f(n)}{da^2} = (\log n)^2 \left( \frac{1}{n^{1/2-a}} - \frac{1}{n^{1/2+a}} \right) \geq 0 \tag{51}$$

Then  $f(3)$  and  $f(2)$  are monotonically increasing and districtly convex functions regarding  $a$  i.e.  $f(3)$  and  $f(2)$  do not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$ .  $f(3) - f(2)$  also does not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$  from the above property of  $f(3)$  and  $f(2)$ . Therefore (Graph 5) shows  $f(3) - f(2)$  correctly.

3.3.5 The following (Graph 6) is plotted by calculating  $-f'(r_0)$  for  $a$  every 0.01. If  $-f'(r_0)$  has a convex or a concave in  $a_0 < a < a_0 + 0.01$ , such a convex or a concave is not displayed in (Graph 6). ( $a_0=0, 0.01, 0.02, \dots, 0.48, 0.49$ )



Graph 6 :  $-f'(r_0)$  regarding  $a$

$a$	0	0.05	0.1	0.15	0.2	0.25	0.3	0.35	0.4	0.45	0.5
$-f'(r_0)$	0	0.000601	0.001149	0.001591	0.00188	0.001976	0.001852	0.001504	0.000968	0.000361	0

Table 3 : The values of  $-f'(r_0)$

We have the following (52) from (32) and (37).

$$\begin{aligned}
-f'(r_0) &= (1/2 - a)r_0^{a-3/2} - (1/2 + a)r_0^{-a-3/2} \\
&= (1/2 - a)\left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{1/2-3/(4a)} \\
&\quad - (1/2 + a)\left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{-1/2-3/(4a)} \\
&= \left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{-3/(4a)} \left[ (1/2 - a)\left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{1/2} \right. \\
&\quad \left. - (1/2 + a)\left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{-1/2} \right] \\
&= \left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{-3/(4a)} \left[ \left\{\frac{(1/4 - a^2)(3/2 + a)}{3/2 - a}\right\}^{1/2} \right. \\
&\quad \left. - \left\{\frac{(1/4 - a^2)(3/2 - a)}{3/2 + a}\right\}^{1/2} \right] \\
&= \left\{\frac{(1/2 - a)(3/2 - a)}{(1/2 + a)(3/2 + a)}\right\}^{3/(4a)} (1/4 - a^2)^{1/2} \left\{ \left(\frac{3/2 + a}{3/2 - a}\right)^{1/2} - \left(\frac{3/2 - a}{3/2 + a}\right)^{1/2} \right\} \\
&= 2a \left\{\frac{(1/2 - a)(3/2 - a)}{(1/2 + a)(3/2 + a)}\right\}^{3/(4a)} \left\{\frac{(1/2 + a)(1/2 - a)}{(3/2 + a)(3/2 - a)}\right\}^{1/2} \\
&= 2a \{g(a)\}^{3/(4a)} \{h(a)\}^{1/2} \tag{52}
\end{aligned}$$

$g(a)$  in the above (52) is a monotonically decreasing and districtly convex func-

tion regarding  $a$  in  $0 \leq a \leq 1/2$  from the following (53-1) and (53-2).

$$\frac{dg(a)}{da} = \frac{4a^2 - 3}{(1/2 + a)^2(3/2 + a)^2} < 0 \quad (53-1)$$

$$\frac{d^2g(a)}{da^2} = \frac{2(6 + 9a - 4a^3)}{(1/2 + a)^3(3/2 + a)^3} > 0 \quad (53-2)$$

$h(a)$  in the above (52) is a monotonically decreasing and districtly concave function regarding  $a$  in  $0 < a \leq 1/2$  from the following (53-3) and (53-4).

$$\frac{dh(a)}{da} = \frac{-4a}{(3/2 + a)^2(3/2 - a)^2} \leq 0 \quad (53-3)$$

$$\frac{d^2h(a)}{da^2} = \frac{-3(3 + 4a^2)}{(3/2 + a)^3(3/2 - a)^3} < 0 \quad (53-4)$$

$a, 3/(4a), g(a)$  and  $h(a)$  that compose  $-f'(r_0)$  meet item 3.3.4.1 and they do not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$  respectively. Then  $-f'(r_0)$  also does not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$  from the above property of  $a, 3/(4a), g(a)$  and  $h(a)$ . Therefore (Graph 6) shows  $-f'(r_0)$  correctly.

Now we could confirm that (Graph 5) and (Graph 6) showed  $f(3) - f(2)$  and  $-f'(r_0)$  correctly, respectively. And we can find that (48) holds from (Graph 5) and (Graph 6).

3.3.6 We can confirm that (48) holds also during  $a \rightarrow +0$  from the following (54) and (55).

$f(3) - f(2)$  can be approximated in  $a \rightarrow +0$  by performing Maclaurin expansion for  $2^a, 2^{-a}, 3^a$  and  $3^{-a}$  as the following (54).

$$\begin{aligned} f(3) - f(2) &= (3^{a-1/2} - 3^{-a-1/2}) - (2^{a-1/2} - 2^{-a-1/2}) \\ &= 3^{-1/2}(3^a - 3^{-a}) - 2^{-1/2}(2^a - 2^{-a}) \\ &= 3^{-1/2}[\{1 + a \log 3 + (a \log 3)^2/2 + \dots\} - \{1 - a \log 3 + (a \log 3)^2/2 - \dots\}] \\ &\quad - 2^{-1/2}[\{1 + a \log 2 + (a \log 2)^2/2 + \dots\} - \{1 - a \log 2 + (a \log 2)^2/2 - \dots\}] \\ &= 2 * 3^{-1/2}\{a \log 3 + (a \log 3)^3/3! + (a \log 3)^5/5! + \dots\} \\ &\quad - 2 * 2^{-1/2}\{a \log 2 + (a \log 2)^3/3! + (a \log 2)^5/5! + \dots\} \\ &\sim 2(3^{-1/2} \log 3 - 2^{-1/2} \log 2)a = 0.29a > 0.012a \quad (a \rightarrow +0) \quad (54) \end{aligned}$$

$-f'(r_0)$  can be approximated in  $a \rightarrow +0$  from (32) and (37) by performing Maclaurin expansion for  $(1 + \frac{16}{3}a)^{1/2}$  and  $(1 + \frac{16}{3}a)^{-1/2}$  as the following (55).

$$\begin{aligned} -f'(r_0) &= (1/2 - a)r_0^{a-3/2} - (1/2 + a)r_0^{-a-3/2} \\ &= (1/2 - a)\left\{\frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)}\right\}^{1/2-3/(4a)} \end{aligned}$$

$$\begin{aligned}
& - (1/2 + a) \left\{ \frac{(1/2 + a)(3/2 + a)}{(1/2 - a)(3/2 - a)} \right\}^{-1/2-3/(4a)} \\
& = (1/2 - a) \left( 1 + \frac{16}{3}a + \frac{128}{9}a^2 + \dots \right)^{1/2-3/(4a)} \\
& \quad - (1/2 + a) \left( 1 + \frac{16}{3}a + \frac{128}{9}a^2 + \dots \right)^{-1/2-3/(4a)} \\
& \sim (1/2 - a) \left( 1 + \frac{16}{3}a \right)^{1/2-3/(4a)} - (1/2 + a) \left( 1 + \frac{16}{3}a \right)^{-1/2-3/(4a)} \\
& = \left( 1 + \frac{16}{3}a \right)^{-3/(4a)} \left\{ (1/2 - a) \left( 1 + \frac{16}{3}a \right)^{1/2} - (1/2 + a) \left( 1 + \frac{16}{3}a \right)^{-1/2} \right\} \\
& = \left( 1 + \frac{16}{3}a \right)^{-3/(4a)} \left\{ (1/2 - a) \left( 1 + \frac{8}{3}a - \frac{32}{9}a^2 + \dots \right) \right. \\
& \quad \left. - (1/2 + a) \left( 1 - \frac{8}{3}a + \frac{32}{9}a^2 + \dots \right) \right\} \\
& \sim \left( 1 + \frac{16}{3}a \right)^{-3/(4a)} \left\{ (1/2 - a) \left( 1 + \frac{8}{3}a \right) - (1/2 + a) \left( 1 - \frac{8}{3}a \right) \right\} \\
& = \left\{ \left( 1 + \frac{16}{3}a \right)^{3/(16a)} \right\}^{-4} \left( \frac{8}{3} - 2 \right) a \\
& \sim \frac{8/3 - 2}{e^4} a = 0.012a < 0.29a \quad (a \rightarrow +0) \tag{55}
\end{aligned}$$

### 3.4. Verification of $B < A$ ( $n_{max}$ is odd number.)

$n_{max}$  is odd number as follows.

$$\begin{aligned}
F(a) & = f(2) - f(3) + f(4) - f(5) + f(6) - \dots \\
& = \{f(2) - f(3)\} + \{f(4) - f(5)\} + \dots + \{f(n_{max} - 3) - f(n_{max} - 2)\} + \{f(n_{max} - 1) - \color{yellow}{f(n_{max})}\} \\
& \quad + \{f(n_{max} + 1) - f(n_{max} + 2)\} + \{f(n_{max} + 3) - f(n_{max} + 4)\} + \{f(n_{max} + 5) - f(n_{max} + 6)\} + \dots
\end{aligned}$$

We can have  $A$  and  $B$  as follows.  $A$  and  $B$  are defined in item 3.2.

$$\begin{aligned}
B & = \{f(3) - f(2)\} + \{f(5) - f(4)\} + \{f(7) - f(6)\} + \dots + \{f(n_{max} - 2) - f(n_{max} - 3)\} + \{\color{yellow}{f(n_{max})} - f(n_{max} - 1)\} \\
A & = \{f(n_{max} + 1) - f(n_{max} + 2)\} + \{f(n_{max} + 3) - f(n_{max} + 4)\} + \{f(n_{max} + 5) - f(n_{max} + 6)\} + \dots
\end{aligned}$$

#### 3.4.1. Condition for $B$

We define as follows.

$\{\color{yellow}\square\}$  : the term which is included within  $B$ .

$\{\color{gray}\square\}$  : the term which is not included within  $B$ .

We have the following (56).

$$\begin{aligned}
f(n_{max}) - f(2) & = \{\color{yellow}{f(n_{max}) - f(n_{max} - 1)}\} + \{\color{gray}{f(n_{max} - 1) - f(n_{max} - 2)}\} + \{\color{yellow}{f(n_{max} - 2) - f(n_{max} - 3)}\} \\
& \quad + \dots + \{\color{yellow}{f(7) - f(6)}\} + \{\color{gray}{f(6) - f(5)}\} + \{\color{yellow}{f(5) - f(4)}\} + \{\color{gray}{f(4) - f(3)}\} + \{\color{yellow}{f(3) - f(2)}\} \tag{56}
\end{aligned}$$

And we have the following (57) from (Table 1) in item 3.1, (Graph 3) and (Graph 4).

$$\begin{aligned}
\{\color{yellow}{f(3) - f(2)}\} & > \{\color{gray}{f(4) - f(3)}\} > \{\color{yellow}{f(5) - f(4)}\} > \{\color{gray}{f(6) - f(5)}\} > \{\color{yellow}{f(7) - f(6)}\} > \dots \\
& > \{\color{yellow}{f(n_{max} - 2) - f(n_{max} - 3)}\} > \{\color{gray}{f(n_{max} - 1) - f(n_{max} - 2)}\} > \{\color{yellow}{f(n_{max}) - f(n_{max} - 1)}\} > 0 \tag{57}
\end{aligned}$$

From the above (56) and (57) we have the following (58).

$$\begin{aligned}
 & f(n_{max}) - f(2) + \{f(3) - f(2)\} \\
 &= \{f(3) - f(2)\} + \{f(5) - f(4)\} + \{f(7) - f(6)\} + \dots + \{f(n_{max} - 2) - f(n_{max} - 3)\} + \{f(n_{max}) - f(n_{max} - 1)\} \\
 & \quad \parallel \quad \wedge \quad \wedge \quad \wedge \quad \leftarrow \text{Value comparison} \rightarrow \quad \wedge \\
 &+ \{f(3) - f(2)\} + \{f(4) - f(3)\} + \{f(6) - f(5)\} + \dots + \{f(n_{max} - 3) - f(n_{max} - 4)\} + \{f(n_{max} - 1) - f(n_{max} - 2)\} \\
 &> 2B \tag{58}
 \end{aligned}$$

The above (58) shows the following inequality.

$$\{\text{Total sum of upper row of (58)}\} = B < \{\text{Total sum of lower row of (58)}\}$$

Then we have the following (59).

$$2B < f(n_{max}) - f(2) + \{f(3) - f(2)\} \tag{59}$$

**3.4.2. Condition for A ( $\{q_{max}\}$  is included within A.)**

We abbreviate  $\{f(n_{max} + q) - f(n_{max} + q + 1)\}$  to  $\{q\}$  for easy description. ( $q = 0, 1, 2, 3, \dots$ ) All  $\{q\}$  has positive value as shown in item 3.3.2.

We define as follows.

- $\{\text{yellow}\}$  : the term which is included within A.
- $\{\text{grey}\}$  : the term which is not included within A.

$\{q_{max}\}$  has the maximum value in all  $\{q\}$ . And  $\{q_{max}\}$  is included within A. Then value comparison of  $\{q\}$  is as follows from item 3.3.2.

$$\{1\} < \{2\} < \{3\} < \dots < \{q_{max} - 3\} < \{q_{max} - 2\} < \{q_{max} - 1\} < \{q_{max}\} > \{q_{max} + 1\} > \{q_{max} + 2\} > \{q_{max} + 3\} > \dots$$

We have the following (60).

$$\begin{aligned}
 & f(n_{max} + 1) = \{f(n_{max} + 1) - f(n_{max} + 2)\} + \{f(n_{max} + 2) - f(n_{max} + 3)\} + \{f(n_{max} + 3) - f(n_{max} + 4)\} \\
 & \quad + \{f(n_{max} + 4) - f(n_{max} + 5)\} + \dots \\
 &= \{1\} + \{2\} + \{3\} + \{4\} + \dots + \{q_{max} - 3\} + \{q_{max} - 2\} + \{q_{max} - 1\} + \{q_{max}\} + \{q_{max} + 1\} + \{q_{max} + 2\} + \{q_{max} + 3\} + \dots \tag{60}
 \end{aligned}$$

From the above (60) we have the following (61).

$$\begin{aligned}
 & f(n_{max} + 1) - \{q_{max} - 1\} \\
 &= \{1\} + \{2\} + \{3\} + \{4\} + \dots + \{q_{max} - 3\} + \{q_{max} - 2\} + \{q_{max}\} + \{q_{max} + 1\} + \{q_{max} + 2\} + \{q_{max} + 3\} + \dots \tag{61} \\
 & \quad \leftarrow \dots \dots \dots \text{Range 1} \dots \dots \dots \rightarrow \leftarrow \dots \dots \dots \text{Range 2} \dots \dots \dots
 \end{aligned}$$

(Range 1) and (Range 2) are determined as above. In (Range 1) value comparison is as follows.

$$\{1\} < \{2\} < \{3\} < \{4\} < \dots < \{q_{max} - 4\} < \{q_{max} - 3\} < \{q_{max} - 2\}$$

And we can find the following.



And we can find the following.

$$\begin{aligned} \text{Total sum of } \{\text{■}\} &= \{q_{max} + 1\} + \{q_{max} + 3\} + \{q_{max} + 5\} + \{q_{max} + 7\} + \dots \\ &\quad \quad \quad \downarrow \quad \quad \quad \downarrow \quad \quad \quad \downarrow \quad \quad \quad \downarrow \quad \quad \quad \leftarrow \text{Value comparison} \\ \text{Total sum of } \{\text{■}\} &= \{q_{max} + 2\} + \{q_{max} + 4\} + \{q_{max} + 6\} + \{q_{max} + 8\} + \dots \end{aligned}$$

Therefore [Total sum of {■} > Total sum of {■}] holds.

In (Range 1)+(Range 2) we have [Total sum of {■} =  $A$  > Total sum of {■}].

We have the following (65).

$$f(n_{max} + 1) - \{q_{max}\} < 2A \quad (65)$$

#### 3.4.4. Condition for $B < A$

From (62) and (65) we have the following inequality.

$$f(n_{max} + 1) - [\{q_{max}\} \text{ or } \{q_{max} - 1\}] < 2A$$

Then the following inequalities hold from (49).

$$\begin{aligned} [\{q_{max}\} \text{ or } \{q_{max} - 1\}] &< f(3) - f(2) \\ f(n_{max}) - f(n_{max} + 1) &< f(3) - f(2) \end{aligned}$$

We have the following (66) from the above 3 inequalities.

$$\begin{aligned} 2A &> f(n_{max} + 1) - [\{q_{max}\} \text{ or } \{q_{max} - 1\}] > f(n_{max} + 1) - \{f(3) - f(2)\} \\ &> f(n_{max}) - \{f(3) - f(2)\} - \{f(3) - f(2)\} = f(n_{max}) - 2\{f(3) - f(2)\} \end{aligned} \quad (66)$$

We have the following (67) for  $B < A$  from (59) and (66).


$$2A > f(n_{max}) - 2\{f(3) - f(2)\} > f(n_{max}) - f(2) + \{f(3) - f(2)\} > 2B \quad (67)$$

From the above (67) we can have the final condition for  $B < A$  as follows.

$$f(3) < (4/3)f(2) \quad (68)$$

The following (Graph 7) is plotted by calculating the following (71) for  $a$  every 0.01.

$$J(a) = (4/3)f(2) - f(3) = (4/3)\left(\frac{1}{2^{1/2-a}} - \frac{1}{2^{1/2+a}}\right) - \left(\frac{1}{3^{1/2-a}} - \frac{1}{3^{1/2+a}}\right) \quad (71)$$

Graph 7 :  $J(a) = (4/3)f(2) - f(3)$ 

a	0	0.05	0.1	0.15	0.2	0.25	0.3	0.35	0.4	0.45	0.5
$(4/3)f(2)-f(3)$	0	0.001903	0.003694	0.005257	0.00648	0.007246	0.007437	0.006933	0.005611	0.003343	0

Table 4 : The values of  $J(a)$ 

$f(2)$  and  $f(3)$  do not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$  as shown in item 3.3.4. ( $a_0=0, 0.01, 0.02, \dots, 0.48, 0.49$ )  $J(a)$  also does not have a convex or a concave in  $a_0 \leq a \leq a_0 + 0.01$  from the above property of  $f(2)$  and  $f(3)$ . Therefore (Graph 7) shows  $J(a)$  correctly. We can confirm that  $0 < J(a)$  holds also during  $a \rightarrow +0$  and  $a \rightarrow 1/2 - 0$  as shown in the following item 3.4.4.1 and 3.4.4.2. From (Graph 7) and item 3.4.4.1 and 3.4.4.2 we can find that  $0 < J(a)$  holds in  $0 < a < 1/2$ . Therefore  $B < A$  i.e.  $0 < F(a)$  holds in  $0 < a < 1/2$  from (44).

3.4.4.1  $J(a)$  can be approximated in  $a \rightarrow +0$  by performing Maclaurin expansion for  $2^a, 2^{-a}, 3^a$  and  $3^{-a}$  as the following (71-1).

$$\begin{aligned}
J(a) &= (4/3)f(2) - f(3) \\
&= (4/3)(2^{a-1/2} - 2^{-a-1/2}) - (3^{a-1/2} - 3^{-a-1/2}) \\
&= (4/3)2^{-1/2}(2^a - 2^{-a}) - 3^{-1/2}(3^a - 3^{-a}) \\
&= (4/3)2^{-1/2}[\{1 + a \log 2 + (a \log 2)^2/2 + \dots\} - \{1 - a \log 2 + (a \log 2)^2/2 - \dots\}] \\
&\quad - 3^{-1/2}[\{1 + a \log 3 + (a \log 3)^2/2 + \dots\} - \{1 - a \log 3 + (a \log 3)^2/2 - \dots\}] \\
&= 2 * (4/3)2^{-1/2}\{a \log 2 + (a \log 2)^3/3! + (a \log 2)^5/5! + \dots\} \\
&\quad - 2 * 3^{-1/2}\{a \log 3 + (a \log 3)^3/3! + (a \log 3)^5/5! + \dots\} \\
&\sim (4/3)2^{-1/2}(2a \log 2) - 3^{-1/2}(2a \log 3) = 0.038a > 0 \quad (a \rightarrow +0) \quad (71-1)
\end{aligned}$$

3.4.4.2 Let  $(1/2 - a)$  be  $t$ .  $J(a)$  can be approximated in  $a \rightarrow 1/2 - 0$  by performing Maclaurin expansion for  $2^t, 2^{-t}, 3^t$  and  $3^{-t}$  as the following (71-2).

$$J(a) = (4/3)f(2) - f(3)$$

$$\begin{aligned}
&= (4/3)(2^{a-1/2} - 2^{-a-1/2}) - (3^{a-1/2} - 3^{-a-1/2}) \\
&= (4/3)(2^{-t} - 2^{t-1}) - (3^{-t} - 3^{t-1}) = (4/3)(2^{-t} - 2^t/2) - (3^{-t} - 3^t/3) \\
&= (4/3)[\{1 - t \log 2 + (t \log 2)^2/2 - \dots\} \\
&\quad - (1/2)\{1 + t \log 2 + (t \log 2)^2/2 + \dots\}] \\
&- [\{1 - t \log 3 + (t \log 3)^2/2 - \dots\} \\
&\quad - (1/3)\{1 + t \log 3 + (t \log 3)^2/2 + \dots\}] \\
&\sim (4/3)\{(1 - t \log 2) - (1 + t \log 2)/2\} - \{(1 - t \log 3) - (1 + t \log 3)/3\} \\
&= (4/3)\{1/2 - (3/2)t \log 2\} - \{2/3 - (4/3)t \log 3\} = 0.0785t \\
&= 0.0785(1/2 - a) > 0 \quad (t \rightarrow +0 \quad a \rightarrow 1/2 - 0) \tag{71-2}
\end{aligned}$$

### 3.5. Verification of $B < A$ ( $n_{max}$ is even number.)

$n_{max}$  is even number as follows.

$$\begin{aligned}
F(a) &= f(2) - f(3) + f(4) - f(5) + f(6) - \dots \\
&= \{f(2) - f(3)\} + \{f(4) - f(5)\} + \dots + \{f(n_{max} - 4) - f(n_{max} - 3)\} + \{f(n_{max} - 2) - f(n_{max} - 1)\} \\
&\quad + \{f(n_{max}) - f(n_{max} + 1)\} + \{f(n_{max} + 2) - f(n_{max} + 3)\} + \{f(n_{max} + 4) - f(n_{max} + 5)\} + \dots
\end{aligned}$$

We can have  $A$  and  $B$  as follows.

$$\begin{aligned}
B &= \{f(3) - f(2)\} + \{f(5) - f(4)\} + \{f(7) - f(6)\} \\
&\quad + \dots + \{f(n_{max} - 3) - f(n_{max} - 4)\} + \{f(n_{max} - 1) - f(n_{max} - 2)\} \\
A &= \{f(n_{max}) - f(n_{max} + 1)\} + \{f(n_{max} + 2) - f(n_{max} + 3)\} + \{f(n_{max} + 4) - f(n_{max} + 5)\} + \dots \\
f(n_{max}) &= \{f(n_{max}) - f(n_{max} + 1)\} + \{f(n_{max} + 1) - f(n_{max} + 2)\} + \{f(n_{max} + 2) - f(n_{max} + 3)\} \\
&\quad + \{f(n_{max} + 3) - f(n_{max} + 4)\} + \dots \\
&= \{0\} + \{1\} + \{2\} + \{3\} + \{4\} \\
&\quad + \dots + \{q_{max} - 3\} + \{q_{max} - 2\} + \{q_{max} - 1\} + \{q_{max}\} + \{q_{max} + 1\} + \{q_{max} + 2\} + \{q_{max} + 3\} + \dots
\end{aligned}$$

After the same process as in item 3.4.1 we can have the following (73).

$$f(n_{max} - 1) - f(2) + \{f(3) - f(2)\} > 2B \tag{73}$$

The following inequality holds from (49).

$$[\{q_{max}\} \text{ or } \{q_{max} - 1\}] < f(3) - f(2)$$

We have the following (74) from the above inequality and the same process as in item 3.4.2 and item 3.4.3.

$$\begin{aligned}
2A &> f(n_{max}) - [\{q_{max}\} \text{ or } \{q_{max} - 1\}] > f(n_{max}) - \{f(3) - f(2)\} \\
&> f(n_{max} - 1) - \{f(3) - f(2)\} \tag{74}
\end{aligned}$$

We have the following (75) for  $B < A$  from (73) and (74).

$$2A > f(n_{max} - 1) - \{f(3) - f(2)\} > f(n_{max} - 1) - f(2) + \{f(3) - f(2)\} > 2B \quad (75)$$

From the above (75) we can have the final condition for  $B < A$  as follows.

$$f(3) < (3/2)f(2) \quad (76)$$

In the following (77),  $(4/3)f(2) < (3/2)f(2)$  is true self-evidently and in item 3.4.4 we already confirmed that the following (68) was true in  $0 < a < 1/2$ .

$$0 < f(3) < (4/3)f(2) < (3/2)f(2) \quad (77)$$

$$f(3) < (4/3)f(2) \quad (68)$$

Therefore the above (76) is true in  $0 < a < 1/2$ . Now we can confirm  $0 < F(a)$  in  $0 < a < 1/2$ .

### 3.6. Conclusion

$0 < F(a)$  holds in  $0 < a < 1/2$  as shown in the above item 3.4 and item 3.5.

**Appendix 4. Graph of  $F(a)$**


4.1 We can approximate  $F(a)$  as the following (81) from (38). We have the following (82) and (83) from (81).

$$F(a)_n = \frac{F(a, n) + F(a, n + 1)}{2} \tag{81}$$

$$\lim_{n \rightarrow \infty} F(a)_n = F(a) \tag{82}$$

$$F(a)_{n+1} = F(a)_n - (-1)^n \frac{f(n + 1) - f(n + 2)}{2} \tag{83}$$

The following (Graph 8) is plotted by calculating  $F(a)_{500}$  and  $F(a)_{501}$  for  $a$  every 0.01.



Graph 8 :  $F(a)_{500}$  and  $F(a)_{501}$

$a$	0	0.01	0.1	0.2	0.3	0.4	0.5
$F(a)_{500}$	0	0.0038667	0.038666	0.077326	0.115971	0.154587	0.193146
$F(a)_{501}$	0	0.0038648	0.038647	0.077289	0.115919	0.154537	0.193148
$F(a)$	0	0.00386	0.0386	0.077	0.1159	0.1545	-

Table 5 : The values of  $F(a)_{500}$  and  $F(a)_{501}$

The range of  $a$  is  $0 \leq a < 1/2$ .  $a = 1/2$  is not included in the range. But we added  $F(1/2)_n$  to calculation due to the following reason.  $f(n)$  at  $a = 1/2$  is  $(1 - 1/n)$  and  $F(1/2)$  fluctuates due to  $\lim_{n \rightarrow \infty} f(n) = 1$ . The above (83) shows that  $F(a)_n$  is partial sum of alternating series which has the term of  $\frac{f(n+1)-f(n+2)}{2}$ . Then  $\lim_{n \rightarrow \infty} F(1/2)_n$  can converge to the fixed value on the condition of  $\lim_{n \rightarrow \infty} \{f(n + 1) - f(n + 2)\} = 0$ . The condition holds due to  $f(n + 1) - f(n + 2) = -1/(n^2 + 3n + 2)$ .

4.2  $r_0$  in (37) has the value of 217 at  $a = 0.49$ . Then  $f(n+1) - f(n+2)$  has positive value

and decreases monotonically with increase of  $n$  in  $217 < n$  and  $0 < a \leq 0.49$ .  $F(a)_n$  converges to  $F(a)$  with  $n \rightarrow \infty$  as (82) shows. Then we can have the following (84) from (83).

$$F(a)_{501} < F(a) < F(a)_{500} \quad (0 < a \leq 0.49) \quad (84)$$

Therefore (Graph 8) shows  $F(a)$  as well as  $F(a)_{500}$  and  $F(a)_{501}$  in  $0 \leq a \leq 0.49$ .

### References

- [1] Yukio Kusunoki, Introduction to infinite series, Asakura syoten, (1972), page 22, (written in Japanese)

Toshihiko ISHIWATA  
E-mail: toshihiko.ishiwata@gmail.com