

The prime Number Theorem and Prime Gaps

Hyeon Jun Ahn

June 1, 2022

Abstract: The prime number theorem; PNT shows the n th prime is asymptotically $n \log n$ where \log is the natural logarithm. By using PNT, This paper proves that

$$\forall k > 0, \exists M \in \mathbb{N} \quad \text{s.t.} \quad n \geq M \Rightarrow g_n = p_{n+1} - p_n < p_n^k$$

where g_n is the prime gap, p_n is the n th prime, and introduces a corollary about the Andrica conjecture, the Cramer conjecture, and the Oppermann conjecture.

1. Introduction

By the prime number theorem, primes less than n are asymptotically $\frac{n}{n \log n}$, so the average gap between primes less than n is $\log n$. Hence, n th prime is asymptotically $n \log n$.

$$\text{i.e.} \quad \lim_{n \rightarrow \infty} \frac{p_n}{n \log n} = 1$$

This is equivalent to

$$p_n \sim n \log n$$

This means that $n \log n$ approximates p_n in the sense that the relative error of this approximation approaches 0 as n increases without bound. So,

$$p_{n+1} + p_n \sim (n+1) \log(n+1) + n \log n$$

because

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{p_{n+1} + p_n}{(n+1) \log(n+1) + n \log n} \\ = & \lim_{n \rightarrow \infty} \left(\frac{1}{(n+1) \log(n+1)/p_{n+1} + n \log n/p_{n+1}} + \frac{1}{(n+1) \log(n+1)/p_n + n \log n/p_n} \right) \\ = & \lim_{n \rightarrow \infty} \left(\frac{1}{1+1} + \frac{1}{1+1} \right) = 1 \end{aligned}$$

This result shows it is possible to add $p_n \sim n \log n$ and $p_{n+1} \sim (n+1) \log(n+1)$. But

$$p_n - p_{n+1} \asymp (n+1) \log(n+1) - n \log n \quad (1)$$

Rather,

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(n+1)\log(n+1) - n\log n} = \infty \text{ and } \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(n+1)\log(n+1) - n\log n} = 0$$

proof. Note that

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} = \infty \text{ and } \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} = 0 \quad (2)$$

E. Westzynthius proved the former in 1931^{1,2}, Daniel Goldston, János Pintz and Cem Yıldırım proved the latter in 2005³. And note that the formula

$$\lim_{n \rightarrow \infty} \frac{\log(n\log n)}{\log p_n} = 1 \quad (3)$$

holds. Because, for every $3 \leq n \in \mathbb{N}$,

$$\frac{\log(n\log n)}{\log p_n} = \log_{p_n}(n\log n) = k(n) \in \mathbb{R}$$

Then,

$$p_n = (n\log n)^{k(n)} \Rightarrow \frac{p_n}{(n\log n)^{k(n)}} = 1$$

Since $p_n \sim n\log n$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{n\log n}{(n\log n)^{k(n)}} &= \lim_{n \rightarrow \infty} \frac{p_n}{(n\log n)^{k(n)}} \frac{n\log n}{p_n} = 1 \times 1 = 1 \\ \therefore \lim_{n \rightarrow \infty} k(n) &= 1 \end{aligned}$$

And note that the formula

$$\lim_{n \rightarrow \infty} \frac{\log(n\log n)}{(n+1)\log(n+1) - n\log n} = 1 \quad (4)$$

holds. Because, by L'Hospital's rule,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\log(n\log n)}{(n+1)\log(n+1) - n\log n} &= \lim_{n \rightarrow \infty} \frac{\log n + \log(\log n)}{(n+1)\log(n+1) - n\log n} \\ &\stackrel{L'H}{=} \lim_{n \rightarrow \infty} \frac{1/n + 1/n\log n}{\log(n+1) - \log n} = \lim_{n \rightarrow \infty} \frac{\log n + 1}{n\log n(\log(n+1) - \log n)} \\ &= \lim_{n \rightarrow \infty} \frac{\log n + 1}{\log n \log(1 + 1/n)^n} = 1 \blacksquare \end{aligned}$$

Now, let $F(n) = \frac{\log p_n}{\log(n\log n)} \frac{\log(n\log n)}{(n+1)\log(n+1) - n\log n}$, then, due to (3),(4),

$$\lim_{n \rightarrow \infty} F(n) = 1$$

$$\begin{aligned}
& \text{i.e. } \exists M \in \mathbb{N} \quad \text{s.t. } n \geq M \Rightarrow \frac{1}{2} < F(n) < \frac{3}{2} \\
& \Rightarrow \frac{1}{2} \frac{p_{n+1} - p_n}{\log p_n} < \frac{p_{n+1} - p_n}{\log p_n} F(n) < \frac{3}{2} \frac{p_{n+1} - p_n}{\log p_n} \\
& \therefore \limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} F(n) = \infty \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} F(n) = 0
\end{aligned}$$

$$\text{Since } \frac{p_{n+1} - p_n}{\log p_n} F(n) = \frac{p_{n+1} - p_n}{(n+1)\log(n+1) - n \log n},$$

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(n+1)\log(n+1) - n \log n} = \infty \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(n+1)\log(n+1) - n \log n} = 0$$

Therefore we need another method to find the approximate expression of $p_{n+1} - p_n$. $(n+1)\log(n+1) - n \log n$ is not appropriate though $p_n \sim n \log n$.

2. Prime gap

Remark 1. For every $k > 0$,

$$\lim_{n \rightarrow \infty} \frac{p_n^k}{(n \log n)^k} = \lim_{n \rightarrow \infty} \left(\frac{p_n}{n \log n} \right)^k = 1 \quad (5)$$

Remark 2.

$$\lim_{n \rightarrow \infty} \frac{\log p_n}{\log(n \log n)} = 1$$

See (3).

Remark 3.

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} = \infty$$

See (2).

Lemma 2. $\forall n, a_n, b_n > 0$,

$$\lim_{n \rightarrow \infty} a_n b_n = 0, \quad \limsup_{n \rightarrow \infty} b_n = \infty \Rightarrow \lim_{n \rightarrow \infty} a_n = 0 \quad (6)$$

proof. Let

$$A_n = \{b_k | k \leq n\}, \quad s_n = \sup A_n$$

Then

$$\limsup_{n \rightarrow \infty} \frac{s_n}{b_n} = 1 \Rightarrow \limsup_{n \rightarrow \infty} \frac{a_n s_n}{a_n b_n} = 1$$

$$\begin{aligned}
\Leftrightarrow \forall \epsilon_1 > 0, \exists N_1 \in \mathbb{N} \quad \text{s.t. } n \geq N_1 & \Rightarrow \frac{a_n s_n}{a_n b_n} < 1 + \epsilon_1 \\
& \Rightarrow a_n s_n < (1 + \epsilon_1) a_n b_n
\end{aligned}$$

Meanwhile,

$$\forall \epsilon_2 > 0, \exists N_2 \in \mathbb{N} \quad \text{s.t. } n \geq N_2 \Rightarrow |a_n b_n| < \epsilon_2$$

So, let $\epsilon = (1 + \epsilon_1)\epsilon_2$, $N = \max(N_1, N_2)$. Then,

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \quad \text{s.t. } n \geq N \Rightarrow a_n s_n < \epsilon$$

$$\therefore \limsup_{n \rightarrow \infty} a_n s_n \leq 0$$

Since $a_n, s_n > 0$,

$$\lim_{n \rightarrow \infty} a_n s_n = 0$$

And since $\lim_{n \rightarrow \infty} s_n = \infty$,

$$\lim_{n \rightarrow \infty} a_n = 0 \blacksquare$$

Lemma 3.

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log(n \log n))^2} = 0 \quad (7)$$

proof. Note that

$$\lim_{n \rightarrow \infty} \frac{\log p_n}{(\log(n \log n))^2} = \lim_{n \rightarrow \infty} \frac{1}{\log(n \log n)} \frac{\log p_n}{\log(n \log n)} = 0 \quad (\because (3))$$

Therefore,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log(n \log n))^2} \frac{\log p_n}{p_{n+1} - p_n} = \lim_{n \rightarrow \infty} \frac{\log p_n}{(\log(n \log n))^2} = 0$$

Let $a_n = \frac{p_{n+1} - p_n}{(\log(n \log n))^2}$, $b_n = \frac{\log p_n}{p_{n+1} - p_n}$ ($n \geq 2$), then

$$a_n, b_n > 0, \quad \lim_{n \rightarrow \infty} a_n b_n = 0, \quad \limsup_{n \rightarrow \infty} b_n = \infty \quad (\because (2))$$

$$\Rightarrow \lim_{n \rightarrow \infty} a_n = 0 \quad (\because (6))$$

Hence,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log(n \log n))^2} = 0 \blacksquare$$

Lemma 4. For every $k > 0$,

$$\lim_{n \rightarrow \infty} \frac{(\log(n \log n))^2}{(n \log n)^k} = 0 \quad (8)$$

proof. Let $x = n \log n$, By L'Hospital's rule,

$$\lim_{x \rightarrow \infty} \frac{(\log(x))^2}{x^k} \stackrel{\text{L'H}}{=} \lim_{x \rightarrow \infty} \frac{2 \log x}{k x^k} \stackrel{\text{L'H}}{=} \lim_{x \rightarrow \infty} \frac{2}{k^2 x^{k+1}} = 0 \blacksquare$$

Due to (5),(7), for every $k > 0$,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{p_n^k} \frac{(n \log n)^k}{(\log(n \log n))^2} = \lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log(n \log n))^2} \frac{(n \log n)^k}{p_n^k} = 0 \times 1 = 0 \quad (9)$$

Hence,

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{p_n^k} &= (8) \times (9) = 0 \\ \Leftrightarrow \lim_{n \rightarrow \infty} \frac{p_n^k}{p_{n+1} - p_n} &= \infty\end{aligned}$$

By epsilon-delta argument,

$$\begin{aligned}\forall k > 0, \exists N \in \mathbb{N} \quad s.t. \quad n \geq N &\Rightarrow g_n := p_{n+1} - p_n < p_n^k \\ &\Rightarrow p_n < p_{n+1} < p_n + p_n^k\end{aligned}\tag{10}$$

corollary 1.

$$g_n := p_{n+1} - p_n = O(p_n^k) \quad \forall k \in \mathbb{R}^+$$

Where O is big O notation.

3. About Andrica conjecture

Andrica conjecture is a conjecture regarding the gaps between prime numbers. The conjecture states that the inequality

$$\sqrt{p_{n+1}} - \sqrt{p_n} < 1$$

holds for all $n \in \mathbb{N}$. And a strong version of Andrica conjecture is as follows: Except for $p_n \in \{3, 7, 13, 23, 31, 113\}$, that is $n \in \{2, 4, 6, 9, 11, 30\}$, one has

$$\sqrt{p_{n+1}} - \sqrt{p_n} < \frac{1}{2}; \quad \text{equivalently} \quad g_n := p_{n+1} - p_n < p_n^{1/2} + \frac{1}{4}$$

And This paper proves that

$$\lim_{n \rightarrow \infty} (\sqrt{p_{n+1}} - \sqrt{p_n}) = 0$$

proof. Let $\epsilon \in (0, \frac{1}{2})$, $k \in (0, \frac{1}{2})$, Then

$$\lim_{n \rightarrow \infty} \frac{p_n^k}{(\sqrt{p_n} + \epsilon)^2 - p_n} = \lim_{n \rightarrow \infty} \frac{p_n^k}{2\epsilon\sqrt{p_n} + \epsilon^2} = 0$$

Thus,

$$\begin{aligned}\exists N_1 \in \mathbb{N} \quad s.t. \quad n > N_1 &\Rightarrow p_n^k < (\sqrt{p_n} + \epsilon)^2 - p_n \\ &\Rightarrow p_n + p_n^k < (\sqrt{p_n} + \epsilon)^2\end{aligned}$$

Meanwhile,

$$\exists N_2 \in \mathbb{N} \quad s.t. \quad n > N_2 \Rightarrow p_{n+1} < p_n + p_n^k \quad (\because (10))$$

Let $N = \max(N_1, N_2)$, Then

$$n > N \Rightarrow p_{n+1} < (\sqrt{p_n} + \epsilon)^2 \Rightarrow \sqrt{p_{n+1}} - \sqrt{p_n} < \epsilon$$

By epsilon-delta argument,

$$\lim_{n \rightarrow \infty} (\sqrt{p_{n+1}} - \sqrt{p_n}) = 0 \blacksquare \quad (11)$$

Furthermore, let $y > 1$, $x < \frac{y-1}{y}$. Then, since $\forall \epsilon > 0$, $\exists M \in \mathbb{N}$ s.t. $n > M \Rightarrow |p_n^{1/y}| > |\epsilon|$, by generalized binomial theorem,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{p_n^x}{(p_n^{1/y} + \epsilon)^y - p_n} \\ &= \lim_{n \rightarrow \infty} \frac{p_n^x}{(p_n + \binom{y}{1} p_n^{(y-1)/y} \epsilon + \binom{y}{2} p_n^{(y-2)/y} \epsilon^2 + \dots) - p_n} \\ &= \lim_{n \rightarrow \infty} \frac{p_n^x}{\left(\binom{y}{1} p_n^{(y-1)/y} \epsilon + \binom{y}{2} p_n^{(y-2)/y} \epsilon^2 + \dots\right)} = 0 \quad (\because x < \frac{y-1}{y}) \end{aligned}$$

In the same method as the proof of (11),

$$\lim_{n \rightarrow \infty} (p_{n+1}^{1/y} - p_n^{1/y}) = 0 \quad \text{for } y > 1$$

3-1. The arithmetic mean, the geometric mean and harmonic mean of primes

The relation between the arithmetic mean and the geometric mean of n th prime and $(n+1)$ th prime is as follows:

$$\frac{p_{n+1} + p_n}{2} \sim \sqrt{p_{n+1} p_n}$$

proof.

$$\begin{aligned} & \lim_{n \rightarrow \infty} (\sqrt{p_{n+1}} - \sqrt{p_n}) = 0 \\ & \Rightarrow \lim_{n \rightarrow \infty} (\sqrt{p_{n+1}} - \sqrt{p_n})^2 = 0 \\ & \Rightarrow \lim_{n \rightarrow \infty} (p_{n+1} + p_n - 2\sqrt{p_{n+1} p_n}) = 0 \end{aligned} \quad (12)$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} + p_n}{2\sqrt{p_{n+1} p_n}} = \lim_{n \rightarrow \infty} \left(\frac{p_{n+1} + p_n - 2\sqrt{p_{n+1} p_n}}{2\sqrt{p_{n+1} p_n}} + 1 \right) = 1 \blacksquare \quad (13)$$

Furthermore,

$$\lim_{n \rightarrow \infty} \left(\frac{p_{n+1} + p_n}{2} - \sqrt{p_{n+1} p_n} \right) = 0$$

trivially holds by (12). And similarly, the relation between the arithmetic mean and the harmonic mean of n th prime and $(n+1)$ th prime is as follows:

$$\frac{p_{n+1} + p_n}{2} \sim \frac{2p_{n+1} p_n}{p_{n+1} + p_n}$$

proof. By (13)

$$\lim_{n \rightarrow \infty} \frac{2p_{n+1}p_n}{p_{n+1} + p_n} \frac{2}{p_{n+1} + p_n} = \lim_{n \rightarrow \infty} \left(\frac{2\sqrt{p_{n+1}p_n}}{p_{n+1} + p_n} \right)^2 = 1 \blacksquare$$

Furthermore,

$$\lim_{n \rightarrow \infty} \left(\frac{p_{n+1} + p_n}{2} - \frac{2p_{n+1}p_n}{p_{n+1} + p_n} \right) = 0$$

proof. Note that

$$\lim_{n \rightarrow \infty} \frac{(\log(n \log n))^4}{4n \log n} = 0 \quad (14)$$

And

$$\lim_{n \rightarrow \infty} \left(\frac{4n \log n}{4p_n} \right) \left(\frac{p_{n+1} - p_n}{(\log(n \log n))^4} \right)^2 = 0 \quad (15)$$

The former is trivial because of (8), and The latter is trivial because of the prime number theorem and (7). Thus,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left(\frac{p_{n+1} + p_n}{2} - \frac{2p_{n+1}p_n}{p_{n+1} + p_n} \right) \\ &= \lim_{n \rightarrow \infty} \frac{(p_{n+1} + p_n)^2 - 4p_{n+1}p_n}{2(p_{n+1} + p_n)} = \lim_{n \rightarrow \infty} \frac{(p_{n+1} - p_n)^2}{2(p_{n+1} + p_n)} \\ &\leq \lim_{n \rightarrow \infty} \frac{(p_{n+1} - p_n)^2}{4p_n} = (14) \times (15) = 0 \end{aligned}$$

By the relation between the geometric mean and the harmonic mean,

$$\lim_{n \rightarrow \infty} \left(\frac{p_{n+1} + p_n}{2} - \frac{2p_{n+1}p_n}{p_{n+1}p_n} \right) = 0 \blacksquare$$

Hence,

$$\frac{p_{n+1} + p_n}{2} \sim \sqrt{p_{n+1}p_n} \sim \frac{2p_{n+1}p_n}{p_{n+1}p_n}$$

Furthermore, the arithmetic mean, the geometric mean, and the harmonic mean of n th prime and $(n+1)$ th prime become asymptotically the same as n increases without bound.

4. About Cramer conjecture

Cramer conjecture is a conjecture regarding the gaps between prime numbers. The conjecture states that

$$p_{n+1} - p_n = O((\log p_n)^2)$$

where O is big O notation. And sometimes the following formulation is called Cramer's conjecture:

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log p_n)^2} = 1$$

which is stronger than the former. And this paper proves that

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log p_n)^2} = 0$$

i.e. Cremer conjecture is true, while the strong version is false.
proof. Note that **Lemma 3;(7)**

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log(n \log n))^2} = 0$$

And **Remark 2;(3)**

$$\lim_{n \rightarrow \infty} \frac{\log p_n}{\log(n \log n)} = 1$$

Then,

$$\lim_{n \rightarrow \infty} \frac{(\log(n \log n))^2}{(\log p_n)^2} = 1 \quad (16)$$

Hence,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log p_n)^2} = (7) \times (16) = 0$$

corollary 2.

$$g_n := p_{n+1} - p_n = O((\log p_n)^2) \blacksquare$$

5. About Oppermann conjecture

Oppermann conjecture is a conjecture regarding the distribution of prime numbers. It is closely related to but stronger than Legendre conjecture, Andrica conjecture, and Brocard conjecture. The conjecture states that for every integer $n \geq 1$,

$$\pi(n^2 - n) < \pi(n^2) < \pi(n^2 + n)$$

Definition 1. Let $\hat{p}(x)$ is the nearest prime less than x , $\hat{P}(x)$ is the nearest prime more than x .

$$e.g. \hat{p}(10) = 7, \hat{P}(10) = 11$$

Lemma 5. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is an increasing function and m is constant, then

$$p_n < p_{n+1} < f(p_n) \Rightarrow \exists p \in \mathbb{P} \text{ with } x < p < f(x)$$

proof (by contradiction). Let $\exists x \in \mathbb{R}$ such that $\nexists p \in \mathbb{P}$ in $(x, f(x))$, then $\hat{P}(x) > f(x)$. And By definition, $\hat{p}(x) \leq x$ and $\hat{P}(x)$ is the next prime of $\hat{p}(x)$. thus,

$$\hat{p}(x) < \hat{P}(x) < f(\hat{p}(x))$$

But, because f is an increasing function, $\hat{p}(x) \leq x \Rightarrow f(\hat{p}(x)) \leq f(x) < \hat{P}(x)$. It's contradiction. \blacksquare

Lemma 6. By **Lemma 5**, (10) is equivalent to

$$\forall k > 0, \exists M_1 \in \mathbb{R}, \quad s.t. \exists p \in \mathbb{P} \text{ with } x < p < x + x^k \text{ for } x \geq M_1 \quad (17)$$

Lemma 7.

$$\forall k > 0, \exists M_2 \in \mathbb{R}, \quad s.t. \exists p \in \mathbb{P} \text{ with } x - x^k < p < x \text{ for } x \geq M_2 \quad (18)$$

proof. In **Lemma 6**, let $x = m + m^k$, then there is a prime in the open interval (m, x) . Since $x > m \Rightarrow x^k > m^k$, $(m, x) \subset (x - x^k, x)$. Hence, there is a prime in the open interval $(x - x^k, x)$. (c.f. $M_1 < M_2$) ■

This psaper proves that for every $k > 0$, there exists $M \in \mathbb{R}$ such that

$$x \geq M \Rightarrow \pi(x^k - x) < \pi(x) < \pi(x^k + x) \quad (19)$$

proof. By (17),(18),

$$\forall k > 0, \exists M_2 \in \mathbb{R}, \quad s.t. \exists p, q \in \mathbb{P} \text{ with } x - x^k < p < x < q < x + x^k \text{ for } x \geq M_2$$

Let $x = t^m$ where $m = \frac{1}{k}$, then

$$\forall m > 0, \exists M' \in \mathbb{R}, \quad s.t. \exists p, q \in \mathbb{P} \text{ with } t^m - t < p < t^m < q < t^m + t \text{ for } t \geq M'$$

(c.f. $x = t^m \Rightarrow M' = M_2^k$) This fomula implies that

$$\forall m > 0, \exists M' \in \mathbb{R} \quad s.t. \quad t \geq M' \Rightarrow \pi(t^m - t) < \pi(t^m) < \pi(t^m + t) \quad \blacksquare$$

Furthermore, how many primes exist in $(x^k, x^k + x)$? In other word, what is the result of $\lim_{x \rightarrow \infty} (\pi(x^k + x) - \pi(x^k))$?

Remark 4.

$$f_1 \sim g_1 \wedge f_2 \sim g_2 \rightarrow f_1 - f_2 \sim g_1 - g_2$$

doesn't always hold. (1) is a counterexample. Due to this,

$$\lim_{x \rightarrow \infty} \frac{\pi(x^m + x) - \pi(x^m)}{(x^m + x)/\log(x^m + x) - x^m/\log(x^m)} = 1$$

may not hold. We need other method.

Lemma 8. for function f and g such that $\forall x \in \mathbb{R}, g(x) > f(x) > 0$, if $\lim_{x \rightarrow \infty} (g(x) - f(x)) = \infty$ and there exists $k \in (0, 1)$ such that $g(x)^k < g(x) - f(x)$ for sufficiently large x , then

$$\lim_{x \rightarrow \infty} (\pi(g(x)) - \pi(f(x))) = \infty$$

proof. Because of (18),

$$\begin{aligned} \forall j \in (0, k), \exists N \in \mathbb{R} \quad s.t. \quad x \geq N &\Rightarrow \exists p \in \mathbb{P} \text{ with } g(x) - g(x)^j < p < g(x) \\ &\Rightarrow \exists p \in \mathbb{P} \text{ with } f(x) < p < g(x) \end{aligned}$$

Let $a_1 = g(x)$, $a_{n+1} = a_n - a_n^j$, then there exists a prime in the open interval $(a_n - a_n^j, a_n) = (a_{n+1}, a_n)$ and for every $n \in \mathbb{N}$, $a_1 \geq a_n$.
Let $f(x) < a_m$, $f(x) > a_{m+1}$, then $\pi(g(x)) - \pi(f(x)) \geq m - 1$. Therefore, for sufficiently large x ,

$$g(x) - f(x) < \sum_{n=1}^m (a_n - a_{n+1}) = \sum_{n=1}^m a_n^j < \sum_{n=1}^m a_1^j = ma_1^j$$

$$\Rightarrow m > \frac{g(x) - f(x)}{a_1^j} = \frac{g(x) - f(x)}{g(x)^j} > \frac{g(x)^k}{g(x)^j}$$

Note that

$$\lim_{x \rightarrow \infty} \frac{g(x)^k}{g(x)^j} = \infty \quad (\because j \in (0, k))$$

Hence,

$$\lim_{x \rightarrow \infty} (\pi(g(x)) - \pi(f(x))) = \infty \quad \blacksquare$$

Since $\forall x \in \mathbb{R}$, $(x + x^m) > x^m > 0$ and for sufficiently large x , every $m > 0$, there exists $k \in (0, 1)$ such that $(x^m + x)^k < (x^m + x) - x^m = x$,

$$\forall m > 0 \quad \lim_{x \rightarrow \infty} (\pi(x^m + x) - \pi(x^m)) = \infty$$

References

- [1] E. Westzynthius. Über die Verteilung der Zahlen, die zu den n ersten Primzahlen teilerfremd sind. *Commentationes Helsingfors* 5, Nr. 25, 37 S. (1931)., 1931.
- [2] Erik Westzynthius. On the distribution of numbers that are not divisible by the first n primes. *Commentat. Phys.-Math.*, 5(25):1–37, 1931.
- [3] DA Goldston and J Pintz. Y. yıldırım, c.: Primes in tuples i. *Ann. Math.*, *AIM Preprint*, 19, 2005.

Keyword: Prime; Prime gap; Andrica conjecture; Cramer conjecture; Oppermann conjecture; Arithmetic mean of primes; Geometric mean of primes; Harmonic mean of primes

E-mail: fghver1051@gmail.com