

The Distribution Of Prime Numbers And The Continued Fractions

Mohammed Bouras

mohammed.bouras33@gmail.com

Abstract. In this paper, we discovered a new sequence contains only ones and the prime numbers, which can be calculated in two different ways that give the same result, the first way using the greatest common divisor (gcd) and Kurepa left factorial function, the second way consisting of using the denominator of the continued fraction defined by

$$\frac{mb(n-3) - nb(n-4)}{n(m-n+2) - m} = \cfrac{1}{2 - \cfrac{3}{3 - \cfrac{4}{4 - \cfrac{5}{\ddots (n-1) - \cfrac{n}{m}}}}}$$

Our sequence defined by

$$a_m(n) = \frac{|n(m-n+2) - m|}{\gcd(n(m-n+2) - m, mb(n-3) - nb(n-4))}$$

Where $|x|$ denotes the absolute value of x .

Keywords. Prime numbers, continued fraction, left factorial, sequence.

1. Introduction and preliminaries

A continued fraction is an expression of the form

$$a_0 + \frac{b_0}{a_1 + \frac{b_1}{a_2 + \frac{b_2}{\ddots}}}$$

Other notation

$$a_0 + \frac{b_0}{a_1 +} \frac{b_1}{a_2 +} \frac{b_2}{a_3 +} \dots$$

Where a_i and b_i are either rational numbers, real numbers or complex numbers.

In 1971, Kurepa introduced the left factorial function, with the symbol $!n$. For more details and formulas see [4], the Kurepa function is defined by

$$\frac{mb(n-3) - nb(n-4)}{n(m-n+2) - m} = \frac{1}{2 - \frac{3}{3 - \frac{4}{4 - \frac{5}{\ddots (n-1) - \frac{n}{m}}}}} \quad (1)$$

Where m is a polynomial in term n .

Proof. Let

$$a_1 = 2a_2 - 3a_3; a_2 = 3a_3 - 4a_4; a_3 = 4a_4 - 5a_5; a_4 = 5a_5 - 6a_6$$

Then we have

$$\begin{aligned} \frac{a_2}{a_1} &= \frac{a_2}{2a_2 - 3a_3} = \frac{1}{\frac{2a_2 - 3a_3}{a_2}} = \frac{1}{2 - \frac{3a_3}{a_2}} = \frac{1}{2 - \frac{3}{\frac{3a_3 - 4a_4}{a_3}}} \\ &= \frac{1}{2 - \frac{3}{3 - \frac{4a_4}{a_3}}} = \frac{1}{2 - \frac{3}{3 - \frac{4}{\frac{4a_4 - 5a_5}{a_4}}}} = \frac{1}{2 - \frac{3}{3 - \frac{4}{4 - \frac{5a_5}{a_4}}}} \end{aligned}$$

After some simplification, we find

$$\frac{a_2}{a_1} = \frac{1}{2 - \frac{3}{3 - \frac{4}{4 - \frac{5}{\ddots (n-1) - \frac{na_n}{a_{n-1}}}}} \quad (2)$$

From (1) and (2), we have

$$ma_n = a_{n-1} \quad (3)$$

We write a_1 in terms of a_{n-1} and a_n

$$a_1 = 2a_2 - 3a_3 = \dots = (n-1)a_{n-1} - (n^2 - 2)a_n \quad (4)$$

Substituting (3) in (4), we find

$$a_1 = (n(m-n+2) - m)a_n$$

Using the same procedure for a_2 , we have

$$a_2 = 3a_3 - 4a_4 = 8a_4 - 15a_5 = 25a_5 - 48a_6 = \dots$$

We observe that

$$a_2 = b(n-3)a_{n-1} - nb(n-4)a_n \quad (5)$$

Substituting (3) in (5), we get

$$a_2 = (mb(n-3) - nb(n-4))a_n$$

Returning to (2), we obtain

$$\frac{a_2}{a_1} = \frac{mb(n-3) - nb(n-4)}{n(m-n+2) - m} = \frac{1}{2 - \frac{3}{3 - \frac{4}{4 - \frac{5}{\ddots (n-1) - \frac{n}{m}}}}} \quad (6)$$

This completes the proof.

Theorem 1.2. For all integers $n \geq 3$. The denominator of the continued fraction is as follows

$$n(m-n+2) - m = 2(mb(n-3) - nb(n-4)) - 3(mc(n-3) - nc(n-4))$$

Where m is a polynomial in term n .

Proof. Similarly, using the same procedure as that of proving the theorem 1.

We have

$$a_3 = 4a_4 - 5a_5 = 15a_5 - 24a_6 = 66a_6 - 105a_7 = \dots$$

We observe that

$$a_3 = c(n-3) \cdot a_{n-1} - nc(n-4) \cdot a_n \quad (7)$$

Substituting (3) in (7), we find

$$a_3 = (mc(n-3) - nc(n-4))a_n$$

Then, we have

$$a_1 = 2a_2 - 3a_3$$

$$(n(m-n+2) - m)a_n = [2(mb(n-3) - nb(n-4)) - 3(mc(n-3) - nc(n-4))] \cdot a_n$$

Then, we get

$$n(m-n+2) - m = 2(mb(n-3) - nb(n-4)) - 3(mc(n-3) - nc(n-4))$$

This completes the proof.

Theorem 1.3. For all integers $n \geq 3$. The continued fraction

$$\frac{2 \cdot (mb(n-3) - nb(n-4))}{n(m-n+1)} = \frac{1}{1 - \frac{1}{2 - \frac{2}{3 - \frac{3}{\ddots (n-1) - \frac{n-1}{m}}}}} \quad (8)$$

Where m is a polynomial in term n .

Proof. Similarly, Using the same procedure of proof the theorem (1.1)

Putting

$$a_1 = a_2 - a_3 ; a_2 = 2a_3 - 2a_4 ; a_3 = 3a_4 - 3a_5 ; a_4 = 4a_5 - 4a_6 ; \dots$$

And we obtain the desired result.

Remark 1

For $m = n$, the Kerupa left factorial function is as continued fraction

$$K(n) = !n = \frac{1}{1 - \frac{1}{2 - \frac{2}{3 - \frac{3}{\ddots (n-1) - \frac{n-1}{n}}}}}$$

The sequence which is actually important is the next one.

2. The sequence $a_m(n)$

The sequence of the unreduced denominator of the continued fraction (theorem 1.1) is as follows

$$a_m(n) = \frac{|n(m - n + 2) - m|}{gcd(n(m - n + 2) - m, mb(n - 3) - nb(n - 4))}$$

Where $gcd(x, y)$ denotes the greatest common divisor of x and y .

3. Main results

In this section we present some new results for our sequence in the following conjectures

Conjecture 3.1. For all integers $n \geq 3$ and $m = n + 1$. The sequence of the unreduced denominator is as follows

$$a(n) = \frac{2n - 1}{gcd(2n - 1, b(n - 2) + b(n - 3))}; n \geq 2$$

The values of $a(n)$

3, 5, 7, 3, 11, 13, 1, 17, 19, 1, 23, 1, 1, 29, 31, 1, 1, 37, 1, 41, 43, 1, 47, 1, 1, 53, 1, 1, 59, 61, 1, 1, 67, 1, 71, 73, 1, 1, 79, 1, 83, 1, 1, 89, 1, 1, 1, 97, 1, 101, 103, 1, 107, 109, 1, 113, 1, 1, 1, 1, 1, 1, 127, 1, 131, 1, 1, 137, 139, 1, 1, 1, 1, 149, 151, 1, 1, 157, 1, 1, 163, 1, 167,...

Every term of this sequence is either a prime number or 1.

For $n \geq 2$, $a(n) = 2n - 1$ if $2n - 1$ is prime (except for $n=5$), 1 otherwise .

Conjecture 3.2. For all integers $n \geq 4$ and $m = n - 3$. The sequence of the unreduced denominator is as follows

$$a(n) = \frac{2n - 3}{gcd(2n - 3, 3b(n - 3) - b(n - 2))}; n \geq 2$$

The values of $a(n)$

1, 1, 5, 7, 1, 11, 13, 1, 17, 19, 1, 23, 1, 1, 29, 31, 1, 1, 37, 1, 41, 43, 1, 47, 1, 1, 53, 1, 1, 59, 61, 1, 1, 67, 1, 71, 73, 1, 1, 79, 1, 83, 1, 1, 89, 1, 1, 1, 97, 1, 101, 103, 1, 107, 109, 1, 113, 1, 1, 1, 1, 1, 1, 127, 1, 131, 1, 1, 137, 139, 1, 1, 1, 1, 149, 151, 1, 1, 157, 1, 1, 163, 1, 167,...

Every term of this sequence is either a prime number or 1.

For $n \geq 4$, $a(n) = 2n - 3$ if $2n - 3$ is prime, 1 otherwise .

Conjecture 3.3. For all integers $n \geq 3$ and $m = -1$. The sequence of the unreduced denominator is as follows

$$a(n) = \frac{n^2 - n - 1}{\gcd(n^2 - n - 1, b(n - 3) + nb(n - 4))} ; \text{ for } n \geq 2$$

The values of $a(n)$

1, 5, 11, 19, 29, 41, 11, 71, 89, 109, 131, 31, 181, 19, 239, 271, 61, 31, 379, 419, 461, 101, 29, 599, 59, 701, 151, 811, 79, 929, 991, 211, 59, 41, 1259, 1, 281, 1481, 1559, 149, 1721, 1, 61, 1979, 2069, 2161, 1, 2351, 79, 2549, 241, 1, 2861, 2969, 3079, 3191, ... (see A356247)

We conjectured that :

* Every term of this sequence is either a prime number or 1.

* Except for 5, the primes all appear exactly twice, such that

$$a(n) = a(a(n) - n + 1)$$

Consequently, let us consider the values of n and m such that we get:

$$a(n) = a(m) = n + m - 1$$

And

$$a(n) = a(m) = \gcd(n^2 - n - 1, m^2 - m - 1)$$

Conjecture 3.4. For all integers $n \geq 3$ and $m = -2$. The expression of the sequence $a(n)$ is as follows

$$a(n) = \frac{n^2 - 2}{\gcd(n^2 - 2, 2b(n - 3) + nb(n - 4))} ; \text{ for } n \geq 3$$

The values of $a(n)$.

7, 7, 23, 17, 47, 31, 79, 7, 17, 71, 167, 97, 223, 127, 41, 23, 359, 199, 439, 241, 31, 41, 89, 337, 727, 1, 839, 449, 137, 73, 1087, 577, 1223, 647, 1367, 103, 1, 47, 73, 881, 1, 967, 1, 151, 2207, 1151, 2399, 1249, 113, 193, 401, 1, 3023, 1567, 191, 41, 71...

The sequence $a(n)$ takes only 1's and primes.

Conjecture 3.5. For all integers $n \geq 3$ and $m = n + 2$. The expression of the sequence $a(n)$ is as follows

$$a(n) = \frac{3n - 2}{\gcd(3n - 2, (n + 1)b(n - 3) - b(n - 4) - (n - 1)b(n - 5))} ; \text{ for } n \geq 3$$

The values of $a(n)$ for $n \geq 3$

7, 5, 13, 2, 19, 11, 5, 1, 31, 17, 37, , 1, 43, 23, 1, 1, 1, 29, 61, 1, 67, 1, 73, 1, 79, 41, 1, 1, 1, 47, 97, 1, 103, 53, 109, 1, 1, 59, 1, 1, 127, 1, 1, 1, 139, 71, 1, 1, 151, 1, 157, 1, 163, 83, 1, 1, 1, 89, 181, 1, 1, 1, 193, 1, 199, 101, 1, 1, 211, ...

The sequence $a(n)$ contains only ones and the primes.

Conjecture 3.6. For all integers $n \geq 3$ and $m = n + 3$. The expression of the sequence $a(n)$ is as follows

$$a(n) = \frac{4n - 3}{\gcd(4n - 3, (n + 2)b(n - 3) - b(n - 4) - (n - 1)b(n - 5))} ; \text{ for } n \geq 3$$

The values of $a(n)$ for $n \geq 3$

3, 13, 17, 7, 5, 29, 11, 37, 41, 1, 7, 53, 19, 61, 1, 23, 73, 1, 1, 1, 89, 31, 97, 101, 1, 109, 113, 1, 1, 1, 43, 1, 137, 47, 1, 149, 1, 157, 1, 1, 1, 173, 59, 181, 1, 1, 193, 197, 67, 1, 1, 71, 1, 1, 1, 229, 233, 79, 241, 1, 83, 1, 257, 1, 1, 269, 1, 277, ...

The sequence $a(n)$ takes only 1's and primes.

Remark 2

There are many sequences that contain's only ones and the primes related to the values of m .

Acknowledgements

I would like to thank Bill Mceachen for the numerous comments and suggestions. Thanks go also to Jon. E. Schoenfeld, Alois. P. Heinz, Michael De Vlieger and the other editor-in-chief of the on-line encyclopedia of integers sequences (Oeis).

References

- [1] Eric S. Rowland, A Natural Prime-Generating Recurrence, Journal of Integer Sequences, Vol. 11 (2008).
- [2] Benoit Cloitre, 10 conjectures in additive number theory, <https://arxiv.org/abs/1101.4274>
- [3] David Angell, A family of continued fractions, Journal of Number Theory 130 (2010) 904–911.
- [4] Bernd C. Kellner, Some remarks on Kurepa's left factorial, arXiv:math/0410477v1 [math.NT] 21 Oct 2004.
- [5] N. J. A. Sloane, The On-line Encyclopedia of integers sequences, <https://oeis.org>
- [6] Plouffe Simon, A set of formulas for primes, <https://vixra.org/abs/1812.0494>