

## **Bond Lengths in Gaseous Alkali Metal Chlorides as Sums of the Golden Ratio Based Cationic Radii and Covalent Radius of Chlorine.**

**Raji Heyrovská**, Academy of Sciences of the Czech Republic (Emer.), Prague, Czech Republic,

Email: [rheyrovs@hotmail.com](mailto:rheyrovs@hotmail.com)

(Paper to mark the 85<sup>th</sup> anniversary of the paper on this subject in Phys. Rev., 51, 1000, 1937)

**Abstract:** Alkali metal halides are known to have shorter bond lengths in the gaseous state than in their crystalline state. It is also known that the bonds are polar covalent. However, an exact interpretation of these bond lengths has remained an open question for the last 85 years. The present author has shown for the first time in 2004 that the crystal ionic distances in alkali halides are exact sums of the Golden ratio based radii of the alkali metal cations and halide anions. In this short report, it is shown that in gaseous alkali metal chlorides, alkali metal cations are bound to covalent chlorine as in the partially ionic bond in HCl.

### **Introduction**

Experimental results by electron diffraction [1] showed that gaseous alkali halides have shorter bond lengths than in their crystalline forms. This result has intrigued many minds ever since and various explanations have been suggested. It was recognized that the bonds are polar covalent and an account of these can be found in [2]. The present author established [3, 4] that the crystal ionic distances,  $d(\text{MX})$  in alkali halides ( $\text{M}^+ \text{X}^-$ , where  $\text{M}^+$  is metal cation and  $\text{X}^-$  is the halogen anion) are exact sums of the Golden ratio based ionic radii of the metal and halogen ions. In general, it was also established over the years that interatomic bonding distances in all molecules including hydrogen bonds, are sums of the covalent atomic and or the Golden ratio based ionic

radii of the adjacent atoms or ions, whether the bond is fully or partially covalent or ionic. For the collected work and a meeting presentation, see [5, 6]. Here, it shown that the additivity of Golden ratio based ionic radii of alkali metals and covalent radii of halogens holds even for the gaseous molecules.

### **Bond lengths in alkali metal chlorides.**

It is found here that the bond lengths in gaseous alkali metal chlorides are given by,

$$d(M^+X)_{\text{gas}} = R(M^+) + R(X)$$

where the Golden ratio based ionic radius [3, 4],  $R(M^+)_{\text{ion}} = d(MM)/\phi^2$  and  $R(X)_{\text{cov}} = d(XX)/2$ .

The bond length  $d(MM)$  is the distance between center to corner atoms of the body centered cubic alkali metals (which is shorter by a factor of 0.866 than the edge length which is involved in the crystalline alkali halides).

The experimental bond lengths  $d(MX)_{\text{gas}}$  for MCl, where  $M = \text{Na, K, Rb and Cs}$ , from [1] reported in [2] are given in column 2 below (in Angstroms). Other values of bond lengths, for example 2.36 Å (< 2.51 Å) for NaCl have been reported [7].

**Table 1** Bond lengths of gaseous alkali chlorides and their correlation with the sum of Golden ratio-based metal ion radius and chlorine covalent radius.

MX	$d(MX)_{\text{obs}}$	$d(MM)$	$R(M^+)$	$R(X)$	$d(M^+X)_{\text{sum}}$
NaCl	2.51	3.66	1.40	0.99	2.39
KCl	2.79	4.52	1.73	0.99	2.72
RbCl	2.89	4.74	1.81	0.99	2.80
CsCl	3.06	5.00	1.91	0.99	2.90

It can be seen that the calculated sum in the last column is slightly less than the observed value in the column 2 by about 0.1 Å. It is interesting to point out that in [1], the covalent radii of halogens are used as ionic radii to interpret the bond lengths. For values calculated by other methods, see [2].

## References

1. S. B. Hendricks and V. M. Mosley, *Phys. Rev.*, 51, 1000 (1937); L. R. Maxwell, S. B. Hendricks and V. M. Mosley, *ibid.*, 52, 968 (1937)
2. An Interpretation of Bond Lengths in Alkali Halide Gas Molecules, R. T. Sanderson *JACS* vol 74, pp 272-4. Jan. 5, 1952.
3. Hydrogen as an atomic condenser; The decisive role of the golden ratio in atomic dimensions; R. Heyrovská, 35th Meeting of DAMOP, APS, Tucson, AZ, May 2004, Poster, Abstracts: P-132 and P-133.
4. The Golden ratio, ionic and atomic radii and bond lengths, R. Heyrovská, *Molecular Physics*, 103 (2005) 877 – 882.
5. Bohr radius as the sum of Golden sections pertaining to the electron and proton, covalent bond lengths between the same two atoms as exact sums of their cationic and anionic radii and additivity of atomic and or ionic radii in bond lengths – Collected work dedicated to J. Kepler (1571- 1630). R. Heyrovská, <http://vixra.org/abs/1709.0066>; <http://vixra.org/pdf/1709.0066v1.pdf>

6. 113th Statistical Mechanics Conference in honor of three renowned Physicists, Rutgers University, Hill Center, May 10-12, 2015,  
<https://www.researchgate.net/publication/28127035722>
7. <http://hyperphysics.phy-astr.gsu.edu/hbase/molecule/NaCl.html>

**Acknowledgement:** I am thankful to Prof. B. Friedrich for asking me about the bond length of gaseous sodium chloride, which made me do this work.