

Investigating Quantum Mechanics In 5th Dimensional Embedding via Deterministic Structure, Small Scale Factor, And Initial Inflaton Field

Andrew Walcott Beckwith¹

Qazi Abdul Ghafoor²

¹ Physics Department, Chongqing University, College of Physics, Chongqing University Huxi Campus, No. 44 Daxuechen Nanlu, Shapinba District, Chongqing 401331, People's Republic of China

² Mathematics Department, Hazara University Manshera, Pakistan

Email: Rwill9955b@gmail.com, qaziabdulghafoor@hu.edu.pk

Abstract: We consider if a generalized HUP set greater than or equal to Planck's constant divided by the square of a scale factor, as well as an inflaton field, yields the result that ΔE times Δt is embedded in a 5 dimensional field which is within a deterministic structure. Our proof concludes with Δt as of Planck time, resulting in enormous potential energy. If that potential energy is induced by a repeating universe structure, we get a free value of ΔE Δt that is almost infinite, supporting a prior conclusion.

1. Introduction'

In this document we are revisiting the following statement made earlier [1] [2]

Quote

Using the following

$$T_{ii} = \text{diag}(\rho, -p, -p, -p) \quad (1)$$

Then

$$\Delta T_{tt} \sim \Delta \rho \sim \frac{\Delta E}{V^{(3)}} \quad (2)$$

Then, Eq. (1) and Eq. (2) together yield

$$\delta t \Delta E \geq \frac{\hbar}{\delta g_{tt}} \neq \frac{\hbar}{2} \quad (3)$$

$$\text{Unless } \delta g_{tt} \sim O(1)$$

By the initial idea given in Eq.(3), the GUP is look complicated at the initial variation, we make the following treatment at the start of expansion of the Universe[1][2][3]

$$\delta g_{tt} \sim a^2(t) \cdot \phi \ll 1 \quad \text{Goes to become effectively almost ZERO.} \quad (4)$$

If this is effectively almost zero, the effect would be to embed Quantum mechanics within a 5 dimensional structure

Snip

I.e. this deterministic embedding is in part in spirit similar to what is given by Wesson [3]

End of quote

We add more context to this through using the Wesson result directly in our own work and further use it to prove a deterministic contribution in line with Eq. (3) and Eq. (4).

2. Modus operandi statement in terms of an inflaton field

Before proceeding, we state that the inflaton field used in Eq. (3) and Eq. (4) satisfies the following properties, [4][5][6]

$$\begin{aligned}
 a(t) &= a_{initial} t^\nu \\
 \Rightarrow \phi &= \ln \left(\sqrt{\frac{8\pi G V_0}{\nu \cdot (3\nu - 1)}} \cdot t \right)^{\sqrt{\frac{\nu}{16\pi G}}} \\
 \Rightarrow \dot{\phi} &= \sqrt{\frac{\nu}{4\pi G}} \cdot t^{-1} \\
 \Rightarrow \frac{H^2}{\dot{\phi}} &\approx \sqrt{\frac{4\pi G}{\nu}} \cdot t \cdot T^4 \cdot \frac{(1.66)^2 \cdot g_*}{m_p^2} \approx 10^{-5}
 \end{aligned} \tag{5}$$

In the spirit of use of the inflaton field what we will propose is that

$$\phi = \ln \left(\sqrt{\frac{8\pi G V_0}{\nu \cdot (3\nu - 1)}} \cdot t \right)^{\sqrt{\frac{\nu}{16\pi G}}} \approx \sqrt{\frac{\nu}{16\pi G}} \cdot \left(\sqrt{\frac{8\pi G V_0}{\nu \cdot (3\nu - 1)}} \cdot t - 1 \right) \tag{6}$$

i.e. assuming that if the initial time step is near Planck time which is normalized to 1 that

$$V_0 \approx \text{initial} - \text{energy} \tag{7}$$

In addition we will go to Wesson [7] and to make the following adjustments

3. Wesson's treatment of embedding of the HUP in deterministic structure[7]

$$|dp_\alpha dx^\alpha| \approx \frac{L}{l} \cdot \frac{h}{c} \cdot \left[\frac{dl}{l} \right]^2 \quad (8)$$

Where we will define l and Λ as follows

First, we define L in terms of the cosmological “constant” by [7]

$$\Lambda = \frac{1}{3L^2} \quad (9)$$

Also[7]

$$dS_{5-d}^2 = \frac{L^2}{l^2} dS_{4-d}^2 - \frac{L^4}{l^4} dl^2 \quad (10)$$

Also 5 dimensional wave number is defined via

$$K_l = 1/l \quad (11)$$

In the case of Pre Planckian space-time the idea is to do the following[7]

$$\begin{aligned} |dp_\alpha dx^\alpha| &\approx \frac{L}{l} \cdot \frac{h}{c} \cdot \left[\frac{dl}{l} \right]^2 \\ \xrightarrow{\alpha=0} |dp_0 dx^0| &\approx |\Delta E \Delta t| \approx (h / a_{init}^2 \phi(t)) \\ \Rightarrow \frac{L}{l} \cdot \frac{h}{c} \cdot \left[\frac{dl}{l} \right]^2 &\approx (h / a_{init}^2 \phi(t_{init})) \end{aligned} \quad (12)$$

Use of all this leads to below equation, [7]

$$\int_{l_1}^{l_2} dl / l^{3/2} \approx \frac{(l_2 - l_1)}{l^{3/2} (c)} \approx \frac{(3\Lambda)^{1/4}}{a_{init} \cdot \left(\frac{\nu}{16\pi G} \right)^{1/4} \cdot \left(\sqrt{\frac{8\pi G V_0}{\nu \cdot (3\nu - 1)}} \cdot t - 1 \right)^{1/2}} \quad (13)$$

4. Extracting time initially from Eq. (13) and assuming time equal to Planck time? Extract V_0

Our approximation is to set $G = 1 = \hbar$ (Planck units) with Planck time normalized to 1. Then

$$t = t_{planck} \rightarrow 1 = \sqrt{\frac{\nu(3\nu-1)}{8\pi V_0}} + \sqrt{\frac{2 \cdot (3\nu-1)}{V_0} \cdot \frac{a_{init}^2 \cdot (l_2 - l_1)^2}{l^3(c) \cdot (3\Lambda)^{1/2}}} \quad (14)$$

Then we have that at Planck time, normalized to 1 we look at

$$V_0 = \left(\sqrt{\frac{\nu(3\nu-1)}{8\pi}} + \sqrt{2 \cdot (3\nu-1) \cdot \frac{a_{init}^2 \cdot (l_2 - l_1)^2}{l^3(c) \cdot (3\Lambda)^{1/2}}} \right)^2 \quad (15)$$

5. At initial configuration in Planck time make the following assumption

We assume that we have an emergent space-time. If so, and

$$V_0 = \left(\sqrt{\frac{\nu(3\nu-1)}{8\pi}} + \sqrt{2 \cdot (3\nu-1) \cdot \frac{a_{init}^2 \cdot (l_2 - l_1)^2}{l^3(c) \cdot (3\Lambda)^{1/2}}} \right)^2 \approx \Delta E \quad (16)$$

Implication is, that if we use the present value of the cosmological constant Λ , that the initial energy, as induced by Eq. (16) becomes almost infinite, thereby confirming by default what is brought up by Eq. (4)

6. Discussion

Our value of the initial energy specifies an almost infinite value, so does this confirm deterministic embedding of the HUP initially in 5 dimensions, in a deterministic structure?

We argue it does, because Eq. (16) still uses the 5 dimensional inputs specified by l which is one over a wave number in an additional dimension of space time. Furthermore we can also compare this expression in (16) with[4]

$$V_0 = \left(\frac{.022}{\sqrt{qN_{efolds}}} \right)^4 = \frac{\nu(\nu-1)\lambda^2}{8\pi Gm_p^2} \quad (17)$$

' λ ' as a dimensionless parameter. From [4] we have a Chamelon mechanism for fifth force as [4]

$$F_{5th-force} = -\frac{\tilde{\beta} \cdot (\vec{\nabla} \phi)}{m_p} \quad (18)$$

We use here Pre Planckian conditions

$$t = \frac{r}{\omega c} \quad (19)$$

First, r is almost Planck in length, if so then

Using this instead of the ω_{gw}^6 expression, then write the rest of it as follows which would have a minimum value as[4][8]

$$\begin{aligned}
\omega_{gw}^6 &\approx c^7 \times \frac{\tilde{\beta}}{2m_p r} \cdot \sqrt{\frac{v}{\pi G}} \times \frac{1}{Gc \cdot (M_{mass})^2 \langle r^2 \rangle^2} \\
\Rightarrow \omega_{gw} &\approx G, m_p, r \approx \ell_p \xrightarrow{\text{Planck-normalization}} 1 \\
M_{mass} &\approx \zeta \cdot m_p \xrightarrow{\text{Planck-normalization}} \zeta \\
\langle r^2 \rangle^2 &\approx \ell_p^4 \xrightarrow{\text{Planck-normalization}} 1 \\
\therefore \omega_{gw} &\xrightarrow{\text{Planck-normalization}} \left(\sqrt{\frac{v}{4\pi}} \times \frac{\tilde{\beta}}{(\zeta)^2} \right)^{1/6}
\end{aligned} \tag{20}$$

$$\omega_{gw} \approx \frac{c^{7/6} \tilde{\beta}^{1/6}}{(2m_p r)^{1/6}} \cdot \left(\frac{v}{\pi G} \right)^{1/12} \cdot \frac{1}{\left(GcM_{Mass}^2 \cdot \langle r^2 \rangle^2 \right)^{1/6}} \text{ so if } G = m_p = \ell_p = 1 \tag{21}$$

We then will conclude this by stating the connection with Eq. (16) and (17) so

$$\lambda = \left(\sqrt{\frac{v(3v-1)}{8\pi}} + \sqrt{2 \cdot (3v-1)} \cdot \frac{a_{init}^2 \cdot (l_2 - l_1)^2}{l^3(c) \cdot (3\Lambda)^{1/2}} \right) \cdot \sqrt{\frac{8\pi}{v(v-1)}} \tag{22}$$

In doing so, we have thoroughly planted 5 dimensional lengths as given by l into our analysis , with the caveat the value of Eq. (16) and Eq. (17) can become enormous with a small enough value of the cosmological constant. Note that the expression $\frac{(l_2 - l_1)^2}{l^3(c)}$ has two lengths, l_2 and l_1 in 5 dimensions, with the first length, l_2 larger in magnitude than l_1 and $l^3(c)$ being the cube of the length,

$$l_2 > l(c) > l_1 \tag{23}$$

And

$$a_{init} \approx 10^{-40} \quad (24)$$

Finally the power of initial gravitational waves at the start of the universe is such that, [4][8][9]

$$P_{GW} \approx \frac{Gc \cdot (M_{mass})^2 \omega_{gw}^6 \langle r^2 \rangle^2}{c^6} \quad (25)$$

$$\approx c \times |F_{5th-force}| = \left| -c \times \frac{\tilde{\beta} \cdot (\vec{\nabla} \phi)}{m_p} \right| \approx c \times \frac{\tilde{\beta}}{2m_p r} \cdot \sqrt{\frac{v}{\pi G}}$$

This will allow for setting $\tilde{\beta}$.

7. Conclusion

We investigated the quantum mechanics along with the concepts of Heisenberg Uncertainty Principle, small scale factor and Inflaton field in the context of 5 dimensional Embedding. Using Wesson's treatment of Embedding we conclude that delta t is planck time and resulted in enormous induced Energy and becomes almost infinite, by a repeating universe structure. This induced energy is basically potential energy and it confirms the Embedding of the HUP initially in 5 dimensions within Deterministic Structure. We found that QM is embedded in 5th dimensional embedding come about due to deterministic setting implying small scale factor and Inflaton field values initially. The reason for this inquiry is to determine an onset of when quantum conditions apply. A summary of what we have tried to prove is that quantum mechanics as represented by the HUP is embedded in a deterministic setup at the onset of the universe, with the results that Corda's results of no firewalls, for black holes due to a mix of

quantum physics, and relativistic effects applies after the start of the expansion of the universe as given in [10]

Acknowledgements

This work is supported in part by National Nature Science Foundation of China grant No.

11375279

References

[1] Beckwith, Andrew, “Does QM embedded in 5th dimensional embedding allow for classical black hole ideas only in early universe, whereas Corda special relativity plus QM may eliminate Event horizons for black holes after big bang?” PAPER ACCEPTED BY jhepgc, TO BE PUBLISHED October , 2023; <https://vixra.org/abs/2308.0126>

[2] Beckwith, A. and Moskaliuk, S. (2017) Generalized Heisenberg Uncertainty Principle in Quantum Geometrodynamics and General Relativity. Ukrainian Journal of Physics, 62, 727-740

[3] Giovannini, M., “*A Primer on the Physics of the Cosmic Microwave Background*” World Press Scientific, Hackensack, New Jersey, USA, 2008

[4] Beckwith, A. ,“New Conservation Law as to Hubble Parameter, Squared Divided by Time Derivative of Inflaton in Early and Late Universe, Compared with Discussion of HUP in Pre Planckian to Planckian Physics, and Relevance of Fifth Force Analysis to Gravitons and GW”, pp 1 -18 in, Gravitational Waves - Theory and Observations ,edited by Prof. Carlos Frajuca, published by Intechopen, London UK, 2023 Submitted: October 24th, 2022 Reviewed: October 30th, 2022 Published: December 22nd, 2022 DOI: 10.5992/intechopen.1000577, <https://www.intechopen.com/online-first/1125889>

[5] Padmanabhan, T., “An Invitation to Astrophysics”, World Press Scientific, World Scientific Series in Astronomy and Astrophysics: Volume 8, Singapore, Republic of Singapore, 2006

[6]Sarkar, U., “ Particle and Astroparticle Physics”, Taylor and Francis, New York City, New York, USA, 2008

[7] Wesson, Paul, “**Five-dimensional Physics: Classical And Quantum Consequences Of Kaluza-klein Cosmology**” World Press Scientific, Singapore, Republic of Singapore 2006

[8] Beckwith, A. and Ghafoor, Q. (2023) Using Model of a Universe as Similar to a Black Hole, Ask If We Have to Have Singularities, If We Are Looking at Initial Time Step and Entropy, from the Beginning. *Journal of High Energy Physics, Gravitation and Cosmology*, **9**, 708-719. doi: [10.4236/jhepgc.2023.93058](https://doi.org/10.4236/jhepgc.2023.93058).

[9] Fischbach, Ephraim; Talmadge, Carrick (19 March 1992). "Six years of the fifth force". *Nature*. **356** (6366): 207–215. Bibcode:1992Natur.356..207F. doi:10.1038/356207a0. S2CID 21255315.

[10] Corda, Christian, “Schrödinger and Klein-Gordon theories of black holes from the quantization of the Oppenheimer and Snyder gravitational collapse” arXiv:2307.05554v1 [gr-qc] 9 Jul 2023